

UC Berkeley

UC Berkeley Previously Published Works

Title

Measurement of time-dependent CP asymmetries in $B^0 \rightarrow K_S^0 \eta \gamma$ decays

Permalink

<https://escholarship.org/uc/item/19d4p3gr>

Journal

Physical Review D, 97(9)

ISSN

2470-0010

Authors

Nakano, H
Ishikawa, A
Sumisawa, K
[et al.](#)

Publication Date

2018-05-01

DOI

10.1103/physrevd.97.092003

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Measurement of the Time-Dependent CP Asymmetry in the $B^0 \rightarrow \phi K^0$ Decay

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ C. LeClerc,⁵ G. Lynch,⁵ A. M. Merchant,⁵ L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ A. V. Telnov,⁵ W. A. Wenzel,⁵ K. Ford,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ S. E. Morgan,⁶ A. T. Watson,⁶ M. Fritsch,⁷ K. Goetzen,⁷ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ M. Steinke,⁷ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸ M. P. Kelly,⁸ T. E. Latham,⁸ F. F. Wilson,⁸ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ P. Kyberd,¹⁰ L. Teodorescu,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹² C. Buchanan,¹³ B. L. Hartfiel,¹³ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵ E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ B. Dahmes,¹⁶ S. L. Levy,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. A. Heusch,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ R. E. Schmitz,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ P. Spradlin,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸ A. Dvoretzskii,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸ S. Jayatilake,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰ S. Chen,²⁰ P. J. Clark,²⁰ W. T. Ford,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. G. Smith,²⁰ L. Zhang,²⁰ A. Chen,²¹ J. L. Harton,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ Q. L. Zeng,²¹ D. Altenburg,²² T. Brandt,²² J. Brose,²² T. Colberg,²² M. Dickopp,²² E. Feltresi,²² A. Hauke,²² H. M. Lacker,²² E. Maly,²² R. Müller-Pfefferkorn,²² R. Nogowski,²² S. Otto,²² A. Petzold,²² J. Schubert,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² J. E. Sundermann,²² D. Bernard,²³ G. R. Bonneaud,²³ F. Brochard,²³ P. Grenier,²³ S. Schrenk,²³ Ch. Thiebaut,²³ G. Vasileiadis,²³ M. Verderi,²³ D. J. Bard,²⁴ A. Khan,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ M. Andreotti,²⁵ V. Azzolini,²⁵ D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,²⁵ G. Cibinetto,²⁵ E. Luppi,²⁵ M. Negrini,²⁵ L. Piemontese,²⁵ A. Sarti,²⁵ E. Treadwell,²⁶ R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ M. Piccolo,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Capra,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ S. Bailey,²⁹ G. Brandenburg,²⁹ M. Morii,²⁹ E. Won,²⁹ R. S. Dubitzky,³⁰ U. Langenegger,³⁰ W. Bhimji,³¹ D. A. Bowerman,³¹ P. D. Dauncey,³¹ U. Egede,³¹ J. R. Gaillard,³¹ G. W. Morton,³¹ J. A. Nash,³¹ G. P. Taylor,³¹ G. J. Grenier,³² U. Mallik,³² J. Cochran,³³ H. B. Crawley,³³ J. Lamsa,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ J. Yi,³³ M. Davier,³⁴ G. Grosdidier,³⁴ A. Höcker,³⁴ S. Laplace,³⁴ F. Le Diberder,³⁴ V. Lepeltier,³⁴ A. M. Lutz,³⁴ T. C. Petersen,³⁴ S. Plaszczynski,³⁴ M. H. Schune,³⁴ L. Tantot,³⁴ G. Wormser,³⁴ C. H. Cheng,³⁵ D. J. Lange,³⁵ M. C. Simani,³⁵ D. M. Wright,³⁵ A. J. Bevan,³⁶ J. P. Coleman,³⁶ J. R. Fry,³⁶ E. Gabathuler,³⁶ R. Gamet,³⁶ R. J. Parry,³⁶ D. J. Payne,³⁶ R. J. Sloane,³⁶ C. Touramanis,³⁶ J. J. Back,³⁷ C. M. Cormack,³⁷ P. F. Harrison,^{37,*} G. B. Mohanty,³⁷ C. L. Brown,³⁸ G. Cowan,³⁸ R. L. Flack,³⁸ H. U. Flaecher,³⁸ M. G. Green,³⁸ C. E. Marker,³⁸ T. R. McMahon,³⁸ S. Ricciardi,³⁸ F. Salvatore,³⁸ G. Vaitsas,³⁸ M. A. Winter,³⁸ D. Brown,³⁹ C. L. Davis,³⁹ J. Allison,⁴⁰ N. R. Barlow,⁴⁰ R. J. Barlow,⁴⁰ P. A. Hart,⁴⁰ M. C. Hodgkinson,⁴⁰ G. D. Lafferty,⁴⁰ A. J. Lyon,⁴⁰ J. C. Williams,⁴⁰ A. Farbin,⁴¹ W. D. Hulsbergen,⁴¹ A. Jawahery,⁴¹ D. Kovalskyi,⁴¹ C. K. Lae,⁴¹ V. Lillard,⁴¹ D. A. Roberts,⁴¹ G. Blaylock,⁴² C. Dallapiccola,⁴² K. T. Flood,⁴² S. S. Hertzbach,⁴² R. Kofler,⁴² V. B. Koptchev,⁴² T. B. Moore,⁴² S. Saremi,⁴² H. Staengle,⁴² S. Willocq,⁴² R. Cowan,⁴³ G. Sciolla,⁴³ F. Taylor,⁴³ R. K. Yamamoto,⁴³ D. J. J. Mangeol,⁴⁴ P. M. Patel,⁴⁴ S. H. Robertson,⁴⁴ A. Lazzaro,⁴⁵ F. Palombo,⁴⁵ J. M. Bauer,⁴⁶ L. Cremaldi,⁴⁶ V. Eschenburg,⁴⁶ R. Godang,⁴⁶ R. Kroeger,⁴⁶ J. Reidy,⁴⁶ D. A. Sanders,⁴⁶ D. J. Summers,⁴⁶ H. W. Zhao,⁴⁶ S. Brunet,⁴⁷ D. Côté,⁴⁷ P. Taras,⁴⁷ H. Nicholson,⁴⁸ N. Cavallo,⁴⁹ F. Fabozzi,^{49,†} C. Gatto,⁴⁹ L. Lista,⁴⁹ D. Monorchio,⁴⁹ P. Paolucci,⁴⁹ D. Piccolo,⁴⁹ C. Sciacca,⁴⁹ M. Baak,⁵⁰ H. Bulten,⁵⁰ G. Raven,⁵⁰ L. Wilden,⁵⁰ C. P. Jessop,⁵¹ J. M. LoSecco,⁵¹ T. A. Gabriel,⁵² T. Allmendinger,⁵³ B. Brau,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵

M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ G. Tiozzo,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Ocariz,⁵⁶ M. Pivk,⁵⁶ L. Roos,⁵⁶ S. T'Jampens,⁵⁶ G. Therin,⁵⁶ P. F. Manfredi,⁵⁷ V. Re,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ F. Anulli,^{27,59} M. Biasini,⁵⁹ I. M. Peruzzi,^{27,59} M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ M. Bondioli,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ V. Del Gamba,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ F. Martinez-Vidal,^{60,‡} M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ F. Sandrelli,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ K. Paick,⁶¹ D. E. Wagoner,⁶¹ N. Danielson,⁶² P. Elmer,⁶² C. Lu,⁶² V. Miftakov,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² F. Bellini,⁶³ G. Cavoto,^{62,63} R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ M. Pierini,⁶³ G. Piredda,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ S. Christ,⁶⁴ G. Wagner,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ N. I. Geddes,⁶⁵ G. P. Gopal,⁶⁵ E. O. Olaiya,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ P.-F. Giraud,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Langer,⁶⁶ M. Legendre,⁶⁶ G. W. London,⁶⁶ B. Mayer,⁶⁶ G. Schott,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ F. X. Yumiceva,⁶⁷ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ G. De Nardo,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ E. E. Elsen,⁶⁸ S. Fan,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ C. Hast,⁶⁸ T. Hryn'ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ S. Petrak,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ G. Simi,⁶⁸ A. Snyder,⁶⁸ A. Soha,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ J. Va'vra,⁶⁸ S. R. Wagner,⁶⁸ M. Weaver,⁶⁸ A. J. R. Weinstein,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ T. I. Meyer,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ M. Saleem,⁷⁰ F. R. Wappler,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² H. Kim,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ C. Borean,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ P. Poropat,^{75,§} L. Vitale,⁷⁵ G. Vuagnin,⁷⁵ R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ H. R. Band,⁷⁸ S. Dasu,⁷⁸ M. Datta,⁷⁸ F. Di Lodovico,⁷⁸ A. M. Eichenbaum,⁷⁸ J. J. Hollar,⁷⁸ J. R. Johnson,⁷⁸ P. E. Kutter,⁷⁸ H. Li,⁷⁸ R. Liu,⁷⁸ A. Mihalyi,⁷⁸ A. K. Mohapatra,⁷⁸ Y. Pan,⁷⁸ R. Prepost,⁷⁸ S. J. Sekula,⁷⁸ P. Tan,⁷⁸ J. H. von Wimmersperg-Toeller,⁷⁸ J. Wu,⁷⁸ S. L. Wu,⁷⁸ Z. Yu,⁷⁸ and H. Neal⁷⁹

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at Riverside, Riverside, California 92521, USA

¹⁵University of California at San Diego, La Jolla, California 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁸California Institute of Technology, Pasadena, California 91125, USA

¹⁹University of Cincinnati, Cincinnati, Ohio 45221, USA

²⁰University of Colorado, Boulder, Colorado 80309, USA

²¹Colorado State University, Fort Collins, Colorado 80523, USA

²²Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²³Ecole Polytechnique, LLR, F-91128 Palaiseau, France

²⁴University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

- ²⁵Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- ²⁶Florida A&M University, Tallahassee, Florida 32307, USA
- ²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ²⁹Harvard University, Cambridge, Massachusetts 02138, USA
- ³⁰Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
- ³¹Imperial College London, London SW7 2AZ, United Kingdom
- ³²University of Iowa, Iowa City, Iowa 52242, USA
- ³³Iowa State University, Ames, Iowa 50011-3160, USA
- ³⁴Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
- ³⁵Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- ³⁶University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³⁷Queen Mary, University of London, London E1 4NS, United Kingdom
- ³⁸University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ³⁹University of Louisville, Louisville, Kentucky 40292, USA
- ⁴⁰University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴¹University of Maryland, College Park, Maryland 20742, USA
- ⁴²University of Massachusetts, Amherst, Massachusetts 01003, USA
- ⁴³Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- ⁴⁴McGill University, Montréal, Quebec, Canada H3A 2T8
- ⁴⁵Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁶University of Mississippi, University, Mississippi 38677, USA
- ⁴⁷Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Quebec, Canada H3C 3J7
- ⁴⁸Mount Holyoke College, South Hadley, Massachusetts 01075, USA
- ⁴⁹Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁵⁰NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵¹University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵²Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA
- ⁵³The Ohio State University, Columbus, Ohio 43210, USA
- ⁵⁴University of Oregon, Eugene, Oregon 97403, USA
- ⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵⁶Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France
- ⁵⁷Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
- ⁵⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁵⁹Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
- ⁶⁰Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- ⁶¹Prairie View A&M University, Prairie View, Texas 77446, USA
- ⁶²Princeton University, Princeton, New Jersey 08544, USA
- ⁶³Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶⁴Universität Rostock, D-18051 Rostock, Germany
- ⁶⁵Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, United Kingdom
- ⁶⁶DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- ⁶⁷University of South Carolina, Columbia, South Carolina 29208, USA
- ⁶⁸Stanford Linear Accelerator Center, Stanford, California 94309, USA
- ⁶⁹Stanford University, Stanford, California 94305-4060, USA
- ⁷⁰State University of New York, Albany, New York 12222, USA
- ⁷¹University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁷²University of Texas at Austin, Austin, Texas 78712, USA
- ⁷³University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁷⁴Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁵Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷⁶Vanderbilt University, Nashville, Tennessee 37235, USA
- ⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- ⁷⁸University of Wisconsin, Madison, Wisconsin 53706, USA
- ⁷⁹Yale University, New Haven, Connecticut 06511, USA
- (Received 15 March 2004; published 10 August 2004)

We present a measurement of the time-dependent CP asymmetry for the neutral B -meson decay $B^0 \rightarrow \phi K^0$. We use a sample of approximately 114×10^6 B -meson pairs taken at the $Y(4S)$ resonance with the BABAR detector at the PEP-II B -meson factory at SLAC. We reconstruct the CP eigenstates ϕK_S^0 and ϕK_L^0 , where $\phi \rightarrow K^+ K^-$, $K_S^0 \rightarrow \pi^+ \pi^-$, and K_L^0 is observed via its

hadronic interactions. The other B meson in the event is tagged as either a B^0 or \bar{B}^0 from its decay products. The values of the CP -violation parameters are $S_{\phi K} = 0.47 \pm 0.34(\text{stat})_{-0.06}^{+0.08}(\text{syst})$ and $C_{\phi K} = 0.01 \pm 0.33(\text{stat}) \pm 0.10(\text{syst})$.

DOI: 10.1103/PhysRevLett.93.071801

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

Decays of B mesons into charmless hadronic final states with a ϕ meson are dominated by $b \rightarrow s\bar{s}s$ gluonic penguin amplitudes, possibly with smaller contributions from electroweak penguins, while other standard model (SM) amplitudes are strongly suppressed [1]. In the SM, CP violation arises from a single complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [2]. Neglecting CKM-suppressed contributions, the time-dependent CP -violating asymmetries in the decays $B^0 \rightarrow \phi K^0$ and $B^0 \rightarrow J/\psi K^0$ are proportional to the same parameter $\sin 2\beta$ [3], where the latter decay is dominated by tree diagrams. Since many scenarios of physics beyond the SM introduce additional diagrams with heavy particles in the penguin loops and new CP -violating phases, comparison of CP -violating observables with SM expectations is a sensitive probe for new physics. Measurements of $\sin 2\beta$ in B decays to charmonium such as $B^0 \rightarrow J/\psi K_S^0$ have been reported by the BABAR [4] and Belle [5] Collaborations, and the world average for $\sin 2\beta$ is 0.731 ± 0.056 [6]. The Belle Collaboration states [7] that their result for $B^0 \rightarrow \phi K_S^0$, $\sin 2\beta = -0.96 \pm 0.50_{-0.11}^{+0.09}$, suggests that there is a large CP -violating phase in its decay amplitude, which cannot be explained by the SM.

In this Letter we report a measurement of the time-dependent CP asymmetry in the final state ϕK^0 based on an integrated luminosity of approximately 108 fb^{-1} collected at the $\Upsilon(4S)$ resonance with the BABAR detector [8] at the PEP-II asymmetric e^+e^- collider [9] located at the Stanford Linear Accelerator Center.

From a $B^0\bar{B}^0$ meson pair we fully reconstruct one meson, B_{CP} , in the final state ϕK^0 , and partially reconstruct the recoil B meson, B_{tag} . We examine B_{tag} for evidence that it decayed either as B^0 or \bar{B}^0 (flavor tag). The asymmetric beam configuration in the laboratory frame provides a boost of $\beta\gamma = 0.56$ to the $\Upsilon(4S)$, which allows the determination of the proper decay time difference $\Delta t = t_{CP} - t_{\text{tag}}$ from the vertex separation of the two neutral B mesons along the beam (z) axis. The decay rate f_+ (f_-) when the tagging meson is a B^0 (\bar{B}^0) is given by

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} [1 \pm S_{\phi K} \sin(\Delta m_d \Delta t) \mp C_{\phi K} \cos(\Delta m_d \Delta t)], \quad (1)$$

where τ_{B^0} is the neutral B meson mean lifetime, and Δm_d is the $B^0\bar{B}^0$ oscillation frequency. The time-dependent CP -violating asymmetry is defined as $A_{CP} \equiv (f_+ - f_-)/(f_+ + f_-)$. In the SM, decays that proceed purely via the $b \rightarrow s\bar{s}s$ penguin transitions have CP parameters

$S_{\phi K} = -\eta_f \sin 2\beta$ and $C_{\phi K} = 0$, where $\beta \equiv \arg[-V_{cd}V_{cb}^*/V_{td}V_{tb}^*]$. Here V_{ik} is the CKM matrix element for quarks i and k , and the CP eigenvalue is $\eta_f = -1$ ($+1$) for ϕK_S^0 (ϕK_L^0).

The B_{CP} candidate is reconstructed in the decay mode ϕK^0 with $\phi \rightarrow K^+K^-$; the K^0 is either a K_L^0 or a $K_S^0 \rightarrow \pi^+\pi^-$. We combine pairs of oppositely charged tracks extrapolated to a common vertex to form ϕ and K_S^0 candidates. For the charged tracks from the ϕ decay we require at least 12 measured drift-chamber (DCH) coordinates and a minimal transverse momentum p_T of $0.1 \text{ GeV}/c$. The tracks must also originate within 1.5 cm in xy and $\pm 10 \text{ cm}$ along the z axis of the nominal beam spot. Tracks used to reconstruct the ϕ mesons are distinguished from pion and proton tracks via a requirement on a likelihood ratio that combines dE/dx information from the silicon vertex tracker (SVT) and the DCH for tracks with momentum $p < 0.7 \text{ GeV}/c$. For tracks with higher p , dE/dx in the DCH and the Cherenkov angle and the number of photons as measured by the internally reflecting ring-imaging Cherenkov detector are used in the likelihood. The two-kaon invariant mass must be within $16 \text{ MeV}/c^2$ of the nominal ϕ mass [6].

For tracks corresponding to K_S^0 and B_{tag} daughters our requirements are less restrictive. A $K_S^0 \rightarrow \pi^+\pi^-$ candidate is accepted if its two-pion invariant mass is within $15 \text{ MeV}/c^2$ of the nominal K^0 mass [6], its reconstructed decay vertex is separated from the ϕ decay vertex by at least 3 standard deviations, and the angle between the line connecting the ϕ and K_S^0 decay vertices and the K_S^0 momentum direction is less than 45 mrad .

We identify a K_L^0 candidate as in our $B^0 \rightarrow J/\psi K_L^0$ analysis [10] either as a cluster of energy deposits in the electromagnetic calorimeter (EMC) or as a cluster of hits in two or more layers of the instrumented flux return (IFR) that cannot be associated with any charged track in the event. The K_L^0 energy is not well measured. Therefore, we determine the K_L^0 laboratory momentum from its flight direction as measured from the EMC or IFR cluster and the constraint that the invariant ϕK_L^0 mass agrees with the known B^0 mass. In those cases where the K_L^0 is detected in both the IFR and EMC we use the angular information from the EMC, as it has a higher precision. In order to reduce background from π^0 decays, we reject an EMC K_L^0 candidate cluster if it forms an invariant mass between 100 and $150 \text{ MeV}/c^2$ with any other cluster in the event under the $\gamma\gamma$ hypothesis, or if it has energy greater than 1 GeV and contains two shower maxima consistent with two photons from a π^0 decay. The remain-

ing background of photons and overlapping showers is further reduced with the use of a neural network constructed from cluster shape variables, trained on Monte Carlo (MC) simulated $B^0 \rightarrow \phi K_L^0$ and measured radiative Bhabha events, and tested on measured $e^+e^- \rightarrow \phi(\rightarrow K_S^0 K_L^0)\gamma$ and $B^0 \rightarrow J/\psi K_L^0$ events. The final ϕK_L^0 sample consists of approximately equal numbers of IFR and EMC K_L^0 candidates.

The results are extracted from an extended unbinned maximum likelihood fit for which we parametrize the distributions of several kinematic and topological variables for signal and background events in terms of probability density functions (PDFs) [11]. The background arises primarily from random combinations of tracks produced in events of the type $e^+e^- \rightarrow q\bar{q}$, where $q = u, d, s, c$ (continuum). Background from other B decay final states with and without charm is estimated with MC simulations. Opposite CP contributions from the $K^+K^-K^0$ final state (K^+K^-S wave) are estimated with a moment analysis [12] on data to be less than 6.6% and treated as a systematic error. The shapes of event variable distributions are obtained from signal and background MC samples and high-statistics data control samples.

Each B_{CP} candidate is characterized by the energy difference $\Delta E = E_B^* - \frac{1}{2}\sqrt{s}$ and, except for $B^0 \rightarrow \phi K_L^0$, the beam-energy-substituted mass $m_{ES} = \sqrt{(\frac{1}{2}s + \vec{p}_0 \cdot \vec{p}_B)^2/E_0^2 - p_B^2}$ [8]. The subscripts 0 and B refer to the initial $Y(4S)$ and the B_{CP} candidate, respectively, and the asterisk denotes the $Y(4S)$ rest frame. For signal events, ΔE is expected to peak at zero, and m_{ES} at the nominal B mass. We require $\Delta E < 0.08$ GeV for $B^0 \rightarrow \phi K_L^0$ and $|\Delta E| < 0.2$ GeV and $m_{ES} > 5.2$ GeV/ c^2 for $B^0 \rightarrow \phi K_S^0$. In the fit we also use the helicity angle θ_H , which is defined as the angle between the directions of the K^+ and the parent B_{CP} in the K^+K^- rest frame. The $\cos\theta_H$ distribution for pseudoscalar-vector B decay modes is $\cos^2\theta_H$, and for the combinatorial background it is nearly uniform.

In continuum events, particles appear bundled into jets. This topology can be characterized with several variables computed in the c.m. frame. One such quantity is the angle θ_T between the thrust axis of the B_{CP} candidate and the thrust axis formed from the other charged and neutral particles in the event. We also use the angle θ_B between the B_{CP} momentum and the beam axis, and the sum of the momenta p_i of the other charged and neutral particles in the event weighted by the Legendre polynomials $L_0(\theta_i)$ and $L_2(\theta_i)$ where θ_i is the angle between the momentum of particle i and the thrust axis of the B_{CP} candidate. For $B^0 \rightarrow \phi K_S^0$ candidates, we combine these variables into a Fisher discriminant \mathcal{F} [13] after requiring $|\cos\theta_T| < 0.9$. In this mode background from other B decays is negligible, as demonstrated in MC simulation studies.

More stringent criteria must be applied to suppress backgrounds in the case of $B^0 \rightarrow \phi K_L^0$ candidates, and we require $|\cos\theta_T| < 0.8$ and $|\cos\theta_B| < 0.85$. We define the missing momentum \vec{p}_{miss} , calculated in the laboratory frame from the sum of beam momenta and all tracks and EMC clusters, excluding the K_L^0 candidate. We require the polar angle θ_{miss} of the missing momentum with respect to the beam direction to be greater than 0.3 rad. The cosine of the angle between \vec{p}_{miss} and the K_L^0 direction, θ_K , must satisfy $\cos\theta_K > 0.6$. In the plane transverse to the beam direction, the difference between the missing momentum projected along the K_L^0 direction and the calculated K_L^0 momentum must be greater than -0.75 GeV/ c . In the Fisher discriminant we replace $|\cos\theta_B|$ by the cosine of the angle between the missing momentum and the K^+ from the ϕ decay. In the ϕK_L^0 sample about 1.4% of the events originate from charm B decays; 0.7% originate from charmless B decays. The dominant CP contamination is the mode $B \rightarrow \phi K^{*0}$, where the K^{*0} decays to $K_L^0\pi^0$; we expect four events in the region $\Delta E < 0.01$ GeV. In the likelihood fit we explicitly parametrize backgrounds from both charm and charmless B decays as derived from MC simulations.

All the other tracks and clusters that are not associated with the reconstructed $B^0 \rightarrow \phi K^0$ decay are used to form the B_{tag} , and its flavor is determined with a multivariate tagging algorithm [4]. The tagging efficiency ϵ and mistag probability w in four hierarchical and mutually exclusive categories is measured from fully reconstructed B^0 decays into the $D^{(*)-}X^+$ ($X^+ = \pi^+, \rho^+, a_1^+$) and $J/\psi K^{*0}$ ($K^{*0} \rightarrow K^+\pi^-$) flavor eigenstates (B_{flav} sample). The analyzing power $\epsilon(1 - 2w)^2$ is $(28.7 \pm 0.7)\%$.

A detailed description of the Δt reconstruction algorithm is given in Ref. [10]. The B_{CP} vertex resolution is dominated by the ϕ vertex. The average Δz resolution is 190 μm and is dominated by the tagging vertex in the event. Thus, we can characterize the resolution with the much larger B_{flav} sample, which we fit simultaneously with the CP samples. The amplitudes for the B_{CP} asymmetries and for the B_{flav} flavor oscillations are reduced by the same factor due to wrong tags. Both distributions are convoluted with a common Δt resolution function, and the backgrounds are accounted for by adding terms to the likelihood, incorporated with different assumptions about their Δt evolution and resolution function [10].

Since we measure the correlations among the observables to be small in the data samples entering the fit, we take the probability density function $\mathcal{P}_{i,c}^j$ for each event j to be a product of the PDFs for the separate observables. For each event hypothesis i (signal, backgrounds) and tagging category c , we define $\mathcal{P}_{i,c}^j = \mathcal{P}_i(m_{ES}) \cdot \mathcal{P}_i(\Delta E) \cdot \mathcal{P}_i(\mathcal{F}) \cdot \mathcal{P}_i(\cos\theta_H) \cdot \mathcal{P}_i(\Delta t; \sigma_{\Delta t}, c)$, where for the ϕK_L^0 mode $\mathcal{P}_i(m_{ES}) = 1$ and for the flavor sample $\mathcal{P}_i(\mathcal{F}) \cdot \mathcal{P}_i(\cos\theta_H) = 1$. The $\sigma_{\Delta t}$ is the error on Δt for a given event. The likelihood function for each decay chain is

then

$$\mathcal{L} = \prod_c \exp\left(-\sum_i N_{i,c}\right) \prod_j \left[\sum_i N_{i,c} \mathcal{P}_{i,c}^j\right], \quad (2)$$

where $N_{i,c}$ is the yield of events of hypothesis i found by the fitter in category c , and N_c is the number of category c events in the sample. The total sample consists of 86 200 B_{flav} , 2138 ϕK_S^0 , and 4730 ϕK_L^0 candidates. We find 70 ± 9 ϕK_S^0 and 52 ± 16 ϕK_L^0 signal events. The signal yields in both the ϕK^0 channels agree well with our determination of the branching fraction for $B^0 \rightarrow \phi K^0$ [14]. Figure 1 shows the m_{ES} (ΔE) distribution of ϕK_S^0 (ϕK_L^0) events together with the result from the fit after a requirement on the likelihood (computed without the variable plotted) to enhance the sensitivity.

We determine the CP parameters $S_{\phi K}$ and $C_{\phi K}$ along with an additional 38 free parameters: the efficiency per tagging category (4 parameters), the average mistag fraction and the difference between B^0 and \bar{B}^0 mistags for each tagging category (8), and the signal Δt resolution (9). For the background we parametrize time dependence (6), Δt resolution (3), and mistag fractions (8). We fix τ_{B^0} and Δm_d to the world averages [6]. The determination of the mistag fractions and Δt -resolution parameters is dominated by the high-statistics B_{flav} sample. The fit was tested with a parametrized simulation of a large number of data-sized experiments and full detector simulated events for the different signal and background samples. The likelihood of our data fit agrees with the likelihoods from fits to the simulated data. The expected error for $S_{\phi K}$ is 0.40 and for $C_{\phi K}$ is 0.29. Compared to the measured values 27% of the fits to the simulated data have a smaller error value for $S_{\phi K}$ and 12% have a higher error value for $C_{\phi K}$. The fit was also verified with our $J/\psi K_S^0$ data sample and a control sample of 232 ϕK^+ candidates where one expects $S_{\phi K^+} = C_{\phi K^+} = 0$. We measure $S_{\phi K^+} = 0.23 \pm 0.24$ and $C_{\phi K^+} = -0.14 \pm 0.18$ with sta-




FIG. 1 (color online). Distribution of the event variable (a) m_{ES} for the ϕK_S^0 final state and (b) ΔE for the ϕK_L^0 final state after reconstruction and a requirement for the likelihood with total signal efficiency of 32% and 5%, respectively. The solid line represents the fit result for the total event yield and the dotted line for the background.

tistical errors only. The simultaneous fit to the ϕK^0 and flavor decay modes yields

$$S_{\phi K} = 0.47 \pm 0.34(\text{stat})^{+0.08}_{-0.06}(\text{syst}),$$

$$C_{\phi K} = 0.01 \pm 0.33(\text{stat}) \pm 0.10(\text{syst}).$$

The result in the dominant channel $B^0 \rightarrow \phi K_S^0$ is $S_{\phi K} = 0.45 \pm 0.43$ and $C_{\phi K} = -0.38 \pm 0.37$ with statistical errors only. Figure 2 shows the Δt distributions of the B^0 - and the \bar{B}^0 -tagged subsets together with the raw asymmetry for the ϕK_S^0 and ϕK_L^0 events with the result of the combined time-dependent CP -asymmetry fit superimposed.

We consider systematic uncertainties in the CP coefficients $S_{\phi K}$ and $C_{\phi K}$ due to the event-yield determination in the two channels (± 0.01 for $S_{\phi K}$, ± 0.05 for $C_{\phi K}$), contributions from B^0 final states with opposite CP ($+0.06$, ± 0.02), the parametrization of PDFs for the event yield in signal and background (± 0.02 , ± 0.05), composition and CP asymmetry of the background (± 0.03 , ± 0.03), the assumed parametrization of the Δt resolution function (± 0.02 , ± 0.01), the m_{ES} background parametrization (± 0.02 , ± 0.05), a possible difference in




FIG. 2 (color online). Plots (a) and (b) show the Δt distributions of B^0 - and \bar{B}^0 -tagged ϕK_S^0 events. The solid lines refer to the fit for all events; the dashed lines correspond to the background. Plot (c) shows the asymmetry. A requirement for the event likelihood is applied. Plots (d)–(f) are the corresponding plots for ϕK_L^0 events.

the efficiency for B^0 and \bar{B}^0 ($\pm 0.01, \pm 0.02$), the fixed values for Δm_d and τ_B ($\pm 0.00, \pm 0.01$), the beam-spot position ($\pm 0.01, \pm 0.01$), and uncertainties in the SVT alignment ($\pm 0.01, \pm 0.01$). The bias in the coefficients due to the fit procedure ($\pm 0.03, \pm 0.01$) is included in the uncertainty without making corrections to the final results. We estimate errors due to the effect of doubly CKM-suppressed decays [15] to be ($\pm 0.01, \pm 0.03$). We add these contributions in quadrature to obtain the total systematic uncertainty.

In summary, we have measured the time-dependent CP asymmetries in the combined B -meson final states ϕK_S^0 and ϕK_L^0 . We obtain values for the CP -violation parameters $S_{\phi K}$ and $C_{\phi K}$ that agree within 1 standard deviation with the ones measured in the charmonium channels [4,5]; the central value of $S_{\phi K}$ is also consistent with no CP asymmetry at the 1.3σ level. Our value of $S_{\phi K}$ differs by 2.3 standard deviations from that measured by the Belle Collaboration [7].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), by NSERC (Canada), by IHEP (China), by CEA and CNRS-IN2P3 (France), by BMBF and DFG (Germany), by INFN (Italy), by FOM (The Netherlands), by NFR (Norway), by MIST (Russia), and by PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, from Research Corporation, and from Alexander von Humboldt Foundation.

*Now at Department of Physics, University of Warwick, Coventry, United Kingdom.

[†]Also with Università della Basilicata, Potenza, Italy.

[‡]Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain.

[§]Deceased.

- [1] N. G. Deshpande and J. Trampetic, *Phys. Rev. D* **41**, 895 (1990); N. G. Deshpande and G. He, *Phys. Lett. B* **336**, 471 (1994); R. Fleischer, *Z. Phys. C* **62**, 81 (1994); Y. Grossman *et al.*, *Phys. Rev. D* **68**, 015004 (2003).
- [2] N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
- [3] A. B. Carter and A. I. Sanda, *Phys. Rev. D* **23**, 1567 (1981); I. I. Bigi and A. I. Sanda, *Nucl. Phys.* **B193**, 85 (1981); Y. Grossman and M. P. Worah, *Phys. Lett. B* **395**, 241 (1997); R. Fleischer, *Int. J. Mod. Phys. A* **12**, 2459 (1997); D. London and A. Soni, *Phys. Lett. B* **407**, 61 (1997).
- [4] *BABAR* Collaboration, B. Aubert *et al.*, *Phys. Rev. Lett.* **89**, 201802 (2002).
- [5] Belle Collaboration, K. Abe *et al.*, *Phys. Rev. D* **66**, 071102 (2002).
- [6] Particle Data Group, K. Hagiwara *et al.*, *Phys. Rev. D* **66**, 010001 (2002).
- [7] Belle Collaboration, K. Abe *et al.*, *Phys. Rev. Lett.* **91**, 261602 (2003).
- [8] *BABAR* Collaboration, B. Aubert *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 1 (2002).
- [9] PEP-II Conceptual Design Report, SLAC Report No. SLAC-R-418, 1993.
- [10] *BABAR* Collaboration, B. Aubert *et al.*, *Phys. Rev. D* **66**, 032003 (2002).
- [11] *BABAR* Collaboration, B. Aubert *et al.*, *Phys. Rev. Lett.* **87**, 151801 (2001).
- [12] S. U. Chung, *Phys. Rev. D* **56**, 7299 (1997).
- [13] R. A. Fisher, *Ann. Eugen.* **7**, 179 (1936).
- [14] *BABAR* Collaboration, B. Aubert *et al.*, *Phys. Rev. D* **69**, 011102 (2004).
- [15] O. Long, M. Baak, R. N. Cahn, and D. Kirkby, *Phys. Rev. D* **68**, 034010 (2003).