

UC Irvine

ICTS Publications

Title

Gas signatures from Escherichia coli and Escherichia coli-inoculated human whole blood

Permalink

<https://escholarship.org/uc/item/1b86w5tk>

Authors

Umber, Brandon J
Shin, Hye-Won
Meinardi, Simone
et al.

Publication Date

2013-07-10

Peer reviewed

RESEARCH

Open Access

Gas signatures from *Escherichia coli* and *Escherichia coli*-inoculated human whole blood

Brandon J Umber^{1†}, Hye-Won Shin^{2,3*†}, Simone Meinardi¹, Szu-Yun Leu^{2,3}, Frank Zaldivar^{2,3}, Dan M Cooper^{2,3} and Donald R Blake¹

Abstract

Background: The gaseous headspace above naïve *Escherichia Coli* (*E. coli*) cultures and whole human blood inoculated with *E. coli* were collected and analyzed for the presence of trace gases that may have the potential to be used as novel, non-invasive markers of infectious disease.

Methods: The naïve *E. coli* culture, LB broth, and human whole blood or *E. coli* inoculated whole blood were incubated in hermetically sealable glass bioreactors at 37°C for 24 hrs. LB broth and whole human blood were used as controls for background volatile organic compounds (VOCs). The headspace gases were collected after incubation and analyzed using a gas chromatographic system with multiple column/detector combinations.

Results: Six VOCs were observed to be produced by *E. coli*-infected whole blood while there existed nearly zero to relatively negligible amounts of these gases in the whole blood alone, LB broth, or *E. coli*-inoculated LB broth. These VOCs included dimethyl sulfide (DMS), carbon disulfide (CS₂), ethanol, acetaldehyde, methyl butanoate, and an unidentified gas S. In contrast, there were several VOCs significantly elevated in the headspace above the *E. coli* in LB broth, but not present in the *E. coli*/blood mixture. These VOCs included dimethyl disulfide (DMDS), dimethyl trisulfide (DMTS), methyl propanoate, 1-propanol, methylcyclohexane, and unidentified gases R2 and Q.

Conclusions: This study demonstrates 1) that cultivated *E. coli* in LB broth produce distinct gas profiles, 2) for the first time, the ability to modify *E. coli*-specific gas profiles by the addition of whole human blood, and 3) that *E. coli*-human whole blood interactions present different gas emission profiles that have the potential to be used as non-invasive volatile biomarkers of *E. coli* infection.

Keywords: Gas biomarker; Volatile organic compounds (VOCs); *Escherichia coli*; Cell culture; Human whole blood culture; Dimethyl sulfide (DMS); Carbon disulfide (CS₂); Ethanol; Acetaldehyde; Methyl butanoate

Background

Over the past decade, the interest in using exhaled gases as non-invasive biomarkers in clinical diagnostics and therapeutic monitoring has steadily increased [1]. Considerable efforts have been made to determine clinically applicable breath biomarkers for such conditions as asthma, diabetes, and various cancers [2-7]. Evidence also exists that cultivated microorganisms can generate species or genus-specific profiles of VOCs [8-13]. Additionally, unique gas signatures have been observed in the exhaled

breath of patients with specific infections, including bronchopneumonia [14] and *Helicobacter pylori* [15].

Identifying and quantifying gases in the exhaled human breath provide a non-invasive means of obtaining information about host responses to infection. However, with the exception of nitric oxide, the cellular sources or mechanisms responsible for the observed gases have not been fully investigated. It is conceivable that the host immune response may yield gases unique to one pathogen or another. *Escherichia Coli* (*E. coli*) was used as a model infectious organism to 1) determine the unique gas profile of bacteria in a “naïve” state and 2) investigate the VOCs that result from the bacteria and whole blood interactions. VOC profiles found in the interaction of bacteria and immune cells containing whole blood may

* Correspondence: hyewons@uci.edu

†Equal contributors

²Department of Pediatrics, University of California, Irvine, CA 92697, USA

³Institute for Clinical and Translational Sciences, University of California, Irvine, CA 92697, USA

Full list of author information is available at the end of the article



then provide a beginning interpretation of the exhaled breath gases from infected subjects.

A hermetically sealable bioreactor was used to grow the cultures and collect the gaseous headspace. The headspace was then analyzed for trace gases produced and released by the microorganisms growing in culture using a multi-column and multi-detector analytical gas chromatography (GC) methodology [16].

This approach will soon be extended to additional infectious agents and it is anticipated that the technology will eventually demonstrate the capability to discriminate between multiple infectious pathogens by means of their unique VOC profiles.

Methods

Blood Sampling. The blood was drawn using EDTA-treated vacutainers from adult male subjects through the Institute for Clinical Translational Sciences (ICTS) general blood donor program. Subjects having a history of any chronic medical conditions or use of any medications were excluded from the study. The Institutional Review Board at the University of California, Irvine, approved the study.

E. Coli culture and headspace Gas collection

E. coli was purchased from Invitrogen (One Shot TOP10 Chemically Competent, Invitrogen, NY). The genotype of this *E. coli* is similar to DH10B strain. The complete genome sequence has been reported elsewhere (<http://www.ncbi.nlm.nih.gov/pubmed/18245285>). Gibco LB broth from Invitrogen was used for the study. The vials of *E. coli*-inoculated whole blood (1×10^6 *E. coli*/10 mL whole blood) and naïve *E. coli* inoculated LB broth (1×10^6 *E. coli*/10 mL LB) were placed inside bioreactors specifically designed to collect the gaseous headspace above aqueous cultures [16]. For comparison, vials of LB broth only (10 mL) or vials of whole blood only (10 mL) were also placed inside bioreactors. To minimize ambient gaseous contaminants, each bioreactor was flushed with whole air containing low levels of VOCs and 5% CO₂ for 10 minutes at 125 mL per minute. The bioreactors were then placed in an incubator at 37°C for 24 hrs. The headspace gases were collected after incubation and analyzed using a gas chromatographic system with multiple column/detector combinations [16-18]. Following sample collection, the bioreactor was disassembled and the cells were immediately collected and counted.

GC system

The analyses of the headspace gases were performed on a system previously developed to measure trace atmospheric gases [17]. Briefly, a 233 cm³ (at STP) sample was cryogenically pre-concentrated and injected into a multi-column/detector GC system. The system consists of

three Hewlett-Packard 6890 GC units (Wilmington, DE) using a combination of columns and detectors capable of separating and quantifying hundreds of gases, including, but not limited to, non-methane hydrocarbons (NMHCs), alkyl nitrates, and halocarbons in the ppbv to pptv range (10^{-9} - 10^{-12}). The detectors include flame ionization detectors (FIDs), electron capture detectors (ECDs), and a mass selective detector (MSD). Preliminary identifications of the unknown compounds were made using GC-MSD and verification was obtained by injecting the diluted headspace of pure compounds (Sigma-Aldrich, St. Louis, MO) to ensure that both the elution time and the mass spectrum matched that of the unknown.

Data analysis

Four study conditions, Broth only, Broth + *E. coli*, Blood only, and Blood + *E. coli* were evaluated in each of the 10 experimental days. The same blood sample and *E. coli* sample were used in the same day. A total of 117 gases were examined for each sample. Thirty-eight gases were excluded; of these, 31 gases were below the limit of detection in over 80% of the samples and seven gases had concentrations less than 100 pptv in all samples. For any gas that could not be detected in a sample, the level was set to zero. Wilcoxon signed-rank test, a non-parametric method for paired data, was used to compare the difference between Broth only and Broth + *E. coli*, between Blood only and Blood + *E. coli*, between Broth + *E. coli* and Blood + *E. coli*, and between Broth only and Blood only. The method was chosen because the sample size was small, the distributions of most gases were highly skewed, and the experiments performed on the same day were considered more correlated than those performed on different days. Multiple-comparison adjustment was further applied using Benjamini and Hochberg's false discover rate (FDR) method [19]. A comparison was considered significant if FDR < 0.01. All analyses were performed using SAS 9 (Cary, NC) and all data were presented with median and range (minimum, maximum).

Results

After statistical analysis, 40 gases were allocated into five categories based on the pattern of gas release: category 1: VOCs elevated from *E. coli*-inoculated human whole blood, (see Table 1 and Figure 1); category 2: VOCs elevated from *E. coli* in LB broth (see Table 1 and Figure 2); category 3: VOCs elevated from naïve *E. coli* in broth, but decreased by *E. coli*-inoculated whole blood (see Table 1, and Figure 3); category 4: VOCs mainly from pure LB broth only (see Table 2 and Figure 4); category 5: VOCs mainly from human whole blood only (see Table 3 and Figure 5).

In category 1, there were six VOCs observed to be significantly increased from *E. coli*-inoculated whole blood

Table 1 VOCs of interest

Gas	Stat. diff.	Broth		Blood	
		+ <i>E. coli</i>	-	+ <i>E. coli</i>	-
Category 1: VOCs elevated in <i>E. coli</i>-infected blood					
Dimethyl sulfide [pptv]	2,4	970 (310,2310)	16010 (180,29500)	3770 (1250,8820)	95920 (25740,234400)
Carbon disulfide [pptv]	2,3,4	85 (55,470)	295 (80,790)	4390 (2450,10040)	8940 (2620,16550)
Acetaldehyde [ppbv]		736 (558,1089)	645 (44,1359)	45 (22,145)	1158 (208,3493)
Ethanol [ppbv]	1,2,3	202 (28,564)	8388 (2037,10867)	218 (5,604)	15486 (7199,40441)
Unk S [§]	2,3	31 (0,316)	74 (12,267)	675 (0,3311)	15001 (2208,37426)
Methyl Butanoate [pptv]		0 (0,25)	0 (0,40)	60 (0,405)	2675 (330,9720)
Category 2: VOCs elevated in naïve <i>E. coli</i> in LB broth					
Isoprene [pptv]	1,4	0 (0,50)	1590 (665,2175)	1040 (350,2785)	1200 (455,2050)
Carbonyl sulfide (OCS) [pptv]	1	2240 (1440,4850)	8360 (4360,20180)	3850 (1970,22040)	6540 (3810,30340)
Unk R1 [§]		7474 (1568,42622)	116738 (15070,1005677)	3758 (105,20243)	2435 (785,10277)
Category 3: VOCs elevated in naïve <i>E. coli</i> but substantially lower from co-culture of <i>E. coli</i> and whole blood					
Dimethyl disulfide [ppbv]	1,3,4	164 (133,185)	354 (182,537)	0.7 (0,3,2)	2 (0,6,4)
Methyl Propanoate [pptv]		3300 (2100,4300)	14500 (5660,21000)	400 (0,1400)	600 (0,1100)
Dimethyl trisulfide [pptv]	1,3	2900 (1900,9700)	160400 (22700,253300)	800 (10,6900)	400 (150,5400)
Unk_Q [§]		2052 (0,13997)	57320 (6317,467654)	620 (0,6232)	289 (0,3787)
1-Propanol [pptv]		175 (0,90200)	757300 (472900,1920800)	700 (0,93600)	33400 (4100,90200)
Methylcyclohexane [pptv]		220 (60,990)	1600 (850,3200)	130 (0,1720)	170 (80,980)
Unk_R2 [§]		416 (0,5269)	15852 (2228,152789)	767 (0,9829)	674 (0,8522)

1: significantly different between Broth only and Broth + *E. coli*; 2: significantly different between Blood only and Blood + *E. coli*; 3: significantly different between Broth + *E. coli* i and Blood + *E. coli*; 4: significantly different between Broth only and blood only.

Data were presented with median and range (minimum, maximum).

[§]unknown peaks were presented as area under the curve.

when compared to VOCs from whole blood alone. These VOCs include dimethyl sulfide, carbon disulfide, ethanol, acetaldehyde, methyl butanoate and an unidentified gas S (see Table 1 and Figure 1 for examples of this category). Ethanol was also emitted from the *E. coli* cultured in the LB broth; however, the concentrations were relatively small when compared to the *E. coli*-inoculated whole blood (see Figure 1B).

In contrast, ten VOCs (see Table 1, category 2 and 3) were significantly elevated from *E. coli* in LB broth (Figures 2 and 3 show several examples). Isoprene, carbonyl sulfide (OCS), and unidentified gas peak R1 were significantly elevated in the headspace of *E. coli* in LB broth. These gases detected in the headspace of whole blood, and *E. coli*-inoculated whole blood. However, emitted isoprene, carbonyl sulfide and unidentified gas peak R1 levels from whole blood and *E. coli*-inoculated whole blood were not different (see Figure 2). Interestingly, seven out of these ten VOCs were substantially lower in the presence of human whole blood (see category 3 and Figure 3). These VOCs included dimethyl disulfide, dimethyl trisulfide, methylpropanoate, 1-propanol, methylcyclohexane, and unidentified gas peaks R2, and Q. The emitted levels from whole blood

and *E. coli*-inoculated whole blood were not different in these seven gases.

Table 2 lists 16 VOCs that were observed to be significantly higher in LB broth when compared to the levels in whole blood (category 4). The emitted levels of 14 out of total 16 gases in this category (except n-Propylbenzene and unidentified gas peak M) were decreased in the presence of *E. coli*. Human whole blood emitted trace amount of these gases, and the levels of these gases were not altered in the presence of *E. coli*. Butanone levels in four different conditions (in the LB broth, in the *E. coli* in the LB broth, human whole blood, and *E. coli*-inoculated whole blood) are presented in Figure 4 as an example.

Table 3 lists eight VOCs that mainly originated from human whole blood (category 5). The emitted levels of these eight VOCs were substantially higher in the whole blood only when compared to the levels in LB broth. Interestingly, co-incubation of *E. coli* in human blood did not alter the concentrations of the gases.

Many elevated whole blood compounds were also detected in the headspace of the blood-collecting vacutainer and were rejected as carried-over contaminants. These include: styrene, 3-methylpentane, n-hexane, cyclohexane, methylcyclopentane, and 2-methyl-2-butene.




Figure 1 Representative patterns of VOCs in Category 1. (A) DMS, (B) Ethanol, and (C) Methyl Butanoate emission from 4 different conditions: broth only (Br), *E. coli* in LB broth (Br + *E. coli*), whole blood only (B), and *E. coli*-inoculated whole blood (B + *E. coli*). 1. significantly different between Broth only and Broth + *E. coli*; 2: significantly different between Blood only and Blood + *E. coli*; 3: significantly different between Broth + *E. coli* and Blood + *E. coli*; 4: significantly different between Broth only and blood only.

Discussion

Specific VOCs released by *E. coli* in either LB broth or human blood were collected and compared to the levels emitted from LB broth or blood alone. Headspace gas samples were collected after incubation for 24 hr at 37°C from 10 mL samples inoculated with (or without) 10^6 *E. coli*. This study demonstrates 1) that cultivated *E. coli* in LB broth produces distinct gas profiles, 2) for the first time the ability to modify *E. coli*-specific gas profiles by the addition of whole human blood (i.e., Methyl Propanoate was highly emitted from *E. coli* in LB broth, but barely detected in human whole blood), and 3) of most significance, that the data suggests that *E. coli*-human whole blood interaction has different gas emission profiles

that can be used as non-invasive markers of *E. coli* infection (i.e., methyl butanoate).

The co-culture of *E. coli* in human whole blood emitted appreciable amounts of dimethyl sulfide when compared to the level from naive *E. coli* in the LB broth (Figure 1A). Dimethyl sulfide is a sulfur-containing odorous gas known to be released from marine phytoplankton and algae, and its release has been reported to protect algae and phytoplankton against environmental changes in salinity, radiation, and temperature [20,21]. Dimethyl sulfide released from damaged phytoplankton has also been proposed to act as a deterrent, or chemical weapon, against consumption by zooplankton [22]. In addition, it has been proposed that some microorganisms are able




Figure 2 Representative patterns of VOCs in Category 2. (A) Isoprene and (B) carbonyl sulfide (OCS) emission from 4 different conditions: only (Br), *E. coli* in LB broth (Br + *E. coli*), whole blood only (B), and *E. coli*-inoculated whole blood (B + *E. coli*). 1: significantly different between Broth only and Broth + *E. coli*; 2: significantly different between Blood only and Blood + *E. coli*; 3: significantly different between Broth + *E. coli* and Blood + *E. coli*; 4: significantly different between Broth only and blood only.

to generate gaseous dimethyl sulfide from dimethylsulphoniopropionate [23-25]. A significant concentration of dimethyl sulfide was observed in the headspace of *E. coli*-inoculated whole blood, whereas the headspace of *E. coli* in LB broth contained much less dimethyl sulfide (Figure 1A). This is the first observation that *E. coli*-

inoculated human whole blood produces dimethyl sulfide. Allardyce and colleagues reported that dimethyl sulfide was detected from *E. coli* culture using BacT/ALERT, which contains a mixture of media with blood [8,26]. It is interesting to speculate the possible roles for dimethyl sulfide. Considering that dimethyl sulfide is a




Figure 3 Representative patterns of VOCs in Category 3. (A) Dimethyl disulfide (DMDS) and (B) 1-propanol emission from 4 different conditions such as broth only (Br), *E. coli* in LB broth (Br + *E. coli*), whole blood only (B), and *E. coli*-inoculated whole blood (B + *E. coli*). 1: significantly different between Broth only and Broth + *E. coli*; 2: significantly different between Blood only and Blood + *E. coli*; 3: significantly different between Broth + *E. coli* and Blood + *E. coli*; 4: significantly different between Broth only and blood only.

Table 2 Gases mainly from pure broth

Gas	Category 4 Gases mainly from LB broth (significantly higher in broth only compared to whole blood only)			
	Broth	+ E. coli	Blood	
		+ E. coli	+ E. coli	
Ethyl nitrate [pptv]	429 (371, 448)	233 (32,284)	28 (17,38)	28 (21,35)
Methyl nitrate [pptv]	705 (670, 825)	446 (38,534)	29 (27,35)	29 (26,36)
2-Propyl nitrate [pptv]	162 (147,177)	103 (30,125)	20 (14,29)	20 (18,33)
2-Methylpropanal [pptv]	90900 (56900,203300)	4300 (700,8900)	2010 (100,600)	600 (200,1300)
3-Methylhexane [pptv]	2000 (1120, 3290)	1020 (660,1660)	510 (400,1850)	530 (220,1410)
2,2,4-Trimethylpentane [pptv]	9140 (8030, 9940)	6100 (1660,7140)	700 (360,1220)	810 (620,1160)
n-Decane [pptv]	100 (70,140)	50 (30,50)	60 (30,70)	60 (30,80)
Benzene [pptv]	93640 (50700,114580)	25910 (300,72270)	1100 (0,2070)	1030 (440,1550)
Toluene [pptv]	98930 (54460,129080)	30870 (1850,69270)	1210 (670,2420)	1680 (720,1880)
Ethylbenzene [pptv]	640 (250,950)	220 (40,530)	150 (80,280)	170 (120,290)
Styrene [pptv]	1130 (670,1950)	800 (290,1180)	280 (140,1110)	360 (200,820)
n-Propylbenzene [pptv]	40 (20,70)	40 (20,110)	20 (5,40)	30 (15,70)
Butanal [pptv]	169900 (60100,238800)	0 (0,8900)	700 (200,17100)	600 (0,34500)
Hexanaldehyde [pptv]	1250 (700,3800)	485 (275,1600)	450 (250,3050)	575 (175,2150)
Butanone [pptv]	478800 (374600,707000)	339600 (219900,555700)	5600 (2300,15900)	9500 (6400,31200)
Unk M ^s	10530 (5841,20041)	12854 (7971,44834)	1095 (263,2664)	1633 (359,3345)

Data were presented with median and range (minimum, maximum).

^sunknown peaks were presented as area under the curve.

cryoprotectant for algae, dimethyl sulfide could be a mediator of survival of blood cells in response to the stresses of bacterial infection or the response of *E. coli* in the face of the bactericidal components of whole blood. Other studies utilizing different media/broth to

culture *E. coli* did not observe dimethyl sulfide in the headspace above the culture [9,13].

Various strains of *E. coli* have been reported to metabolize glucose (and other sugars) to generate ethanol regardless of culture medium or broth used [3,27-34]. In the current study, ethanol has been detected in *E. coli* in LB broth and in the presence of human whole blood (Figure 1B). *E. coli* in LB broth emitted greater than 8 ppmv ethanol. The emission level was even greater (>15 ppmv) when *E. coli* was mixed with human blood. It is likely that the ethanol emission from *E. coli* depends on the available glucose in the LB broth or human whole blood.

Acetaldehyde has previously been detected in exhaled human breath [35,36] and in human lung cancer cell line cultures [37-39]. We previously demonstrated that both malignant myelocytic white blood cells such as HL60 and neutrophils are capable of producing acetaldehyde (16;18). The current study also demonstrates that *E. coli*-inoculated whole blood obtained from healthy subjects emitted appreciable amounts of acetaldehyde into the gas phase greater than the emission from whole blood (Table 1). In addition, acetaldehyde has been detected from other strains of *E. coli* and other microorganisms such as *Pseudomonas* cultured in human blood and medium mixture [8,26]. The LB broth itself also emits some acetaldehyde. However, the levels were not altered in the presence of *E. coli* in LB broth. This observation




Figure 4 Butanone emission from 4 different conditions such as broth only (Br), *E. coli* in LB broth (Br + *E. coli*), whole blood only (B), and *E. coli*-inoculated whole blood (B + *E. coli*). Butanone is an exemplary gas that originates from LB broth.

Table 3 Gases mainly from whole human blood

Gas	Category 5 Gases mainly from whole human blood (significantly higher in whole blood only compared to broth only)			
	Broth		Blood	
		+ E. coli		+ E. coli
n-Pentane [pptv]	50 (0,160)	10 (0,165)	1220 (835,1900)	1625 (815,2440)
Methanol [ppbv]	48 (34,132)	62 (41,110)	300 (210,346)	378 (226,704)
n-Heptane [pptv]	0 (0,315)	150 (30,165)	370 (260,915)	700 (210,4305)
2,3,4-Trimethylpentane [pptv]	35 (0,130)	55 (0,205)	125 (105,275)	145 (130,365)
alpha-Pinene [pptv]	5 (0,15)	5 (0,60)	40 (20,220)	60 (15,370)
Methyl chloride (CH ₃ Cl) [pptv]	1045 (800,1390)	1140 (675,1815)	2190 (1215,2670)	1890 (1235,3025)
Dibromomethane (CH ₂ Br ₂) [pptv]	1 (0,2)	1 (0,2)	47 (11,75)	54 (10,106)
i-Pentane [pptv]	225 (0,870)	210 (0,510)	1400 (545,2395)	1310 (0,4770)

Data were presented with median and range (minimum, maximum).

and others may suggest that acetaldehyde is likely a metabolic byproduct of the interaction between *E. coli* and the various components of whole blood [40,41].

Other VOCs generated only in the *E. coli*-inoculated whole blood were methyl butanoate and a gas that remains unidentified, unknown S. Methyl butanoate has been detected in the exhaled breath of patients with liver disorders [42]. These gases have the potential to be breath biomarkers of *E. coli* infection since these VOCs only appear in the *E. coli*-inoculated whole blood.

There were four VOCs found in the headspace above cultured *E. coli* in LB broth that were significantly lower

in the presence of human whole blood. These VOCs include 1-propanol and the unknown VOCs R1, R2, and Q (see Table 1). Note that both LB broth and human whole blood generate negligible amounts of these VOCs. It remains unclear whether these VOCs are produced by *E. coli* and metabolized by certain components in the blood or the gas production itself was suppressed in the blood. One such possible mechanism for 1-propanol would be the production of 1-propanol by *E. coli* [43] and then, when co-cultured with whole blood, 1-propanol is metabolized by alcohol dehydrogenase in the human whole blood [44]. These compounds may prove interesting in cellular mechanistic studies as they have potentially been catabolised by *E. coli* and/or whole blood.

Isoprene, carbonyl sulfide, and unidentified gas peak R1 were higher in the headspace of *E. coli* in LB broth when compared to the LB broth only. Carbonyl sulfide has been detected in the human breath [14]. Breath carbonyl sulfide concentrations were significantly higher in subjects with cystic fibrosis and it has been suggested that it originates from respiratory bacterial colonization. In effect, subjects with poorer pulmonary function tended to have greater breath carbonyl sulfide concentrations [14]. Isoprene has also been detected in exhaled human breath, and its possible association with blood cholesterol levels [45,46] and end-stage renal failure [47] has been suggested. In addition, various bacterial species, including *E. coli*, have been reported to produce isoprene, though emitted isoprene levels varied depending on the culture medium [48]. In the current study, *E. coli* contributes to isoprene emission with median value of 1600 pptv when cultured in the LB broth. Human whole blood obtained from healthy subjects emitted a median value of 1000 pptv of isoprene. Interestingly, *E. coli* inoculation into the whole blood does not affect emitted isoprene levels. This observation may suggest that isoprene release may be governed by different




Figure 5 i-Pentane emission from 4 different conditions such as broth only (Br), *E. coli* in LB broth (Br + *E. coli*), whole blood only (B), and *E. coli*-inoculated whole blood (B + *E. coli*). i-Pentane is an exemplary gas that originates from human whole blood.

mechanisms when *E. coli* is cultured in the LB broth and human whole blood.

In contrast, there are certain VOCs that originated from LB broth only or human whole blood only that are not increased by the addition of *E. coli* (Table 2, category 4 and Figure 4; Table 3, category 5 and Figure 5). For example, butanone (an exemplary gas, category 4) and *i*-Pentane (an exemplary gas, category 5) emissions were observed either in LB broth only or human whole blood only, respectively. However, the emission levels were not altered in the presence of *E. coli*; the VOCs under these categories were not considered of interest in searching for *E. coli* -associated gas biomarkers.

Conclusions

The current study demonstrates that cultivated *E. coli* produces distinct gas profiles depending on the medium used (i.e., LB broth or human whole blood). In addition, and of most significance, the data suggests that the interaction between *E. coli* and human whole blood presents a different gas emission profile that may have the potential to be used as non-invasive marker of *E. coli* infection. Additional investigations will be required to provide a more specific understanding of how the immune response of whole blood specifically alters the gas signature of invading *E. coli*. These data also provide the beginning of an inventory of VOCs associated with *E. coli* –the necessary first step from which an exploration of other pathogenic microorganisms can follow.

Abbreviations

VOCs: Volatile organic compounds; *E. coli*: Escherichia Coli; DMS: Dimethyl sulfide; CS₂: Carbon disulfide; OCS: Carbonyl sulfide; DMDS: Dimethyl disulfide; DMTS: Dimethyl trisulfide.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HWS and BJU designed and performed experiments and wrote the manuscript. SM participated in chemical analysis of volatile gases obtained from *E. coli* cultures. FPZ contributed experimental design and participated in culturing *E. coli*. SYL carried out statistical analysis. DMC and DRB participated in the design of the experiments and provided a review of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by grants from the Pacific Southwest Regional Center of Excellence for Biodefense & Emerging Infectious Diseases Research (U54AI065359) and from the National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH), and the NIH Roadmap for Medical Research (UL1 RR031985). We thank Alan Barbour MD and the Institute for Clinical Translational Research Center (ICTS) at the University of California, Irvine.

Author details

¹Department of Chemistry, University of California, Irvine, CA 92697, USA.

²Department of Pediatrics, University of California, Irvine, CA 92697, USA.

³Institute for Clinical and Translational Sciences, University of California, Irvine, CA 92697, USA.

Received: 4 April 2013 Accepted: 24 June 2013
Published: 10 July 2013

References

1. Amann A, Spanel P, Smith D: Breath analysis: the approach towards clinical applications. *Mini Rev Med Chem* 2007, **7**(2):115–129.
2. Bajtarevic A, Ager C, Pienz M, Klieber M, Schwarz K, Ligor M, et al: Noninvasive detection of lung cancer by analysis of exhaled breath. *BMC Cancer* 2009, **9**:348.
3. Galassetti PR, Novak B, Nemet D, Rose-Gottron C, Cooper DM, Meinardi S, et al: Breath ethanol and acetone as indicators of serum glucose levels: an initial report. *Diabetes Technol Ther* 2005, **7**(1):115–123.
4. Lee J, Ngo J, Blake D, Meinardi S, Pontello AM, Newcomb R, et al: Improved predictive models for plasma glucose estimation from multi-linear regression analysis of exhaled volatile organic compounds. *J Appl Physiol* 2009, **107**(1):155–160.
5. Ligor M, Ligor T, Bajtarevic A, Ager C, Pienz M, Klieber M, et al: Determination of volatile organic compounds in exhaled breath of patients with lung cancer using solid phase microextraction and gas chromatography mass spectrometry. *Clin Chem Lab Med* 2009, **47**(5):550–560.
6. Novak BJ, Blake DR, Meinardi S, Rowland FS, Pontello A, Cooper DM, et al: Exhaled methyl nitrate as a noninvasive marker of hyperglycemia in type 1 diabetes. *Proc Natl Acad Sci U S A* 2007, **104**(40):15613–15618.
7. Shin HW, Rose-Gottron CM, Cooper DM, Newcomb RL, George SC: Airway diffusing capacity of nitric oxide and steroid therapy in asthma. *J Appl Physiol* 2004, **96**(1):65–75.
8. Allardyce RA, Langford VS, Hill AL, Murdoch DR: Detection of volatile metabolites produced by bacterial growth in blood culture media by selected ion flow tube mass spectrometry (SIFT-MS). *J Microbiol Methods* 2006, **65**(2):361–365.
9. Bunge M, Araghipour N, Mikoviny T, Dunkl J, Schnitzhofer R, Hansel A, et al: On-line monitoring of microbial volatile metabolites by proton transfer reaction-mass spectrometry. *Appl Environ Microbiol* 2008, **74**(7):2179–2186.
10. Lechner M, Fille M, Hausdorfer J, Dierich MP, Rieder J: Diagnosis of bacteria *in vitro* by mass spectrometric fingerprinting: a pilot study. *Curr Microbiol* 2005, **51**(4):267–269.
11. Lechner M, Tilg H, Rieder J: Analysis of volatile compounds emitted by the *Helicobacter pylori* reference strain NCTC 11637 *in vitro*. *Helicobacter* 2006, **11**(1):66.
12. Moularat S, Robine E, Ramalho O, Oturan MA: Detection of fungal development in a closed environment through the identification of specific VOC: demonstration of a specific VOC fingerprint for fungal development. *Sci Total Environ* 2008, **407**(1):139–146.
13. Zhu J, Bean HD, Kuo YM, Hill JE: Fast detection of volatile organic compounds from bacterial cultures by secondary electrospray ionization-mass spectrometry. *J Clin Microbiol* 2010, **48**(12):4426–4431.
14. Kamboures MA, Blake DR, Cooper DM, Newcomb RL, Barker M, Larson JK, et al: Breath sulfides and pulmonary function in cystic fibrosis. *Proc Natl Acad Sci U S A* 2005, **102**(44):15762–15767.
15. Lechner M, Karlsseder A, Niederseer D, Lirk P, Neher A, Rieder J, et al: *H. pylori* infection increases levels of exhaled nitrate. *Helicobacter* 2005, **10**(5):385–390.
16. Shin HW, Umber BJ, Meinardi S, Leu SY, Zaldivar F, Blake DR, et al: Acetaldehyde and hexanaldehyde from cultured white cells. *J Transl Med* 2009, **7**:31.
17. Colman JJ, Swanson AL, Meinardi S, Sive BC, Blake DR, Rowland FS: Description of the analysis of a wide range of volatile organic compounds in whole air samples collected during PEM-tropics A and B. *Anal Chem* 2001, **73**(15):3723–3731.
18. Shin HW, Umber BJ, Meinardi S, Leu SY, Zaldivar F, Blake DR, et al: Gas signatures from cultured neutrophils and peripheral blood mononuclear cells obtained from healthy humans. *J Mol Biomark Diagnosis* 2012, **2**(4):112.
19. Benjamini Y, Hochberg Y: Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J Royal Stat Soc B* 1995, **57**:289–300.
20. Cañavate JP, Lubian LM: Relationship between cooling rates, cryoprotectant concentrations and salinities in the cryopreservation of marine microalgae. *Mar Biol* 1995, **124**(2):325–334.
21. Sunda W, Kieber DJ, Kiene RP, Huntsman S: An antioxidant function for DMSP and DMS in marine algae. *Nature* 2002, **418**(6895):317–320.

22. Dacey JW, Wakeham SG: Oceanic dimethylsulfide: production during zooplankton grazing on phytoplankton. *Science* 1986, **233**(4770):1314–1316.
23. Johnston AW, Todd JD, Sun L, Nikolaidou-Katsaridou MN, Curson AR, Rogers R: Molecular diversity of bacterial production of the climate-changing gas, dimethyl sulphide, a molecule that impinges on local and global symbioses. *J Exp Bot* 2008, **59**(5):1059–1067.
24. Todd JD, Rogers R, Li YG, Wexler M, Bond PL, Sun L, et al: Structural and regulatory genes required to make the gas dimethyl sulfide in bacteria. *Science* 2007, **315**(5812):666–669.
25. Curson AR, Todd JD, Sullivan MJ, Johnston AW: Catabolism of dimethylsulphoniopropionate: microorganisms, enzymes and genes. *Nat Rev Microbiol* 2011, **9**(12):849–859.
26. Allardice RA, Hill AL, Murdoch DR: The rapid evaluation of bacterial growth and antibiotic susceptibility in blood cultures by selected ion flow tube mass spectrometry. *Diagn Microbiol Infect Dis* 2006, **55**(4):255–261.
27. Alterthum F, Ingram LO: Efficient ethanol production from glucose, lactose, and xylose by recombinant *Escherichia coli*. *Appl Environ Microbiol* 1989, **55**(8):1943–1948.
28. Boumba VA, Economou V, Kourkoumelis N, Gousia P, Papadopoulos C, Vougiouklakis T: Microbial ethanol production: experimental study and multivariate evaluation. *Forensic Sci Int* 2011, **215**(1–3):189–198.
29. Donnelly MI, Millard CS, Clark DP, Chen MJ, Rathke JW: A novel fermentation pathway in an *Escherichia coli* mutant producing succinic acid, acetic acid, and ethanol. *Appl Biochem Biotechnol* 1998, **70**–**72**:187–198.
30. Huerta-Beristain G, Utrilla J, Hernandez-Chavez G, Bolivar F, Gosset G, Martinez A: Specific ethanol production rate in ethanologenic *Escherichia coli* strain KO11 is limited by pyruvate decarboxylase. *J Mol Microbiol Biotechnol* 2008, **15**(1):55–64.
31. Martin GJ, Knepper A, Zhou B, Pamment NB: Performance and stability of ethanologenic *Escherichia coli* strain FBR5 during continuous culture on xylose and glucose. *J Ind Microbiol Biotechnol* 2006, **33**(10):834–844.
32. Padukone N, Evans KW, McMillan JD, Wyman CE: Characterization of recombinant *E. coli* ATCC 11303 (pLOI 297) in the conversion of cellulose and xylose to ethanol. *Appl Microbiol Biotechnol* 1995, **43**(5):850–855.
33. Salaspuro V, Nyfors S, Heine R, Siitonen A, Salaspuro M, Jousimies-Somer H: Ethanol oxidation and acetaldehyde production *in vitro* by human intestinal strains of *Escherichia coli* under aerobic, microaerobic, and anaerobic conditions. *Scand J Gastroenterol* 1999, **34**(10):967–973.
34. Samy T, Arnaldos M, Kunkel SA, Pagilla KR, Stark BC: Engineering of ethanolic *E. coli* with the *Vitreoscilla hemoglobin* gene enhances ethanol production from both glucose and xylose. *Appl Microbiol Biotechnol* 2010, **88**(5):1103–1112.
35. Turner C, Spanel P, Smith D: A longitudinal study of ethanol and acetaldehyde in the exhaled breath of healthy volunteers using selected-ion flow-tube mass spectrometry. *Rapid Commun Mass Spectrom* 2006, **20**(1):61–68.
36. Fuchs P, Loeseken C, Schubert JK, Miekisch W: Breath gas aldehydes as biomarkers of lung cancer. *Int J Cancer* 2010, **126**(11):2663–2670.
37. Sponring A, Filipiak W, Mikoviny T, Ager C, Schubert J, Miekisch W, et al: Release of volatile organic compounds from the lung cancer cell line NCI-H2087 *in vitro*. *Anticancer Res* 2009, **29**(1):419–426.
38. Filipiak W, Sponring A, Filipiak A, Ager C, Schubert J, Miekisch W, et al: TD-GC-MS analysis of volatile metabolites of human lung cancer and normal cells *in vitro*. *Cancer Epidemiol Biomark Prev* 2010, **19**(1):182–195.
39. Filipiak W, Sponring A, Mikoviny T, Ager C, Schubert J, Miekisch W, et al: Release of volatile organic compounds (VOCs) from the lung cancer cell line CALU-1 *in vitro*. *Cancer Cell Int* 2008, **8**:17.
40. Hazen SL, Hsu FF, D'Avignon A, Heinecke JW: Human neutrophils employ myeloperoxidase to convert alpha-amino acids to a battery of reactive aldehydes: a pathway for aldehyde generation at sites of inflammation. *Biochemistry* 1998, **37**(19):6864–6873.
41. Hazen SL, D'Avignon A, Anderson MM, Hsu FF, Heinecke JW: Human neutrophils employ the myeloperoxidase-hydrogen peroxide-chloride system to oxidize alpha-amino acids to a family of reactive aldehydes. Mechanistic studies identifying labile intermediates along the reaction pathway. *J Biol Chem* 1998, **273**(9):4997–5005.
42. Manolis A: The diagnostic potential of breath analysis. *Clin Chem* 1983, **29**(1):5–15.
43. Jain R, Yan Y: Dehydratase mediated 1-propanol production in metabolically engineered *Escherichia coli*. *Microb Cell Fact* 2011, **10**:97.

doi:10.1186/2001-1326-2-13

Cite this article as: Umber et al.: Gas signatures from *Escherichia coli* and *Escherichia coli*-inoculated human whole blood. *Clinical and Translational Medicine* 2013 **2**:13.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com