

UC Davis

UC Davis Previously Published Works

Title

Deletions in the pyruvate pathway of Salmonella Typhimurium alter SPI1-mediated gene expression and infectivity

Permalink

<https://escholarship.org/uc/item/3dz306xg>

Journal

Journal of Animal Science and Biotechnology, 4(1)

ISSN

2049-1891

Authors

Abernathy, Jason
Corkill, Carolina
Hinojosa, Carolee
et al.

Publication Date

2013-02-25

DOI

<http://dx.doi.org/10.1186/2049-1891-4-5>

Peer reviewed

RESEARCH

Open Access

Deletions in the pyruvate pathway of *Salmonella* Typhimurium alter SPI1-mediated gene expression and infectivity

Jason Abernathy¹, Carolina Corkill², Carolee Hinojosa², Xianyao Li³ and Huaijun Zhou^{1*}

Abstract

Background: *Salmonella enterica* serovar Typhimurium is a major foodborne pathogen worldwide. *S. Typhimurium* encodes type III secretion systems via *Salmonella* pathogenicity islands (SPI), producing the major effector proteins of virulence. Previously, we identified two genes of *Salmonella* pyruvate metabolism that were up-regulated during chicken cell infection: pyruvate formate lyase I (*pflB*) and bifunctional acetaldehyde-CoA/alcohol dehydrogenase (*adhE*). We were therefore interested in examining the role these genes may play in the transmission of *Salmonella* to humans.

Methods: Mutant strains of *Salmonella* with single gene deletions for *pflB* and *adhE* were created. Invasion and growth in human HCT-8 intestinal epithelial cells and THP-1 macrophages was examined. Quantitative PCR was performed on 19 SPI-1 genes.

Results: In HCT-8 cells, both mutant strains had significantly higher intracellular counts than the wild-type from 4 to 48 h post-infection. Various SPI-1 genes in the mutants were up-regulated over the wild-type as early as 1 h and lasting until 24 h post-infection. In THP-1 cells, no significant difference in internal *Salmonella* counts was observed; however, SPI-1 genes were largely down-regulated in the mutants during the time-course of infection. We also found five SPI-1 genes - *hilA*, *hilC*, *hilD*, *sicP* and *rtsA* - which were up-regulated in at least one of the mutant strains in log-phase broth cultures alone. We have therefore identified a set of SPI-1 virulence genes whose regulation is effected by the central metabolism of *Salmonella*.

Keywords: *adhE*, Metabolism, *pflB*, Pyruvate, *Salmonella* pathogenicity island, *Salmonella* Typhimurium, Virulence

Background

Salmonella serovars cause an array of diseases, ranging from gastroenteritis to systemic infections [1,2]. While there are thousands of characterized serovars, few have been demonstrated as virulent to humans [3,4]. *Salmonella enterica* serovar Typhimurium is one such strain, with a broad spectrum of hosts including pig, chicken and humans [1,5,6]. *S. Typhimurium* can be ubiquitous in the environment and within certain animal reservoirs; it can act as a commensal, produce an asymptomatic infection, and/or trigger disease in various hosts [7]. Serovar Typhimurium thus remains one of the top infectious food-borne pathogens worldwide.

The most common route of entry for *S. Typhimurium* is by oral transmission. Once contact is made with a human host through this route, the bacterium initiates infection by invading intestinal epithelial cells using a type III secretion system (T3SS) [8-10]. This system is encoded by a 40 kilobase region of the genome called *Salmonella* pathogenicity island (SPI) 1 [11-18]. The T3SS of *S. Typhimurium* consists of structural components (i.e. needle complex) that make up the apparatus used to inject effector proteins into the cytoplasm of host cells [12,19-21]. The effector proteins then promote infection by altering the physiology of the host cell. Specifically, *Salmonella* can seize the host cell machinery and induce rearrangements in the actin cytoskeleton and cause alterations in signal transduction that helps facilitate further systemic infection [22-24]. In mammals, the intracellular phase of infection proceeds in a coordinated effort with an additional type III secretion

* Correspondence: hzhou@ucdavis.edu

¹Department of Animal Science, University of California, Davis, CA 95616, USA

Full list of author information is available at the end of the article

system encoded by SPI-2 [25-27]. Overall, the T3SS-1 encoded by SPI-1 promotes invasion of host intestinal epithelial cells [28,29], initiation of inflammation [27,30,31], and intracellular survival and persistence [32-34], while TTSS-2 encoded by SPI-2 is needed to promote intracellular growth [35-39].

While the T3SS and its genetic regulators are a central component in the ability of *Salmonella* to establish a systemic infection, the capability of *Salmonella* to invade and persist in a host has been partly linked to the central metabolism of the bacterium (reviewed in [40]) and environmental cues [41-44]. Alterations in *Salmonella* metabolic pathways, such as glycolysis and the tricarboxylic acid (TCA) cycle, using both *in vitro* and *in vivo* models have demonstrated the importance of factors outside of the T3SS in its pathogenicity [40]. For instance, deletions in the pyruvate dehydrogenase subunit E1 (*aceE*) gene of *S. Enteritidis* has been shown to attenuate the strain in a chicken model [45]. In murine models, the loss of functional phosphofructokinase (Pfk) activity prevents growth of *S. Typhimurium* [46], while deletions in other TCA cycle genes hyper-activate the mutants in macrophages though interestingly attenuates the response in BALB/c mice [47]. As the SPI genes and the T3SS have been extensively studied as key pathogen modulators in *Salmonella*, determining the ability of additional metabolic genes to affect pathogenicity remains to be explored.

In our previous study, we measured global gene expression of the *Salmonella enterica* serotype Typhimurium genome in chicken heterophils (data not shown). Two *Salmonella* genes, pyruvate formate lyase I (*pflB*) and bifunctional acetaldehyde-CoA/alcohol dehydrogenase (*adhE*), were significantly up-regulated after infection. Since these genes are not typical virulence genes but play a role in cellular metabolism (Figure 1), we sought to understand their role during infection. Therefore, the overall aim of this study is to further characterize the importance these *Salmonella* genes play in its pathogenesis. Deletion mutants of these two genes were created, and gentamicin protection assays and quantitative PCR of SPI-1 genes were performed on human epithelial and macrophage cell lines. The resulting experiments show that these two genes of the pyruvate metabolic pathway affect both infectivity of epithelial cells and SPI-1 gene expression.

Methods

Creation of *Salmonella* mutants

The *Salmonella enterica* serovar Typhimurium ATCC 14028 strain was used as the wild-type in this study. This strain was also used in the construction of mutant gene knock-out isolates. Two genes, pyruvate formate lyase I (*pflB*) and bifunctional acetaldehyde-CoA/alcohol




Figure 1 A simplified model of the pyruvate metabolic pathway in *Salmonella Typhimurium*. Genes producing metabolic enzymes (in italics) were deleted in the creation of mutant strains, including pyruvate formate lyase I (*pflB*) and bifunctional acetaldehyde-CoA/alcohol dehydrogenase (*adhE*).

dehydrogenase (*adhE*), produce enzymes of the pyruvate metabolic pathway as illustrated (Figure 1). To assess the effects of these genes during invasion, two mutant strains, $\Delta pflB::kan$ (*S. Typhimurium* gene number STM0973) and $\Delta adhE::kan$ (*S. Typhimurium* gene number STM1749), were created by the lambda red recombination method with adaptations as described by Santiviago et.al. [48]. Gene-specific primers used for the creation of each isolate are listed in Table 1. The recombination event was confirmed by polymerase chain reaction (PCR) using primers to the flanking genomic regions.

Cell culture

Human cell lines HCT-8 and THP-1 were used in this study. HCT-8 is an ileocecal colorectal adenocarcinoma epithelia cell line, while THP-1 is a monocytic leukemia cell line that can be stimulated to differentiate into macrophages. HCT-8 was grown using RPMI-1640 with 2.05 mmol/L L-Glutamine media (HyClone, Thermo Scientific, Rockford, IL) supplemented with 10% fetal bovine serum (Atlanta Biologicals, Lawrenceville, GA), 1% of 100 mmol/L sodium pyruvate (Thermo Scientific), and 1% of 10 000 U penicillin/10 mg streptomycin per mL (Thermo Scientific). THP-1 was grown in suspension using RPMI-1640 with 2.05 mmol/L L-Glutamine

Table 1 A list of the *Salmonella* primers used in this study

| Gene | Forward sequence (5'-3') | Reverse sequence (5'-3') | Locus or reference | Technique |
|-------------|--|---|--------------------|------------|
| <i>pflB</i> | GAAGGTAGGTGTTACATGCCGAGCTT AATGAAAAGTTAGCCACAGTGCAGGCT GGAGCTGCTC | TTTCAGTCAAACCCATTACAT GGTCTGCGTGAAGGTACGAGT AATCATATGAATATCCTCCTTAG | STM0973 | Lambda red |
| <i>adhE</i> | TATCAGGAGAGCATTATGGCTGTTACT AATGTCGCTGAACCTAACGTGCAGGCT GGAGCTGCTC | AAAACAGATAACGAATTAAGC GGATTTTTTCGCTTTTTTCTC TGCCATATGAATATCCTCCTTAG | STM1749 | |
| 16S rRNA | AGGCCTTCGGGTTGTAAAGT | GTTAGCCGGTGCTTCTCTG | [49] | qPCR |
| <i>hilA</i> | ATAGCAAACCTCCCGACGATG | ATTAAGGCGACAGAGCTGG | [50] | |
| <i>hilC</i> | CTCACCTCTTCAGCGGCCAGT | CACCCGCAAATGGTCACAGGCT | STM2867 | |
| <i>hilD</i> | GGTAGTTAACGTGACGCTTG | GATCTTCTGCGCTTCTCTG | [50] | |
| <i>invC</i> | TCGGTCGATCGCTGCAA | CCCTGGTCGCGAAAATATTC | STM2894 | |
| <i>invE</i> | CGAATGACGCCAGCTGTTC | TGCGTCAGGCGTCGTAAG | STM2897 | |
| <i>invF</i> | TGAAAGCCGACACAATGAAAT | GCCTGCTCGCAAAAAGC | STM2899 | |
| <i>invH</i> | GGTGCCCTCCCTTCT | TGCGTTGGCCAGTTGCT | STM2900 | |
| <i>invJ</i> | CGCCGCGGTTAATTG | GCTTCCGCTGCAACCAA | STM2892 | |
| <i>orgA</i> | AGGCAGGGAGCCTTGCTT | CCCTGATGCATTGCCAAAA | STM2870 | |
| <i>orgB</i> | ACCATCCCGAAACGCTTTTA | TTGCCCTCAGGCTTATCG | STM2869 | |
| <i>prgH</i> | TGAACGGCTGTGAGTTTCCA | GCGCATCACTCTGACCTACCA | STM2874 | |
| <i>prgJ</i> | GGTCTATGGAACGGACATTGTC | CGCCGAACCAGAAAAAGC | STM2872 | |
| <i>prgK</i> | GGGTGGAATAGCGCAGATG | TCAGCTCGCGGAGACGATA | STM2871 | |
| <i>rtsA</i> | ACCCGTGGTGAGCTTGATGAGT | CCTGTCCAGGTGGGGAGCAT | SEE_03946 | |
| <i>sicA</i> | GATGAGTCTCTCGGGGCAA | GCTCTGTCTCCGCGTTTT | STM2886 | |
| <i>sicP</i> | AGATGATATCTGTTATTGAACGGTATG | CTGCCGCCAGATAGAATCG | STM2879 | |
| <i>sipA</i> | CGTCTTCGCTCAGGAGAAT | TGCCGGGCTCTTTCGTT | STM2882 | |
| <i>sipB</i> | GGCGCGCTGCTAACCAT | TCGCCCCACCGGTAAGAA | STM2885 | |
| <i>sipC</i> | ATCAGGCTGGTCGATTACG | GTACGCCGCTACTCAGGAAC | [50] | |

media (HyClone) supplemented with 10% fetal bovine serum (Atlanta Biologicals), 1% of 10,000 U penicillin/10 mg streptomycin per mL (Thermo Scientific), and 1.75 μ L of 14.3 mM β -Mercaptoethanol (Sigma-Aldrich, St. Louis, MO). Both cell lines were cultured at 37°C in an air-jacketed incubator (NuAire, Plymouth, MN) with 5% CO₂ and constant humidity.

Invasion assays

The infectivity of *Salmonella* on both HCT-8 and THP-1 was defined as the number of internal *Salmonella* cells per viable human cell at each unit of time. Internal *Salmonella* counts on both cell lines were determined using a gentamicin protection assay (GPA). Viable human cells were assayed by trypan blue exclusion. Invasion assays on each cell line were performed in at least three independent experiments, using triplicate wells at each time-point. Probabilities for statistical significance were calculated using Student's *t*-test.

For HCT-8 cells, 2 \times 10⁵ cells were seeded in 12-well plates with 1 mL complete media (without antibiotics) and grown until confluent. Overnight ATCC 14028, Δ *pflB*::*kan*, and Δ *adhE*::*kan* strains of *Salmonella* grown at 37°C

in Luria-Bertani (LB) broth were sub-cultured for 3 h in pre-warmed (37°C) LB broth before challenge to ensure log-phase cultures. HCT-8 cells were challenged with each *Salmonella* strain at a multiplicity of infection (MOI) of 100 by replacing media with *Salmonella* infectious media. Immediately upon challenge, plates were centrifuged at 500 \times g for 5 min and placed in a CO₂ incubator and allowed to incubate for 30 min. After this time, cells were washed with phosphate buffered saline (PBS, pH 7.4) and infectious media was replaced with 1 mL complete media supplemented with 100 μ g/mL gentamicin sulfate (Mediatech, Manassas, VA). After 30 min, media was replaced with 2 mL complete media supplemented with 10 μ g/mL gentamicin sulfate (Mediatech) and kept through the duration of the study. The first time-point of 1 h post-infection was collected by washing the infected HCT-8 cells with PBS and lysing with 1 mL PBS + 1% Triton X-100 (Sigma-Aldrich). Internalized *Salmonella* were then counted by serial dilutions on pre-warmed LB agar in triplicate. At the same time, infected HCT-8 cells were counted on a TC10 automated cell counter (Bio-Rad, Hercules, CA) using a 1:1 ratio of cells to 0.4% trypan blue dye (Bio-Rad). Additional time-

points of 4, 18, 24, and 48 h post-infection were also collected.

For THP-1 cells, 8×10^5 THP-1 cells were seeded in 12-well plates with 1 mL complete media (without β -Mercaptoethanol or antibiotics) and 200 nmol/L phorbol 12-myristate 13-acetate (PMA; Millipore, Billerica, MA) for 24 h to allow differentiation of suspension monocytes to adherent macrophages. After 24 h, media was exchanged with pre-warmed media (without β -Mercaptoethanol or antibiotics) omitting PMA and cells were allowed to develop for an additional 24 h. Overnight cultures from ATCC 14028, $\Delta pflB::kan$, and $\Delta adhE::kan$ strains of *Salmonella* were sub-cultured for 2.5 h in pre-warmed LB broth and then incubated with 10% human AB serum (Mediatech) for 30 min prior to challenge. THP-1 cells were challenged 48 h after seeding at a MOI10 from each *Salmonella* strain, and invasion was assessed similarly as with HCT-8 challenge. To assess the role that opsonization may play on infectivity using these *Salmonella* strains, separate GPA experiments were also performed where strains were not incubated with human AB serum prior to challenge. Time-points of 1, 2.5, 4, and 6 h post-infection were collected from non-opsonized challenge experiments. Lower time-points and MOI were selected for non-opsonized challenge as the THP-1 cells were largely damaged and apoptotic at MOI 100 and with longer incubations (data not shown), and has previously been observed [51].

Quantitative PCR

Real-time reverse-transcription quantitative PCR (RT-qPCR) was performed to assess mRNA expression levels of 19 selected SPI-1 genes. The primers used for RT-qPCR are listed in Table 1. SPI-1 transcripts were measured from both *Salmonella* grown in cultures and after invasion of host cells.

To generate complementary DNA (cDNA) for RT-qPCR from *Salmonella* cultures, 5 mL of ATCC14028, $\Delta pflB::kan$, and $\Delta adhE::kan$ strains were grown overnight at 37°C with agitation, then sub-cultured 1:25 in pre-warmed LB broth for 4 h. The three strains were equalized by OD₆₀₀ spectrophotometry and RNA was isolated by the RNeasy Protect Bacteria Kit (Qiagen, Valencia, CA) according to the manufacturer's protocol. Potential contaminating genomic DNA was removed by the TURBO DNA-free™ kit (Ambion, Carlsbad, CA), as directed by the manufacturer. Reverse transcription reactions were then performed by the ThermoScript™ RT-PCR System for First-Strand cDNA Synthesis (Invitrogen, Carlsbad, CA) from 4 μ g total RNA from each sample, using the supplied random hexamers as RT-primers, in 20 μ L total volume reactions. Reactions for each sample were diluted 1:10 with sterile nanopure water and used as templates for RT-qPCR.

To generate cDNA for RT-qPCR from *Salmonella* after invasion of host cells, HCT-8 and THP-1 cells were again infected with ATCC 14028, $\Delta pflB::kan$, or $\Delta adhE::kan$ strains by the GPA in additional independent experiments, with the following modifications. At each time-point post infection, cells were washed with PBS and lysed directly by the addition of 400 μ L TRI-reagent (Ambion) to each well and gentle pipetting. Total RNA was then extracted using the RiboPure™-Bacteria Kit (Ambion) according to the manufacturer's protocol. The TURBO DNA-free™ kit (Ambion) was used to remove potential genomic DNA contamination. Total RNA from each sample (representing one well of a 12-well plate) was quantified using a spectrophotometer (NanoDrop 1000, Wilmington, DE) and equimolar amounts from each of three replicate wells were pooled. RT-reactions were performed using the ThermoScript™ RT-PCR System for First-Strand cDNA Synthesis (Invitrogen) from 800 ng total RNA from each sample, using the supplied random hexamers as RT-primers, in 20 μ L total volume reactions. Reactions for each sample were diluted 1:5 with sterile nanopure water and used as templates for RT-qPCR.

All RT-qPCR reactions were performed using the Power SYBR Green PCR Master Mix (Applied Biosystems, Foster City, CA), with modifications. A total reaction volume of 10 μ L was used, to include: 5 μ L of 2X SYBR Green, 2 μ L of [1 μ mol/L] forward primer, 2 μ L of [1 μ mol/L] reverse primer, and 1 μ L template cDNA. Quantitative PCR was performed on an ABI 7900HT system (Applied Biosystems), with cycling and dissociation curve parameters used according to the recommendations from the manufacturer. Each RT-qPCR reaction was performed using at least three technical replicates for each sample. After real-time PCR cycling, each product was assessed for quality and specificity by the corresponding dissociation curve. Raw data was edited using the SDS v2.4 Software (Applied Biosystems), and cycle threshold (Ct) values were recorded. Ct values were uploaded into the Relative Expression Software Tool (REST2009; Qiagen) [52] for comparisons. Data was considered significant at p -value < 0.05 after permutation testing using randomizations. The standard error of the mean was estimated from confidence intervals.

Results

Salmonella invasion of HCT-8 epithelial cells

After HCT-8 cells were challenged with either *Salmonella* wild-type or mutant strains, the number of intracellular bacteria as assessed by counting colony forming units (CFUs) on LB agar were averaged from repeated independent GPA experiments. At the earliest time post-infection of 1 h, there was no significant difference in average CFUs between the wild-type and mutant strains




Figure 2 Invasion of HCT-8 cells by *Salmonella* Typhimurium strains. (A) Time-points of 1 h and 4 h post-infection; (B) Time-points of 18 h, 24 h, and 48 h post-infection. The X-axis represents time (h) while the Y-axis represents average colony forming units (CFU) of bacteria per viable HCT-8 cell. The different *Salmonella* strains are color-coded according to the legend. Asterisk indicates significance of the mutant strain from the wild-type at $p < 0.01$ assessed at each time.

(Figure 2A). However, by 4 h, there was a significant increase in average CFUs in the mutant strains, $\Delta pflB::kan$ and $\Delta adhE::kan$, compared to the wild-type strain, ATCC 14028 (Figure 2A). At 4 h, there were approximately 4–5 CFU of either mutant per 10 HCT-8 cells, while ATCC 14028 remained near 1 CFU/10 HCT-8 cells, thus representing an approximate 4–5 fold increase in *Salmonella* mutants over the wild-type. Increased numbers of mutant strains over the wild-type observed at 4 h post-infection also lasted through the duration of the study at 48 h post-infection, measured at increments of 18, 24, and 48 h (Figure 2B). At 18 h post-infection, there were 4.2 and 3.9 CFU per HCT-8 cell from $\Delta pflB::kan$ and $\Delta adhE::kan$ strains, respectively, while the wild-type ATCC 14028 strain remained at near 1 CFU/cell, thus representing an approximate 4-fold significant difference. Similarly at 24 h post-infection, there were 3.9 and 4.3 CFU per viable epithelial cell from $\Delta pflB::kan$ and $\Delta adhE::kan$ strains, respectively, while the wild-type again remained around 1 CFU/HCT-8 cell at this time-point. The wild-type strain ATCC 14028 was mostly consistent in the level of infectivity and growth measured by GPA with an average of 6.4 CFU per 10 HCT-8 cells over the 48 h course of study, ranging from 3 CFU/100 HCT-8 cells at 1 h to 1.3 CFU/HCT-8 cell at 48 h post-infection. However, in the mutant strains there was an approximate 8-fold significant increase in the number of CFUs between the 1 h and 4 h time-points and, likewise, an approximate 8-fold increase between the 4 h and 18 h time-points.

Average CFU/HCT-8 cell was also increased in the mutants, with an average of 3.9 and 2.5 CFU/HCT-8 cell in $\Delta pflB::kan$ and $\Delta adhE::kan$, respectively. CFUs of both mutant strains remained consistent between 18 h and 24 h post-infection; however, at 48 h post-infection a spike in $\Delta pflB::kan$ was observed (Figure 2B). At 4 h through 48 h post-infection, internal growth of the mutants was found to be highly significant (p -value < 0.01). Overall there was either a steady or an upward trend in the numbers of internalized *Salmonella* through the 48 h course of infection in epithelial cells with the magnitude of replication being significantly greater by the mutant strains beginning a 4 h post-infection (Figure 2).

Salmonella invasion of THP-1 macrophage cells

For challenge of THP-1 cells with *Salmonella* strains opsonized by treatment with human AB serum, time increments were chosen the same as with HCT-8 challenge. Again, average numbers of internalized *Salmonella* strains were assessed by counting CFUs. The numbers of internalized *Salmonella* ranged from approximately 1.5 to 21 CFU per viable THP-1 cell (Figure 3). ATCC 14028 wild-type strain had an average of 7.4 CFU/THP-1 cell, $\Delta pflB::kan$ strain had an average of 11.4 CFU/THP-1 cell, and $\Delta adhE::kan$ strain had an average of 9.9 CFU/THP-1 cell over the course of the 48 h study. There was a general trend that the mutant strains, in particular $\Delta pflB::kan$, had higher CFUs per differentiated THP-1 cell than ATCC 14028. However, no significant difference in average




Figure 3 Invasion of THP-1 cells by opsonized *Salmonella* Typhimurium strains. Time-points of 1, 4, 18, 24, and 48 h post-infection are shown. The X-axis represents time (h) while the Y-axis represents average colony forming units (CFU) of bacteria per viable THP-1 cell. The different *Salmonella* strains are color-coded according to the legend.

CFU counts was observed between either of the mutant or wild-type strains after repeated GPA experiments. This was the case at all of the five time-points tested, from 1 h to 48 h post-infection (Figure 3). Overall, there was a bell-curved trend in the numbers of internalized *Salmonella* throughout this study, where bacteria peaked at 18 h post-infection, and then began to decline as the viable THP-1 cells declined (Figure 3).

For challenge of THP-1 cells with non-opsonized *Salmonella* strains, time-points of 1, 2.5, 4, and 6 h post-infection were selected to assess infectivity. The

numbers of internalized *Salmonella* cells ranged from approximately 1.4 to 10.4 CFU per viable THP-1 cell (Figure 4). ATCC 14028 wild-type strain had an average of 3.3 CFU/THP-1 cell, $\Delta pflB::kan$ strain had an average of 5.4 CFU/THP-1 cell, and $\Delta adhE::kan$ strain had an average of 4.1 CFU/THP-1 cell over the course of the 6 h study. There was a general trend that the mutant strains, in particular $\Delta pflB::kan$, had higher CFUs per differentiated THP-1 cell than ATCC 14028. Again however, no significant difference in average CFU counts was observed between either of the mutant or




Figure 4 Invasion of THP-1 cells by non-opsonized *Salmonella* Typhimurium strains. Time-points of 1, 2.5, 4, and 6 h post-infection are shown. The X-axis represents time (h) while the Y-axis represents average colony forming units (CFU) of bacteria per viable THP-1 cell. The different *Salmonella* strains are color-coded according to the legend.

wild-type strains after repeated GPA experiments. This was the observation at any of the four time-points tested, from 1 h to 6 h post-infection (Figure 4). Overall, there was an upward trend in the numbers of internalized *Salmonella* throughout this study as viable THP-1 cells had not yet become noticeably apoptotic (Figure 3).

After the THP-1 GPA trials, a direct comparison was made between 1 h and 4 h post-infection to assess the role opsonization may play between the *Salmonella* parent and mutant strains. For the opsonized challenge, ATCC 14028 had an average internalization of 7.3 and 7.2 CFU/THP-1 cell at 1 h and 4 h, respectively. $\Delta pflB::kan$ had an average internalization of 12.4 and 12.8 CFU/THP-1 cell at 1 h and 4 h, respectively. $\Delta adhE::kan$ had an average internalization of 7.7 and 10.6 CFU/THP-1 cell at 1 h and 4 h, respectively (Figure 3). For the non-opsonized challenge, ATCC 14028 had an average internalization of 1.4 and 3.6 CFU/THP-1 cell at 1 h and 4 h, respectively. $\Delta pflB::kan$ had an average internalization of 1.9 and 5.4 CFU/THP-1 cell at 1 h and 4 h, respectively. $\Delta adhE::kan$ had an average internalization of 1.4 and 4.0 CFU/THP-1 cell at 1 h and 4 h, respectively (Figure 4). Thus, larger average CFU counts were observed by challenge of THP-1 cells with opsonized bacteria.

SPI-1 gene expression in culture

To determine if the significant increase in numbers of *Salmonella* mutants to HCT-8 cells was potentially due

to the type III secretion system, gene expression was measured in 19 SPI-1 genes (Table 1) by qPCR and compared to the expression in the wild-type strain. Differential regulation was measured from SPI-1 transcripts in log-phase cultures. A total of five genes were found significantly up-regulated in at least one of the mutant strains (Figure 5). In the $\Delta pflB::kan$ strain, *hilA*, *hilD*, *rtsA*, and *sicP* genes were up-regulated compared to ATCC 14028. Relative expression values ranged from 1.36 to 1.96 over the wild-type in genes found significant by qPCR ($p < 0.05$). In the $\Delta adhE::kan$ strain, *hilA*, *hilC*, *hilD*, and *sicP* genes were up-regulated compared to ATCC 14028. Relative expression values ranged from 1.64 to 2.58 over the wild-type in genes found significant by qPCR (Figure 5).

SPI-1 gene expression after invasion of HCT-8 epithelial cell line

Since a set of SPI-1 genes were up-regulated in cultures alone, we were interested to determine if a larger response involving a greater number of genes or a greater magnitude of the fold-change was observed through interaction with epithelial cells. The response of SPI-1 after infection with HCT-8 cells was determined by using qPCR on the same panel of 19 genes. SPI-1 genes were found to be up-regulated over the wild-type as early as 1 h post-infection (Table 2).




Figure 5 Quantitative RT-PCR of *Salmonella* Typhimurium strains from cultures. Gene expression of select SPI-1 invasion genes measured by RT-qPCR from *Salmonella* mutants $\Delta pflB::kan$ and $\Delta adhE::kan$ and comparing expression to the wild-type strain. The 16S rRNA gene was used for normalization. Genes are listed along the X-axis while relative expression level is listed along the Y-axis. Asterisk indicates significance of the mutant strain from the wild-type at $p < 0.05$ assessed for each gene.

Table 2 Quantitative RT-PCR of *Salmonella* during invasion of HCT-8 cells

| Time post-infection | Strain | <i>Salmonella</i> gene | Expression* | Regulation** | | | |
|---------------------|--------------------|------------------------|--------------------|--------------|-------------|------|----|
| 1 h | $\Delta pflB::kan$ | <i>hilC</i> | 1.28 | UP | | | |
| | | <i>hilD</i> | 1.60 | | | | |
| | | <i>prgJ</i> | 1.48 | | | | |
| | | <i>prgK</i> | 1.92 | | | | |
| | | <i>sicA</i> | 1.31 | | | | |
| | $\Delta adhE::kan$ | <i>hilC</i> | 1.26 | | | | |
| | | <i>invE</i> | 3.16 | | | | |
| | | <i>invH</i> | 1.56 | | | | |
| | | <i>orgB</i> | 1.25 | | | | |
| | | <i>prgJ</i> | 1.60 | | | | |
| | | <i>prgK</i> | 1.96 | | | | |
| | | <i>sicA</i> | 1.15 | | | | |
| | | 4 h | $\Delta pflB::kan$ | | <i>rtsA</i> | 2.52 | UP |
| | | | | | <i>sicA</i> | 1.64 | |
| | | | | | <i>sipA</i> | 1.88 | |
| $\Delta adhE::kan$ | <i>hilA</i> | | 3.12 | | | | |
| | <i>orgA</i> | | 3.70 | | | | |
| 18 h | $\Delta pflB::kan$ | <i>hilD</i> | 2.15 | UP | | | |
| | | <i>prgH</i> | 1.85 | | | | |
| | | <i>prgJ</i> | 3.08 | | | | |
| | | <i>sicA</i> | 2.78 | | | | |
| | | <i>sipA</i> | 1.88 | | | | |
| | | <i>sipC</i> | 3.21 | | | | |
| | $\Delta adhE::kan$ | <i>invC</i> | 1.61 | | | | |
| | 24 h | $\Delta pflB::kan$ | <i>prgH</i> | | -1.92 | DOWN | |
| | | | <i>prgJ</i> | | -2.43 | | |
| | | | <i>prgK</i> | | -1.75 | | |
| <i>sicP</i> | | | -3.41 | | | | |
| <i>sipA</i> | | | -3.36 | | | | |
| <i>sipB</i> | | -3.39 | | | | | |
| $\Delta adhE::kan$ | | <i>invC</i> | -2.66 | | | | |
| <i>sipA</i> | -2.07 | | | | | | |

*Values listed are relative fold-change values as compared to the wild-type strain ATCC 14028; **All genes listed are significant at p -value < 0.05.

In $\Delta pflB::kan$, five SPI-1 genes were up-regulated at 1 h, ranging from 1.28 to 1.92 fold change over the wild-type. At 4 h, three SPI-1 genes were up-regulated with relative expression values ranging from 1.64 to 2.52 fold over the wild-type. At 18 h post-infection, six SPI-1

genes were found to be up-regulated, with fold changes of 1.85 to 3.21 over the wild-type. In $\Delta adhE::kan$, seven SPI-1 genes were up-regulated at 1 h post-infection, with relative expression values ranging from 1.15 to 3.16 fold over the wild-type. At 4 h, seven SPI-1 genes were up-regulated with relative expression values ranging from 1.62 to 3.70 fold over the wild-type. At 18 h post-infection, *invC* was up-regulated 1.61 fold in $\Delta adhE::kan$ over ATCC 14028. By 24 h post-infection, SPI-1 genes were either down-regulated in both mutant strains, or not significantly different from the wild-type strain.

SPI-1 gene expression after invasion of THP-1 macrophage cell line

Salmonella invasion gene expression from bacteria invaded into differentiated macrophages, THP-1 cell line, was also examined for non-opsonized wild-type and mutant strains. Similarly to non-opsonized *Salmonella* GPA, time-points of 1, 2.5, 4, and 6 h post-infection were chosen for testing of SPI-1 gene expression by qPCR.

In strain $\Delta pflB::kan$, several SPI-1 genes were differentially expressed between the mutant and wild-type (Table 3). At 1 h, there was no significant differential gene expression. By 2.5 h, 6 SPI-1 genes were down-regulated with expression values ranging from -1.34 to -1.70, while *invE* was up-regulated 1.7-fold over the wild-type at the same time. At 4 h, *hilA*, *hilD*, *invF*, and *prgJ* were slightly up-regulated, with expression values of 1.18 to 1.72 fold over the wild-type. The largest degree of differential expression was observed at 6 h post-infection. A total of 17 out of the 19 SPI-1 genes tested (89%) were down-regulated, ranging from -1.3 to -2 fold change over the wild-type.

In strain $\Delta adhE::kan$, SPI-1 gene expression was down-regulated at every time-point examined (Table 4). At 1 and 2.5 h post-infection, all genes tested were significantly down-regulated as compared to ATCC 14028. At 4 h post-infection, 14 of the 19 SPI-1 genes tested (74%) were down-regulated, with *prgJ* having the lowest magnitude of the fold-change at -1.67 and *sipB* having the highest magnitude of the fold-change at -4.44 under the wild-type. At 6 h post-infection, 16 of the 19 SPI-1 genes tested (84%) were down-regulated, with *orgB* having the lowest magnitude of the fold-change at -1.31 and *sipB* having the highest magnitude of the fold-change at -3.58 under the wild-type.

Discussion

SPI-1 of *Salmonella enterica* serovars is an important virulence factor that has been well-studied in several animal models and in disease outbreaks [53,54]. SPI-1-mediated infectivity of *Salmonella* Typhimurium was examined by creation of null mutations in genes that were up-regulated in our previous host-pathogen microarray study (data not shown). Those *Salmonella*

Table 3 Quantitative RT-PCR of non-opsonized *Salmonella* strain $\Delta pflB::kan$ during invasion of THP-1 cells

| Time post-infection | <i>Salmonella</i> gene | Expression* | Regulation** |
|---------------------|------------------------|-------------|--------------|
| 1 h | No change | | |
| 2.5 h | <i>invE</i> | 1.71 | UP |
| | <i>invC</i> | -1.46 | DOWN |
| | <i>invH</i> | -1.41 | |
| | <i>rtsA</i> | -1.70 | |
| | <i>sicA</i> | -1.52 | |
| | <i>sicP</i> | -1.34 | |
| | <i>sipC</i> | -1.36 | |
| 4 h | <i>hilA</i> | 1.72 | UP |
| | <i>hilD</i> | 1.36 | |
| | <i>invF</i> | 1.19 | |
| | <i>prgJ</i> | 1.18 | |
| 6 h | <i>hilA</i> | -1.61 | DOWN |
| | <i>hilC</i> | -1.97 | |
| | <i>hilD</i> | -1.55 | |
| | <i>invC</i> | -1.54 | |
| | <i>invE</i> | -1.51 | |
| | <i>invF</i> | -1.52 | |
| | <i>invH</i> | -1.61 | |
| | <i>orgA</i> | -1.66 | |
| | <i>orgB</i> | -1.30 | |
| | <i>prgH</i> | -1.43 | |
| | <i>prgJ</i> | -1.46 | |
| | <i>rtsA</i> | -1.83 | |
| | <i>sicA</i> | -1.89 | |
| | <i>sicP</i> | -1.63 | |
| | <i>sipA</i> | -1.58 | |
| | <i>sipB</i> | -2.00 | |
| <i>sipC</i> | -1.70 | | |

*Values listed are relative fold-change values as compared to the wild-type strain ATCC 14028; **All genes listed are significant at p -value < 0.05.

genes, pyruvate formate lyase I (*pflB*) and bifunctional acetaldehyde-CoA/alcohol dehydrogenase (*adhE*), play a major role in the pyruvate metabolism of the organism (Figure 1). Here we show that these two genes can also affect virulence and SPI-1 gene expression.

For the HCT-8 intestinal epithelial cell line, internalized mutant *Salmonella* CFUs were significantly increased over the wild-type at 4 h post-infection and lasted through 48 h post-infection at which time the experiment was terminated (Figure 2). Since the T3SS encoded by SPI-1 is canonically involved in early contact with non-phagocytic cells such as fibroblasts and intestinal epithelia [27,30,31], a selected group of SPI-1 genes (Table 1) in mutant *Salmonella* were measured by qPCR and compared to the wild-type expression. Several SPI-1 genes were up-

Table 4 Quantitative RT-PCR of non-opsonized *Salmonella* strain $\Delta adhE::kan$ during invasion of THP-1 cells

| Time post-infection | <i>Salmonella</i> gene | Expression* | Regulation** | |
|---------------------|--|-------------|--------------|------|
| 1 h | All genes tested were significantly down-regulated | | | |
| 2.5 h | All genes tested were significantly down-regulated | | | |
| 4 h | <i>hilC</i> | -2.53 | DOWN | |
| | <i>invC</i> | -2.63 | | |
| | <i>invE</i> | -2.46 | | |
| | <i>invF</i> | -2.14 | | |
| | <i>invH</i> | -1.98 | | |
| | <i>orgB</i> | -1.71 | | |
| | <i>prgH</i> | -1.79 | | |
| | <i>prgJ</i> | -1.67 | | |
| | <i>prgK</i> | -1.91 | | |
| | <i>rtsA</i> | -2.58 | | |
| | <i>sicA</i> | -3.91 | | |
| | <i>sicP</i> | -1.73 | | |
| | <i>sipB</i> | -4.44 | | |
| | <i>sipC</i> | -2.91 | | |
| | 6 h | <i>hilA</i> | -2.08 | DOWN |
| | | <i>hilC</i> | -2.76 | |
| <i>hilD</i> | | -1.56 | | |
| <i>invC</i> | | -2.76 | | |
| <i>invE</i> | | -2.07 | | |
| <i>invF</i> | | -2.44 | | |
| <i>invH</i> | | -2.30 | | |
| <i>orgA</i> | | -2.36 | | |
| <i>orgB</i> | | -1.31 | | |
| <i>prgH</i> | | -1.71 | | |
| <i>rtsA</i> | | -2.33 | | |
| <i>sicA</i> | | -3.31 | | |
| <i>sicP</i> | | -1.72 | | |
| <i>sipA</i> | -2.22 | | | |
| <i>sipB</i> | -3.58 | | | |
| <i>sipC</i> | -2.28 | | | |

*Values listed are relative fold-change values as compared to the wild-type strain ATCC 14028; **All genes listed are significant at p -value < 0.05.

regulated after HCT-8 cell challenge, from 1 h post-infection lasting to 24 h post-infection when SPI-1 expression in the mutants decreased compared to the wild-type (Table 2). These up-regulated genes included *hilC*, *hilD*, *rtsA*, *sipA*, - all major genes involved in the early response to infection [55]. Some SPI-1 genes were up-regulated at multiple times post-infection and in both mutants, such as *sicA*, a gene encoding a complex chaperone protein required for *Salmonella* entry into host cells [56,57]. Overall, the results indicated that increased internalized growth

of strains of *S. Typhimurium* deficient for transcripts of the pyruvate pathway in HCT-8 cells may in part be due to a differential regulation of genes in the T3SS.

Additional evidence in supporting this notion included qPCR results on SPI-1 genes in log-phase cultures alone. We found that the major regulators of SPI-1 were up-regulated when compared to the wild-type. These included *hilA* and *hilD* in both of the mutant strains, and *hilC* and *rtsA* in at least one of the mutant strains (Figure 5). *HilA* is the major regulator of SPI-1 invasion [41]. The *hilA* gene encodes an OmpR/ToxR family transcriptional regulator that activates the expression of invasion genes in response to both environmental and genetic factors [41]. Further, the proteins encoded by *rtsA*, *hilC* and *hilD* bind to a DNA region upstream of *hilA* and induce *hilA* expression [58-60]. Therefore, these genes could be expected to be induced at an early stage, as they are key modulators of invasive *Salmonella* infection and *hilD* appears to bridge communication between SPI-1 and SPI-2 regulation (reviewed in [55]). These results further indicate the increased virulence of the mutant strains to HCT-8 is occurring via a differential regulation of genes involved in the T3SS. Interestingly, the *sicP* gene was also up-regulated in both mutant strains measured from cell culture alone. *SicP* serves as a chaperone for the SPI-1 effector *Salmonella* protein tyrosine phosphatase (SptP). SptP has been shown to have several functions in the establishment of *Salmonella* infection, in both epithelial cells and macrophages [61,62]. While the gene expression of SptP was not examined, the up-regulation of the chaperone *sicP* transcripts could indicate early formation of *SicP*-SptP complexes even before contact with a host, as these two genes appear to be transcriptionally coupled [63]. The full significance of the gene expression response of *sicP* to alterations in the pyruvate metabolism pathway remains to be further explored.

We further examined the effects of the mutant strains on the infectivity of THP-1 macrophage cell line. We observed no significant difference in the internalization of *Salmonella* mutants over the wild-type strain, whether the invasion assays were using opsonized or non-opsonized bacteria (Figures 2 & 3). Since we were interested in addressing the effects of our mutations on SPI-1-dependent virulence, we chose to examine SPI-1 gene expression using invasion assays with non-opsonized bacteria. Interestingly, while no significant internalization between strains was observed, SPI-1 gene expression was mostly down-regulated in the mutants after a challenge of THP-1 cells (Tables 3 & 4). During the systemic phase of macrophage infection by *Salmonella*, SPI-1 expression is typically repressed to allow internal replication of the bacterium and suppress excess apoptosis [64]. Therefore, as differentiated THP-1 are phagocytic-type cells and

infection is largely established via SPI2-encoded T3SS, low SPI-1 expression of invaded macrophages could therefore be expected based on previous efforts [65-70].

The specific *Salmonella* knockout genes and the role they may play in its pathogenesis were further examined. Our data supports the overall results of Huang et al. [50] that a *pflB* mutation in *Salmonella* can effect SPI-1 gene expression as measured from cultures alone. In addition to the previous work [50], we identified multiple SPI-1 genes that are differentially expressed both in cultures and at several time-points post-infection of an *in vitro* challenge using human HCT-8 cells. Knockout of the *pflB* gene likely produces an imbalance of metabolites in which the presence mimics the conditions in the host gastrointestinal tract [50]. We further show that another gene of the *Salmonella* pyruvate pathway (*adhE*) can have a similar effect on intestinal epithelia, suggesting that gut metabolites in addition to formate and/or other feedback mechanisms due to environmental cues may be at work. As *AdhE* is a multifunctional enzyme, another possibility for increased invasion of a Δ *adhE* mutant in addition to an SPI1-mediated virulence mechanism could be formulated. A lack of *AdhE* appears to stimulate type I fimbrial adhesion [71,72], important adhesions on the surface of *Salmonella* that may mediate attachment and internalization of bacteria [73]. Further experimentation would be warranted to determine the exact mechanism and co-factors involved with the pyruvate pathway involvement in *Salmonella* virulence.

For the generation of mutant strains, each recombination event was confirmed by PCR using primers to the flanking genomic regions. These confirmations, in addition to the evidence provided in this study, give confidence in the specificity of our deletions. By using lambda-red for site-specific mutagenesis, evidence suggests any potential downstream effects are reduced or eliminated with this system over that of random mutagenesis/transposons, including issues of polarity (e.g. [48]). We do acknowledge however that complementation assays to restore the phenotype could provide additional evidence, but these were not performed here.

We also acknowledge the role the host microbiome plays in *Salmonella* pathogenicity [74], particularly since microbiota may contribute to pyruvate pathway metabolites utilized in the establishment of infection [50]. The mammalian gastrointestinal tract contains high levels of short chain fatty acids as the result of the breakdown of foodstuffs by the digestive processes and the action of resident bacteria [75]. We therefore would be highly interested in future *in vivo* studies to help determine the mechanisms involved in the interplay between the host genetics, the gut environment, and the microbiome on *Salmonella* pathogenesis when dealing with mutant strains produced through metabolic pathway mutations.

Conclusions

Deletions of two genes, *pflB* and *adhE*, involved in the pyruvate metabolism pathway of *Salmonella* Typhimurium were shown to alter SPI-1 gene expression. In bacterial cell culture alone, five SPI-1 genes from the two mutant strains were up-regulated from the wild-type. In a challenge with human HCT-8 and THP-1 cell lines, SPI-1 gene expression was also significantly altered from the wild-type at various times post-infection. We therefore have identified a set of SPI-1 genes that crosstalk between *Salmonella* pyruvate metabolism and virulence. This work provides a further framework for examining the role that central metabolism plays in the infection of *Salmonella* Typhimurium.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JA and XL designed assays and performed the necessary experiments to generate *Salmonella* mutants. JA, CC and CH performed cell culture and gene expression experiments. HZ conceived, planned and guided this study. JA and HZ drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors would like to thank Dr. Helene Andrews-Polymeris at Texas A&M University for helpful discussions and technical assistance in the generation of *Salmonella* mutants. The authors also thank Drs. Palmy Jesudhasan and Suresh Pillai at Texas A&M University for use of equipment and providing some of the *Salmonella* qPCR primers used in this study. This work was supported in part by a USDA NIFA Postdoctoral Fellowship Grant (Award No. 2011-67012-30684).

Author details

¹Department of Animal Science, University of California, Davis, CA 95616, USA. ²Department of Poultry Science, Texas A&M University, College Station, TX 77845, USA. ³College of Animal Science and Technology, Shandong Agricultural University, Taian, Shandong 271018, China.

Received: 12 October 2012 Accepted: 20 February 2013

Published: 25 February 2013

References

1. Ellermeier C, Slauch J: **The Genus *Salmonella***. In *The Prokaryotes*. Edited by Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E. New York: Springer; 2006:123–158.
2. Raffatellu M, Tukul C, Chessa D, Wilson RP, Baumler AJ: **The Intestinal Phase of *Salmonella* Infections**. In *Salmonella: Molecular Biology and Pathogenesis*. Edited by Rhen M, Maskell D, Mastroeni P, Threlfall J. Norfolk UK: Taylor & Francis; 2007:31–52.
3. Litrup E, Torpdahl M, Malorny B, Huehn S, Christensen H, Nielsen EM: **Association between phylogeny, virulence potential and serovars of *Salmonella enterica***. *Infect Genet Evol* 2010, **10**(7):1132–1139.
4. Brenner FW, Villar RG, Angulo FJ, Tauxe R, Swaminathan B: ***Salmonella* nomenclature**. *J Clin Microbiol* 2000, **38**(7):2465–2467.
5. Fedorka-Cray PJ, Whipp SC, Isaacson RE, Nord N, Lager K: **Transmission of *Salmonella typhimurium* to swine**. *Vet Microbiol* 1994, **41**(4):333–344.
6. Harvey PC, Watson M, Hulme S, Jones MA, Lovell M, Berchieri A Jr, Young J, Bumstead N, Barrow P: ***Salmonella enterica* serovar typhimurium colonizing the lumen of the chicken intestine grows slowly and upregulates a unique set of virulence and metabolism genes**. *Infect Immun* 2011, **79**(10):4105–4121.
7. Callaway T, Edrington T, Anderson R, Byrd J, Nisbet D: **Gastrointestinal microbial ecology and the safety of our food supply as related to *Salmonella***. *J Anim Sci* 2008, **86**(14 suppl):E163–E172.
8. Hansen-Wester I, Hensel M: ***Salmonella* pathogenicity islands encoding type III secretion systems**. *Microbes Infect* 2001, **3**(7):549–559.
9. Shea JE, Hensel M, Gleeson C, Holden DW: **Identification of a virulence locus encoding a second type III secretion system in *Salmonella typhimurium***. *Proc Natl Acad Sci USA* 1996, **93**(6):2593–2597.
10. Ellermeier JR, Slauch JM: **Adaptation to the host environment: regulation of the SPI1 type III secretion system in *Salmonella enterica* serovar Typhimurium**. *Curr Opin Microbiol* 2007, **10**(1):24–29.
11. Morgan E: ***Salmonella* Pathogenicity Islands**. In *Salmonella: Molecular Biology and Pathogenesis*. Edited by Rhen M, Maskell D, Mastroeni P, Threlfall J. Norfolk UK: Taylor & Francis; 2007:67–88.
12. Altier C: **Genetic and environmental control of *Salmonella* invasion**. *J Microbiol* 2005, **43**(Spec No):85–92.
13. Kimbrough TG, Miller SI: **Contribution of *Salmonella typhimurium* type III secretion components to needle complex formation**. *Proc Natl Acad Sci USA* 2000, **97**(20):11008–11013.
14. Kimbrough TG, Miller SI: **Assembly of the type III secretion needle complex of *Salmonella typhimurium***. *Microbes Infect* 2002, **4**(1):75–82.
15. Kubori T, Matsushima Y, Nakamura D, Uralil J, Lara-Tejero M, Sukhan A, Galan JE, Aizawa SI: **Supramolecular structure of the *Salmonella typhimurium* type III protein secretion system**. *Science* 1998, **280**(5363):602–605.
16. Lee CA, Jones BD, Falkow S: **Identification of a *Salmonella typhimurium* invasion locus by selection for hyperinvasive mutants**. *Proc Natl Acad Sci USA* 1992, **89**(5):1847–1851.
17. Mills DM, Bajaj V, Lee CA: **A 40 kb chromosomal fragment encoding *Salmonella typhimurium* invasion genes is absent from the corresponding region of the *Escherichia coli* K-12 chromosome**. *Mol Microbiol* 1995, **15**(4):749–759.
18. Sukhan A, Kubori T, Wilson J, Galan JE: **Genetic analysis of assembly of the *Salmonella enterica* serovar Typhimurium type III secretion-associated needle complex**. *J Bacteriol* 2001, **183**(4):1159–1167.
19. Collazo CM, Galan JE: **Requirement for exported proteins in secretion through the invasion-associated type III system of *Salmonella typhimurium***. *Infect Immun* 1996, **64**(9):3524–3531.
20. Collazo CM, Galan JE: **The invasion-associated type III system of *Salmonella typhimurium* directs the translocation of Sip proteins into the host cell**. *Mol Microbiol* 1997, **24**(4):747–756.
21. Cornelis GR: **The type III secretion injectisome**. *Nat Rev Microbiol* 2006, **4**(11):811–825.
22. Guiney DG, Lesnick M: **Targeting of the actin cytoskeleton during infection by *Salmonella* strains**. *Clin Immunol* 2005, **114**(3):248–255.
23. Galan JE: ***Salmonella* interactions with host cells: type III secretion at work**. *Annu Rev Cell Dev Biol* 2001, **17**:53–86.
24. Abrahams GL, Hensel M: **Manipulating cellular transport and immune responses: dynamic interactions between intracellular *Salmonella enterica* and its host cells**. *Cell Microbiol* 2006, **8**(5):728–737.
25. McLaughlin LM, Govoni GR, Gerke C, Gopinath S, Peng K, Laidlaw G, Chien YH, Jeong HW, Li Z, Brown MD, et al: **The *Salmonella* SPI2 effector SseI mediates long-term systemic infection by modulating host cell migration**. *PLoS Pathog* 2009, **5**(11):e1000671.
26. Coombes BK, Brown NF, Valdez Y, Brumell JH, Finlay BB: **Expression and secretion of *Salmonella* pathogenicity island-2 virulence genes in response to acidification exhibit differential requirements of a functional type III secretion apparatus and SsaL**. *J Biol Chem* 2004, **279**(48):49804–49815.
27. Hapfelmeier S, Stecher B, Barthel M, Kremer M, Muller AJ, Heikenwalder M, Stallmach T, Hensel M, Pfeffer K, Akira S, et al: **The *Salmonella* pathogenicity island (SPI)-2 and SPI-1 type III secretion systems allow *Salmonella* serovar typhimurium to trigger colitis via MyD88-dependent and MyD88-independent mechanisms**. *J Immunol* 2005, **174**(3):1675–1685.
28. Lostroh CP, Lee CA: **The *Salmonella* pathogenicity island-1 type III secretion system**. *Microbes Infect* 2001, **3**(14–15):1281–1291.
29. Schlumberger MC, Hardt WD: ***Salmonella* type III secretion effectors: pulling the host cell's strings**. *Curr Opin Microbiol* 2006, **9**(1):46–54.
30. Coombes BK, Coburn BA, Potter AA, Gomis S, Mirakhor K, Li Y, Finlay BB: **Analysis of the contribution of *Salmonella* pathogenicity islands 1 and 2 to enteric disease progression using a novel bovine ileal loop model and a murine model of infectious enterocolitis**. *Infect Immun* 2005, **73**(11):7161–7169.
31. Hapfelmeier S, Ehrbar K, Stecher B, Barthel M, Kremer M, Hardt WD: **Role of the *Salmonella* pathogenicity island 1 effector proteins SipA, SopB, SopE, and SopE2 in *Salmonella enterica* subspecies 1 serovar Typhimurium colitis in streptomycin-pretreated mice**. *Infect Immun* 2004, **72**(2):795–809.

32. Brawn LC, Hayward RD, Koronakis V: *Salmonella* SPI1 effector SipA persists after entry and cooperates with a SPI2 effector to regulate phagosome maturation and intracellular replication. *Cell Host Microbe* 2007, **1**(1):63–75.
33. Lawley TD, Chan K, Thompson LJ, Kim CC, Govoni GR, Monack DM: Genome-wide screen for *Salmonella* genes required for long-term systemic infection of the mouse. *PLoS Pathog* 2006, **2**(2):e11.
34. Steele-Mortimer O, Brumell JH, Knodler LA, Meresse S, Lopez A, Finlay BB: The invasion-associated type III secretion system of *Salmonella enterica* serovar Typhimurium is necessary for intracellular proliferation and vacuole biogenesis in epithelial cells. *Cell Microbiol* 2002, **4**(1):43–54.
35. Buckner MM, Croxen MA, Arena ET, Finlay BB: A comprehensive study of the contribution of *Salmonella enterica* serovar Typhimurium SPI2 effectors to bacterial colonization, survival, and replication in typhoid fever, macrophage, and epithelial cell infection models. *Virulence* 2011, **2**(3):208–216.
36. Lilic M, Stebbins CE: Re-structuring the host cell: up close with *Salmonella*'s molecular machinery. *Microbes Infect* 2004, **6**(13):1205–1211.
37. Patel JC, Galan JE: Manipulation of the host actin cytoskeleton by *Salmonella*—all in the name of entry. *Curr Opin Microbiol* 2005, **8**(1):10–15.
38. Ly KT, Casanova JE: Mechanisms of *Salmonella* entry into host cells. *Cell Microbiol* 2007, **9**(9):2103–2111.
39. Brown NF, Vallance BA, Coombes BK, Valdez Y, Coburn BA, Finlay BB: *Salmonella* pathogenicity island 2 is expressed prior to penetrating the intestine. *PLoS Pathog* 2005, **1**(3):e32.
40. Dandekar T, Astrid F, Jasmin P, Hensel M: *Salmonella enterica*: a surprisingly well-adapted intracellular lifestyle. *Front Microbiol* 2012, **3**:164.
41. Bajaj V, Lucas RL, Hwang C, Lee CA: Co-ordinate regulation of *Salmonella typhimurium* invasion genes by environmental and regulatory factors is mediated by control of *hilA* expression. *Mol Microbiol* 1996, **22**(4):703–714.
42. Lee CA, Falkow S: The ability of *Salmonella* to enter mammalian cells is affected by bacterial growth state. *Proc Natl Acad Sci USA* 1990, **87**(11):4304–4308.
43. Lundberg U, Vinatzer U, Berdnik D, von Gabain A, Baccarini M: Growth phase-regulated induction of *Salmonella*-induced macrophage apoptosis correlates with transient expression of SPI-1 genes. *J Bacteriol* 1999, **181**(11):3433–3437.
44. Hernandez SB, Cota I, Ducret A, Aussel L, Casadesus J: Adaptation and preadaptation of *Salmonella enterica* to bile. *PLoS Genet* 2012, **8**(1):e1002459.
45. Pang E, Tien-Lin C, Selvaraj M, Chang J, Kwang J: Deletion of the *aceE* gene (encoding a component of pyruvate dehydrogenase) attenuates *Salmonella enterica* serovar Enteritidis. *FEMS Immunol Med Microbiol* 2011, **63**(1):108–118.
46. Bowden SD, Rowley G, Hinton JC, Thompson A: Glucose and glycolysis are required for the successful infection of macrophages and mice by *Salmonella enterica* serovar typhimurium. *Infect Immun* 2009, **77**(7):3117–3126.
47. Bowden SD, Ramachandran VK, Knudsen GM, Hinton JC, Thompson A: An incomplete TCA cycle increases survival of *Salmonella* Typhimurium during infection of resting and activated murine macrophages. *PLoS One* 2010, **5**(11):e13871.
48. Santiviago CA, Reynolds MM, Porwollik S, Choi SH, Long F, Andrews-Polymenis HL, McClelland M: Analysis of pools of targeted *Salmonella* deletion mutants identifies novel genes affecting fitness during competitive infection in mice. *PLoS Pathog* 2009, **5**(7):e1000477.
49. Lee SH, Jung BY, Rayamahji N, Lee HS, Jeon WJ, Choi KS, Kweon CH, Yoo HS: A multiplex real-time PCR for differential detection and quantification of *Salmonella* spp., *Salmonella enterica* serovar Typhimurium and Enteritidis in meats. *J Vet Sci* 2009, **10**(1):43–51.
50. Huang Y, Suyemoto M, Garner CD, Cicconi KM, Altier C: Formate acts as a diffusible signal to induce *Salmonella* invasion. *J Bacteriol* 2008, **190**(12):4233.
51. Valle E, Guiney DG: Characterization of *Salmonella*-induced cell death in human macrophage-like THP-1 cells. *Infect Immun* 2005, **73**(5):2835–2840.
52. Pfaffl MW, Horgan GW, Dempfle L: Relative expression software tool (REST) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Res* 2002, **30**(9):e36.
53. Ibarra JA, Steele-Mortimer O: *Salmonella*—the ultimate insider. *Salmonella* virulence factors that modulate intracellular survival. *Cell Microbiol* 2009, **11**(11):1579–1586.
54. Hu Q, Coburn B, Deng W, Li Y, Shi X, Lan Q, Wang B, Coombes BK, Finlay BB: *Salmonella enterica* serovar Senftenberg human clinical isolates lacking SPI-1. *J Clin Microbiol* 2008, **46**(4):1330–1336.
55. Srikanth CV, Mercado-Lubo R, Hallstrom K, McCormick BA: *Salmonella* effector proteins and host-cell responses. *Cell Mol Life Sci* 2011, **68**(22):3687–3697.
56. Tucker SC, Galan JE: Complex function for SicA, a *Salmonella enterica* serovar typhimurium type III secretion-associated chaperone. *J Bacteriol* 2000, **182**(8):2262–2268.
57. Kaniga K, Tucker S, Trollinger D, Galan JE: Homologs of the Shigella IpaB and IpaC invasins are required for *Salmonella typhimurium* entry into cultured epithelial cells. *J Bacteriol* 1995, **177**(14):3965–3971.
58. Schechter LM, Lee CA: AraC/XylS family members, HilC and HilD, directly bind and derepress the *Salmonella typhimurium* *hilA* promoter. *Mol Microbiol* 2001, **40**(6):1289–1299.
59. Olekhnovich IN, Kadner RJ: DNA-binding activities of the HilC and HilD virulence regulatory proteins of *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 2002, **184**(15):4148–4160.
60. Ellermeier CD, Slauch JM: RtsA and RtsB coordinately regulate expression of the invasion and flagellar genes in *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 2003, **185**(17):5096.
61. Lin SL, Le TX, Cowen DS: SptP, a *Salmonella typhimurium* type III-secreted protein, inhibits the mitogen-activated protein kinase pathway by inhibiting Raf activation. *Cell Microbiol* 2003, **5**(4):267–275.
62. Haraga A, Miller SI: A *Salmonella enterica* serovar typhimurium translocated leucine-rich repeat effector protein inhibits NF- κ B-dependent gene expression. *Infect Immun* 2003, **71**(7):4052–4058.
63. Button JE, Galan JE: Regulation of chaperone/effector complex synthesis in a bacterial type III secretion system. *Mol Microbiol* 2011, **81**(6):1474–1483.
64. Kage H, Takaya A, Ohya M, Yamamoto T: Coordinated regulation of expression of *Salmonella* pathogenicity island 1 and flagellar type III secretion systems by ATP-dependent ClpXP protease. *J Bacteriol* 2008, **190**(7):2470–2478.
65. Cirillo DM, Valdivia RH, Monack DM, Falkow S: Macrophage-dependent induction of the *Salmonella* pathogenicity island 2 type III secretion system and its role in intracellular survival. *Mol Microbiol* 1998, **30**(1):175–188.
66. Deiwick J, Nikolaus T, Erdogan S, Hensel M: Environmental regulation of *Salmonella* pathogenicity island 2 gene expression. *Mol Microbiol* 1999, **31**(6):1759–1773.
67. Beuzon CR, Meresse S, Unsworth KE, Ruiz-Albert J, Garvis S, Waterman SR, Ryder TA, Boucrot E, Holden DW: *Salmonella* maintains the integrity of its intracellular vacuole through the action of SifA. *EMBO J* 2000, **19**(13):3235–3249.
68. Eriksson S, Lucchini S, Thompson A, Rhen M, Hinton JC: Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica*. *Mol Microbiol* 2003, **47**(1):103–118.
69. Pfeifer CG, Marcus SL, Steele-Mortimer O, Knodler LA, Finlay BB: *Salmonella typhimurium* virulence genes are induced upon bacterial invasion into phagocytic and nonphagocytic cells. *Infect Immun* 1999, **67**(11):5690–5698.
70. Drecktrah D, Knodler LA, Ireland R, Steele-Mortimer O: The mechanism of *Salmonella* entry determines the vacuolar environment and intracellular gene expression. *Traffic* 2006, **7**(1):39–51.
71. Chuang YC, Wang KC, Chen YT, Yang CH, Men SC, Fan CC, Chang LH, Yeh KS: Identification of the genetic determinants of *Salmonella enterica* serotype Typhimurium that may regulate the expression of the type 1 fimbriae in response to solid agar and static broth culture conditions. *BMC Microbiol* 2008, **8**:126.
72. Umitsuki G, Wachi M, Takada A, Hikichi T, Nagai K: Involvement of RNase G in *in vivo* mRNA metabolism in *Escherichia coli*. *Genes Cells* 2001, **6**(5):403–410.
73. Guo A, Lasaro MA, Sirard JC, Kraehenbuhl JP, Schifferli DM: Adhesin-dependent binding and uptake of *Salmonella enterica* serovar Typhimurium by dendritic cells. *Microbiology* 2007, **153**(Pt 4):1059–1069.
74. Sekirov I, Gill N, Jogova M, Tam N, Robertson M, de Llanos R, Li Y, Finlay BB: *Salmonella* SPI-1-mediated neutrophil recruitment during enteric colitis is associated with reduction and alteration in intestinal microbiota. *Gut Microbes* 2010, **1**(1):30–41.
75. Lawhon SD, Maurer R, Suyemoto M, Altier C: Intestinal short-chain fatty acids alter *Salmonella typhimurium* invasion gene expression and virulence through BarA/SirA. *Mol Microbiol* 2002, **46**(5):1451–1464.

doi:10.1186/2049-1891-4-5

Cite this article as: Abernathy et al.: Deletions in the pyruvate pathway of *Salmonella* Typhimurium alter SPI1-mediated gene expression and infectivity. *Journal of Animal Science and Biotechnology* 2013 **4**:5.