

# UC Davis

## UC Davis Previously Published Works

### Title

Exposure to umbilical cord management approaches and death or neurodevelopmental impairment at 22-26 months corrected age after extremely preterm birth.

### Permalink

<https://escholarship.org/uc/item/3x42h9kv>

### Journal

Archives of Disease in Childhood. Fetal and Neonatal Edition, 108(3)

### Authors

Handley, Sara  
Kumbhat, Neha  
Eggleston, Barry  
[et al.](#)

### Publication Date

2023-05-01

### DOI

10.1136/archdischild-2022-324565

Peer reviewed



Published in final edited form as:

*Arch Dis Child Fetal Neonatal Ed.* 2023 May ; 108(3): 224–231. doi:10.1136/archdischild-2022-324565.

## Exposure to Umbilical Cord Management Approaches and Death or Neurodevelopmental Impairment at 22–26 Months Corrected Age after Extremely Preterm Birth

Sara C. Handley, MD MSCE<sup>1,2,\*</sup>, Neha Kumbhat, MD MS Epi<sup>3,\*</sup>, Barry Eggleston, MS<sup>4</sup>, Elizabeth E. Foglia, MD MSCE<sup>1</sup>, Alexis S. Davis, MD MS Epi<sup>3</sup>, Krisa P. Van Meurs, MD<sup>3</sup>, Satyan Lakshminrusimha, MD<sup>5</sup>, Michele C. Walsh, MD MS<sup>6</sup>, Kristi L. Watterberg, MD<sup>7</sup>, Myra H. Wyckoff, MD<sup>8</sup>, Abhik Das, PhD<sup>9</sup>, Sara B. DeMauro, MD MSCE<sup>1</sup>,  
Generic Database Subcommittee of the NICHD Neonatal Research Network

<sup>1</sup>Children's Hospital of Philadelphia, Philadelphia, PA,

<sup>2</sup>Leonard Davis Institute of Health Economics, The University of Pennsylvania, Philadelphia, PA,

<sup>3</sup>Stanford University, Stanford, CA,

<sup>4</sup>RTI International, Research Triangle Park, NC,

<sup>5</sup>University of California, Davis, Sacramento, CA,

<sup>6</sup>University Hospitals Rainbow Babies and Children's Hospital, Case Western Reserve University, Cleveland, OH,

<sup>7</sup>University of New Mexico Health Sciences Center, Albuquerque, NM,

<sup>8</sup>University of Texas, Southwestern Medical Center, Dallas, TX,

<sup>9</sup>RTI International, Rockville, MD

### Abstract

**Objective:** To compare death or severe neurodevelopmental impairment (NDI) at 22–26 months' corrected age (CA) among extremely preterm infants following exposure to different forms of umbilical cord management.

**Design:** Retrospective study.

---

**Corresponding author:** Sara B. DeMauro, MD MSCE, The Children's Hospital of Philadelphia, Main Hospital, Division of Neonatology, 2<sup>nd</sup> Floor, 3401 Civic Center Blvd, Philadelphia, PA 19104, Telephone: 215-913-8931, Fax: 215-590-3051, [demauro@chop.edu](mailto:demauro@chop.edu).

\*Indicates co-first authorship

**Contributorship statement:** SH, NK, and SD designed the project and the main conceptual ideas. BE completed the data analysis. SH, NK, BE, AD, KVM, EF, SL, MW, KW, MW, AD and SD interpreted the analysis. SH and NK equally contributed to the article draft, which was critically revised by AD, KVM, EF, SL, MW, KW, MW, AD, and SD. National Institute of Child Health and Human Development, Eunice Kennedy Shriver provided the Generic Database for this study. SH, NK, BE, AD, KVM, EF, SL, MW, KW, MW, AD, SD, and the Generic Database subcommittee gave the final approval to the version submitted for publication.

**ClinicalTrials.gov ID:** Generic Database: [NCT00063063](https://clinicaltrials.gov/ct2/show/study/NCT00063063)

**Research Ethics Approval:** This study was performed utilizing the Neonatal Research Network (NRN) Generic Database (GDB) of the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD). Institutional review board approval for the NRN GDB registry was obtained for each center.

**Setting:** *Eunice Kennedy Shriver* National Institute of Child Health and Human Development Neonatal Research Network registry.

**Patients:** Infants born <27 weeks' gestation in 2016–2018 without severe congenital anomalies who received active treatment after birth and underwent neurodevelopmental assessments between 22–26 months' CA.

**Exposures:** Immediate cord clamping (ICC), delayed cord clamping (DCC) or umbilical cord milking (UCM).

**Main Outcomes and Measure:** Primary composite outcome of death or severe NDI at 22–26 months' CA, defined as severe cerebral palsy, Bayley-III cognitive/motor composite score <70, bilateral deafness or blindness; individual components were examined as secondary outcomes. Multivariable regression examined associations, adjusting for risk factors identified *a priori* and potential confounders. Mediation analysis explored the effect of severe intraventricular hemorrhage (IVH) on the exposure-outcome relationship.

**Results:** Among 1,900 infants, 64.1% were exposed to ICC, 27.8% to DCC, and 8.1% to UCM. Compared to ICC-exposed infants, DCC-exposed infants had lower odds of death or severe NDI (aOR 0.64, 95% CI 0.50–0.83). No statistically significant differences were observed when comparing UCM with either ICC or DCC, or between secondary outcomes across groups. Association between cord management and the primary outcome was not mediated by severe IVH.

**Conclusion:** Compared to ICC, DCC exposure was associated with lower death or severe NDI at 22–26 months' CA among extremely preterm infants, which was not mediated by severe IVH.

## Keywords

Delayed cord clamping (DCC); umbilical cord milking (UCM); Immediate cord clamping (ICC); Neurodevelopmental impairment (NDI); Neonatal Research Network (NRN); Generic Database (GDB)

## Introduction

The prognosis of extremely low birth weight (ELBW) infants surviving to hospital discharge remains guarded, as a large proportion experience neurodevelopmental impairment (NDI). [1]–[2] Severe intraventricular hemorrhage (IVH) may contribute to NDI in the ELBW population.[3]–[4] Forms of placental transfusion, such as delayed cord clamping (DCC), offer a potential neuro-protective strategy for ELBW infants.[5] Studies evaluating the relationships of DCC, umbilical cord milking (UCM), and immediate cord clamping (ICC) with severe IVH have reported mixed findings, leaving questions about neuroprotection and neurodevelopment unanswered.

Most studies examining umbilical cord management and neurodevelopmental outcomes are limited by small sample size, single center cohorts with heterogeneous neurodevelopmental measures, and variable assessment time points. [6] [7] [8] Recent data from the Australian Placental Transfusion Study (APTS) showed a 30% relative risk reduction of death with no difference in major disabilities at two years among DCC-exposed infants, compared to ICC-exposed infants.[9] The International Liaison Committee of Resuscitation systematic review

and meta-analysis, published in 2015, recommends DCC following extremely preterm birth, but does not recommend UCM due to low quality evidence and limited long-term outcome data.[10] Evidence about neurodevelopmental outcomes comparing all three approaches to cord management, DCC, ICC, and UCM, remains limited.

We have examined short-term outcomes following different approaches to umbilical cord management among extremely preterm infants and demonstrated that DCC is the preferred approach, as UCM was associated with increases in severe IVH. [11] [12] Using the detailed follow-up data collected by the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD) Neonatal Research Network (NRN), the objective of the current study was to compare rates of death or severe NDI at 22–26 months' corrected age (CA) after exposure to ICC, DCC, or UCM among infants born <27 weeks' gestation. Individual components of the primary composite outcome were examined as secondary outcomes. This study also explored whether the relationship between cord management and the primary outcome was mediated by severe IVH.

## Methods

### Patient selection

This retrospective analysis included infants 22<sup>0/7</sup> to 26<sup>6/7</sup> weeks' gestation born at NRN centers from January 1<sup>st</sup> 2016 to December 31<sup>st</sup> 2018. The NRN Generic Database (GDB) registry collects sociodemographic, maternal, and neonatal information prospectively during the birth hospitalization using previously defined, pre-specified definitions.[12] [13] The Follow-up (FU) database collects neurodevelopmental assessments of eligible survivors born <27 weeks' gestation at 22–26 months' CA. Each participating center obtained institutional review board (IRB) approval for both databases; per local IRB guidance a waiver of consent was approved or parental consent obtained.

Infants with missing exposure documentation, those with severe congenital malformations, including congenital heart disease and/or genetic syndromes, and those who did not receive active treatment after birth (any ventilatory support, chest compressions, epinephrine administration, surfactant, mechanical ventilation, or parental nutrition) were excluded [14]. Infants with neurodevelopmental assessments completed more than four months outside of the target window of 22–26 months CA were excluded. The final cohort included infants who died prior to follow-up and survivors with complete neurodevelopmental assessments. Survivors without NDI data were deemed lost to follow-up. The total number of infants exposed to the three cord management forms was used as a denominator to calculate lost-to-follow-up rates.

### Exposures

The exposure was the documented approach to umbilical cord management performed in the delivery room as decided by the clinical team and center practice guidelines. Exposure to DCC, UCM, or ICC was determined using two yes/no questions answered for each infant in the GDB: 1) Is there documentation of at least 30 seconds of DCC? and 2) Is there documentation of UCM? Infants with answers to both questions documented as 'no' were

classified as exposed to ICC, functionally defined as cord clamping <30 seconds after birth. Infants with missing data and those with answers to both questions documented as ‘yes’ were excluded, thus the analytic cohort only includes infants with one documented exposure.

## Outcomes

The primary outcome was a composite of death before follow-up evaluation or severe NDI at follow-up (defined below). The neurodevelopmental assessment included a physical examination with standardized neurological examination and developmental evaluation using the Bayley Scales of Infant and Toddler Development, 3<sup>rd</sup> Edition (Bayley-III), administered by certified examiners.[15][16] A score <70 on any of the Bayley-III composite scales indicates development that is two standard deviations below the expected population mean score. The language composite score includes assessment of both receptive and expressive language. Motor abilities were also classified using the Gross Motor Function Classification System (GMFCS) and severity of cerebral palsy (CP) was classified using the GMFCS level: mild (level 1), moderate (level 2–3), and severe CP (level 4–5). [17] Bilateral blindness was defined as vision <20/200 despite corrective lenses bilaterally. Hearing impairment was defined as permanent hearing loss with or without amplification and obtained by parental report. The NRN follow up subcommittee defines severe NDI as the presence of any one of the following: severe CP, Bayley-III cognitive composite score <70, Bayley-III motor composite score <70, bilateral blindness or hearing impairment.[18] Secondary outcomes were: 1) death prior to follow-up evaluation, and among survivors: 2) severe NDI, 3) moderate or severe CP (GMFCS 2), 4) severity of CP based on GMFCS level, 5) Bayley-III cognitive score <70, 6) Bayley-III cognitive score <85, 6) Bayley-III motor score <70 7) Bayley-III composite language score <70, 8) bilateral blindness and 9) hearing impairment.

## Statistical analysis

We examined three comparisons: DCC versus ICC, UCM versus ICC, and DCC versus UCM. Unadjusted comparisons were made between groups for sociodemographic and perinatal-neonatal characteristics using t-tests and Fischer’s exact test for continuous and categorical variables, respectively. Risk-adjusted associations of cord management technique with the primary composite outcome were estimated using complete case analysis and multivariable logistic regression. The regression model incorporated the following variables: 1) risk factors for death or severe NDI identified *a priori*: gestational age (in weeks), sex, race/ethnicity, maternal education, and antenatal steroids; [19] 2) covariates occurring prior to the exposure that were statistically imbalanced across the three exposure groups (p-value < 0.1) in unadjusted analyses; 3) birth year; and 4) NRN center as a random effect. The same model examined secondary outcomes. Models assessing severe NDI among survivors also included age at follow-up assessment. No adjustments were made for multiple comparisons. To examine the effect of the coronavirus 2019 (COVID-19) pandemic on disruptions to healthcare delivery, specifically, the effect on follow-up assessment timing, we performed a sensitivity analysis limiting the study population to infants who reached 26 months’ CA by March 15<sup>th</sup>, 2020.

A prespecified mediation analysis examined whether severe IVH mediated the relationship between cord management and the primary outcome. Given that severe IVH is a binary variable, a model-based causal mediation analysis using differences in proportions was used to report direct and indirect effects of the mediator on the outcome of interest. This analysis quantifies the association between the exposure and the outcome without (direct) and through (indirect) the mediator, respectively.[20] The mediation analysis adjusted for all variables in the primary model.

Analyses were conducted by the NRN Data Coordinating Center (RTI International) using R statistical software version 3.5.1 (Feather Spray, Vienna, Austria).

## Results

### Study population

During the study period, 2,277 infants were born between 22<sup>0/7</sup> and 26<sup>6/7</sup> weeks' gestation and met initial inclusion criteria. Infants seen more than four months outside of the target neurodevelopmental assessment window of 22–26 months CA were excluded (n=77) and those without follow-up data were deemed lost to follow-up (n=300, 13.2%). The final analytic cohort included 1,900 infants; 583 died before follow-up assessment and 1,317 had complete neurodevelopmental assessments. Compared to infants in the analytic cohort, those lost to follow-up were slightly older, had higher birth weights, higher Apgar scores, and lower rates of severe brain injury (Supplemental table 1).

In the final cohort, 1,218 (64.1%) infants were exposed to ICC, 528 (27.8%) to DCC, and 154 (8.1%) to UCM (Figure 1). Baseline characteristics differed among the three groups (Table 1). For example, the proportion of infants exposed to antenatal steroids was lower in the ICC group and the proportion of infants who received PPV or intubation was lowest in the DCC group.

### Primary and secondary outcomes

Compared to ICC, DCC-exposed infants had significantly lower adjusted odds of death or severe NDI (DCC: 36.3% versus ICC: 50.3%; aOR 0.64, 95% CI 0.50–0.83) (Table 2). A statistically significant difference was not observed in the primary outcome for the remaining two comparisons (UCM versus ICC and DCC versus UCM) (Table 2).

DCC-exposed infants had a significantly lower adjusted odds of death prior to follow-up compared to ICC-exposed infants (DCC: 22.4% versus ICC: 34.3%; aOR 0.61, 95% CI 0.46–0.81) (Table 3). There was no statistically significant difference in severe NDI between infants exposed to DCC versus ICC (16.7% versus 23.0%; aOR 0.74, 95% CI 0.52–1.05). There were no differences in secondary outcomes when comparing UCM to ICC or DCC. The results of the COVID-19-informed sensitivity analysis found no change in the primary outcome (Supplemental table 2).

### Mediation analysis

Compared to ICC, DCC directly reduced death or severe NDI (average direct effect –0.063 after adjustment, p=0.012) with no indirect effect on the primary outcome via severe IVH

(average indirect effect  $-0.009$ ,  $p=0.52$ ). No statistically significant direct or indirect effects were found in the remaining comparisons.

## Discussion

In this large, contemporary observational study, exposure to DCC compared to ICC was associated with a significantly decreased odds of the composite outcome of death or severe NDI. The association was not mediated via severe IVH.

We have previously reported associations between umbilical cord management approaches and short-term outcomes. These studies highlighted practice variation across the NRN and demonstrated a benefit of DCC. [12] [13] In the current cohort, only 27.8% of infants were exposed to DCC, which may reflect provider preferences, literature interpretation, or desire for additional follow-up data. The impetus for the current study was sparse literature describing neurodevelopmental outcomes following different forms of cord management, particularly in the extremely preterm infant population. Comparing DCC and ICC, Mercer et al evaluated outcomes of 161 infants <32 weeks' gestation at 18–22 months' CA (77% of the original cohort) using the Bayley-III and found improved motor function following DCC.[6] Recently published data from APTS reported a reduced risk of death or disability at two years CA after DCC compared to ICC (RR 0.83 95% CI 0.72, 0.95) and a 39% reduction in the composite outcome of death or severe NDI.[9] Although our study is observational, the results similarly support DCC over ICC. The other two published trials report long-term outcomes comparing DCC and UCM among preterm infants. Rabe et al assessed children at two (n=39, 67% of the original cohort) and three and a half years (n=29, 50% of the original cohort) and found no differences in Bayley-III scores.[7] However, the inclusion of infants born from 27<sup>0/7</sup> to 32<sup>6/7</sup> weeks' gestation, a more mature population compared to our study, and small sample size may contribute to these findings. Katheria et al reported that among 135 toddlers assessed at 18 months (69% of the original cohort), infants randomized to UCM had higher language and cognitive scores than infants randomized to DCC.[8] Neurodevelopmental outcomes from Katheria's multicenter randomized controlled trial of UCM versus DCC are not yet known.[11] Aside from the APTS findings, published studies of neurodevelopmental outcomes are from trials limited by small samples sizes (29–263 infants) from single institutions, high attrition rates, variation in neurodevelopmental assessment measures, and a wide age range at the time of assessment, making comparisons challenging. The current study uses a large, multicenter cohort with higher rates (>86%) of in-person, standardized neurodevelopmental assessments to demonstrate the effectiveness of umbilical cord management approaches in premature infants outside the context of a trial.

Data regarding cord management techniques and IVH are evolving. The largest trial comparing DCC to ICC and associated meta-analysis reported comparable rates of severe IVH following either exposure.[21] [22] In contrast, the largest trial comparing DCC and UCM was stopped early due to increased severe IVH rates among UCM randomized infants. [11] We reported an association between UCM and severe IVH in the NRN.[12] While a recent meta-analysis comparing umbilical cord management strategies reported that DCC may be beneficial, particularly with regard to hematologic measures, it found no differences between UCM and ICC or DCC in survival or severe morbidities, including IVH.[23]



The increased severe IVH rates with UCM exposure, which have not been appreciated following DCC exposure, have informed guidelines supporting DCC.[24] We completed a pre-specified mediation analysis to further understand the relationship between cord management, severe IVH, and death or severe NDI, specifically to examine whether severe IVH mediates the exposure-outcome relationship. Relative to ICC, we found that exposure to DCC reduced death or severe NDI by 6.3 percentage points, which was not mediated by severe IVH. Within the other exposure comparisons (UCM versus ICC and DCC versus UCM), there was no difference in death or severe NDI and severe IVH was not a mediator.

This study has limitations. As a retrospective observational study, causality cannot be assigned and results are hypothesis generating. In the unadjusted comparisons, DCC-exposed infants had more favorable baseline characteristics, such as higher antenatal steroid exposure rates and lower receipt of resuscitative interventions. Despite model adjustments for variables prior to the exposure, residual confounding may influence study results. Those exposed to ICC may have been sicker at birth or experienced bradycardia after ICC, either of which may prompt more resuscitation. Comparisons with UCM should be interpreted cautiously given the small number of exposed infants. Additionally, this study lacks information surrounding the reason a specific cord management approach was chosen as these decisions are informed by individual clinical scenarios and local center policies. Studies examining long-term outcomes are at risk for attrition bias. In the current study this bias was compounded by the COVID-19 pandemic and associated impact on follow-up assessments, which was evident in the sensitivity analysis cohort that had a loss to follow-up rate of 9.9%, a lower rate compared to the primary cohort. Notably, the sensitivity analysis did not change the results.

Strengths of this study include detailed information on infants born at 15 academic centers across the United States. During the study period, 1,900 infants met inclusion criteria, making this the largest observational study to compare long-term neurodevelopmental outcomes following the three approaches to cord management. Previous studies of long-term outcomes have not included all possible pairwise comparisons of approaches to cord management to discern which approach is most beneficial for ELBW infants. Additionally, this study utilizes in-person, comprehensive, standardized neurodevelopmental assessments by trained examiners. These data further support the effectiveness of DCC in the real-world, clinical context.

In conclusion, in this large, contemporary, observational study of infants born <27 weeks' gestation, DCC was associated with improvement in the primary composite outcome of death or severe NDI at 22–26 months' CA compared to ICC. The protective effect of DCC on death or severe NDI was not mediated by severe IVH.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.



## Acknowledgements

The National Institutes of Health and the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD) provided grant support for the Neonatal Research Network for the Generic Database and the Follow-up Study. NICHD staff provided input into the study design, conduct, analysis, and manuscript drafting; NCATS cooperative agreements provided infrastructure support to the NRN. While NICHD staff had input into the study design, conduct, analysis, and manuscript drafting, the comments and views of the authors do not necessarily represent the views of NICHD, the National Institutes of Health, the Department of Health and Human Services, or the U.S. Government.

Data collected at participating sites of the NICHD Neonatal Research Network were transmitted to RTI International, the data coordinating center (DCC) for the network, which stored, managed and analyzed the data included in this study. On behalf of the NRN, RTI International had full access to all the data in the study and take responsibility for the integrity of the data and accuracy of the data analysis. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

We are indebted to our medical and nursing colleagues and the infants and their parents who agreed to take part in this study. The following investigators, in addition to those listed as authors, participated in this study:

NRN Steering Committee Chair: Richard A. Polin, MD, Division of Neonatology, College of Physicians and Surgeons, Columbia University, (2011-present).

Alpert Medical School of Brown University and Women & Infants Hospital of Rhode Island (UG1 HD27904) – Abbot R. Laptook, MD; Martin Keszler, MD; Betty R. Vohr, MD; Angelita M. Hensman, PhD RNC-NIC; Elisa Vieira, BSN RN; Lucille St. Pierre, BS; Barbara Alksninis, RNC PNP; Andrea Knoll; Mary L. Keszler, MD; Teresa M. Leach, MEd CAES; Elisabeth C. McGowan, MD; Victoria E. Watson, MS CAS.

Case Western Reserve University, Rainbow Babies & Children’s Hospital (UG1 HD21364) – Anna Maria Hibbs, MD MSCE; Nancy S. Newman, RN; Deanne E. Wilson-Costello, MD; Bonnie S. Siner, RN; Harriet G. Friedman, MA.

Children’s Mercy Hospital (UG1 HD68284) – William E. Truog, MD; Eugenia K. Pallotto, MD MSCE; Howard W. Kilbride MD; Cheri Gauldin, RN BS CCRC; Anne Holmes RN MSN MBA-HCM CCRC; Allison Scott, BSN RNC-NIC; Prabhu S. Parimi, MD; Lisa Gaetano, RN MSN.

Cincinnati Children’s Hospital Medical Center, University Hospital, and Good Samaritan Hospital (UG1 HD27853, UL1 TR77) – Stephanie L. Merhar, MD MS; Brenda B. Poindexter, MD MS; Kurt Schibler, MD; Tanya E. Cahill, MD; Cathy Grisby, BSN CCRC; Kristin Kirker, CRC; Sandra Wuertz, RN BSN CLC; Juanita Dudley, RN BSN; Julia Thompson, RN BSN; Lisa Henkes, RN BSN; Sara Stacey, BA; Devan Hayes, BS.

Duke University School of Medicine, University Hospital, University of North Carolina, Duke Regional Hospital, and WakeMed Health & Hospitals (UG1 HD40492, UL1 TR1117) – C. Michael Cotten, MD MHS; Ronald N. Goldberg, MD; William F. Malcolm, MD; Patricia L. Ashley, MD; Deesha Mago-Shah, MD; Mollie Warren, MD; Joanne Finkle, RN JD; Kimberley A. Fisher, PhD FNP-BC IBCLC; Kathryn E. Gustafson, PhD; Matthew M. Laughon, MD MPH; Carl L. Bose, MD; Janice Bernhardt, MS RN; Gennie Bose, RN; Janice Wereszczak, CPNP-AC/PC; Andrea Trembath, MD MPH; Jennifer Talbert, MS RN; Stephen D. Kicklighter, MD; Alexandra Bentley, MD; Laura Edwards, MD; Ginger Rhodes-Ryan, ARNP MSN, NNP-BC; Donna White, RN-BC BSN.

Emory University, Children’s Healthcare of Atlanta, Grady Memorial Hospital, and Emory University Hospital Midtown (UG1 HD27851, UL1 TR454) – Ravi M. Patel, MD MSc; David P. Carlton, MD; Ira Adams-Chapman, MD (deceased); Yvonne Loggins, RN; Diane Bottcher, RN; Sheena L. Carter, PhD; Salathiel Kendrick-Allwood, MD; Maureen Mulligan LaRossa, RN; Judith Laursen, RN; Colleen Mackie, RRT; Amy Sanders, PsyD; Gloria Smikle, PNP; Lynn Wineski, NNP.

*Eunice Kennedy Shriver* National Institute of Child Health and Human Development – Andrew A. Bremer, MD PhD; Rosemary D. Higgins, MD; Stephanie Wilson Archer, MA.

Indiana University, Methodist Hospital, and Riley Hospital for Children (UG1 HD27856, UL1 TR6) – Gregory M. Sokol, MD; Abbey C. Hines, PsyD; Dianne E. Herron, RN CCRC; Donna Watkins, MSN NNP-BC; Susan Gunn, NNP-BC CCRC; Jeff Joyce, CCRC (deceased).

McGovern Medical School at The University of Texas Health Science Center at Houston, Children’s Memorial Hermann Hospital, and Memorial Hermann Southwest Hospital (U10 HD21373, UG1 HD87229) – Jon E. Tyson, MD MPH; Amir M. Khan, MD; Kathleen A. Kennedy, MD MPH; Barbara J. Stoll, MD; Ricardo A. Mosquera, MD MS; Andrea F. Duncan, MD MS; Elizabeth Allain, PhD; Julie Arldt-McAlister, MSN APRN; Fatima Boricha, MD;

Allison G. Dempsey, PhD; Carmen Garcia, RN BSN; Donna J. Hall, RN; Janice John, CPNP; M. Layne Lillie, RN BSN; Karen Martin, RN; Georgia E. McDavid, RN; Shannon L. McKee, EdS; Michelle Poe, PhD RN, Kimberly Rennie, PhD; Tina Reddy, MD; Shawna Rodgers, RNC-NIC BSN; Daniel K. Sperry, RN; Emily Stephens, BSN RNC-NIC; Sharon L. Wright, MT (ASCP).

Nationwide Children's Hospital, The Abigail Wexner Research Institute at Nationwide Children's Hospital, Center for Perinatal Research, The Ohio State University College of Medicine, The Ohio State University Wexner Medical Center, and Riverside Methodist Hospital (UG1 HD68278) – Pablo J. Sánchez, MD; Leif D. Nelin, MD; Jonathan L. Slaughter, MD MPH; Sudarshan R. Jadcherla, MD; Nathalie L. Maitre, MD PhD; Christopher Timan, MD; Keith O. Yeates, MD PhD; Patricia Luzader, RN; Julie Gutentag, RN BSN; Jennifer L. Grothaus, BA RN BSN; Melanie Stein, RRT BBS; Rox Ann Sullivan, RN BSN; Cole D. Hague, BA MS; Helen Carey, PT DHSc PCS; Michelle Chao, BS; Stephanie Burkhardt, BS MPH; Margaret Sullivan, BS; Lina Yossef-Salameh, MD; Mary Ann Nelin, MD; Erna Clark, BA; Julie C. Shadd, BSN RD; Courtney Park, RN BSN; Courtney Cira, BS; Erin Fearnis; Kristi Small, BS; Sarah A. Keim, PhD MA MS; Christine A. Fortney, RN PhD; Aubrey Fowler, BS, Jacqueline McCool; Lindsay Pietruszewski, PT DPT; Jessica Purnell, BS CCRC; Kyrstin Warnimont, BS; Laura Marzec, MD; Bethany Miller, RN BSN; Demi R. Beckford, MHS; Hallie Baugher, BS MSN; Julia Newton, MPH; Katelyn Levengood, PT DPT; Nancy Batterson, OT/L; Brittany DeSantis, BS.

RTI International (UG1 HD36790) – Carla M. Bann, PhD; Dennis Wallace, PhD; Marie G. Gantz, PhD; Jeanette O'Donnell Auman, BS; Margaret Crawford, BS; Jenna Gabrio, BS MPH; Jamie E. Newman, PhD MPH; Lindsay Parlberg, BS; Carolyn M. Petrie Huitema, MS; Kristin M. Zaterka-Baxter, RN BSN.

Stanford University, El Camino Hospital, and Lucile Packard Children's Hospital (UG1 HD27880, UL1 TR93) – David K. Stevenson, MD; Susan R. Hintz, MD MS Epi; M. Bethany Ball, BS CCRC; Valerie Chock, MD MS Epi; Marian M. Adams, MD; Dona Bahmani, CRC; Barbara Bentley, PsychD MEd; Maria Elena DeAnda, PhD; Anne M. DeBattista, RN PNP PhD; Beth Earhart, PhD; Lynne C. Huffman, MD; Casey E. Krueger, PhD; Ryan E. Lucash, PhD; Melinda S. Proud, RCP; Elizabeth N. Reichert, MA CCRC; Heather Taylor, PhD; Hali E. Weiss, MD; R. Jordan Williams, MD.

University of Alabama at Birmingham Health System and Children's Hospital of Alabama (UG1 HD34216) – Waldemar A. Carlo, MD; Namasivayam Ambalavanan, MD; Myriam Peralta-Carcelen, MD MPH; Kirstin J. Bailey, PhD; Fred J. Biasini, PhD; Stephanie A. Chopko, PhD; Monica V. Collins, RN BSN MaEd; Shirley S. Cosby, RN BSN; Kristy A. Domnanovich, PhD; Chantel J. Jno-Finn, PT DPT; Morissa Ladinsky, MD; Mary Beth Moses, PT MS PCS; Tara E. McNair, RN BSN; Vivien A. Phillips, RN BSN; Julie Preskitt, MSOT MPH; Richard V. Rector, PhD; Kimberlly Stringer, MD MPH; Sally Whitley, MA OTR-L FAOTA; Sheree York Chapman, PT DPT PCS.

University of California - Los Angeles, Mattel Children's Hospital, Santa Monica Hospital, Los Robles Hospital and Medical Center, and Olive View Medical Center (UG1 HD68270) – Uday Devaskar, MD; Meena Garg, MD; Isabell B. Purdy, PhD CPNP; Teresa Chanlaw, MPH; Rachel Geller, RN BSN.

University of Iowa, Mercy Medical Center, and Sanford Health (UG1 HD53109, UL1 TR442) – Edward F. Bell, MD; Tarah T. Colaizy, MD MPH; Jane E. Brumbaugh, MD; Heidi M. Harmon, MD; Karen J. Johnson, RN BSN; Mendi L. Schmelzel, RN MSN; Jacky R. Walker, RN; Claire A. Goeke, RN; Diane L. Eastman, RN CPNP MA; Michelle L. Baack, MD; Laurie A. Hogden, MD; Megan M. Henning, RN; Chelsey Elenkwich, BSN RN; Megan Broadbent, RN BSN; Sarah Van Muyden, RN BSN; Dan L. Ellsbury, MD; Tracy L. Tud, RN.

University of New Mexico Health Sciences Center (UG1 HD53089, UL1 TR41) – Janell Fuller, MD; Robin K. Ohls, MD; Conra Backstrom Lacy, RN; Carol Hartenberger, BSN MPH; Sandra Sundquist Beauman, MSN RNC; Mary Ruffner Hanson, RN BSN; Jean R. Lowe, PhD; Elizabeth Kuan, RN BSN.

University of Pennsylvania, Hospital of the University of Pennsylvania, Pennsylvania Hospital, Children's Hospital of Philadelphia, and Virtua Voorhees Hospital (UG1 HD68244) – Eric C. Eichenwald, MD; Barbara Schmidt, MD MSc; Haresh Kirpalani, MB MSc; Soraya Abbasi, MD; Aasma S. Chaudhary, BS RRT; Toni Mancini, RN BSN CCRC; Dara M. Cucinotta, RN; Judy C. Bernbaum, MD; Marsha Gerdes, PhD; Sarvin Ghavam, MD; Hallam Hurt, MD; Jonathan Snyder, RN BSN, Kristina Ziolkowski, CMA(AAMA) CCRP.

University of Rochester Medical Center, Golisano Children's Hospital, and the University of Buffalo Women's and Children's Hospital of Buffalo (UG1 HD68263, UL1 TR42) – Carl T. D'Angio, MD; Ronnie Guillet, MD PhD; Gary J. Myers, MD; Anne Marie Reynolds, MD; Holly I.M. Wadkins; Michael G. Sacilowski, BS; Rosemary L. Jensen; Joan Merzbach, LMSW; William Zorn, PhD; Osman Farooq, MD; Stephanie Guilford, BS; Kelley Yost, PhD; Mary Rowan, RN; Diane Prinzing; Ann Marie Scorsone, MS CCRC; Michelle Hartley-McAndrew, MD; Kyle Binion, BS; Constance Orme; Premini Sabaratnam, MPH; Alison Kent, BMBS FRACP MD; Rachel Jones; Elizabeth Boylin, BA; Daisy Rochez, BS MHA; Emily Li, BA; Jennifer Kachelmeyer, BS; Kimberly G. McKee, BS; Kelly R. Coleman, PsyD; Deanna Maffett, RN.

University of Texas Southwestern Medical Center, Parkland Health & Hospital System, and Children's Medical Center Dallas (UG1 HD40689) – Luc P. Brion, MD; Roy J. Heyne, MD; Diana M. Vasil, MSN BSN RNC-NIC; Maria M. De Leon, RN BSN; Frances Eubanks, RN BSN; E. Rebecca McDougald, MSN APRN CPNP-PC/AC; Lara Pavageau, MD; Polliciana Sepulveda, RN; Alicia Guzman; Elizabeth Heyne, PsyD PA-C; Lizette E. Lee, RN; Azucena Vera, AS; Jillian Waterbury, DNP RN CPNP-PC; Cathy Twell Boatman, MS CIMI; Kristine Tolentino-Plata, MS.

University of Utah Medical Center, Intermountain Medical Center, McKay-Dee Hospital, Utah Valley Hospital, and Primary Children's Medical Center (UG1 HD87226, UL1 TR105) – Bradley A. Yoder, MD; Mariana Baserga, MD MSCI; Roger G. Faix, MD; Sarah Winter, MD; Stephen D. Minton, MD; Mark J. Sheffield, MD; Carrie A. Rau, RN BSN CCRC; Shawna Baker, RN; Jill Burnett, RNC BSN; Susan Christensen, RN; Sean D. Cunningham, PhD; Brandy Davis, RN BSN; Jennifer O. Elmont, RN BSN; Becky Hall, APRN; Erika R. Jensen, APRN; Jamie Jordan, RN BSN; Manndi C. Loertscher, BS CCRP; Trisha Marchant, RNC BSN; Earl Maxson, RN CCRN; Kandace M. McGrath, BS; Hena G. Mickelsen, BA; Galina Morshedzadeh, BSN APRN; D. Melody Parry, RN BSN; Susan T. Schaefer, RN BSN RRT; Kelly Stout, PhD; Ashley L. Stuart, PhD; Katherine Tice, RN BSN; Kimberlee Weaver-Lewis, RN MS; Kathryn D. Woodbury, RN BSN.

Wayne State University, Hutzel Women's Hospital, and Children's Hospital of Michigan (UG1 HD21385) – Seetha Shankaran, MD; Girija Natarajan, MD; Beena G. Sood, MD MS; Athina Pappas, MD; Monika Bajaj, MD; Melissa February, MD; Prashant Agarwal, MD; Sanjay Chawla, MD; Rebecca Bara, RN BSN; Kirsten Childs, RN BSN; Eunice Hinz Woldt, RN MSN; Laura Goldston, MA, John Barks, MD; Stephanie A. Wiggins, MS; Mary K. Christensen, BA RRT; Martha Carlson, MD; Diane F. White, RRT CCRP.

#### Funding:

The Clinical Centers of the NRN receive funding from the NIH (5UG1HD068244-12 to SBD).

#### Data Sharing:

Data reported in this paper may be requested through a data use agreement. Further details are available at <https://neonatal.rti.org/index.cfm?fuseaction=DataRequest.Home>.

#### References

1. Ishii N, Kono Y, Yonemoto N, et al. Outcomes of Infants Born at 22 and 23 Weeks' Gestation. *Pediatrics* 2013;132:62–71. doi:10.1542/peds.2012-2857 [PubMed: 23733804]
2. Adams-Chapman I, Heyne RJ, DeMauro SB, et al. Neurodevelopmental Impairment Among Extremely Preterm Infants in the Neonatal Research Network. *Pediatrics* 2018;141:e20173091. doi:10.1542/peds.2017-3091 [PubMed: 29666163]
3. Hintz SR, Kendrick DE, Vohr BR, et al. Changes in neurodevelopmental outcomes at 18 to 22 months' corrected age among infants of less than 25 weeks' gestational age born in 1993–1999. *Pediatrics* 2005;115:1645–51. doi:10.1542/peds.2004-2215 [PubMed: 15930228]
4. Hollebrandse NL, Spittle AJ, Burnett AC, et al. School-age outcomes following intraventricular haemorrhage in infants born extremely preterm. *Arch Dis Child - Fetal Neonatal Ed* 2021;106:4–8. doi:10.1136/archdischild-2020-318989 [PubMed: 32732377]
5. Lea CL, Smith-Collins A, Luyt K. Protecting the premature brain: current evidence-based strategies for minimising perinatal brain injury in preterm infants. *Arch Dis Child - Fetal Neonatal Ed* 2017;102:F176–82. doi:10.1136/archdischild-2016-311949 [PubMed: 28011793]
6. Mercer JS, Erickson-Owens DA, Vohr BR, et al. Effects of Placental Transfusion on Neonatal and 18 Month Outcomes in Preterm Infants: A Randomized Controlled Trial. *J Pediatr* 2016;168:50–55.e1. doi:10.1016/j.jpeds.2015.09.068 [PubMed: 26547399]
7. Rabe H, Sawyer A, Amess P, et al. Neurodevelopmental Outcomes at 2 and 3.5 Years for Very Preterm Babies Enrolled in a Randomized Trial of Milking the Umbilical Cord versus Delayed Cord Clamping. *Neonatology* 2016;109:113–9. doi:10.1159/000441891 [PubMed: 26650133]
8. Katheria A, Garey D, Truong G, et al. A Randomized Clinical Trial of Umbilical Cord Milking vs Delayed Cord Clamping in Preterm Infants: Neurodevelopmental Outcomes at 22–26 Months of Corrected Age. *J Pediatr* 2018;194:76–80. doi:10.1016/j.jpeds.2017.10.037 [PubMed: 29246467]

9. Robledo KP, Tarnow-Mordi WO, Rieger I, et al. Effects of delayed versus immediate umbilical cord clamping in reducing death or major disability at 2 years corrected age among very preterm infants (APTS): a multicentre, randomised clinical trial. *Lancet Child Adolesc Health* 2021;;S2352464221003734. doi:10.1016/S2352-4642(21)00373-4
10. Wyckoff MH, Singletary EM, Soar J, et al. 2021 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations: Summary From the Basic Life Support; Advanced Life Support; Neonatal Life Support; Education, Implementation, and Teams; First Aid Task Forces; and the COVID-19 Working Group. *Circulation* 2022;145. doi:10.1161/CIR.0000000000001017 [PubMed: 35861771]
11. Katheria A, Reister F, Essers J, et al. Association of Umbilical Cord Milking vs Delayed Umbilical Cord Clamping With Death or Severe Intraventricular Hemorrhage Among Preterm Infants. *JAMA* 2019;322:1877. doi:10.1001/jama.2019.16004 [PubMed: 31742630]
12. Kumbhat N, Eggleston B, Davis AS, et al. Umbilical Cord Milking vs Delayed Cord Clamping and Associations with In-Hospital Outcomes among Extremely Premature Infants. *J Pediatr* 2021;;S0022347620315821. doi:10.1016/j.jpeds.2020.12.072
13. Kumbhat N, Eggleston B, Davis AS, et al. Placental transfusion and short-term outcomes among extremely preterm infants. *Arch Dis Child - Fetal Neonatal Ed* 2020;;fetalneonatal-2019-318710. doi:10.1136/archdischild-2019-318710
14. Rysavy MA, Li L, Bell EF, et al. Between-Hospital Variation in Treatment and Outcomes in Extremely Preterm Infants. *N Engl J Med* 2015;372:1801-11. doi:10.1056/NEJMoa1410689 [PubMed: 25946279]
15. Newman JE, Bann CM, Vohr BR, et al. Improving the Neonatal Research Network Annual Certification for Neurologic Examination of the 18-22 month Child. *J Pediatr* 2012;161:1041-1046.e2. doi:10.1016/j.jpeds.2012.05.048 [PubMed: 22748517]
16. Albers CA, Grieve AJ. Test Review: Bayley, N. (2006). *Bayley Scales of Infant and Toddler Development- Third Edition*. San Antonio, TX: Harcourt Assessment. *J Psychoeduc Assess* 2007;25:180-90. doi:10.1177/0734282906297199
17. Palisano RJ, Avery L, Gorter JW, et al. Stability of the Gross Motor Function Classification System, Manual Ability Classification System, and Communication Function Classification System. *Dev Med Child Neurol* 2018;60:1026-32. doi:10.1111/dmcn.13903 [PubMed: 29726578]
18. Bell EF, Hintz SR, Hansen NI, et al. Mortality, In-Hospital Morbidity, Care Practices, and 2-Year Outcomes for Extremely Preterm Infants in the US, 2013-2018. 2022;;29.
19. Givens M, Teal EN, Patel V, et al. Preterm birth among pregnant women living in areas with high social vulnerability. *Am J Obstet Gynecol MFM* 2021;3:100414. doi:10.1016/j.ajogmf.2021.100414 [PubMed: 34082172]
20. Imai K, Keele L, Tingley D. A general approach to causal mediation analysis. *Psychol Methods* 2010;15:309-34. doi:10.1037/a0020761 [PubMed: 20954780]
21. Tarnow-Mordi W, Morris J, Kirby A, et al. Delayed versus Immediate Cord Clamping in Preterm Infants. *N Engl J Med* 2017;377:2445-55. doi:10.1056/NEJMoa1711281 [PubMed: 29081267]
22. Fogarty M, Osborn DA, Askie L, et al. Delayed vs early umbilical cord clamping for preterm infants: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2018;218:1-18. doi:10.1016/j.ajog.2017.10.231 [PubMed: 29097178]
23. Seidler AL, Gyte GML, Rabe H, et al. Umbilical Cord Management for Newborns <34 Weeks' Gestation: A Meta-analysis. *Pediatrics* 2021;147. doi:10.1542/peds.2020-0576
24. Perlman JM, Wyllie J, Kattwinkel J, et al. Part 7: Neonatal Resuscitation: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations (Reprint). *PEDIATRICS* 2015;136:S120-66. doi:10.1542/peds.2015-3373D [PubMed: 26471381]

### Key Points

#### What is already known on this topic


Comparisons of short-term in-hospital outcomes following exposure to different forms of umbilical cord management in extremely preterm infants have shown delayed cord clamping to be beneficial compared to immediate cord clamping or umbilical cord milking. Published data about neurodevelopmental outcomes among extremely preterm infants following exposure to different forms of cord management are limited.

#### What this study adds

Compared to immediate cord clamping, delayed cord clamping may be beneficial for outcomes at 22 to 26 months' corrected age.

#### How this study might affect research, practice, or policy

This retrospective study, which captures clinical care and outcomes outside the context of a randomized clinical trial, provides further evidence supporting delayed cord clamping as a beneficial practice and hospital policy for extremely preterm infants.



**Figure 1.** Patient identification flowchart. \*Infants seen but outside of the 18–30-month CA window. †Infants seen and examination window closed on or before 31/12/2020. DCC, delayed cord clamping; UCM, umbilical cord milking.

**Table 1.**

## Sociodemographic and perinatal-neonatal characteristics

Characteristics	Exposure (%)			p value <sup>+</sup>		
	ICC (N=1218)	DCC (N=528)	UCM (N=154)	DCC vs. ICC	UCM vs. ICC	DCC vs. UCM
<b>Neonatal characteristics</b>						
Gestational age in weeks [mean weeks, (SD)]	24.9 (1.2)	25.1 (1.2)	25.0 (1.2)	0.005	0.97	0.10
Birth weight in grams [mean grams, (SD)]	702 (171)	732(173)	722 (159)	0.001	0.10	0.67
Small for gestational age	9.7%	6.8%	7.1%	0.05	0.38	0.86
Male sex	51.7%	51.0%	52.6%	0.79	0.86	0.78
Multiple births	26.0%	24.6%	30.5%	0.55	0.25	0.14
Age at follow up assessment [mean months, (SD)]	24.5 (2.1)	24.3 (2.2)	25.2 (2.1)	0.05	0.0004	< 0.0001
<b>Sociodemographic characteristics</b>						
Race/Ethnicity				0.55	< 0.0001	< 0.0001
Asian	2.6%	1.9%	7.1%			
Black, non-Hispanic	42.6%	44.3%	8.4%			
Hispanic	20.2%	10.8%	14.9%			
Other	2.0%	3.6%	4.6%			
White, non-Hispanic	32.2%	39.0%	64.3%			
Unknown/Not reported	0.5%	0.4%	0.7%			
Maternal Insurance				0.001	<0.0001	<0.0001
Private	36.5%	47.8%	59.7%			
Public	60.2%	50.7%	33.1%			
Other	3.4%	1.5%	7.1%			
Maternal education				0.04	0.96	0.23
Less than high school	15.2%	10.2%	6.5%			
High school degree	25.5%	23.3%	17.5%			
Partial college/trade/tech	24.5%	31.8%	28.6%			
College degree or higher	18.1%	22.7%	36.4%			
Missing	16.8%	11.9%	11.0%			
Multilingual at home	1.5%	2.9%	0%	< 0.0001	0.52	0.01
Missing due to death	34.4%	22.4%	30.7%			
Under state supervision	2.6%	3.4%	2.0%	< 0.0001	0.45	0.03
Missing due to death	34.4%	22.4%	30.7%			
<b>Perinatal characteristics</b>						
Limited or no prenatal care	14.5%	11.2%	7.8%	0.07	0.02	0.29
Diabetes prior to pregnancy	5.1%	2.7%	3.3%	0.02	0.43	0.78
Gestational diabetes	3.6%	2.9%	3.3%	0.47	1.00	0.79



Characteristics	Exposure (%)			p value <sup>+</sup>		
	ICC (N=1218)	DCC (N=528)	UCM (N=154)	DCC vs. ICC	UCM vs. ICC	DCC vs. UCM
Hypertension during pregnancy	26.9%	25.2%	21.4%	0.48	0.17	0.39
PIH	9.8%	12.7%	9.1%	0.08	0.89	0.26
Antenatal steroids	88.2%	96.6%	98.7%	< 0.0001	< 0.0001	0.28
Antenatal MgSO <sub>4</sub>	77.7%	88.8%	94.2%	< 0.0001	< 0.0001	0.07
<b>Neonatal delivery characteristics</b>						
1-minute Apgar score 4	75.5%	62.5%	67.3%	<0.0001	0.04	0.29
5-minute Apgar score 4	33.6%	23.3%	29.4%	<0.0001	0.32	0.14
Delivery room interventions						
PPV	91.8%	87.0%	97.3%	0.003	0.01	0.0001
Intubation	77.4%	70.4%	91.3%	0.003	< 0.0001	< 0.0001
Chest Compressions	5.7%	3.7%	4.7%	0.11	0.85	0.63
Epinephrine	2.8%	1.9%	3.4%	0.40	0.60	0.35
<b>Among survivors beyond 12 hours after birth</b>						
Severe brain injury	28.4%	24.0%	34.7%	0.08	0.12	0.01
Severe IVH <sup>a</sup>	22.9%	19.8%	29.9%	0.19	0.06	0.01
Cystic PVL	6.5%	5.3%	7.5%	0.37	0.60	0.31
Porencephalic cyst	2.1%	1.9%	0.7%	1.00	0.35	0.47
Ventriculomegaly	11.7%	10.3%	12.1%	0.45	0.89	0.55
Late onset sepsis	27.1%	24.7%	28.2%	0.30	0.77	0.39
Necrotizing enterocolitis	11.6%	12.2%	11.4%	0.74	1.00	0.89
Severe ROP <sup>b</sup>	17.3%	16.3%	14.0%	0.69	0.43	0.66
<b>Among survivors at 36 weeks PMA</b>						
Severe BPD <sup>c</sup>	11.6%	13.3%	11.9%	0.40	0.87	0.87

Abbreviations: BPD = bronchopulmonary dysplasia, ICC = Immediate cord clamping, DCC = Delayed cord clamping, UCM = Umbilical cord milking, PIH = Pregnancy induced hypertension, IVH = Intraventricular hemorrhage, PVL = Periventricular leukomalacia, PPV = Positive pressure ventilation, PMA = post menstrual age, ROP = retinopathy of prematurity.

<sup>a</sup>Severe IVH - grade III or IV

<sup>b</sup>Severe ROP- undergone ophthalmologic intervention for ROP or having retinal detachment

<sup>c</sup>Grade 3 BPD – requiring mechanical ventilation at 36 weeks corrected age

<sup>+</sup>p-values based on t-test/Wilcoxon rank sum test for continuous variables and Fisher's exact test for categorical variables. Data presented as n (%) for categorical variables and median (SD) for continuous variables.

**Table 2.**

Associations between cord management practices and death or severe NDI

	Death or severe NDI (%) <sup>a</sup>		Adjusted OR (95% CI)
	DCC (N=528)	ICC (N=1218)	
<b>DCC vs. ICC</b>	36.3%	50.3%	0.64 (0.50, 0.83)
	UCM (N=154)	ICC (N=1218)	
	<b>UCM vs. ICC</b>	46.3%	
	DCC (N=528)	UCM (N=152)	
	<b>DCC vs. UCM</b>	36.4%	

Abbreviations: CI =confidence interval, DCC = Delayed cord clamping, ICC = Immediate cord clamping, NDI = Neurodevelopmental impairment, OR = Odds Ratio, UCM = Umbilical cord milking.

<sup>a</sup>The N indicates the total number of infants in the cohort exposed to the umbilical cord management practice listed. The fractional N reports the number of infants with death or severe NDI over the number of infants in the cohort in whom the primary outcome could be assessed.

Variables in the model include a priori variables associated with death or NDI (gestational age, race/ethnicity, maternal education, and antenatal steroids), those that differed in the bivariate comparison of DCC vs ICC and occurred before the exposure (insurance, limited or no prenatal care, diabetes prior to pregnancy, pregnancy induced hypertension and antenatal magnesium), sex, SGA, and birth year, with NRN center as a random effect.

Table 3.

Associations of cord management practices with secondary outcomes

Characteristics	Exposure N (%)			DCC/ICC aOR (95% CI)	UCM/ICC aOR (95% CI)	DCC/UCM aOR (95% CI)
	ICC (%)	DCC N (%)	UCM (%)			
Mortality <sup>a</sup>	418/1218 (34.3%)	118/528 (22.4%)	47/154 (30.5%)	0.61 (0.46, 0.81)	0.66 (0.39, 1.11)	0.77 (0.42, 1.42)
Severe NDI <sup>b</sup>	175/761 (23.0%)	64/383 (16.7%)	21/100 (21.0%)	0.74 (0.52, 1.05)	1.12 (0.62, 2.03)	0.72 (0.33, 1.55)
Moderate or severe CP <sup>c</sup>	97/796 (12.2%)	43/406 (10.6%)	9/106 (8.5%)	0.85 (0.58, 1.25) <sup>d</sup>	0.67 (0.33, 1.37) <sup>d</sup>	1.28 (0.60, 2.71)
CP classification				1.01 (0.73, 1.40)	1.02 (0.59, 1.76)	0.91 (0.41, 2.02)
None	608/800 (76.0%)	317/406 (78.1%)	81/106 (76.4%)			
Mild	115/800 (14.4%)	52/406 (12.8%)	17/106 (16.0%)			
Moderate	45/800 (5.6%)	24/406 (5.9%)	6/106 (5.7%)			
Severe	32/800 (4.0%)	13/406 (3.2%)	2/106 (1.9%)			
Cognitive Bayley-3 score <70	156/778 (20.1%)	57/398 (14.3%)	20/104 (19.2%)	0.72 (0.50, 1.04)	1.23 (0.65, 2.31)	0.62 (0.28, 1.36)
Cognitive Bayley-3 score <85	391/778 (50.3%)	164/398 (41.2%)	44/104 (42.3%)	0.88 (0.68, 1.18)	1.08 (0.62, 1.91)	0.94 (0.50, 1.75)
Motor Bayley-3 score <70	165/764 (21.6%)	67/384 (17.5%)	19/102 (18.6%)	0.76 (0.53, 1.09)	0.85 (0.45, 1.62)	0.94 (0.41, 2.16)
Moderate to severe language delay (Language Bayley-3 <70)	257/770 (33.4%)	111/386 (28.8%)	28/103 (27.2%)	0.92 (0.67, 1.25)	1.03 (0.60, 1.77)	1.03 (0.50, 2.10)
Bilateral blindness	11/800 (1.4%)	5/408 (1.2%)	1/106 (0.9%)	0.89 (0.31, 2.58) <sup>d</sup>	0.68 (0.09, 5.34) <sup>d</sup>	1.30 (0.15, 11.27) <sup>d</sup>
Hearing impairment	24/778 (3.1%)	11/390 (2.8%)	2/103 (1.9%)	0.91 (0.44, 1.88) <sup>d</sup>	0.62 (0.14, 2.67) <sup>d</sup>	1.47 (0.32, 6.72) <sup>d</sup>

Abbreviations: aOR=adjusted odds ratio, Bayley-3 = Bayley Scales of Infant Development, CP = Cerebral palsy, DCC = Delayed cord clamping, ICC = Immediate cord clamping, NDI = Neurodevelopmental impairment, UCM = Umbilical cord milking.

<sup>a</sup>Mortality before assessment at 22–26 months' corrected age.

<sup>b</sup>Severe NDI-severe cerebral palsy (CP) [Gross Motor Function Classification System (GMFCS) >=4], cognitive BSID III <70, motor BSID scores <70, bilateral blindness despite corrective lens, or bilateral hearing impairment with or without amplification

<sup>c</sup>Moderate or severe CP-GMFCS >=2

<sup>d</sup>Model did not converge, was nearly unidentifiable, or variance-covariance matrix was not positive-definite, so raw Odds Ratios are report.

Variables in the model include a priori variables associated with death or NDI (gestational age, race/ethnicity, maternal education, and antenatal steroids), those that differed in the bivariate comparison of DCC vs ICC and occurred before the exposure (insurance, multilingual at home, limited or no prenatal care, diabetes prior to pregnancy, pregnancy induced hypertension and antenatal magnesium) and age at follow-up assessment and birth year, with NRN center as a random effect. Age at follow up assessment and multilingual at home were not included in the model assessing mortality