

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### Title

PRODUCTION OF THE NEUTRON-RICH NUCLIDES  $^{20}\text{C}$  AND  $^{27}\text{F}$  BY FRAGMENTATION OF 213 MeV/NUCLEON  $^{48}\text{Ca}$

### Permalink

<https://escholarship.org/uc/item/5bd0v4xw>

### Author

Stevenson, J.D.

### Publication Date

1980-12-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Submitted to Physical Review Letters

PRODUCTION OF THE NEUTRON-RICH NUCLIDES  $^{20}\text{C}$  AND  $^{27}\text{F}$   
BY FRAGMENTATION OF 213 MeV/NUCLEON  $^{48}\text{Ca}$

J.D. Stevenson and P.B. Price

December 1980

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

FEB 17 1981

LIBRARY AND  
DOCUMENTS SECTION

## TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 6782.*

LBL-11737 c. 2

## DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Production of the Neutron-Rich Nuclides  $^{20}\text{C}$  and  $^{27}\text{F}$   
by Fragmentation of 213 MeV/nucleon  $^{48}\text{Ca}$

J.D. Stevenson<sup>†</sup> and P.B. Price<sup>†</sup>

Lawrence Berkeley Laboratory  
and  
Department of Physics  
University of California  
Berkeley, California 94720

Abstract

Production cross sections of neutron-rich projectile fragments have been measured for the reaction 213 MeV/nucleon  $^{48}\text{Ca} + \text{Be}$ . The neutron-rich nuclides  $^{20}\text{C}$  and  $^{27}\text{F}$  have been observed for the first time. A method to strengthen our evidence for the possible particle-stability of  $^{21}\text{C}$ ,  $^{23}\text{N}$ , and  $^{25}\text{O}$  is presented.

<sup>†</sup>Also Space Sciences Laboratory

Of the more than 7000 nuclides predicted to be particle-stable, only about one-fourth have ever been observed. The actual limits of particle-stability, the so-called neutron and proton driplines, are known only up to beryllium for neutron-rich nuclides and up to sodium for proton-rich nuclides. A more complete knowledge of the location of the driplines would have widespread implications for both nuclear physics and nuclear astrophysics.

Among the methods used to predict the masses of nuclei far from the valley of stability are the droplet model;<sup>1-3</sup> the shell model;<sup>4,5</sup> self-consistent calculations based on the energy density concept;<sup>6</sup> and the Garvey-Kelson mass relations.<sup>7-10</sup> These models contain parameters chosen to fit known masses of nuclei near the valley of stability. Predictions of the limits of particle-stability of nuclides require considerable extrapolations from the fitted data and therefore provide a good test of the global validity of these models.

Techniques to produce extremely neutron-rich nuclides include proton-induced fragmentation of heavy nuclei,<sup>11</sup> deep inelastic heavy-ion reactions,<sup>12</sup> and fragmentation of relativistic heavy ions.<sup>13,14</sup>

In this Letter we report results of a search for new neutron-rich nuclides produced by fragmenting 213 MeV/nucleon  $^{48}\text{Ca}$  nuclei. These results include the discovery of the particle-stability of  $^{20}\text{C}$  and  $^{27}\text{F}$ , confirmation of a previous report of the particle-stability of  $^{22}\text{N}$  and  $^{26}\text{F}$  (ref. 14), and evidence for the possible particle-stability of  $^{21}\text{C}$ ,  $^{23}\text{N}$ , and  $^{25}\text{O}$ .

The process of relativistic heavy-ion projectile fragmentation has been studied extensively.<sup>15,16</sup> Projectile fragments viewed in

the projectile frame have Gaussian transverse and longitudinal momentum distributions,<sup>15</sup> with momentum widths  $\sigma_{p_{\parallel}} \approx \sigma_{p_{\perp}} \approx 200$  MeV/c. In the laboratory frame the fragments are "focused" within  $\sim 1^\circ$  of the beam direction and have a momentum-per-nucleon within a few percent of that of the beam. By passing this beam of fragments through a spectrometer, neutron-rich nuclei are cleanly separated from stability-line isotopes which are  $\sim 10^6$  times more abundant.

The experimental techniques used in this experiment are described in detail in ref. 17. The zero-degree magnetic spectrometer at the Lawrence Berkeley Laboratory Bevalac was used to focus neutron-rich projectile fragments on a stack of Lexan plastic track detectors<sup>18</sup> thick enough to stop fragments with  $Z > 5$ . Later the detectors were chemically etched and the sizes and positions of the etch pits at the ends of range of the fragments that stopped in the stack were measured with a microscope. The range of each fragment, its deflection in the spectrometer, and its charge at the end of its range were determined from the etch pit measurements. The charge resolution was  $\sigma_Z \approx 0.2$ . Absolute charge assignments were based on calibrations of the plastic detectors with low-energy  $^{12}\text{C}$  and  $^{16}\text{O}$  ions. The energy and rigidity of the beam were determined precisely by measuring its range and deflection in a stack of plastic detectors. Rigidities of the fragments were found by comparing their deflections to that of the beam. The charge, range and rigidity measurements allow the mass of each particle to be calculated. The beam intensity of  $^{48}\text{Ca}$  ions,  $\sim 10^7$  ions/sec, was monitored with a scintillator telescope that counted fragments from the target. The monitor was calibrated at

intensities low enough to count. The target was 0.89 g/cm<sup>2</sup> beryllium.

Figure 1 shows data accumulated over a 40 hour run, for carbon, nitrogen, oxygen, and fluorine nuclei. The mass resolution ranges from  $\sigma_m \approx 0.23$  for carbon to  $\sigma_m \approx 0.33$  for fluorine. The ratios of peak heights do not directly reflect the relative yields of isotopes.

There is clear evidence for the first observation of <sup>20</sup>C (~40 counts) and <sup>27</sup>F (~20 counts). We also confirm the particle-stability of <sup>22</sup>N and <sup>26</sup>F, which were recently reported.<sup>14</sup> In addition, at <sup>21</sup>C, <sup>23</sup>N, and <sup>25</sup>O there are bumps consisting of about 5 counts clearly outside the Gaussian envelopes of the adjacent lighter isotopes. Although the number of counts is too small to convincingly establish their existence, it should be possible with small improvements in sensitivity to obtain definitive evidence.

To determine fragmentation cross sections we calculated the transmission through the spectrometer assuming that the fragment momentum distributions viewed in the projectile frame were Gaussian with  $\sigma$  given by

$$\sigma = 94[M_f(M_p - M_f)/(M_p - 1)]^{1/2} \text{ MeV/c} \quad (1)$$

where  $M_f$  and  $M_p$  are the fragment and projectile masses in amu. This equation was shown<sup>16</sup> to fit the measured momentum widths for the reaction 213 MeV/nucleon <sup>40</sup>Ar + C → projectile fragments. Figure 2 compares our calculated fragmentation cross sections with those of Westfall et al.<sup>14</sup> for exactly the same reaction, 213 MeV/nucleon <sup>48</sup>Ca + Be. The absolute normalization of our results is somewhat uncertain but is in reasonable agreement with the work of Westfall

et al. Because our Lexan detector stack was much larger than the detector telescope used by Westfall et al., we were able to survey a wider rigidity interval and detect nuclides more neutron-rich and with lower production cross sections despite the fact that our total fluence of  $^{48}\text{Ca}$  ions was smaller than theirs.

Table 1 gives the mass excess calculated by several methods for nuclides in the vicinity of the neutron dripline for C, N, O, and F. Single and double asterisks signify nuclides calculated to be unstable against single and double neutron emission respectively. Although the nuclides  $^{20}\text{C}$  and  $^{27}\text{F}$  that we have detected are calculated to be particle-stable by each of the models, it is interesting to note that  $^{21}\text{C}$  and  $^{25}\text{O}$ , for whose existence we have some evidence, are calculated to be unstable against neutron emission by some of the models and particle-stable by others.  $^{24}\text{N}$  and  $^{28}\text{F}$  are also predicted by several models to be particle-unstable.

Judging from the trends in Fig. 2, an experiment capable of discriminating among these models will require about 50 times the present sensitivity, using the 213 MeV/nucleon  $^{48}\text{Ca} + \text{Be}$  reaction. One possible way of achieving this is to use considerably thicker targets. Figure 3 shows the expected yield of carbon isotopes as a function of target thickness, calculated on the assumption that the partial cross section for isotopes of a given element has a Gaussian dependence on fragment mass about a most probable mass, with a mass width taken to be a constant  $\sigma_A = 1.5$  amu. This prescription, together with appropriate normalization factors and a geometrical total cross section, gives reasonable agreement with the production



cross sections of neutron-rich nuclei found by Westfall et al.<sup>14</sup> and in our Fig. 2.

The arrow in Fig. 3 indicates the target thickness for our experiment and that of Westfall et al. The dashed horizontal line assumes present beam intensities and only 10 events. The predicted yield increases somewhat faster than linearly with target thickness because of multiple interactions in the target. One can see that, with current Bevalac beam intensities, experiments with very thick (~1 interaction length) targets may well have the sensitivity to test the limits of particle stability. Improvements in the Bevalac will soon provide higher beam intensities and ions up to Pb or U. Thus, fragmentation of relativistic heavy ions promises to be a powerful tool in the study of nuclei far from stability.

This work was supported by the U.S. Department of Energy Contract W-7405-ENG-48.

## References

1. W.D. Myers, At. Data Nucl. Data Tables 17, 411 (1976).
2. H.v. Groote, E.R. Hilf, and K. Takahashi, At. Data Nucl. Data Tables 17, 418 (1976).
3. P.A. Seeger and W.M. Howard, At. Data Nucl. Data Tables 17, 428 (1976).
4. S. Liran and N. Zeldes, At. Data Nucl. Data Tables 17, 431 (1976).
5. M. Bauer, At. Data Nucl. Data Tables 17, 442 (1976).
6. M. Beiner, R.J. Lombard, and D. Mas, At. Data Nucl. Data Tables 17, 450 (1976).
7. J. Jänecke, At. Data Nucl. Data Tables 17, 455 (1976).
8. E. Comay and I. Kelson, At. Data Nucl. Data Tables 17, 463 (1976).
9. C. Thibault and R. Klapisch, Phys. Rev. C9, 793 (1974).
10. N.A. Jelley, J. Cerny, D.P. Stahel, and K.H. Wilcox, Phys. Rev. C11, 2049 (1975).
11. C. Detraz, D. Guillemaud, G. Huber, R. Klapisch, M. Langevin, F. Naulin, C. Thibault, L.C. Carraz, and F. Touchard, Phys. Rev. C19, 164 (1979).
12. P. Auger, T.H. Chiang, J. Galin, B. Gatty, D. Guerreau, E. Nolte, J. Pouthas, X. Tarrago, and J. Girard, Z. Phys. A 289, 255 (1979).
13. T.J.M. Symons, Y.P. Viyogi, G.D. Westfall, P. Doll, D.E. Greiner, H. Faraggi, P.J. Lindstrom, D.K. Scott, H.J. Crawford, and C. McParland, Phys. Rev. Lett. 42, 40 (1979).
14. G.D. Westfall, T.J.M. Symons, D.E. Greiner, H.H. Heckman, P.J. Lindstrom, J. Mahoney, A.C. Shotter, D.K. Scott, H.J. Crawford, C. McParland, T.C. Awes, C.K. Gelbke, and J.M. Kidd, Phys. Rev.

- Lett. 43, 1859 (1979).
15. D.E. Greiner, P.J. Lindstrom, H.H. Heckman, B. Cork, and F.S. Bieser, Phys. Rev. Lett. 35, 152 (1975).
  16. Y.P. Viyogi, T.J.M. Symons, P. Doll, D.E. Greiner, H.H. Heckman, D.L. Hendrie, P.J. Lindstrom, J. Mahoney, D.K. Scott, K. Van Bibber, G.D. Westfall, H. Wieman, H.J. Crawford, C. McParland, and C.K. Gelbke, Phys. Rev. Lett. 42, 33 (1979).
  17. J.D. Stevenson, P.B. Price, and M.P. Budiansky, Nucl. Instr. and Meth. 171, 93 (1980).
  18. R.L. Fleischer, P.B. Price, and R.M. Walker, Nuclear Tracks in Solids (University of California Press, Berkeley, 1975).
  19. G.M. Raisbeck, P. Boerstling, P. Roesenfeldt, T.D. Thomas, R. Klapisch, and G.T. Garvey, in High Energy Physics and Nuclear Structure, edited by S. Devons (Plenum, New York, 1970), pp. 341-345.

Table 1. Mass Excesses for Some Neutron-Rich Nuclides (MeV)

	Particle- stable?	Calculated Mass Excess				
		ref. 2	ref. 4	ref. 6	ref. 8	ref. 9
$^{19}\text{C}$	yes <sup>a</sup>	33.38 <sup>*</sup>		30.0	33.57 <sup>*</sup>	33.55
$^{20}\text{C}$	yes <sup>b</sup>	37.33		33.9	36.95	37.17
$^{21}\text{C}$	yes? <sup>b</sup>	47.34 <sup>*</sup>		41.1	45.22 <sup>*</sup>	46.01 <sup>*</sup>
$^{22}\text{C}$		53.51 <sup>**</sup>		47.4	50.63	51.72
$^{22}\text{N}$	yes <sup>c</sup>	33.08 <sup>*</sup>			30.72	31.54
$^{23}\text{N}$	yes? <sup>b</sup>	38.06		31.7	36.13	37.27
$^{24}\text{N}$		47.76 <sup>*</sup>			45.08 <sup>*</sup>	46.04 <sup>*</sup>
$^{25}\text{N}$		53.87		49.4 <sup>*</sup>	52.81 <sup>**</sup>	53.17
$^{25}\text{O}$	yes? <sup>b</sup>	29.64 <sup>*</sup>	28.99	24.7 <sup>*</sup>	28.18 <sup>*</sup>	28.91 <sup>*</sup>
$^{26}\text{O}$		34.75	33.66	31.6	34.13	33.97
$^{26}\text{F}$	yes <sup>c</sup>	19.15	16.95		18.60	18.84
$^{27}\text{F}$	yes <sup>b</sup>	23.15	20.91	21.2	23.70	23.06
$^{28}\text{F}$		31.54 <sup>*</sup>	28.71		32.58 <sup>*</sup>	31.06

(a) ref. 19; (b) this Letter; (c) ref. 14

### Figure Captions

- Figure 1. Mass histograms of neutron-rich isotopes of C, N, O, and F observed in this experiment. The peak heights do not directly reflect the relative abundances of the isotopes.
- Figure 2. Measured cross sections for production of neutron-rich isotopes in the 213 MeV/nucleon  $^{48}\text{Ca} + \text{Be}$  reaction. Squares are results of ref. 14; crosses are results presented in this letter.
- Figure 3. Calculated yield of neutron-rich carbon ions per  $^{48}\text{Ca}$  ion as a function of target thickness. The calculation assumes a Gaussian mass yield fitted to measured cross sections. The arrow indicates the target thickness for this experiment and that of ref. 14. The dashed line corresponds to a yield of 10 events with present beam intensities and a detector with collecting power comparable to ours.




Figure 1

XBL 809-11912




Figure 2

XBL 809-11911




Figure 3

XBL 809-11913



