

# UC Irvine

## UC Irvine Previously Published Works

### Title

hESC-derived photoreceptors survive and integrate better in immunodeficient retina

### Permalink

<https://escholarship.org/uc/item/5cs477c3>

### Journal

Stem Cell Investigation, 4(8)

### ISSN

2306-9759

### Author

Seiler, Magdalene J

### Publication Date

2017

### DOI

10.21037/sci.2017.08.05

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# hESC-derived photoreceptors survive and integrate better in immunodeficient retina

Magdalene J. Seiler

Department of Physical Medicine & Rehabilitation, Sue & Bill Gross Stem Cell Research Center, University of California, Irvine, USA

*Correspondence to:* Magdalene J. Seiler, PhD. Assistant Professor, Department of Physical Medicine & Rehabilitation, Sue & Bill Gross Stem Cell Research Center, University of California, 845 Health Science Rd, 2035 Gross Hall, Irvine CA 92697-1705, USA. Email: mseiler@uci.edu.

*Provenance:* This is an invited Editorial commissioned by Editor-in-Chief Zhizhuang Joe Zhao (Pathology Graduate Program, University of Oklahoma Health Sciences Center, Oklahoma City, USA).

*Comment on:* Zhu J, Cifuentes H, Reynolds J, *et al.* Immunosuppression via Loss of IL2 $\gamma$  Enhances Long-Term Functional Integration of hESC-Derived Photoreceptors in the Mouse Retina. *Cell Stem Cell* 2017;20:374-84.e5.

Received: 03 August 2017; Accepted: 11 August 2017; Published: 23 August 2017.

doi: 10.21037/sci.2017.08.05

View this article at: <http://dx.doi.org/10.21037/sci.2017.08.05>

Retinal degenerative diseases resulting in photoreceptor death, such as age-related macular degeneration (AMD) and retinitis pigmentosa (RP) lead to incurable vision loss in millions of patients worldwide. When photoreceptors are lost, the resulting visual deficit is permanent. Currently, there are no effective therapies for these diseases except to delay degeneration in early disease stages. A potential approach in regaining vision for more advanced disease is to replace the degenerated photoreceptors. Particularly, this is a viable strategy for AMD as it has been shown that ganglion cells can survive in a degenerate retina even when there is severe underlying photoreceptor loss (1). Finding a suitable source of transplantable cells to replace the dying host tissue is the main challenge.

There are two different strategies to replace photoreceptors: (I) transplantation of retinal progenitor sheets (fetal retina or stem-cell derived) (2-5). Fetal retinal sheet transplants have resulted in long-term visual improvements in different animal models of retinal degeneration (4,6-8) and in patients (9), and have shown to integrate and synaptically connect with a degenerated retina (6,8). (II) The second approach is injection of dissociated photoreceptor precursors (10-15). Studies have shown that a small percentage of subretinally injected photoreceptor precursor cells can integrate in the photoreceptors layer, form synaptic terminals (10,15) and outer segments (14,15). However, this requires the presence of an outer nuclear layer in the host retina; transplanted photoreceptor

progenitors do not develop proper morphology in recipients with severe loss of photoreceptors (10,11,16). Therefore, most studies have been performed in transgenic mutants where the photoreceptors remain viable although non-functional, such as transgenic knockouts of rhodopsin, CRX or rod transducin (10,11,15). In such very specific models, several groups have shown some vision improvement with transplantation of photoreceptor precursors (11,13,15); however most experiments have been short-term (mostly 6, up to 12 weeks). Even when transplanted within the same species, dissociated photoreceptor precursor transplants disappear over time due to a slow rejection process (12). This is not the case with fetal retinal sheet transplants (2).

The study of Zhu *et al.* injected dissociated retinal cells derived from human pluripotent stem cells which has been done previously in few studies (11,17-20). Testing of human cells in animals requires immunosuppression. Although the retina has a relative immune privilege (21) this does not extend to xenografts. In spite of the so-called “immune privilege” of the eye, xenografts require immunosuppression to survive (21,22). In addition, retinal degeneration causes activation of microglia and macrophages. Therefore, this study of Zhu *et al.* developed an immunodeficient mouse (lacking IL2 $\gamma$ ) for transplantation of hESC-derived photoreceptor precursors (23), in a cross with retinal degenerate *Crx*  $-/-$  mice, a model of Leber’s congenital amaurosis (LCA). *Crx*  $-/-$  mice have non-functional photoreceptors which degenerate very slowly over a long

time frame. This means there was still an outer nuclear layer present at the time of transplantation in their study which is different from severe retinal degeneration.

The title of the Zhu *et al.* paper is misleading: the animals were actually not immunosuppressed, but they were genetically manipulated to be immunodeficient and did not receive immunosuppressive drugs. The side effects of immunosuppressive drugs are difficult to balance against any visual benefits, in contrast to the benefits of immunosuppression for organ replacement (24).

The donor tissue in the Zhu *et al.* study contained mixed hESC-derived retinal cells (3 months of differentiation) not purified photoreceptor precursors (25). The cells were labeled with a lentivirus expressing enhanced green fluorescent protein (EGFP) under an ubiquitous promoter that labels all cells (11). This virus labeled 60–70% of all donor cells. GFP label was restricted to cells of human origin as confirmed by co-staining with human nuclear or cytoplasmic markers.

In the first part of the study, the cells were injected into the subretinal space of mice with a normal retina, either wildtype or IL2r-gamma knockout, without immunosuppression. Very few cells integrated into the outer nuclear layer of normal mice, whereas robust integration was observed in IL2r-gamma knockout mice. This integration represented 2.5–4% of the injected cells. In the wildtype retina, transplantation caused an upregulation of lymphocytes, T-cells, dendritic cells and activation of microglia/macrophages. In contrast, host retinas of IL2r-gamma knockout mice contained more CD3+ cells, but CD4 and CD8a expression (T-cell marker) was almost completely eliminated. Also absent were CD49b natural killer cells, and CD11c dendritic cells. CD68 and F4/80 (marker for activated microglia/macrophages) were significantly reduced. Thus, knockout of IL2r-gamma caused a suppression of the normally occurring immune response. GFP-cells that were integrated in the host outer nuclear layer expressed mature photoreceptor and synaptic markers. To sum up the first part: due to the reduced immune response, transplanted hESC-derived retinal cells survived and integrated better in the IL2r-gamma mice than in mice with normal immune system.

In the second part of the study, hESC-derived retinal cells were transplanted to the Crx  $-/-$  model of LCA, either with normal immune system or IL2r-gamma knockouts. At 3 months, few cells survived in Crx  $-/-$  mice with a normal immune system, but there was significantly higher integration in IL2r-gamma knockout animals. However,

only about 20% of the integration rate (4,000 *vs.* 20,000 cells) was seen compared with IL2r-gamma knockout mice with normal retina. Transplanted cells could still be detected at 9 months post-transplantation which is a significantly longer survival time tested than in previous studies with non-immunodeficient animals.

Improvement of vision in transplanted animals was done by testing for pupillary responses. This test consists of illuminating the transplanted eye, and recording pupillary constriction from the non-transplanted eye. This test has been used in previous studies of photoreceptor precursor transplants (10,16). At 3 and 9 months post-transplant, a partial restoration of pupillary responses was seen in Crx  $-/-$  mice that were also IL2r-gamma knockouts, but not in transplanted Crx  $-/-$  with an intact immune system. However, the intensity of the pupillary reflex does not correlate with the number of photoreceptor cells (26).

Transplanted Crx  $-/-$  mice with IL2r-gamma knockout also had very small detectable B-waves in ERG recordings that were absent in sham controls. However, the responses were very small and close to the noise level, with a small number of animals tested.

Another indication of visual function restoration was the demonstration that the immediate early genes *c-fos* and *Arc* were upregulated in visual brain centers after intense light exposure, in transplanted Crx  $-/-$ , IL2r-gamma knockout mice. The authors did not show a comparative panel of mice with normal retina. Dark-adapted anesthetized mice were exposed for 2 h with a light intensity of 10,000 lux, followed by sacrifice after 2 h. However, this light exposure was much more intense than in other publications investigating the upregulation of *c-fos* expression by light exposure. E.g., Barnard *et al.* (27) exposed mice to 15 min of fluorescent white light of 33  $\mu\text{W}/\text{cm}^2$ , which would approximately correspond to 225 lux. It would have been interesting if they had seen an effect of the transplant under physiological light intensity conditions.

Recently, several laboratories have demonstrated that recipient photoreceptors incorporate GFP label from transplanted photoreceptor precursor cells that were injected into the subretinal space. This means that GFP label alone is insufficient to tell whether donor cells really integrated into the host retina (28–30). However, Zhu *et al.* showed that GFP-labeled cells stain for human specific markers and do not co-express a mouse-specific MHC class I marker which clearly determined that they were all of human origin. Thus, there was no cytoplasmic transfer in this study.

In summary, the study of Zhu *et al.* shows that using an immunodeficient mouse model significantly improves the integration and survival of transplanted retinal cells. One word of caution: Zhu *et al.* argued that immunosuppression may have enhanced transplant integration in the study of Shirai *et al.* (5), but this study already used cyclosporine A as immunosuppressant. The study by Zhu *et al.* confirms that immunodeficient animal models are better for transplant survival and integration than models that need immunosuppression by drugs because of the side effects of immunosuppressant drugs (24). It is unclear however, how this would translate to future clinical trials. In a previous clinical trial with fetal retina-RPE sheet transplants to patients with RP and AMD, no immunosuppression was used, and transplants survived for many years (9).

### Acknowledgements

The author thanks Drs. Robert B. Aramant and Bin Lin (UC Irvine) for their critical comments.

**Funding:** The author is supported by CIRM TR4-06648, and a subcontract of NIH grant R01EY024045 (PI Hans S. Keirstead, AIVITA Biomedical, Irvine CA).

### Footnote

**Conflicts of Interest:** The author has a proprietary interest in the implantation instrument and method to transplant retinal sheets (Ocular Transplantation LLC).

### References

1. Medeiros NE, Curcio CA. Preservation of ganglion cell layer neurons in age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2001;42:795-803.
2. Seiler MJ, Aramant RB. Cell replacement and visual restoration by retinal sheet transplants. *Prog Retin Eye Res* 2012;31:661-87.
3. Assawachananont J, Mandai M, Okamoto S, et al. Transplantation of Embryonic and Induced Pluripotent Stem Cell-Derived 3D Retinal Sheets into Retinal Degenerative Mice. *Stem Cell Reports* 2014;2:662-74.
4. Seiler MJ, Lin RE, McLelland BT, et al. Vision recovery and connectivity by fetal retinal sheet transplantation in an immunodeficient retinal degenerate rat model. *Invest Ophthalmol Vis Sci* 2017;58:614-30.
5. Shirai H, Mandai M, Matsushita K, et al. Transplantation of human embryonic stem cell-derived retinal tissue in two primate models of retinal degeneration. *Proc Natl Acad Sci U S A* 2016;113:E81-90.
6. Seiler MJ, Aramant RB, Thomas BB, et al. Visual restoration and transplant connectivity in degenerate rats implanted with retinal progenitor sheets. *Eur J Neurosci* 2010;31:508-20.
7. Seiler MJ, Jones BW, Aramant RB, et al. Computational molecular phenotyping of retinal sheet transplants to rats with retinal degeneration. *Eur J Neurosci* 2012;35:1692-704.
8. Seiler MJ, Thomas BB, Chen Z, et al. Retinal transplants restore visual responses - Trans-synaptic tracing from visually responsive sites labels transplant neurons. *Eur J Neurosci* 2008;28:208-20.
9. Radtke ND, Aramant RB, Petry HM, et al. Vision Improvement in Retinal Degeneration Patients by Implantation of Retina Together with Retinal Pigment Epithelium. *Am J Ophthalmol* 2008;146:172-82.
10. MacLaren RE, Pearson RA, MacNeil A, et al. Retinal repair by transplantation of photoreceptor precursors. *Nature* 2006;444:203-7.
11. Lamba DA, Gust J, Reh TA. Transplantation of human embryonic stem cell-derived photoreceptors restores some visual function in Crx-deficient mice. *Cell Stem Cell* 2009;4:73-9.
12. West EL, Pearson RA, Barker SE, et al. Long-term survival of photoreceptors transplanted into the adult murine neural retina requires immune modulation. *Stem Cells* 2010;28:1997-2007.
13. Tucker BA, Park IH, Qi SD, et al. Transplantation of adult mouse iPS cell-derived photoreceptor precursors restores retinal structure and function in degenerative mice. *PLoS ONE* 2011;6:e18992.
14. Eberle D, Kurth T, Santos-Ferreira T, et al. Outer segment formation of transplanted photoreceptor precursor cells. *PLoS One* 2012;7:e46305.
15. Pearson RA, Barber AC, Rizzi M, et al. Restoration of vision after transplantation of photoreceptors. *Nature* 2012;485:99-103.
16. Singh MS, Charbel Issa P, Butler R, et al. Reversal of end-stage retinal degeneration and restoration of visual function by photoreceptor transplantation. *Proc Natl Acad Sci U S A* 2013;110:1101-6.
17. West EL, Gonzalez-Cordero A, Hippert C, et al. Defining the integration capacity of embryonic stem cell-derived photoreceptor precursors. *Stem Cells* 2012;30:1424-35.
18. Tucker BA, Mullins RF, Streb LM, et al. Patient-specific iPSC-derived photoreceptor precursor cells as a means to

- investigate retinitis pigmentosa. *Elife* 2013;2:e00824.
19. Laver CR, Metcalfe AL, Szczygiel L, et al. Bimodal in vivo imaging provides early assessment of stem-cell-based photoreceptor engraftment. *Eye (Lond)* 2015;29:681-90.
  20. Yanai A, Laver CR, Gregory-Evans CY, et al. Enhanced functional integration of human photoreceptor precursors into human and rodent retina in an ex vivo retinal explant model system. *Tissue Eng Part A* 2015;21:1763-71.
  21. Streilein JW, Ma N, Wenkel H, et al. Immunobiology and privilege of neuronal retina and pigment epithelium transplants. *Vision Res* 2002;42:487-95.
  22. Warfvinge K, Kiilgaard JF, Klassen H, et al. Retinal progenitor cell xenografts to the pig retina: immunological reactions. *Cell Transplant* 2006;15:603-12.
  23. Zhu J, Cifuentes H, Reynolds J, et al. Immunosuppression via Loss of IL2rgamma Enhances Long-Term Functional Integration of hESC-Derived Photoreceptors in the Mouse Retina. *Cell Stem Cell* 2017;20:374-84 e5.
  24. Diehl R, Ferrara F, Muller C, et al. Immunosuppression for in vivo research: state-of-the-art protocols and experimental approaches. *Cell Mol Immunol* 2017;14:146-79.
  25. Lamba DA, McUsic A, Hirata RK, et al. Generation, purification and transplantation of photoreceptors derived from human induced pluripotent stem cells. *PLoS One* 2010;5:e8763.
  26. Kovalevsky G, DiLoreto D Jr, Wyatt J, et al. The intensity of the pupillary light reflex does not correlate with the number of retinal photoreceptor cells. *Exp Neurol* 1995;133:43-9.
  27. Barnard AR, Appleford JM, Sekaran S, et al. Residual photosensitivity in mice lacking both rod opsin and cone photoreceptor cyclic nucleotide gated channel 3 alpha subunit. *Vis Neurosci* 2004;21:675-83.
  28. Santos-Ferreira T, Llonch S, Borsch O, et al. Retinal transplantation of photoreceptors results in donor-host cytoplasmic exchange. *Nat Commun* 2016;7:13028.
  29. Pearson RA, Gonzalez-Cordero A, West EL, et al. Donor and host photoreceptors engage in material transfer following transplantation of post-mitotic photoreceptor precursors. *Nat Commun* 2016;7:13029.
  30. Ortin-Martinez A, Tsai EL, Nickerson PE, et al. A Reinterpretation of Cell Transplantation: GFP Transfer From Donor to Host Photoreceptors. *Stem Cells* 2017;35:932-9.

doi: 10.21037/sci.2017.08.05

**Cite this article as:** Seiler MJ. hESC-derived photoreceptors survive and integrate better in immunodeficient retina. *Stem Cell Investig* 2017;4:70.