

UC San Diego

UC San Diego Previously Published Works

Title

Visually guided behavior in freely moving mice

Permalink

<https://escholarship.org/uc/item/5vk0k34b>

Journal

BMC Neuroscience, 14(Suppl 1)

ISSN

1471-2202

Authors

Sriram, Balaji
Cruz-Martin, Alberto
DeNardo, Laura
et al.

Publication Date

2013-07-08

DOI

<http://dx.doi.org/10.1186/1471-2202-14-S1-P141>

Peer reviewed

POSTER PRESENTATION

Open Access

Visually guided behavior in freely moving mice

Balaji Sriram^{1*}, Alberto Cruz-Martin¹, Laura DeNardo^{1,2}, Mohit Patel¹, Eui-seok J Kim³, Anirvan Ghosh^{1,4}

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

The ability of neuroscience to ascribe functions to brain regions and to different neuronal subtypes within these regions depends on our ability to identify behavioral paradigms that depend on these functions and to measure these behaviors in a quantitative fashion. Due to their inexpensive nature, extensive similarities in brain architecture, availability of genetic tools and ease of handling, rodents have recently become an important tool in the study of neuronal coding. We train common mice (*Mus musculus*) in a potentially cortex dependent, visually guided task.

Adult C57BL/6 mice were trained to perform an orientation discrimination task in a 2AFC (2 alternate forced choice) behavioral training chamber. Subjects performed trials in a semi-closed economy obtaining water through

the training chamber 5-6 days a week with ad-libitum water provided for the remaining 1-2 days. During the initial training stage, stimuli were full field drifting gratings (100% contrast, ~0.05 cpd, 2 Hz) with randomly interleaved trials showing gratings oriented $\pm 45^\circ$ from the vertical. Mice performed on average ~200 trials every day. Subjects learned the task until they reached a criterion performance (80-85% correct) (Figure 1A).

Past this initial training stage, subjects were tested on a variety of stimuli where the discriminated gratings showed varying spatial frequency (Figure 1B), contrast (Figure 1C), temporal frequency (Figure 1D) and orientation (Figure 1E) from the vertical allowing us to measure precise psychometric curves on the performance of subjects.




Figure 1 Example psychometric curves from a typical subject. (A) Initial training with full contrast, 0.05cpd, 2Hz gratings. After the initial training period, performance was measured with varying spatial frequency (B), contrast (C), temporal frequency (D) and orientation of grating (E). All data show binomial fits to the data with 95% confidence intervals. In B,C,D and E the condition corresponding to the original training is denoted by '*'.

* Correspondence: basriram@ucsd.edu

¹Division of Biology, University of California, San Diego, La Jolla, USA
Full list of author information is available at the end of the article

The role of cortical (Primary visual cortex; V1) and subcortical circuits (Superior colliculus; SC) mediating this behavior will be probed through the use of lesions. We hope that such measurements will provide a basis to constrain and in the future uniquely describe models of the neocortex.

Author details

¹Division of Biology, University of California, San Diego, La Jolla, USA.

²Department of Biology, Stanford University, Stanford, USA. ³SNL-C, Salk Institute for Biological Sciences, La Jolla, USA. ⁴CNS Discovery, F. Hoffmann La Roche, Basel, Switzerland.

Published: 8 July 2013

doi:10.1186/1471-2202-14-S1-P141

Cite this article as: Sriram et al.: Visually guided behavior in freely moving mice. *BMC Neuroscience* 2013 14(Suppl 1):P141.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

