

UCSF

UC San Francisco Previously Published Works

Title

Conditional ablation of brain-derived neurotrophic factor-TrkB signaling impairs striatal neuron development.

Permalink

<https://escholarship.org/uc/item/61k55988>

Journal

Proceedings of the National Academy of Sciences of USA, 109(38)

Authors

Li, Yun
Yui, Daishi
Luikart, Bryan
[et al.](#)

Publication Date

2012-09-18

DOI

10.1073/pnas.1212899109

Peer reviewed

Conditional ablation of brain-derived neurotrophic factor-TrkB signaling impairs striatal neuron development

Yun Li^{a,1}, Daishi Yui^a, Bryan W. Luikart^{a,2}, Renée M. McKay^a, Yanjiao Li^a, John L. Rubenstein^b, and Luis F. Parada^{a,3}

^aDepartment of Developmental Biology and Kent Waldrep Center for Basic Research on Nerve Growth and Regeneration, University of Texas Southwestern Medical Center, Dallas, TX 75390; and ^bNina Ireland Laboratory of Developmental Neurobiology, Department of Psychiatry, University of California, San Francisco, CA 94143

Contributed by Luis F. Parada, August 2, 2012 (sent for review June 21, 2012)

Neurotrophic factors, such as brain-derived neurotrophic factor (BDNF), are associated with the physiology of the striatum and the loss of its normal functioning under pathological conditions. The role of BDNF and its downstream signaling in regulating the development of the striatum has not been fully investigated, however. Here we report that ablation of *Bdnf* in both the cortex and substantia nigra depletes BDNF in the striatum, and leads to impaired striatal development, severe motor deficits, and postnatal lethality. Furthermore, striatal-specific ablation of *TrkB*, the gene encoding the high-affinity receptor for BDNF, is sufficient to elicit an array of striatal developmental abnormalities, including decreased anatomical volume, smaller neuronal nucleus size, loss of dendritic spines, reduced enkephalin expression, diminished nigral dopaminergic projections, and severe deficits in striatal dopamine signaling through DARPP32. In addition, *TrkB* ablation in striatal neurons elicits a non-cell-autonomous reduction of tyrosine hydroxylase protein level in the axonal projections of substantia nigral dopaminergic neurons. Thus, our results establish an essential function for TrkB in regulating the development of striatal neurons.

Huntington disease | neurotrophin

The striatum is a subcortical part of the telencephalon and a major input component of the basal ganglia system. The striatum plays a crucial role in regulating voluntary movements, as well as reward-associated learning and memory. It can be separated into dorsal and ventral parts based on known anatomical and functional distinctions, with the dorsal portion associated with movement regulation and the ventral striatum involved predominantly in mediating neurological functions pertaining to motivation, reward, and emotion. Progressive degeneration of the striatal medium spiny neurons (MSNs) and their main dopaminergic inputs from the substantia nigra are the cardinal symptoms of the neurological disorders Huntington disease (HD) and Parkinson disease, respectively.

Like other telencephalic regions, the striatum contains inhibitory neurons generated from the medial ganglionic eminence. The majority of inhibitory GABAergic neurons are MSNs that project outside of the striatum to neighboring structures, such as the globus pallidus and the substantia nigra. The mechanism regulating development of the striatum involves combinatorial codes of transcription factors and parallels the development of other neural structures; however, the molecular mechanisms underlying the maturation and maintenance of the striatum remain largely unknown. Recent studies have shown that brain-derived neurotrophic factor (BDNF) exerts prosurvival effects on striatal neurons in vitro (1, 2) and delays the onset of degeneration in a transgenic mouse model of HD (3). However, it has been shown that during development and in adulthood, *Bdnf* mRNA is largely absent from striatal neurons and glia. Instead, BDNF is produced in the cortex and substantia nigra and is transported anterogradely to striatal neurons (4). Indeed, cortical ablation of *Bdnf* leads to cellular deficits in the striatum (5) and global gene expression impairment (6). These findings suggest an important role for BDNF in the striatum; however, given BDNF's complex downstream signaling capacities, the nature of its intracellular signaling

casades pertaining to development, maturation, and function of the striatum remains to be delineated.

In this study, we conditionally ablated BDNF or its receptor TrkB in corticostriatal and nigrostriatal neuronal circuits. We found that *Bdnf* deletion in both cortex and substantia nigra led to complete depletion of BDNF protein in the striatum. Mutant mice displayed dramatic developmental abnormalities and neurological impairments. Furthermore, specific deletion of *TrkB* from striatal neurons was sufficient to produce this wide range of developmental deficits. Thus, our results demonstrate that BDNF and TrkB play critical paracrine and cell-autonomous roles, respectively, in the development and maintenance of striatal neurons.

Results

***Bdnf* Ablation in Cortex and Substantia Nigra Depletes Striatal BDNF Protein.** The striatum does not itself produce BDNF, but rather receives BDNF protein through anterograde transport from cortical and nigral neurons (4). To generate conditional knock-out (cKO) mice lacking BDNF protein in the striatum, we crossed mice harboring flox alleles of *Bdnf* with a Cre recombinase line that mediates recombination in both cortex and substantia nigra. It has been shown that mice expressing Cre under the endogenous *Bfl* promoter have widespread recombination in the telencephalon (7). By crossing the *Bfl-Cre* mice with *R26* reporter mice (8), we confirmed that *Bfl-Cre* was active in the cortex and mediated effective recombination in neurons, including those in layer V, where the majority of projections to the striatum originate (*R26*^{Bfl} mice; Fig. 1*A*). X-Gal staining on adult *R26*^{Bfl} mice also revealed positive cells in the midbrain region in the proximity of the substantia nigra (Fig. 1*A*, Right). Double immunostaining for β -gal and tyrosine hydroxylase (TH), a marker for dopaminergic neurons, confirmed that TH-positive neurons also expressed β -gal (Fig. 1*B*, Right). To definitively demonstrate the effectiveness of *Bfl-Cre*-mediated deletion of *Bdnf*, we examined *Bdnf* mRNA expression by in situ hybridization. We found that although *Bdnf* was expressed robustly in the cortex and substantia nigra of control mice, it was absent in the *Bfl-Cre*; *Bdnf*^{flox/flox} mice (BDNF^{Bfl} mice) at age 1 mo (Fig. 1*C*). Deletion of *Bdnf* from these two structures led to complete depletion of BDNF protein from the striatum as measured by ELISA [(Fig. 1*D*); $n = 5-6$ for each; $P < 0.001$ for control-BDNF^{Bfl} comparisons].

BDNF Depletion Leads to Motor Dysfunction and Deficits in Striatal Neuron Development. In contrast to *Bdnf* germline KO mice that die shortly after birth, the BDNF^{Bfl} mice were viable. However,

Author contributions: Yun Li, D.Y., B.W.L., and L.F.P. designed research; Yun Li, D.Y., B.W.L., and Yanjiao Li performed research; J.L.R. contributed new reagents/analytic tools; Yun Li, D.Y., B.W.L., R.M.M., J.L.R., and L.F.P. analyzed data; and Yun Li, R.M.M., and L.F.P. wrote the paper.

The authors declare no conflict of interest.

¹Present address: Whitehead Institute for Biomedical Research, Cambridge, MA 02142.

²Present address: Department of Physiology and Neurobiology, Geisel School of Medicine at Dartmouth, Lebanon, NH 03756.

³To whom correspondence should be addressed. E-mail: luis.parada@utsouthwestern.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1212899109/-DCSupplemental.




Fig. 1. *Bdnf* ablation in the cortex and substantia nigra depletes BDNF from the striatum. (A) X-Gal staining on coronal sections of 5-wk-old R26^{Bfl} mice at the forebrain (Left) and midbrain (Right) levels. Note the robust recombination in the cortex (ctx), striatum (str), and substantia nigra (sn). (Middle) Higher-magnification view of the cortex showing X-Gal staining throughout layers II–VI. (B) Double immunostaining for β -gal (green) and NeuN (red) (Left) or TH (red) (Right) revealed *Bfl*-Cre-mediated recombination in neurons of the cortex and substantia nigra. (Scale bars: 50 μ m.) (C) In situ hybridization for *Bdnf* mRNA in 4-wk-old control and BDNF^{Bfl} mice. Note the ablation of *Bdnf* mRNA from the cortex (Upper) and substantia nigra (Lower) of BDNF^{Bfl} mice. (Scale bars: 500 μ m.) (D) ELISA showed complete removal of BDNF protein from the cortex and striatum of BDNF^{Bfl} mice. $n = 5$ –6 for each. Results are mean \pm SEM. *** $P < 0.001$.

they were runty compared with littermate controls by 3 wk of age (Fig. 2A) and did not survive past 8 wk of age. In addition, the BDNF^{Bfl} mice displayed early-onset behavioral abnormalities, including hindlimb clamping, hyperactivity, and circling (not shown), indicative of neurological dysfunction. Although born with normal-sized brains, the BDNF^{Bfl} mice also showed reduced postnatal brain weight (Fig. 2B and C; $n = 4$ –6 for each genotype and age; $P < 0.05$ at 3 wk and $P < 0.001$ at 5 wk). Analysis of postnatal brain regions revealed significant volume reduction in mutant striatum (Fig. 2D; $n = 4$ –6 for each; $P < 0.05$ at 3 wk and $P < 0.001$ at 5 wk), as well as reduction in dentate gyrus (DG) and cortex volume (Fig. S1; $n = 5$ for each; $P < 0.01$ and $P < 0.05$, respectively), consistent with previous reports (5, 9). Notably, the overall architecture of the striatum in BDNF^{Bfl} mice, as evaluated by Nissl and luxol fast blue staining at age 3 wk, was

comparable to that in controls (Fig. S1), indicating that BDNF depletion did not affect structural formation of the striatum.

To assess whether BDNF depletion resulted in striatum deficits at the cellular level, we evaluated expression levels and distribution of DARPP32 (dopamine- and cAMP-regulated phosphoprotein, 32 kDa), a major target of dopamine-activated adenylyl cyclase and the key mediator of dopamine signaling (10). We saw a dramatic reduction in the intensity of DARPP32 immunoreactivity throughout the striatum of BDNF^{Bfl} mice at age 1 mo (Fig. 2E, Upper). High magnification revealed loss of DARPP32 protein in both the soma and processes of striatal neurons (Fig. 2E, Lower). We further examined striatal MSN morphology using Golgi staining, and found significantly fewer spines on secondary dendritic branches in 3-wk-old BDNF^{Bfl} mice (Fig. 2F and G; $n = 7$ mice for each; $P < 0.01$), as well as shorter spines (Fig. 2H; $n = 7$ mice for each; $P < 0.05$). Taken together, these results indicate impaired striatal neuronal maturation in the BDNF^{Bfl} mice.

TrkB Is Required for Striatal Development. BDNF signals through its high-affinity receptor TrkB, and the pan-neurotrophin, low-affinity receptor p75. Although both receptors are reportedly present in the developing striatum (11), only *TrkB* is robustly expressed in postnatal and adult striatum (12) (Fig. 3A). To evaluate the requirement for TrkB in striatal development in vivo, we examined *TrkB*^{-/-} mice at postnatal days 4 and 8. We previously reported that although the majority of *TrkB*^{-/-} mice die at birth (13), a small percentage are viable in the first postnatal week. Compared with littermate controls, these mice had decreased striatal volume and a dramatic reduction in DARPP32 protein level (Fig. 3B and C). We also generated a line of *TrkB* cKO mice using the *Bfl*-Cre mice, which lack TrkB in the striatum, in addition to other brain regions including cortex, hippocampus, and substantia nigra (*TrkB*^{Bfl} mice). The *TrkB*^{Bfl} mice displayed developmental abnormalities similar to those of the BDNF^{Bfl} mice, including reduced brain and body weight, hindlimb clamping, postnatal lethality, circling behavior, and ataxia (not shown). Cellular analysis revealed reduction in DARPP32 immunostaining and decreased spine density on secondary dendrites at 3 wk postnatal age (Fig. 3D–F; $n = 6$ mice for each; $P < 0.001$). Thus, a loss of TrkB impairs normal striatal development. However, whether the requirement for TrkB lies within the striatum or elsewhere could not be assessed in these mutant mice, owing to global deletion of *TrkB*.

To examine whether striatal-specific deletion of *TrkB* would recapitulate the BDNF^{Bfl} and *TrkB*^{Bfl} phenotypes, we used a well-characterized *Dlx5/6i-cre* transgenic line (14) that mediates robust recombination in the GABAergic cells of the striatum and olfactory bulb (OB) (Fig. 4A). Limited recombination was also observed in the inhibitory neurons of the cortex and




Fig. 2. BDNF depletion impairs striatal development. (A) BDNF^{Bfl} mice had lower body weight at postnatal 3 wk and 5 wk. $n = 4$ –6 for each. (B and C) At postnatal age 3 wk (C) and 5 wk (B and C), the BDNF^{Bfl} mice had significantly lower brain weight compared with littermate controls. $n = 4$ –6 for each. (D) BDNF^{Bfl} mice had lower striatal volumes at postnatal age 3 wk and 5 wk compared with littermate controls. $n = 4$ –6 for each. (E) (Upper) Immunostaining demonstrated reduced levels of DARPP32 protein in the striatum of BDNF^{Bfl} mice. (Lower) Higher-magnification images showing loss of DARPP32 in both the soma and processes of MSNs. ctx, cortex; str, striatum. (Scale bars: Upper, 500 μ m; Lower, 50 μ m.) (F–H) BDNF depletion led to fewer and shorter dendritic spines on striatal MSNs. (Scale bar: 10 μ m.) $n = 7$ mice for each. Results are mean \pm SEM. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

striatal, but not cortical, neurons (TrkB^{Dlx} mice; Fig. 4C). Finally, immunoblotting showed complete absence of the full-length TrkB protein from striatum of 3-wk-old TrkB^{Dlx} mice (Fig. 4D; $n = 3$ for each; $P < 0.001$). It has been reported that a truncated TrkB splice variant is selectively expressed by astrocytes and mediates neurotrophin-evoked calcium signaling (15). We found that truncated TrkB expression in striatum of TrkB^{Dlx} mice was not affected (Fig. 4D and E; $n = 3$ for each; $P > 0.02$ for all control–TrkB^{Dlx} comparisons), further confirming the neuron-specific ablation of TrkB by *Dlx5/6i-cre*. Expression of both forms of TrkB protein was unaffected in cortex and hippocampus (Fig. 4D and E). These results demonstrate a complete and specific ablation of *TrkB* from striatal neurons, permitting us to investigate *TrkB* cell-autonomous function in this cell population.

Like BDNF^{Bfl} mice, TrkB^{Dlx} mice were runty and did not survive past age 4 wk. Significant decreases in body weight and striatal volume were evident at 2 wk and 3 wk of age (Fig. 5A and C; $n = 5–7$ for each; $P < 0.001$), and a significant reduction in brain weight was evident by 3 wk (Fig. 5B; $n = 5–6$ for each; $P < 0.05$). Attesting to the restricted recombination pattern mediated by the *Dlx5/6i-cre*, we found no overt difference in the size of the DG or cortex (Fig. S2; $n = 5–6$ for each). Striatal neurons in the TrkB^{Dlx} mice had reduced soma size, as evaluated by NeuN immunostaining (Fig. 5D and E; $n = 92–98$ neurons from five mice of either genotype; $P < 0.001$). Neurons in the striatum of TrkB^{Dlx} mice were more compact, and quantitative analysis revealed a significant increase in the density of NeuN-positive striatal neurons (Fig. 5D and F; $n = 9$ for each; $P < 0.5$). Along with the reduction in striatal volume (Fig. 5C), TrkB^{Dlx} mice had approximately 13% fewer NeuN-positive striatal neurons compared with their littermate controls at age 3 wk (1.58×10^6 vs. 1.81×10^6). Golgi staining demonstrated reduced spine density and length in TrkB^{Dlx} mice compared with littermate controls (Fig. 5G–I; $n = 8–9$ mice for each; $P < 0.001$ and $P < 0.05$, respectively). We further assessed the development of striatal MSNs by DARPP32 immunostaining and found reduced protein expression

in both the soma and the processes (Fig. 5J). In addition, the density of DARPP32-positive MSNs was reduced in the striatum of TrkB^{Dlx} mice (Fig. 5J). Given the increased density of NeuN-positive cells in the TrkB^{Dlx} mice (Fig. 5F), along with the fact that DARPP32-positive MSNs represent more than 90% of all neurons in the striatum, the loss of DARPP32 staining likely reflects a decline in gene expression rather than cell loss.

BDNF has been implicated in regulating the development of a subgroup of striatal MSNs, including the striatopallidal neurons that express the neuropeptide enkephalin (16). To determine whether TrkB loss impairs the development of these neurons, we examined enkephalin expression using an antibody against the mature form of the peptide. By age 3 wk, TrkB^{Dlx} mice showed a substantial reduction in the intensity of enkephalin immunostaining in both the soma and the surrounding dendrites (Fig. 5K). Similar to the loss of DARPP32, this reduced enkephalin expression was not accompanied by a comparable loss of NeuN-positive cells (Fig. 5F), and thus most likely reflects impaired terminal maturation and maintenance on the level of individual cells.

Taken together, these findings indicate that a lack of striatal TrkB signaling was responsible for the deficits associated with BDNF depletion in the cortex and substantia nigra.

TrkB Ablation Attenuates Intracellular Signaling. Our findings in the TrkB^{Dlx} mice demonstrated impaired cellular striatal development and maturation. The decrease in DARPP32 protein level in the absence of BDNF or TrkB further indicates a disruption in normal dopamine signaling. Consequently, we examined the level of total DARPP32 protein and phospho-Thr³⁴ DARPP32 in striatum of 3-wk-old mice. Recent studies have shown that activation of the D1 class dopamine receptors elicits phosphorylation of DARPP32 on Thr³⁴, which conveys key signal transduction to the protein kinase A cascade (17). Immunoblot analysis revealed a significant loss of total DARPP32 protein in the TrkB^{Dlx} mice, confirming the immunohistochemistry results (Fig. 6A and B; $n = 3$ for each; $P < 0.001$), as well as a loss of phospho-Thr³⁴ DARPP32 (Fig. 6A).




Fig. 5. Removal of TrkB in striatal neurons impairs striatal development. (A–C) TrkB^{Dlx} mice had reduced body weight (A) and brain weight (B), as well as decreased striatal volume (C), at postnatal age 2 wk and 3 wk. $n = 5–7$ for each. (D) NeuN immunostaining showed a comparable striatal structure in 3-wk-old TrkB^{Dlx} mice. (Insets) Higher-magnification views of NeuN-positive striatal cells. (Scale bars: Upper, 200 μm; Lower, 50 μm; Insets, 10 μm.) (E) Quantification revealed significantly smaller neuronal nucleus size in striatum of 3-wk-old TrkB^{Dlx} mice. $n = 92–98$ neurons from five mice of each genotype. (F) The density of NeuN-positive striatal neurons was higher in 3-wk-old TrkB^{Dlx} mice. $n = 9$ for each. (G–I) Golgi staining showed that the TrkB^{Dlx} mice had fewer and shorter spines on the secondary dendrites of striatal MSNs. (Scale bars: 10 μm.) $n = 8–9$ mice for each. (J) Immunostaining revealed significant loss of DARPP32 protein in the striatum of TrkB^{Dlx} mice. (Scale bars: Upper, 200 μm; Lower, 50 μm.) (K) Enkephalin-positive striatopallidal neurons were reduced in TrkB^{Dlx} mice. (Scale bars: Upper, 500 μm; Lower, 50 μm.) Results are mean \pm SEM. * $P < 0.05$; *** $P < 0.001$.

Quantitative analysis demonstrated a significant reduction in the ratio of phospho to total DARPP32 (Fig. 6C; $n = 3$ for each; $P < 0.001$), indicating that the loss of phospho-Thr³⁴ DARPP32 reflects a reduction in total protein, as well as impaired activation of the dopamine signaling pathway in the basal state.

TrkB activation can be propagated through downstream pathways common to receptor tyrosine kinases, including the ERK and PI3-kinase pathways, which promote neuronal growth, differentiation, and survival. We examined the levels of phospho-ERK1/2 at Thr²⁰²/Thr²⁰⁴ and phospho-Ser⁴⁷⁹ AKT in striatum by immunoblot analysis. At age 3 wk, TrkB^{Dlx} mice showed a significant reduction in the levels of both phospho-proteins compared with littermate controls (Fig. 6D and E; $n = 3$ for each; $P < 0.001$ for both). Thus, ablating *TrkB* in striatal neurons leads to deficits in key intracellular signaling.

TrkB Ablation Leads to Non-Cell-Autonomous Loss of Dopaminergic Projections. The BDNF^{±Bfl} and TrkB^{Dlx} mice displayed severe neurological phenotypes, including shortened lifespan and locomotor defects. In particular, the TrkB^{Dlx} mice were hypoactive, reminiscent of phenotypes seen in other mutant mice with impaired nigral dopamine signaling, including dopamine-deficient TH mutant mice (18). A previous study suggested that *Bdnf* ablation in midbrain neurons leads to age-dependent dopaminergic neuron loss (19). We and others have shown that the *Dlx5/6i-cre*

does not mediate recombination in the substantia nigra (Fig. 6F) (14, 20). We examined the substantia nigra dopaminergic neurons in the TrkB^{Dlx} mice by TH immunostaining. We found no overt difference in the number of TH-positive dopaminergic neurons at age 3 wk (Fig. 6G); however, in the striatum, we found a substantial reduction in the area and intensity of TH immunostaining in 3-wk-old TrkB^{Dlx} mice compared with controls (Fig. 6H). We confirmed this finding by immunostaining for dopamine transporter, a protein residing in dopaminergic terminals (Fig. 6I). Finally, we measured TH levels in striatal protein lysates and found a 50% reduction in TrkB^{Dlx} mice (Fig. 6J; $n = 3$ for each; $P < 0.001$). These data demonstrate that ablation of *TrkB* in striatal neurons results in a loss of dopaminergic projections to the striatum through a non-cell-autonomous mechanism.

Discussion

The striatum is the largest CNS structure lacking a detectable endogenous source of BDNF. Instead, striatal neurons receive BDNF from projection neurons of the cortex and substantia nigra via anterograde transport (4). BDNF is important to the survival of striatal neurons in vitro (1, 2), and global ablation of *Bdnf* in the CNS leads to selective impairment in striatal MSN development (2). Furthermore, modulation of BDNF levels by genetic deletion or overexpression in mice harboring HD-related mutations led to acceleration or amelioration, respectively, of the cellular phenotypes associated with HD (3, 16, 21). Here we demonstrate that *Bdnf* ablation in cortex and substantia nigra depletes BDNF in the striatum and leads to developmental deficits. Taken together, these findings suggest that striatal neurons are critically dependent on BDNF.

BDNF and its downstream signaling pathways have been implicated in regulating the development and functioning of the rodent brain (22). Removal of BDNF from the cortex and substantia nigra (regions that supply the striatum with BDNF) affects the physiology of these brain regions (19, 23). We used the TrkB^{Dlx} mice generated in this study to specifically examine the role of BDNF downstream signaling in striatal neurons, while leaving the cortical and nigra neurons largely unaffected. These mutant mice exhibited a panel of molecular, cellular, and behavioral phenotypes demonstrating severe developmental abnormalities, thus establishing a cell-autonomous role for TrkB in striatal neurons.

Our findings support the notion that neurons in the striatum are highly sensitive to the availability of BDNF. We previously reported that ablating *TrkB* in neurons of the cerebral cortex, hippocampus, OB, and cerebellum does not lead to drastic cellular degeneration or postnatal lethality (9, 24, 25). Here we demonstrate that TrkB^{Dlx} mice, in which *TrkB* is selectively ablated in striatal neurons, have severe locomotor deficits, reduced body and brain weight, and postnatal lethality, phenotypes almost identical to those seen in mice with broader ablation of *TrkB* (TrkB^{B1} mice). This finding is consistent with a comparatively prominent role for TrkB in the striatum. In addition, a lack of TrkB in striatal MSNs resulted in reduced cell size, fewer dendritic spines, and diminished expression of a key signaling peptide (enkephalin) and a key protein (DARPP32). We further note that DARPP32 reduction coincided with dramatically impaired activation of DARPP32 in the form of Thr³⁴ phosphorylation, indicating reduced basal dopamine signaling. A recent study using a different *Dlx5/6-Cre* transgenic mouse line reported that deletion of TrkB in the striatum led to profound reduction of striatal volume (by 53%) and NeuN-positive striatal neurons (by 58%) (26). Further investigation revealed normal proliferation but dramatically higher apoptosis. However, we caution that this particular line of *Dlx5/6-Cre* transgenic mice was reported to have higher cell death levels during embryogenesis compared with control mice lacking the transgene (27). Baydyuk et al. (26) used only mice homozygous for the floxed *TrkB* alleles as controls for their TrkB mutant mice; thus, whether the observed cell death was caused by the *Dlx5/6-Cre* transgene alone or was affected by *TrkB* deletion remains unclear. In the present study, we used a Cre line that did not cause cell death or striatal volume reduction (20) and used littermate *TrkB^{fllox/fllox}*, *TrkB^{fllox/wt}*, and *TrkB^{wt/wt};cre/wt* mice as controls to eliminate a potential




Fig. 6. TrkB^{Dlx} mice have striatal molecular and cellular phenotypes. (A–C) Immunoblot analysis of striatal lysates from control and TrkB^{Dlx} mice showed a significant reduction in total DARPP32 levels (A and B), as well as phospho-Thr³⁴ DARPP32 levels (A and C). $n = 3$ for each. (D and E) *TrkB* ablation in striatal neurons led to reduced activity of the ERK and AKT pathways, as shown by immunoblot analysis for phospho-Thr²⁰²/Thr²⁰⁴ ERK and total ERK proteins, as well as phospho-Ser⁴⁷⁹ and total AKT proteins. $n = 3$ for each. (F) Coimmunostaining for TH (red) and β -gal (green) on sections of R26^{Dlx} mice demonstrated lack of recombination in dopaminergic neurons of substantia nigra. (Scale bars: 100 μ m.) (G) TH immunostaining in substantia nigra showed comparable numbers of dopaminergic neurons in control and TrkB^{Dlx} mice. (Scale bars: 200 μ m.) (H and I) (Upper) Three-week-old TrkB^{Dlx} mice displayed reduced dopaminergic terminals in striatum, labeled by TH (H) or dopamine transporter (I). (Lower) Higher-magnification views. (Scale bars: Upper, 200 μ m; Lower, 50 μ m.) (J) Immunoblot analysis showed a 50% reduction in TH protein in the striatum of TrkB^{Dlx} mice. $n = 3$ for each. Results are mean \pm SEM. *** $P < 0.001$.

effect of the *Dlx5/6-Cre* transgene on striatal development. Our results demonstrate that lack of TrkB leads to substantial impairment in striatal cellular functionality.

Our findings are in agreement with previous reports showing that cortical ablation of *Bdnf* using *Emx-Cre* (*Emx-BDNF^{KO}* mice) (5) or global deletion of *Bdnf* in the postmitotic neurons of the CNS using *Tau-Cre* (2) leads to a significant reduction in striatal BDNF and striatal impairment. Interestingly, these lines of *Bdnf* mutant mice were viable and survived into advanced age, whereas the *BDNF^{B1}* mice generated in the current study had an average lifespan of 2 mo. A possible cause of this difference is the complete deletion of *Bdnf* in the substantia nigra pars compacta mediated by the *Bfl-Cre*. It has been demonstrated that dopaminergic neurons of the substantia nigra express *Bdnf* (19) and supply 14% of BDNF protein in the ipsilateral striatum (4). Although this is a smaller contribution compared with the corticostriatal input (66% ipsilateral) (4), the nigrostriatal projection apparently supplies sufficient BDNF to maintain basal striatal function for survival of the *Emx-BDNF^{KO}* mice. Dopaminergic neurons emit extensive axonal arborizations in the striatum, with each neuron forming axonal terminals that cover up to 5.7% of the total striatal volume (28), thus establishing a strong delivery system for neurotransmitters (dopamine) and trophic factors (e.g., BDNF). Thus, by ablating *Bdnf* in neurons of both the cortex and substantia nigra, we created a cKO mouse with more severe neurological impairment and demonstrated that striatal BDNF could play a role in postnatal survival.

Taken together, our results demonstrate that ablation of TrkB signaling in striatal neurons impairs the development of these neurons and their associated neural circuits. Future studies on the role of TrkB signaling in the pathophysiology of the striatum and the potential benefits of restoring TrkB signaling may provide insight into preventing, delaying, and reversing the progression of striatal neurodegeneration.

Methods

Mice. The *TrkB^{fl/fl}*, *TrkB^{fl/wt}*, *BDNF^{fl/wt}*, *Bfl-Cre*, and *Dlx5/6i-cre* mouse lines were generated as described previously (7, 13, 14, 24, 25). *R26* mice were obtained from Jackson Laboratory. For generation of the *BDNF^{B1}* mice, *BDNF^{fl/wt};cre/wt* mice were interbred with *BDNF^{fl/wt}* or *BDNF^{fl/wt}* mice. Littermate *BDNF^{fl/wt}*, *BDNF^{fl/wt}*, and *BDNF^{wt/wt};cre/wt* mice were in-

distinguishable and thus were pooled as the control group. *TrkB* cKO mice (*TrkB^{B1}* and *TrkB^{Dlx}*) were derived by crossing *TrkB^{fl/wt};cre/wt* mice with *TrkB^{fl/wt}* or *TrkB^{fl/wt}* mice. Littermate *TrkB^{fl/wt}*, *TrkB^{fl/wt}*, and *TrkB^{wt/wt}*, *cre/wt* mice served as the control group. All animals were maintained on the 129/SvEv and C57BL/6 mixed background. All mouse protocols used in this study were approved by the University of Texas Southwestern Medical Center's Institutional Animal Care and Research Advisory Committee.

Histology. Mice were anesthetized and perfused transcardially with PBS followed by 4% (wt/vol) paraformaldehyde (PFA), and the dissected brains post-fixed in 4% (wt/vol) PFA at 4 °C. For Golgi staining, mice were perfused with PBS alone, and the dissected brain was incubated in Golgi solution for 12 d, as described previously (24). In situ hybridization for *Bdnf* and *TrkB* mRNA was done using 14- μ m-thick fresh-frozen sections (24, 29). X-Gal staining was performed on 50- μ m-thick floating brain sections prepared on a vibratome (20).

Immunoblot Analysis. Immunoblot analysis was performed as described previously (30) using the following primary antibodies: actin (Sigma-Aldrich), phospho-Ser⁴⁷⁹ and total AKT (Cell Signaling), phospho-Thr³⁴ DARPP32 (Cell Signaling), total DARPP-32 (BD Biosciences), phospho-Thr²⁰²/Thr²⁰⁴ and total ERK (Cell Signaling), TH (Chemicon), and TrkB (Dr. Louis Reichardt, University of California San Francisco). HRP-conjugated secondary antibodies (Santa Cruz Biotechnology) were used to detect primary antibodies, and visualized using the ChemiGlow kit (Alpha Innotech) in accordance with the manufacturer's instructions. Membranes blotted for phosphoproteins were stripped and reprobed with antibodies against total proteins. Quantification of bands was performed using Kodak 1D Image Analysis software (30). Values for phosphoproteins were normalized to total proteins, and the levels of actin protein were used as controls for total DARPP32, and TrkB proteins.

BDNF ELISA. The concentration of BDNF protein in cortical and striatal lysates was measured using a BDNF ELISA Kit (Promega) as described previously (9, 24).

ACKNOWLEDGMENTS. We thank Shawna Kennedy, Linda McClellan, Steven McKinnon, and Ahmed Mahmoud for technical assistance, and Dr. Louis Reichardt for the TrkB antibody. The *Bfl-Cre* was a generous gift from Dr. Susan McConnell. We thank Drs. Mark Lush, James Bibb, Serge Nef, and members of the L.F.P. laboratory for sharing reagents and helpful suggestions. This work was supported by National Institute of Neurological Disorders and Stroke Grant R37NS033199, Conte Center Grant 7P50MH066172-07, and the American Cancer Society.

- Ivkovic S, Ehrlich ME (1999) Expression of the striatal DARPP-32/ARPP-21 phenotype in GABAergic neurons requires neurotrophins in vivo and in vitro. *J Neurosci* 19:5409–5419.
- Rauskolb S, et al. (2010) Global deprivation of brain-derived neurotrophic factor in the CNS reveals an area-specific requirement for dendritic growth. *J Neurosci* 30:1739–1749.
- Gharami K, Xie Y, An JJ, Tonegawa S, Xu B (2008) Brain-derived neurotrophic factor over-expression in the forebrain ameliorates Huntington's disease phenotypes in mice. *J Neurochem* 105:369–379.
- Altar CA, et al. (1997) Anterograde transport of brain-derived neurotrophic factor and its role in the brain. *Nature* 389:856–860.
- Baquet ZC, Gorski JA, Jones KR (2004) Early striatal dendrite deficits followed by neuron loss with advanced age in the absence of anterograde cortical brain-derived neurotrophic factor. *J Neurosci* 24:4250–4258.
- Strand AD, et al. (2007) Expression profiling of Huntington's disease models suggests that brain-derived neurotrophic factor depletion plays a major role in striatal degeneration. *J Neurosci* 27:11758–11768.
- Hébert JM, McConnell SK (2000) Targeting of cre to the Foxg1 (BF-1) locus mediates loxP recombination in the telencephalon and other developing head structures. *Dev Biol* 222:296–306.
- Soriano P (1999) Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet* 21:70–71.
- Li Y, et al. (2008) TrkB regulates hippocampal neurogenesis and governs sensitivity to antidepressant treatment. *Neuron* 59:399–412.
- Greengard P, Allen PB, Nairn AC (1999) Beyond the dopamine receptor: The DARPP-32/protein phosphatase-1 cascade. *Neuron* 23:435–447.
- Klein R, Martin-Zanca D, Barbacid M, Parada LF (1990) Expression of the tyrosine kinase receptor gene *trkB* is confined to the murine embryonic and adult nervous system. *Development* 109:845–850.
- Yan Q, Johnson EM, Jr. (1989) Immunohistochemical localization and biochemical characterization of nerve growth factor receptor in adult rat brain. *J Comp Neurol* 290:585–598.
- Luikart BW, Nef S, Shipman T, Parada LF (2003) In vivo role of truncated *trkb* receptors during sensory ganglion neurogenesis. *Neuroscience* 117:847–858.
- Zerucha T, et al. (2000) A highly conserved enhancer in the *Dlx5/Dlx6* intergenic region is the site of cross-regulatory interactions between *Dlx* genes in the embryonic forebrain. *J Neurosci* 20:709–721.
- Rose CR, et al. (2003) Truncated TrkB-T1 mediates neurotrophin-evoked calcium signaling in glia cells. *Nature* 426:74–78.
- Canals JM, et al. (2004) Brain-derived neurotrophic factor regulates the onset and severity of motor dysfunction associated with encephalimeric neuronal degeneration in Huntington's disease. *J Neurosci* 24:7727–7739.
- Bibb JA, et al. (2000) Severe deficiencies in dopamine signaling in presymptomatic Huntington's disease mice. *Proc Natl Acad Sci USA* 97:6809–6814.
- Zhou QY, Palmiter RD (1995) Dopamine-deficient mice are severely hypoactive, adipic, and aphagic. *Cell* 83:1197–1209.
- Baquet ZC, Bickford PC, Jones KR (2005) Brain-derived neurotrophic factor is required for the establishment of the proper number of dopaminergic neurons in the substantia nigra pars compacta. *J Neurosci* 25:6251–6259.
- Sanchez-Ortiz E, et al. (2012) TrkA gene ablation in basal forebrain results in dysfunction of the cholinergic circuitry. *J Neurosci* 32:4065–4079.
- Pineda JR, et al. (2005) Brain-derived neurotrophic factor modulates dopaminergic deficits in a transgenic mouse model of Huntington's disease. *J Neurochem* 93:1057–1068.
- Huang EJ, Reichardt LF (2003) Trk receptors: Roles in neuronal signal transduction. *Annu Rev Biochem* 72:609–642.
- Gorski JA, Zeiler SR, Tamowski S, Jones KR (2003) Brain-derived neurotrophic factor is required for the maintenance of cortical dendrites. *J Neurosci* 23:6856–6865.
- Luikart BW, et al. (2005) TrkB has a cell-autonomous role in the establishment of hippocampal Schaffer collateral synapses. *J Neurosci* 25:3774–3786.
- He XP, et al. (2004) Conditional deletion of TrkB but not BDNF prevents epileptogenesis in the kindling model. *Neuron* 43:31–42.
- Baydyuk M, et al. (2011) TrkB receptor controls striatal formation by regulating the number of newborn striatal neurons. *Proc Natl Acad Sci USA* 108:1669–1674.
- Fishell G (2006) Loss of a notch activity in the developing central nervous system leads to increased cell death. *Dev Neurosci* 28:517.
- Matsuda W, et al. (2009) Single nigrostriatal dopaminergic neurons form widely spread and highly dense axonal arborizations in the neostriatum. *J Neurosci* 29:444–453.
- Malkovska I, Kernie SG, Parada LF (2006) Differential expression of the four untranslated BDNF exons in the adult mouse brain. *J Neurosci Res* 83:211–221.
- Lush ME, Li Y, Kwon CH, Chen J, Parada LF (2008) Neurofibromin is required for barrel formation in the mouse somatosensory cortex. *J Neurosci* 28:1580–1587.