

UCLA

UCLA Previously Published Works

Title

Substance Use Homophily Among Geosocial Networking Application Using Gay, Bisexual, and Other Men Who Have Sex With Men

Permalink

<https://escholarship.org/uc/item/65v7v8nr>

Journal

Archives of Sexual Behavior, 44(7)

ISSN

0004-0002

Author

Holloway, Ian W

Publication Date

2015-10-01

DOI

10.1007/s10508-015-0581-6

Peer reviewed



HHS Public Access

Author manuscript

Arch Sex Behav. Author manuscript; available in PMC 2016 October 01.

Published in final edited form as:

Arch Sex Behav. 2015 October ; 44(7): 1799–1811. doi:10.1007/s10508-015-0581-6.

Substance Use Homophily Among Geosocial Networking Application Using Gay, Bisexual and Other Men Who Have Sex With Men

Ian W. Holloway, MSW, MPH, PhD¹

¹Department of Social Welfare, Luskin School of Public Affairs, University of California, Los Angeles, Los Angeles, CA

Abstract

Geosocial networking applications (GSN apps) represent important virtual contexts in which gay, bisexual and other men who have sex with men (MSM) seek affiliation. These apps allow users to create and view public profiles, send photos and text messages, and connect with other users based on shared interests and geographic proximity. The present study examined substance use homophily among a sample of 295 MSM recruited via a popular GSN app. Comparisons of social network members met via GSN app versus elsewhere and associations between both individual and network characteristics and recent binge drinking, marijuana use and illicit substance use were explored using bivariate tests of association and multivariate logistic regression analyses. High rates of recent binge drinking (59 %), marijuana use (37 %) and illicit substance use (27 %) were observed among participants. GSN app use greater than one year and showing naked chest or abs in a profile picture were positively associated with recent illicit substance use. In multivariate analyses, the strongest predictors of binge drinking (AOR = 3.81; 95 % CI = 1.86–7.80), marijuana use (AOR = 4.12; 95 % CI = 2.22–7.64) and illicit substance use (AOR = 6.45; 95 % CI = 3.26–12.79) were the presence of a social network member who also engaged in these behaviors. Social network interventions that target binge drinking, marijuana use and illicit substance use may be delivered via GSN apps to reduce the prevalence of substance use and related risks among MSM in these virtual contexts.

Keywords

MSM; networks; technology; binge drinking; substance use

Introduction

Substance use and misuse is a major public health problem among gay, bisexual and other men who have sex with men (hereafter MSM). Alcohol and recreational drug use are highly prevalent in this population (Cochran, Ackerman, Mays, & Ross, 2004; Stall et al., 2001) and have been associated with other health issues including sexual risk behaviors (Celentano

Corresponding Author: Ian W. Holloway, MSW, MPH, PhD, Department of Social Welfare, Luskin School of Public Affairs, University of California, Los Angeles, 3250 Public Affairs Building, Box 951656, Los Angeles, CA 90095-1656, Phone: 310-825-7840, Fax: 310-206-7564, holloway@luskin.ucla.edu.

et al., 2006; Greenwood et al., 2001; Hirschfield, Remien, Humberstone, Walavalkar, & Chiasson, 2004; Operario et al., 2006). Attempts to explain the association between substance use and sexual risk include alcohol- or drug-induced disinhibition (Colfax et al., 2004; Drumright et al., 2006; Semple, Patterson, & Grant, 2002), erotic arousal (Schilder, Lampinen, Miller, & Hogg, 2005), and increased length of sexual sessions leading to greater opportunities for engagement in risk behaviors (Guss, 2000). However, the ways in which MSM meet substance use and sexual partners and the contexts in which substances are used remain underexplored. Understanding the contexts of MSM's substance use (i.e., where and with whom substances are used) may inform the development of tailored substance use and sexual risk reduction interventions for MSM.

MSM are avid Internet users (Groves, Breslow, Newcomb, Rosenberger, & Bauermeister, 2014) and many use the Internet (e.g., websites, chatrooms) to seek sex and substance use partners virtually (Bauermeister, Giguere, Carballo-Diequez, Ventuneac, & Eisenberg, 2010; Halkitis, Fischgrund, & Parsons, 2005; Kubicek, Carpineto, McDavitt, Weiss, & Kipke, 2011). New technologies, such as geosocial networking applications (GSN apps) targeting MSM (e.g., *Grindr*, *Scruff*, *Jack'd*), have also emerged as important avenues through which MSM meet and maintain relationships with sex partners (Landovitz et al., 2013; Rice et al., 2012). These apps allow users to identify proximity of other users in real time, a feature that may increase engagement in risk behaviors (Winetrobe, Rice, Bauermeister, Petering, & Holloway, 2014). As relatively new technologies, much remains unknown about the relationship between GSN apps and MSM's social networks, whose composition is associated with MSM's sexual risk behavior (Berry, Raymond, & McFarland, 2007; Miller, Serner, & Wagner, 2005; Smith, Grierson, Wain, Pitts, & Pattison, 2004). The present study sought to document the social networks of MSM using a popular GSN app and to understand associations between individual and social network characteristics and substance use among GSN app-using MSM in Los Angeles, CA.

Literature Review

Substance Use and Abuse Among MSM

Substance use and sexual risk behaviors among MSM are interconnected and contribute to health disparities among MSM (Stall & Purcell, 2000). MSM are at increased risk for both substance use and substance abuse (Kipke et al., 2007a; Moon, Fornili, & O'Briant, 2007), including the use of alcohol and marijuana (Russell, Driscoll, & Truong, 2002), cocaine, ecstasy and other club drugs (Kipke, et al., 2007b). In a study of 172 MSM recruited online, 49 % endorsed using club drugs (defined as crystal methamphetamine, ecstasy, poppers, cocaine and Viagra) and of those, 51 % used two or more at the same time and 25 % used three or more at the same time (Fernandez et al., 2005). Reviews on substance use and sexual risk behavior among MSM have demonstrated positive associations between alcohol use (Woolf & Maisto, 2009; Shuper, Joharchi, Irving, & Rehm, 2009), erectile dysfunction drugs (e.g., Viagra, Cialis) (Romanelli & Smith, 2004; Swearingen & Klausner, 2005), methamphetamine (Shoptaw & Reback, 2007; Halkitis, Parsons, & Stirratt, 2001) and sexual risk. However, the relationship between substance use and sexual risk behavior has been complicated by variations in how substance use is classified (e.g., "alcohol or drug use",

“multi-drug use” or “other drug use”) and the time period under which substance use is measured (e.g., globally, situationally or at the event-level) (Leigh & Stall, 1993). A review of event-level measurement of substance use and sexual risk undertaken by Waverly Vosburgh and colleagues (2012) demonstrated consistent associations between binge alcohol use and sexual risk behaviors and methamphetamine use and sexual risk behaviors.

Studies of MSM in Los Angeles have documented high rates of substance use. For example, a comparative study of young MSM across 7 U.S. urban areas found that in Los Angeles recent substance use (in the past 6 months) included alcohol use (87 %), illicit drug use (67 %), “upper”/amphetamine use (32 %), and cocaine use (16 %). Of note was that 28 % of MSM reported using 3 or more different drugs in the past 6 months (Thiede et al., 2003). Another study compared substance use and HIV risk among MSM in Chicago and Los Angeles by serostatus and found that among HIV-positive men there were significantly higher rates of Viagra use in Los Angeles (Carey et al., 2009). Also, methamphetamine use was higher among men in Los Angeles compared to Chicago regardless of serostatus. A more recent study of young MSM in Los Angeles (ages 18–24 at baseline) found that 40 % reported frequent binge drinking, 40 % had ever used club drugs (defined as cocaine, crystal/methamphetamine, ecstasy, poppers, GHB, Ketamine, and other forms of speed) and 22 % were frequent or heavy cigarette smokers (Kipke et al., 2007a; Kipke et al., 2007b; Holloway et al., 2012). The association between alcohol use and sexual risk behavior, in particular, has been shown to vary across development and to be dependent upon the context in which it is used by MSM (Mustanski, 2008; Newcomb, 2013; Venable et al., 2004).

Social Networks and Substance Use Among MSM

Over the past two decades, there has been increasing interest in the ways in which social networks influence health behaviors (Smith & Christakis, 2008). Social networks refer to groups of individuals who are connected through personal relationships. Within social networks, members may influence another member’s behavior based on social comparison, social sanctions and rewards, socialization, and information exchange (Fisher, 1988; Latkin et al., 1995). Social network analysis allows researchers to quantitatively document how individuals (i.e., egos) are connected to network members (i.e., alters) and the ways in which processes, such as social support and social influence, are transmitted through networks (Christakis & Fowler, 2008; Berkman & Glass, 2000). Network structure (e.g., size, density) and composition (e.g., proportion of types of social ties – family, friends, etc.) have been shown to influence health behaviors in a variety of populations (Smith & Christakis, 2008; Valente, 2010), including MSM (Smith, Grierson, Wain, Pitts, & Pattison, 2004; Peterson, Rothenberg, Kraft, Beeker, & Trotter, 2009; Tobin & Latkin, 2008).

Social network structure and composition have been linked to substance use. Latkin and colleagues (1995; 2003) demonstrated that social networks, norms, and HIV risk behaviors were linked among urban drug users at risk for HIV, finding that network density and size of drug subnetworks were associated with frequency of drug injection (Latkin, Mandell, Vlahov, Oziemkowska, & Celentano, 1995; Latkin, Forman, Knowlton, & Sherman, 2003). Their results also provided ways to intervene with specific social ties to improve norms around condom use. Other mechanisms through which social networks influence health

include social support to promote coping; engagement and interaction on a particular topic; exposure to new ideas, technologies, and access to other individuals or resources that could be potentially harmful or beneficial (e.g., connections to be able to obtain illegal drugs versus harm reduction strategies).

The notion that norms and behavior are “contagious” among social ties has gained momentum in studying the direct impact social networks have on a number of health behaviors, including binge drinking (Reifman, Watson, & McCourt, 2006), marijuana use (Kobus, & Henry, 2009), and illicit substance use (Schroeder et al., 2001). Social network homophily refers to the clustering of similar individuals within networks (McPherson, Smith-Lovin, & Cook, 2001). Studies of smoking and obesity have demonstrated that health behaviors often cluster in social networks (Christakis & Fowler, 2007; Christakis & Fowler, 2008). Egocentric network studies of young MSM have demonstrated that the presence of network members who engage in sexual risk behaviors is associated with greater sexual risk behavior among participants themselves (Amirkhanian et al., 2006; Tucker et al., 2012; Kapadia et al., 2013). However, to our knowledge, no studies have examined substance use homophily among MSM who use geosocial networking applications.

Geosocial Networking Applications and Substance Use Among MSM

Technology has been recognized as an important avenue for implementing risk behavior prevention and health promotion among MSM, including young MSM (Allison et al., 2012; Holloway et al., 2014). Several studies published in recent years have focused on the use of GSN apps among MSM with a particular focus on motivations for GSN app use (Grosskopf, LaVasseur, & Glaser, 2014; Van De Wiele & Tong, 2014), sexual risk behaviors (Beymer et al., 2014, Rice et al., 2012; Lehmler & Ioegeer, 2014; Winetrobe, Rice, Bauermeister, Petering, & Holloway, 2014), HIV testing (Rendina, Jimenez, Grov, Ventuneac, & Parsons, 2014), and the acceptability of varied HIV prevention strategies among users (Burrell et al., 2012; Holloway et al., 2014; Landovitz et al., 2013). While none of these studies have focused explicitly on substance use, several have reported substance use prevalence among users. For example, among a sample of 146 young MSM (18–24 year old) GSN app users in Los Angeles, 64 % reported binge drinking, 35 % reported marijuana use and 26 % reported “hard drug” use (which included poppers, cocaine, heroin, methamphetamine and ecstasy) (Winetrobe, Rice, Bauermeister, Petering, & Holloway, 2014). Among an older sample of MSM (median age 25) in Los Angeles, Landovitz and colleagues (2013) found that 48 % of participants reported using drugs or alcohol during sex in the past month. Phillips and colleagues (2014) found that MSM who used GSN apps to look for other men in the past year were one and a half times more likely to have used non-injection drugs, including crystal methamphetamine, painkillers and poppers, compared to MSM who did not use GSN apps for partner seeking. To our knowledge, there are no published studies documenting correlates of substance use behaviors among GSN app-using MSM.

Present Study

Given high rates of substance use and abuse among MSM, the important influence of social networks in determining substance use among MSM, and the emergence of new GSN apps to facilitate social networking among MSM, the present study sought to understand the

relationship between GSN app use, social network characteristics and substance use among MSM. Specifically, the following research questions were addressed: (1) “What is the composition of the social networks of GSN app-using MSM?” and (2) “Which individual and social network factors are associated with alcohol, marijuana and illicit substance use in this population?”

Method

Participants

MSM were recruited from two neighborhoods with large populations of gay and bisexually identified men in Los Angeles, CA: West Hollywood and Long Beach. Individuals were eligible to participate if they were users of a popular GSN app and had not previously participated in the study. Utilizing the geo-location feature of the GSN app, research assistants created their own profiles to recruit GSN app users who were within a seven-mile radius of West Hollywood and Long Beach, CA. From August 8, 2011 and October 3, 2011, GSN app users between the ages of 18–24 were recruited (young MSM). From December 5, 2011 and January 3, 2012, GSN app users 25 years of age and over were recruited (older MSM). The recruiters’ profiles contained the study institution’s name and identified the recruiters as researchers; their profile pictures were of the research assistant or a stock photo.

Procedures

Participants were randomly selected based on their location at the time of recruitment. On the GSN app, profiles are organized by geo-location, with the first profiles being closest in proximity to the user. Users appeared on a grid displaying four profile photos in each row and continued for all users within a seven-mile range. Potential participants were selected using a randomization number chart displaying numbers between 1 and 4, to match the app’s profile display. Randomly selected persons were sent a message providing information about the study. Interested participants received a link and unique log-in code to an anonymous, online survey, which took approximately 20–30 minutes to complete. Upon completion, participants received a \$25 downloadable gift card to either iCard or Amazon.com. For every user who was approached, his distance from the recruiter was recorded. Recruiters were available to answer respondents’ questions and to provide minor technical support through the GSN app’s chat feature. Recruitment occurred between 9 a.m. and 8 p.m. on weekdays. Overall, 11.95 % of the men approached via GSN app text message completed the survey resulting in a total sample of 295 participants. All study procedures were approved by the Institutional Review Board of the University of Southern California. Secondary data analysis for the present study was approved by the Institutional Review Board of the University of California, Los Angeles.

Measures

The self-report survey was used to obtain a range of information from participants, including demographic characteristics, GSN app use characteristics, substance use and social network characteristics.

Demographics—Participants were asked to identify their age in years, race/ethnicity (African American, Latino/Hispanic, white, Asian, Native Hawaiian or other Pacific Islander, American Indian or Alaska Native, mixed race, other race). Native Hawaiian/Pacific Islander, American Indian/Alaska Native, and other race were then collapsed to form one “Other” race category. Participants also reported highest level of education (less than high school, high school graduate or GED, some college, 4 year college/university degree, master’s degree or professional degree, and doctorate); current employment status (not currently working, currently working); sexual identity (gay, bisexual, heterosexual, questioning, queer, other); whether they were out (i.e., had “disclosed having sex with other men”) to parents, brothers/sisters, or other family; and their relationship status out of nine options, which were then collapsed to reflect whether the participant was single or not.

GSN App Use—Participants were asked how often they logged on to the GSN app (less than five times per day, five or more times per day); how long ago they started using the GSN app (less than one year, greater than one year); if their profile picture showed their face; what naked body parts were visible in their GSN app profile picture, which was dichotomized to reflect whether their profile picture showed their naked chest or abs or not; what time of day they usually logged on to the GSN app (before midnight, after midnight); and whether they used the GSN app both on weekdays and weekends. Participants were also asked whether they used the GSN app “to find people to drink or use drugs with” and whether “the last time [they] used the GSN app, was it during or immediately after [they] had been drinking alcohol or using drugs?”

Substance use—To assess binge drinking, participants were asked a question adapted from the Centers for Disease Control and Prevention Behavioral Risk Surveillance System (BRSS), “During the past 30 days, [have you] had 5 or more drinks of alcohol in a row (i.e., within a couple of hours) at least once?” To assess illicit substance use, participants were asked to report on the frequency of their past 30 day use of marijuana, poppers, heroin, methamphetamine and ecstasy (0 days, 1 or 2 days, 3 to 5 days, 6 to 9 days, 10 to 19 days, 20 to 29 days, All 30 days). Responses were subsequently collapsed and dichotomized to reflect at least one episode of marijuana use and one episode of recent illicit drug (i.e., poppers, heroin, methamphetamines, ecstasy) use.

Social Networks—A single-item egocentric name generator asked participants to list their top five closest social network members (i.e., alters) using the following prompt: “The next several questions are about the most important people that you regularly communicate with on a social basis. These are people that you interact with, either through face-to-face contact or via the Internet or cell phone and could be family members, friends, sex-partners, co-workers or anyone else who is important to you. Based on this criteria, we ask that you please list the five people you interact with the most and/or who are most important to you in the space provided below.” Participants gave a first name or nickname for each alter; last names were not gathered to preserve confidentiality of nominated alters. Next, participants were asked to describe their relationship to each alter (e.g., life partner/husband, boyfriend, lover/sex partner/hook-up, family, friend, coworker, other) and the age, race/ethnicity, and sexual orientation of that alter. Alters described as husband, boyfriend, lover, sex partner

and hook-up were grouped to represent “intimate partners”. In addition, participants were asked to report whether they had known each network alter for more than a year, whether they provided the participant with emotional support (i.e., “anyone who you can go to if you have an important problem to discuss about your personal life”), and whether they provided the participant with instrumental support (i.e., “anyone who you could borrow \$100 from if you needed it”).

Alters’ substance use was assessed by asking participants to select any of their alters who had engaged in recent alcohol, marijuana or illicit substance use. Specifically, participants were asked the following questions: (1) “In the past month, who has drunk alcohol to the point of drunkenness?”; (2) “In the past month, who smoked marijuana, pot, or weed?”; (3) “In the past month, who used meth, crystal, or Tina?”; (4) “In the past month, who used cocaine?”; and (5) “In the past month, who used heroin?” All responses were scored dichotomously. An additional dichotomous item representing illicit drug use (methamphetamines, cocaine, heroin) was created. Participants were also asked which of their network members would object to them “drinking to the point of drunkenness,” “smoking marijuana” or using any of the other substances named above.

Statistical Analysis

Bivariate tests of association (i.e., chi square, independent sample t-test) determined associations between individual level variables (i.e., demographic variables, GSN app use) and whether the participant had included a GSN app-met partner in his network. Associations between individual and social network characteristics and each of the three substance use outcomes (i.e., binge drinking, marijuana use, illicit substance use) were determined using chi square tests of association. In social network analyses, family member alters were excluded from the total sample of alters because a primary aim of this analysis was to determine differences between GSN app-met and non-GSN app-met alters and family members were unlikely to be met through the GSN app. Due to the large number of tests that were conducted, we employed the false discovery rate controlling procedure described by Benjamini and Hochberg (1995). Multivariate logistic regressions were also performed to simultaneously test for associations between individual- and network-level factors and substance use outcomes. All data were analyzed using SAS 9.2 (2011).

Results

Individual Characteristics

Table 1 presents demographic characteristics, GSN app use and substance use behaviors for the total sample (N = 295) by whether the participant included a GSN app-met partner in his network (N = 62) or not (N = 226). In general, participants were young (mean age = 25.56), educated (i.e., had completed some college or more at the time of the study) (88.82 %), single (81.36 %), gay-identified (90.14 %), and out to their families (81.69 %). White men made up the largest racial/ethnic group (51.53 %); smaller percentages of the sample were Latino/Hispanic (23.39 %), Asian (10.17 %), and Black/African American (4.41 %). Most were employed (70.85 %). About half of the sample reported using the GSN app for over one year (46.53 %), logged on five or more times per day (50.69 %) and logged on after

midnight (52.20 %). Less than one third showed their naked chest or abs in their profile picture (29.62 %), approximately 10 % used the GSN app to find alcohol or substance using partners and approximately 8 % reported using the GSN app immediately after using alcohol or drugs. Over half reported recent binge drinking (59.32 %); over a third reported recent marijuana use (36.81 %); over a quarter reported using one or more illicit drug in the past month (26.44 %) or being under the influence of alcohol or drugs during their last sexual encounter (26.96 %). Additionally, over three-quarters of the sample had at least one network member who drank alcohol to the point of drunkenness (77.14 %), slightly more than half had at least one network member who used marijuana (55.00 %), and less than a quarter had at least one network member who used any illicit substances (22.92 %).

A number of statistically significant differences emerged between participants who included a GSN app-met partner in their social network and those who did not (Table 1). A smaller percentage of participants with a GSN app-met partner in their social network were single compared to those without a GSN app-met partner in their network (72.58 % versus 84.07 %, $\chi^2 = 4.28$, $p < 0.05$). Additionally, a greater percentage of participants with a GSN app-met partner in their network reported having sex under the influence of drugs or alcohol with a GSN app-met partner versus those who did not include a GSN app-met partner in their social network (32.29 % versus 17.04 %, $\chi^2 = 7.31$, $p < 0.01$). Finally, a greater percentage of participants with a GSN app-met partner in their social network also had at least one person who drinks to the point of drunkenness included in their social network compared to those who did not include a GSN app-met partner in their social network (86.67 % versus 74.55 %, $\chi^2 = 3.93$, $p < 0.05$).

Social Network Characteristics

Excluding family members, MSM nominated a total of 1,239 alters, which included friends (76.42 %), intimate partners (15.96 %), co-workers (3.65 %) and others (2.76 %). A greater percentage of GSN app-met partners were male (98.80 % vs. 64.22 %, $\chi^2 = 41.49$, $p < 0.001$), LGBT-identified (98.84 % vs. 60.67 %, $\chi^2 = 50.09$, $p < 0.001$), and intimate partners (52.87 % vs. 13.23 %, $\chi^2 = 94.00$, $p < 0.001$). More non-GSN app-met partners were friends (78.64 % vs. 43.68 %, $\chi^2 = 54.29$, $p < 0.001$) had known participants for longer than one year (78.56 % vs. 27.59 %, $\chi^2 = 111.59$, $p < 0.001$) and provided emotional (65.09 % vs. 42.53 %, $\chi^2 = 17.71$, $p < 0.001$) and instrumental support (54.16 % vs. 41.18 %, $\chi^2 = 5.34$, $p < 0.02$) to participants. There were no statistically significant differences in substance use behavior or objecting to substance use between alters met via GSN app vs. alters met elsewhere (Table 2).

Substance Use Outcomes

Binge Drinking—Bivariate associations between individual characteristics, social network characteristics and substance use outcomes are presented in Table 3. Only two individual-level characteristics were associated with binge drinking: White race/ethnicity and having completed at least some college or university ($p < 0.05$). At the network level, GSN app use and instrumental support were also associated with binge drinking ($p < 0.05$). Additionally, the proportion of alters who objected to participants' drinking to the point of drunkenness, as well as whether alters drank alcohol to the point of drunkenness, smoked marijuana and used

an illicit substance were associated with binge drinking ($p < 0.05$). In multivariate analyses (not shown) greater educational attainment (AOR = 0.51; 95 % CI = 0.29–0.91) and the proportion of network members who would object to the participant getting drunk (AOR = 0.34; 95 % CI = 0.13–0.92) were protective against binge drinking. The strongest predictor of participants' recent binge drinking was having at least one alter who drinks alcohol to the point of drunkenness in their network (AOR = 3.81; 95 % CI = 1.86–7.80).

Marijuana Use—The only individual-level predictor associated with recent marijuana use was GSN app use for more than one year ($p < 0.05$). At the social network level, recent marijuana use was associated with knowing their network members for greater than one year and having network members who provide instrumental support ($p < 0.05$). Recent marijuana use was negative associated with the proportion of network members who objected to them smoking marijuana and positively associated with whether network members drank alcohol to the point of drunkenness, smoked marijuana, and used an illicit substance ($p < 0.05$). In multivariate analyses, the proportion of network members who objected to smoking marijuana was protective against recent marijuana use (AOR = 0.18, 95 % CI = 0.06–0.56). A greater proportion of network members who provided instrumental support increased the odds of recent marijuana use (AOR = 2.52; 95 % CI = 1.04–6.12) as did having at least one alter who smokes marijuana (AOR = 4.12; 95 % CI = 2.22–7.64).

Illicit Drug Use—At the individual level, age and white race/ethnicity were both associated with recent illicit substance use ($p < 0.05$). Additionally, GSN app use greater than one year and displaying their naked chest or abs in their GSN app profile picture were associated with recent illicit substance use ($p < 0.05$). At the network-level, having older, male, white, gay-identified and GSN app-met alters in one's network were also associated with having used an illicit drug in the past 30 days, as was having network members with whom participants “hook up” in their network. Having alters who provided instrumental support was also associated with recent illicit substance use ($p < 0.05$). Recent illicit substance use was negatively associated with the proportion of alters who objected to using illicit substance and was positively associated with whether alters drank to the point of drunkenness and used illicit substances ($p < 0.05$). The only statistically significant predictor of participants' illicit substance use in multivariate analyses was having at least one alter who uses any illicit substance in their social network (AOR = 6.45; 95 % CI = 3.26–12.79).

Discussion

This study is among the first to examine the social networks of GSN app-using MSM in relation to substance use. Similar to other samples of MSM in Los Angeles, rates of binge drinking, marijuana use and illicit substance use were high (Kipke et al., 2007a; Thiede et al., 2003), demonstrating the need for increased substance use/misuse prevention with MSM. Our first research question sought to identify the composition of the social networks of GSN app using MSM. Results indicate diversity in the composition of GSN app users' networks, which are comprised of friends, intimate partners, family members (although excluded for this analysis), co-workers and others. While results indicate that a small subsample of MSM used the GSN app to find partners with whom to drink alcohol and use substances (approximately 10 %), large percentages of the GSN app-users had network

members who engaged in binge drinking, marijuana use and illicit substance use. Results from the present study may inform network-based interventions targeting these behaviors that are staged via GSN apps, which may be especially useful in reducing substance use/misuse among MSM given the popularity and widespread use of these apps.

Over one fifth of participants in the present study included a GSN app-met alter as a member of their closest social network and statistically significantly higher percentages of GSN app-met alters belonged to participants who had used the technology for greater than one year. GSN app-met network members were more likely to be “hook ups” than social network members met elsewhere, demonstrating, as has been shown by others (Rice et al., 2012), that sexual partner seeking is a primary purpose for using the GSN app. It is important to note that 12 % of GSN app-met alters were classified as husbands, life partners or boyfriends and 44 % were classified as friends. The popular press has characterized GSN apps as platforms for casual sex seeking (Kapp, 2011; Wortham, 2013). While this is true for many GSN app-users, it is also true that primary romantic relationships and close friendships are formed via GSN apps. As such, these platforms represent important gay male social context for dating, serious relationship and friendship seeking. While larger percentages of alters providing emotional and instrumental social support were met elsewhere, over 40 % of alters met on the GSN app provided social support to participants.

Our second research question sought to understand the influence of individual and social network factors on engagement in binge drinking, marijuana use and illicit substance use. Despite the pro-social roles that GSN app-met alters held in the lives of participants in this study, there were some individual-level GSN app use patterns that emerged as correlates of substance use outcomes. Specifically, older age, higher education and white race/ethnicity were associated with greater substance use. Furthermore, using the GSN app for longer than one year was correlated with recent use of marijuana and illicit substances. Finally, displaying naked chest or abs in their GSN app profile photos was associated with illicit substance use. These findings at the individual level are similar to those of Winetrobe and colleagues (2014) who analyzed data from the young men (ages 18–24) in this sample and found that those who had used the GSN app longer and displayed sexualized profile photos were more likely to have engaged in unprotected anal intercourse with their last GSN app-met partner. Taken together, these findings may indicate heightened risk behaviors for a subset of GSN app-users. However, the addition of network data presented here suggests that for substance use outcomes this heightened risk is primarily function of social network dynamics, rather than individual-level risk behaviors, as the strongest predictors of binge drinking, marijuana use and illicit substance use were presence of a social network member who also engaged in these behaviors.

Social network results from this study emphasize the importance of network influence on substance use, as has been shown by others. Participants with alcohol users, marijuana users and illicit substance users in their networks were more likely to engage in those behaviors themselves. Described by McPherson (2001), this phenomenon is referred to as homophily and refers to the idea that like individuals are more likely to affiliate with others who are similar to themselves (i.e., “birds of a feather flock together”). These results are intuitive given the powerful influence of peer norms and the fact that substance use among MSM

often occurs in groups and social settings (Halkitis et al., 2011). Because our data is cross sectional it is impossible for us to determine whether substance users seek out other substance users for inclusion into their networks or whether peer influence processes operate within networks to promote greater substance use. Further longitudinal research with GSN app-using MSM should attempt to elucidate these processes over time, as their clarification may have important implications for preventing substance abuse and related risks among GSN app-using MSM.

A key finding for the development of risk behavior prevention and harm reduction interventions is the protective nature of having a social network member who objects to binge drinking, marijuana use and illicit substance use. Network studies of young MSM have demonstrated lower levels of sexual risk behaviors among participants who have a “pro-social” peer as part of their social network. Tucker and colleagues (2012) found that homeless YMSM were less likely to engage in unprotected sex and had fewer sex partners if their networks included fewer sex partners and if the majority of their network members were not heavy drinkers (i.e., greater than 50% had not drunk alcohol to the point of drunkenness in the past 30 months). Social network based interventions that take into consideration the composition of MSM networks, leverage ties to peers who do not engage in risk behaviors and promote diffusion of peer norms that are discouraging of substance use and related risks may be especially effective (Valente, 2012).

Limitations

Limitations of the present research should be taken into consideration when interpreting findings. As mentioned above, this was a cross-sectional study, making it impossible to determine the direction of our findings or causality. Furthermore, our study did not include a comparison group of non-GSN app users, making it impossible to determine whether substance use is comparable between users and non-users. All data were collected via self-report, which may underestimate or overestimate the actual prevalence of binge drinking, marijuana use and illicit substance use. Enabling MSM to take the survey using their private computers, smartphones or tablets likely contributed to veracity in reporting of behaviors; however, it is impossible to know for certain. In addition, we gathered data on the attitudes of participants’ social network members from the participants themselves, without consulting nominated network members. It is quite possible that participants’ perceptions of the attitudes and behaviors of their network members do not correspond to actual behavior. However, several studies have demonstrated the importance of perception of peer behavior on the actual risk behavior of participants, so this may be less of a concern for many readers (Peterson et al., 2009; Hart & Peterson, 2004).

The recruitment methods used (i.e., active recruitment, recruitment of younger MSM and older MSM at different times, and limiting recruitment to just one popular GSN app in Los Angeles, CA) could also have introduced bias and limit the generalizability of our results, especially if there are underlying differences between those who were available or not available to participate in the study during the recruitment periods, for example. The large numbers of white, well-educated participants is likely a function of recruiting in affluent gay neighborhoods in Los Angeles (i.e., West Hollywood, Long Beach). Future research with

racial/ethnic minority and low-income GSN app users is needed. Additionally, because contact with GSN app representatives could not be made, we were unable to develop a fully collaborative research approach. As noted in previous work, collaborative approaches between GSN app companies and public health researchers are warranted to fully elucidate the role that GSN apps play in the lives of MSM (Holloway et al., 2014).

Conclusions

Despite the limitations of this study, the findings may have useful implications for the formulation of substance use/misuse prevention interventions for GSN-app using MSM. First, due to the widespread use of these technologies and high prevalence rates of binge drinking, marijuana use and illicit substance use among GSN app users, it appears that these platforms may be well suited for the dissemination of prevention and harm reduction messaging. Our previous research suggests high levels of acceptability of HIV prevention interventions delivered via smartphone among young MSM (Holloway et al., 2014); however, no research has been conducted on the acceptability of substance abuse prevention interventions in these contexts. Formative research is needed on the feasibility of app-based prevention interventions with this population as little is known about how MSM may respond to these efforts. Banner ads and push notifications can be easily purchased on GSN apps to remind users of relevant information regarding substance use and accompanying sexual risk behaviors (e.g., using substances during sex increases one's risk for contracting HIV); however, it is unclear how effective these approaches may be in capturing the attention of MSM using GSN apps.

Network homophily among GSN app-using MSM points to the possibility of network based interventions that can promote peer norms to discourage substance misuse. Facebook-delivered popular opinion leader models to increase HIV testing have been successful with racially and ethnically diverse MSM previously (Young et al., 2014). Similar interventions implemented via GSN apps may be protective against substance misuse among GSN app-using MSM. Finally, those interested in promoting health behaviors among GSN app-using MSM must bear in mind that GSN apps are used for much more than substance use and/or casual sex partner seeking; instead, GSN apps represent important social contexts for affiliation between gay, bisexual and other MSM. Recognition of the important social role that GSN apps play in the lives of MSM will assist public health practitioners to develop interventions that promote positive affiliation while reducing high-risk behaviors.

Acknowledgments

This work was also supported by the Center for HIV Identification, Prevention, and Treatment (CHIPTS) NIMH grant MH58107; the UCLA Center for AIDS Research (CFAR) grant 5P30AI028697; and the National Center for Advancing Translational Sciences through UCLA CSTI Grant UL1TR000124. The content is solely the responsibility of the authors and does not necessarily represent the official views of NIH. The author would like to thank Dr. Eric Rice for his leadership on the parent study, Diane Tan for her assistance with manuscript preparation, and the community advisory board who helped guide the research.

References

- Allison S, Bauermeister JA, Bull S, Lightfoot M, Mustanski B, Shegog R, Levine D. The intersection of youth, technology, and new media with sexual health: moving the research agenda forward. *Journal of Adolescent Health*. 2012; 51(3):207–212. [PubMed: 22921129]
- Amirkhanian YA, Kelly JA, Kirsanova AV, DiFranceisco W, Khoursine RA, Semenov AV, Rozmanova VN. HIV risk behaviour patterns, predictors, and sexually transmitted disease prevalence in the social networks of young men who have sex with men in St Petersburg, Russia. *International journal of STD & AIDS*. 2006; 17(1):50–56. [PubMed: 16409680]
- Bauermeister JA, Giguere R, Carbello-Diequez A, Ventuneac A, Eisenburg A. Perceived risks and protective strategies employed by young men who have sex with men (YMSM) when seeking online sexual partners. *Journal of Health Communication*. 2010; 15(6):679–690. [PubMed: 20812127]
- Benjamini, Y.; Hochberg, Y. *Journal of Royal Statistical Society, Series B*. New York: Oxford University Press; 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing; p. 289-300.
- Berkman, LF.; Glass, T. Social integration, social networks, social support, and health. In: Berkman, LF.; Kawachi, I., editors. *Social Epidemiology*. 2000. p. 137-173.
- Berry M, Raymond HF, McFarland W. Same race and older partner selection may explain higher HIV prevalence among black men who have sex with men. *AIDS*. 2007; 21(17):2349–2350. [PubMed: 18090287]
- Beymer MR, Weiss RE, Bolan RK, Rudy ET, Bourque LB, Rodriguez JP, Morisky DE. Sex on demand: geosocial networking phone apps and risk of sexually transmitted infections among a cross-sectional sample of men who have sex with men in Los Angeles County. *Sex Transmitted Infections*. 2014 Epub online ahead of print. 10.1136/sextrans-2013-051494
- Burrell ER, Pines HA, Robbie E, Coleman L, Murphy RD, Hess KL, Gorbach PM. Use of the location-based social networking application GSN APP as a recruitment tool in rectal microbicide development research. *AIDS and Behavior*. 2012; 16(7):1816–1820. [PubMed: 22851153]
- Carey JW, Mejia R, Bingham T, Ciesielski C, Gelaude D, Herbst JH, Stall R. Drug use, high-risk sex behaviors, and increased risk for recent HIV infection among men who have sex with men in Chicago and Los Angeles. *AIDS and Behavior*. 2009; 13(6):1084–1096. [PubMed: 18498049]
- Celentano DD, Valleroy LA, Sifakis F, MacKellar DA, Hylton J, Thiede H. Young Men's Survey Study Group. Associations between substance use and sexual risk among very young men who have sex with men. *Sexually Transmitted Diseases*. 2006; 33(4):265–271. [PubMed: 16434886]
- Centers for Disease Control and Prevention. [Accessed on May 29, 2015] Behavioral Risk Factor Surveillance System: Questionnaires. 2014. Retrieved from <http://www.cdc.gov/brfss/questionnaires/index.htm>
- Christakis NA, Fowler JH. The spread of obesity in a large social network over 32 years. *New England Journal of Medicine*. 2007; 357(4):370–379. [PubMed: 17652652]
- Christakis NA, Fowler JH. The collective dynamics of smoking in a large social network. *New England Journal of Medicine*. 2008; 358(21):2249–2258. [PubMed: 18499567]
- Cochran SD, Ackerman D, Mays VM, Ross MW. Prevalence of non-medical drug use and dependence among homosexually active men and women in the US population. *Addiction*. 2004; 99(8):989–998. [PubMed: 15265096]
- Colfax G, Vittinghoff E, Husnik MJ, McKirnan D, Buchbinder S, Koblin B, Coates TJ. Substance use and sexual risk: a participant- and episode-level analysis among a cohort of men who have sex with men. *American Journal of Epidemiology*. 2004; 159(10):1002–1012. [PubMed: 15128613]
- Drumright LN, Little SJ, Strathdee SA, Slymen DJ, Araneta MR, Malcarne VL, Gorbach PM. Unprotected anal intercourse and substance use among men who have sex with men with recent HIV infection. *Journal of Acquired Immune Deficiency Syndrome*. 2006; 43(3):344–350.
- Fernández MI, Perrino T, Collazo JB, Varga LM, Marsh D, Hernandez N, Bowen GS. Surfing new territory: club-drug use and risky sex among Hispanic men who have sex with men recruited on the Internet. *Journal of Urban Health*. 2005; 82(1 Suppl 1):i79–88. [PubMed: 15738317]

- Fisher JD. Possible effects of reference group-based social influence on AIDS-risk behavior and AIDS prevention. *American Psychologist*. 1988; 43:914–920. [PubMed: 3214003]
- Greenwood GL, White EW, Page-Shafer K, Bein E, Osmond DH, Paul J, Stall RD. Correlates of heavy substance use among young gay and bisexual men: The San Francisco Young Men's Health Study. *Drug and Alcohol Dependence*. 2001; 61(2):105–112. [PubMed: 11137274]
- Grosskopf NA, LeVasseur MT, Glaser DB. Use of the Internet and mobile-based "apps" for sex-seeking among men who have sex with men in New York City. *American Journal of Men's Health*. 2014 Epub ahead of print. 10.1177/1557988314527311
- Grov C, Breslow AS, Newcomb ME, Rosenberger JG, Bauermeister JA. Gay and bisexual men's use of the Internet: Research from the 1990s through 2013. *American Journal of Men's Health*. 2014; 51(4):390–409.
- Guss JR. Sex like you can't even imagine: "Crystal," crack and gay men. *Journal of Gay & Lesbian Psychotherapy*. 2000; 3(3/4):105–122.
- Halkitis PN, Fischgrund BN, Parsons JT. Explanations for methamphetamine use among gay and bisexual men in New York City. *Substance Use and Misuse*. 2005; 40(9–10):1331–1345. [PubMed: 16048820]
- Halkitis PN, Parsons JT, Stirratt MJ. A double epidemic: crystal methamphetamine drug use in relation to HIV transmission. *Journal of Homosexuality*. 2001; 41(2):17–35. [PubMed: 11482426]
- Halkitis PN, Pollock JA, Pappas MK, Dayton A, Moeller RW, Siconolfi D, Solomon T. Substance use in the MSM population of New York City during the era of HIV/AIDS. *Substance Use & Misuse*. 2011; 46(2–3):264–273. [PubMed: 21303246]
- Hart T, Peterson JL. Predictors of risky sexual behavior among young African men who have sex with men. *American Journal of Public Health*. 2004; 94(7):1122–1123. [PubMed: 15226130]
- Hirshfield S, Remien RH, Humberstone M, Walavalkar I, Chiasson MA. Substance use and high-risk sex among men who have sex with men: a national online study in the USA. *AIDS Care*. 2004; 16(8):1036–1047. [PubMed: 15511735]
- Holloway IW, Rice E, Gibbs J, Winetrobe H, Dunlap S, Rhoades H. Acceptability of smartphone application-based HIV prevention among young men who have sex with men. *AIDS and Behavior*. 2014; 18(2):285–296. [PubMed: 24292281]
- Holloway IW, Traube DE, Rice E, Schrage SS, Palinkas LA, Richardson J, Kipke MD. Community and individual factors associated with cigarette smoking among young men who have sex with men. *Journal of Research on Adolescence*. 2012; 22(2):199–205. [PubMed: 22661879]
- Leigh BC, Stall R. Substance use and risky sexual behavior for exposure to HIV: Issues in methodology, interpretation, and prevention. *American Psychologist*. 1993; 48(10):1035. [PubMed: 8256876]
- Kapadia F, Siconolfi DE, Barton S, Olivieri B, Lombardo L, Halkitis PN. Social support network characteristics and sexual risk taking among a racially/ethnically diverse sample of young, urban men who have sex with men. *AIDS and Behavior*. 2013; 17(5):1819–1828. [PubMed: 23553346]
- Kapp, M. [Accessed on: July 16, 2014] Welcome to the world's biggest, scariest ga bar. *Vanity Fair*. 2011. Available at: <http://www.vanityfair.com/culture/features/2011/05/GSNapp-201105>
- Kipke MD, Kubicek K, Weiss G, Wong C, Lopez D, Iverson E, Ford W. The health and health behaviors of young men who have sex with men. *Journal of Adolescent Health*. 2007a; 40(4):342–350. [PubMed: 17367727]
- Kipke MD, Weiss G, Ramirez M, Dorey F, Ritt-Olsen A, Iverson E, Ford W. Club drug use in Los Angeles among young men who have sex with men. *Substance Use and Misuse*. 2007b; 42(11):1723–1743. [PubMed: 17934992]
- Kobus K, Henry DB. Interplay of network position and peer substance use in early adolescent cigarette, alcohol, and marijuana use. *Journal of Early Adolescence*. 2009; 30(2):225–245.
- Kubicek K, Carpineto J, McDavitt B, Weiss G, Kipke MD. Use and perceptions of the Internet for sexual information and partners: A study of young men who have sex with men. *Archives of Sexual Behavior*. 2011; 40(4):803–816. [PubMed: 20809373]
- Landovitz RJ, Tseng CH, Weissman M, Haymer M, Mendenhall B, Rogers K, Shoptaw S. Epidemiology, sexual risk behavior, and HIV prevention practices of men who have sex with men

using GSN APP in Los Angeles, California. *Journal of Urban Health*. 2013; 90(4):729–39. [PubMed: 22983721]

- Latkin CA, Forman V, Knowlton A, Sherman S. Norms, social networks, and hiv-related risk behaviors among urban disadvantaged drug users. *Social Science & Medicine*. 2003; 56(3):465–476. [PubMed: 12570967]
- Latkin C, Mandell W, Oziemkowska M, Celentano D, Vlahov D, Ensminger M, Knowlton A. Using social network analysis to study patterns of drug use among urban drug users at high risk for hiv/aids. *Drug and Alcohol Dependence*. 1995; 38(1):1–9. [PubMed: 7648991]
- Lehmiller JJ, Iorger M. Social networking smartphone applications and sexual health outcomes among men who have sex with men. *PLoS One*. 2014; 9(1):e86603. [PubMed: 24466166]
- McPherson M, Smith-Lovin L, Cook JM. Birds of a feather: Homophily in social networks. *Annual Review of Sociology*. 2001; 27(1):415–444.
- Mustanski B. Moderating effects of age on the alcohol and sexual risk taking association: an online daily diary study of men who have sex with men. *AIDS and Behavior*. 2008; 12(1):118–126. [PubMed: 18034298]
- Miller M, Sernver M, Wagner M. Sexual diversity among black men who have sex with men in an inner-city community. *Journal of Urban Health*. 2005; 82(1):i26–i34. [PubMed: 15738323]
- Moon MW, Fornili K, O'Briant AL. Risk comparisons among youth who report sex with same sex vs. both-sex partners. *Youth and Society*. 2007; 38(3):267–284.
- Newcomb ME. Moderating effect of age on the association between alcohol use and sexual risk in MSM: evidence for elevated risk among younger MSM. *AIDS and Behavior*. 2013; 17(5):1746–1754. [PubMed: 23553348]
- Operario D, Choi KH, Chu PL, McFarland W, Secura GM, Behel S, Valleroy L. Prevalence and correlates of substance use among young Asian Pacific Islander men who have sex with men. *Prevention Science*. 2006; 7(1):19–29. [PubMed: 16435077]
- Peterson J, Rothenberg R, Kraft J, Beeker C, Trotter R. Perceived condom norms and HIV risks among social and sexual networks of young African American men who have sex with men. *Health Education Research*. 2009; 24(1):119–27. [PubMed: 18281710]
- Phillips G II, Magnus M, Kuo I, Rawls A, Peterson J, Jia Y, Greenberg AE. Use of geosocial networking (GSN) mobile phone applications to find men for sex by men who have sex with men (MSM) in Washington, D.C. *AIDS and Behavior*. 2014 Epub ahead of print. 10.1007/s10461-014-0760-9
- Reifman A, Watson WK, McCourt A. Social networks and college drinking: Probing processes of social influence and selection. *Personality and Social Psychology Bulletin*. 2006; 32(6):820–832. [PubMed: 16648206]
- Rendina HJ, Jimenez RH, Grov C, Ventuneac A, Parsons JT. Patterns of lifetime and recent HIV testing among men who have sex with men in New York city who use GSN app. *AIDS and Behavior*. 2014; 18(1):41–49. [PubMed: 23925515]
- Rice E, Holloway I, Winetrobe H, Rhoades H, Barman-Adhikari A, Gibbs J, Dunlap S. Sex risk among young men who have sex with men who use GSN app, a smartphone geosocial networking application. *Journal of AIDS & Clinical Research*. 2012; S4:1–8.
- Romanelli F, Smith KM. Recreational use of sildenafil by HIV-positive and-negative homosexual/bisexual males. *Annals of Pharmacotherapy*. 2004; 38(6):1024–1030. [PubMed: 15113986]
- Russell ST, Driscoll AK, Truong N. Adolescent same-sex romantic attractions and relationships: Implications for substance use and abuse. *American Journal of Public Health*. 2002; 92(2):198–202. [PubMed: 11818291]
- SAS Institute Inc. *Procedures Guide*. Cary, NC: SAS Institute Inc; 2011. Base SAS® 9.3.
- Schilder AJ, Lampinen TM, Miller ML, Hogg RS. Crystal methamphetamine and ecstasy differ in relation to unsafe sex among young gay men. *Canadian Journal of Public Health*. 2005; 96(5):340–343. [PubMed: 16238150]
- Schroeder JR, Latkin CA, Hoover DR, Curry AD, Knowlton AR, Celentano DD. Illicit drug use in one's social network and in one's neighborhood predicts individual heroin and cocaine use. *Annals of Epidemiology*. 2001; 11(6):389–394. [PubMed: 11454498]

- Semple SJ, Patterson TL, Grant I. Motivations associated with methamphetamine use among HIV+ men who have sex with men. *Journal of Substance Abuse Treatment*. 2002; 22(3):149–156. [PubMed: 12039618]
- Shoptaw S, Reback CJ. Methamphetamine use and infectious disease - related behaviors in men who have sex with men: implications for interventions. *Addiction*. 2007; 102(s1):130–135. [PubMed: 17493062]
- Shuper PA, Joharchi N, Irving H, Rehm J. Alcohol as a correlate of unprotected sexual behavior among people living with HIV/AIDS: review and meta-analysis. *AIDS and Behavior*. 2009; 13(6): 1021–1036. [PubMed: 19618261]
- Smith AMA, Grierson I, Wain D, Pitts M, Pattison P. Associations between the sexual behaviour of men who have sex with men and the structure and composition of their social networks. *Sexually Transmitted Infections*. 2004; 81(1):455–458. [PubMed: 15572613]
- Smith KP, Christakis NA. Social networks and health. *Annual Review of Sociology*. 2008; 34:405–429.
- Stall R, Paul JP, Greenwood G, Pollack LM, Bein E, Crosby GM, Catania JA. Alcohol use, drug use and alcohol - related problems among men who have sex with men: the Urban Men's Health Study. *Addiction*. 2001; 96(11):1589–1601. [PubMed: 11784456]
- Stall R, Purcell DW. Intertwining epidemics: A review of research on substance use among men who have sex with men and its connection to the AIDS epidemic. *AIDS & Behavior*. 2000; 4(2):181–192.
- Swearingen SG, Klausner JD. Sildenafil use, sexual risk behavior, and risk for sexually transmitted diseases, including HIV infection. *American Journal of Medicine*. 2005; 118(6):571–577. [PubMed: 15922685]
- Thiede H, Valleroy LA, MacKellar DA, Celentano DD, Ford WL, Hagan H, Torian LV. Regional patterns and correlates of substance use among young men who have sex with men in 7 US urban areas. *American Journal of Public Health*. 2003; 93(11):1915–1921. [PubMed: 14600066]
- Tobin KE, Latkin CA. An examination of social network characteristics of men who have sex with men who use drugs. *Sexually Transmitted Infections*. 2008; 84(6):420–424. [PubMed: 19028939]
- Tucker JS, Hu J, Golinelli D, Kennedy DP, Green HD, Wenzel SL. Social network and individual correlates of sexual risk behavior among homeless young men who have sex with men. *Journal of Adolescent Health*. 2012; 51(4):386–392. [PubMed: 22999840]
- Valente, TW. *Social Networks and Health: Models, Methods, and Applications*. New York, NY: Oxford Press; 2010.
- Valente TW. Network interventions. *Science*. 2012; 337(6090):49–53. [PubMed: 22767921]
- Van De Wiele C, Tong ST. Breaking boundaries: The uses and gratifications of GSN app. 2014 Epub online ahead of print. 10.1145/2632048.2636070
- Vanable PA, McKirnan DJ, Buchbinder SP, Bartholow BN, Douglas JM Jr, Judson FN, MacQueen KM. Alcohol use and high-risk sexual behavior among men who have sex with men: the effects of consumption level and partner type. *Health Psychology*. 2004; 23(5):525. [PubMed: 15367072]
- Vosburgh HW, Mansergh G, Sullivan PS, Purcell DW. A review of the literature on event-level substance use and sexual risk behavior among men who have sex with men. *AIDS and Behavior*. 2012; 16(6):1394–1410. [PubMed: 22323004]
- Winetrobe H, Rice E, Bauermeister J, Petering R, Holloway IW. Associations of unprotected anal intercourse with GSN app-met partners among GSN app-using young men who have sex with men in Los Angeles. *AIDS Care*. 2014 Epub ahead of print. 10.1080/09540121.2014.911811
- Woolf SE, Maisto SA. Alcohol use and risk of HIV infection among men who have sex with men. *AIDS and Behavior*. 2009; 13(4):757–782. [PubMed: 18236149]
- Wortham, J. [Accessed on July 16, 2014] How GSN app is changing the way we connect. *New York Times*. 2013. Available at: http://bits.blogs.nytimes.com/2013/02/10/how-GSNapp-is-changing-the-way-we-all-connect?_php=true&_type=blogs&_r=0
- Young SD, Holloway I, Jaganath D, Rice E, Westmoreland D, Coates T. Project HOPE: online social network changes in an HIV prevention randomized controlled trial for African American and Latino men who have sex with men. *American Journal of Public Health*. 2014; 104(9):1707–1712. [PubMed: 25033137]

Table 1

Demographic characteristics by whether the participants included a GSN app-met partner in their social network. (N = 295)^a

Variable	App-met partner in network (n = 62)			No app-met partner in network (n = 226)			Total (n = 295)		X ² or t Statistic	p value
	N	% or Mean (SD)	N	% or Mean (SD)	N	% or Mean (SD)	N	% or Mean (SD)		
Age	62	25.94 (6.56)	226	25.58 (6.61)	295	25.56 (6.56)			0.2404	.6240
Race/Ethnicity ^d										
White	34	54.84	116	51.33	152	51.53				
Black/AA	2	3.23	11	4.87	13	4.41				
Latino/Hispanic	13	20.97	53	23.45	69	23.39				
Asian	5	8.06	24	10.62	30	10.17				
Mixed	7	11.29	20	8.85	28	9.49				
Other	1	1.61	2	0.88	3	1.02				
Sexual Orientation ^b									1.1420	0.2852
Gay	58	93.55	201	88.94	265	90.14				
Bisexual	4	6.45	17	7.56	22	7.48				
Straight	0	0.00	0	0.00	0	0.00				
Questioning	0	0.00	4	1.78	4	1.36				
Queer	0	0.00	1	0.44	1	0.34				
Other	0	0.00	2	0.89	2	0.68				
Education ^c									0.0009	0.9755
< High school	1	1.61	4	1.77	5	1.69				
High school GED	8	12.90	18	7.96	28	9.49				
Some college	24	38.71	87	38.50	115	38.98				
4 year college	20	32.26	91	40.27	112	37.97				
Master's degree	9	14.52	21	9.29	30	10.17				
Doctorate	0	0.00	5	2.21	5	1.69				
Employed	45	72.58	158	69.91	209	70.85			0.1666	0.6831
“Out” to family members	52	83.87	184	81.42	241	81.69				
Single ^{*d}	45	72.58	190	84.07	240	81.36			4.276	0.0386
GSN app Use										

Variable	App-met partner in network (n = 62)			No app-met partner in network (n = 226)			Total (n = 295)			X ² or t Statistic	p value
	N	% or Mean (SD)	N	% or Mean (SD)	N	% or Mean (SD)					
Use 5 + times/day	35	56.45	111	49.12	146	50.69	1.0477	0.3060			
Use > 1 year*	37	59.68	97	42.92	134	46.53	5.4911	0.0191			
Chest/abs showing	21	33.87	64	28.44	85	29.62	0.6866	0.4073			
Use after midnight	35	56.45	117	51.77	152	52.78	0.4279	0.5130			
Use to find substance users*	12	19.35	17	7.52	29	10.07	7.5225	0.0061			
Use after alcohol / drug use [^]	4	6.45	19	8.41	23	7.99	--	--			
Last sex under influence of alcohol/drugs	21	33.87	55	24.55	79	29.96	2.1605	0.1416			
Recent Substance Use											
Binge drinking	35	56.45	135	59.73	175	59.32	0.2168	0.6415			
Marijuana use	19	31.15	82	37.27	106	36.81	0.7782	0.3777			
Illicit ^e drug use	20	32.26	56	24.78	78	26.44	1.4011	0.2365			
Has at least one alter who:											
Drinks alcohol to drunkenness*	52	86.67	164	74.55	216	77.14	3.9282	0.0475			
Smokes marijuana	33	53.23	121	55.50	154	55.00	0.1013	0.7503			
Uses any illicit substance	17	27.42	49	21.68	66	22.92	0.9036	0.3410			

* p < .05

** p < .01

*** p < .001

[^] Contains cells with < 5 expected counts

^a May not add up to total sample size due to missing values

^b X² test: White vs. all others

^c X² test: Gay/Homosexual vs. all others

^d X² test: Some college or university vs. all others

^e X² test: Single vs. all others

^f Includes poppers, cocaine, heroin, methamphetamine and/or ecstasy

Bivariate comparisons of GSN app-met social network members and non-GSN app-met social network members (N=1239, excludes family alters)^a

Table 2

Variable	GSN app-met (n = 87)			Non-GSN app-met (n = 1124)			Total (n = 1239)			χ ² or t Statistic	p value
	N	% or Mean (SD)	N	% or Mean (SD)	N	% or Mean (SD)	N	% or Mean (SD)			
Age	87	29.56 (8.41)	1089	28.49 (8.85)	1204	28.47 (8.81)			-1.0900	0.2763	
Male ^{***}	82	98.80	700	64.22	798	66.44			41.4900	<.0001	
Race/Ethnicity ^b									0.3846	0.5352	
White	43	49.43	581	52.91	641	52.93					
Latino/Hispanic	22	25.29	273	24.86	301	24.86					
Asian	11	12.64	117	10.66	128	10.57					
Pacific Islander	2	2.30	10	0.91	12	0.99					
AI/Alaska Native	0	0.00	7	0.64	8	0.66					
Mixed	2	2.30	34	3.10	37	3.06					
Other	1	1.15	3	0.27	4	0.33					
LGBT ^{***}	85	98.84	668	60.67	770	63.37			50.0939	<.0001	
Relationship											
Intimate ^{***}	46	52.87	148	13.23	197	15.96			94.0038	<.0001	
Life partner [^]	2	2.30	20	1.79	22	1.78			--	--	
Boyfriend ^{**}	10	11.49	46	4.11	57	4.62			9.9387	0.0016	
Hook-up ^{***}	34	39.08	82	7.33	118	9.56			91.2882	<.0001	
Friend ^{***}	38	43.68	880	78.64	943	76.42			54.2864	<.0001	
Coworker [^]	1	1.15	44	3.93	45	3.65			--	--	
Other [^]	2	2.30	32	2.86	34	2.76			--	--	
Known > 1 year ^{***}	24	27.59	883	78.56	921	74.33			111.5889	<.0001	
Uses GSN app ^{***}	87	100.00	269	24.75	354	29.77			154.54	<.0001	
Support provided Emotional ^{***}	37	42.53	727	65.09	783	63.56			17.7088	<.0001	
Instrumental [*]	35	41.18	593	54.16	642	53.15			5.3368	0.0209	

Objects to:

Variable	GSN app-met (n = 87)		Non-GSN app-met (n = 1124)		Total (n = 1239)		X ² or t Statistic	p value
	N	% or Mean (SD)	N	% or Mean (SD)	N	% or Mean (SD)		
Getting drunk	9	10.34	178	16.45	188	15.71	2.2343	0.1350
Marijuana	15	17.44	229	21.34	248	20.96	0.7287	0.3933
Illicit substance ^c	74	85.06	929	82.65	1103	82.82	0.3287	0.5665
Gets drunk	45	52.94	464	42.45	523	43.51	3.5361	0.0600
Smokes marijuana	16	18.39	257	23.60	277	23.18	1.2263	0.2681
Uses Illicit substance ^c	8	9.20	96	8.54	106	8.56	0.0441	0.8337

* p < .05

** p < .01

*** p < .001

[^] Contains cells with < 5 expected counts

^a Sample sizes may vary due to missing data

^b X² test: White vs. all others

^c Includes cocaine, heroin and/or methamphetamine

Table 3
 Bivariate comparisons of recent substance use outcomes by individual- and network-level characteristics

	Binge Drinking % (n)	Marijuana Use % (n)	Illicit Substance Use % (n)
<i>Individual-level (n = 295)</i>			
25 years of age or older	33.71 (59)	33.96 (36)	47.44 (37)*
White race/ethnicity	60.00 (105)*	55.66 (59)	66.67 (52)*
Gay/Homosexual Sexual Orientation	90.86 (159)	86.79 (92)	92.31 (72)
Some college or university	32.00 (56)*	41.51 (44)	38.46 (30)
Employed	72.57 (127)	70.75 (75)	74.36 (58)
"Out" to family	82.29 (144)	84.91 (90)	84.62 (66)
Single	84.00 (147)	76.42 (81)	79.49 (62)
<i>GSN app Use</i>			
Use 5 + times/day	48.57 (85)	48.11 (51)	52.56 (41)
Use > 1 year	49.14 (86)	39.62 (42)*	57.69 (45)*
Chest/abs showing	31.03 (54)	31.43 (33)	41.56 (32)*
Use after midnight	52.57 (92)	53.77 (57)	52.56 (41)
Includes app-met partner in SN	20.59 (35)	18.81 (19)	26.32 (20)
<i>Network-level (n = 1239, excludes family alters)</i>			
Alter met via GSN app	7.07 (51)	6.24 (27)	8.08 (27)
25 years of age or older	57.09 (423)	54.55 (246)	67.25 (230)*
Male	66.34 (477)	65.75 (286)	74.02 (245)*
White race/ethnicity	53.58 (397)	53.22 (240)	60.82 (208)*
LGBT Sexual Orientation	64.75 (474)	60.64 (265)	69.28 (230)*
<i>Relationship with Alter</i>			
Intimate	15.25 (113)	15.74 (71)	18.71 (64)
Life partner/husband	1.62 (12)	1.55 (7)	2.63 (9)
Boyfriend	4.59 (34)	5.76 (26)	3.80 (13)
Hook-up	9.04 (67)	8.43 (38)	12.28 (42)*
Friend	77.33 (573)	75.17 (339)	73.39 (251)
Coworker	3.91 (29)	4.88 (22)	4.09 (14)

	Binge Drinking % (n)	Marijuana Use % (n)	Illicit Substance Use % (n)
Other	2.70 (20)	1.77 (8)	1.75 (6)
Known > 1 year	75.03 (556)	78.27 (353)*	76.02 (260)
Uses GSN app	32.87 (237)*	31.55 (136)	37.69 (124)*
Support provided			
Emotional support	64.91 (481)	65.70 (293)	60.83 (205)
Instrumental support	56.87 (418)*	59.73 (261)*	61.18 (197)*
Alter would object to you:			
Getting drunk	10.80 (77)*	–	–
Smoking marijuana	–	9.43 (40)*	–
Using illicit substance ^d	–	–	77.78 (266)*
Substance use among alters			
Drinks alcohol to Drunkenness	54.84 (402)*	51.47 (228)*	52.69 (176)*
Smokes marijuana	27.25 (197)*	36.16 (158)*	26.77 (87)
Any illicit substance use ^d	11.88 (88)*	12.42 (56)*	21.35 (73)*

* p < .05

^d Includes cocaine, heroin and/or methamphetamines

Note: Independent sample t-test used for continuous predictors; chi-square or Fisher's exact used for categorical predictors. Multiple comparisons adjusted using the Benjamini and Hochman (1995) procedure.