

UC Irvine

UC Irvine Previously Published Works

Title

Ecosystem-level controls on root-rhizosphere respiration

Permalink

<https://escholarship.org/uc/item/6cf6x8tb>

Journal

New Phytologist, 199(2)

ISSN

0028-646X

Authors

Hopkins, Francesca
Gonzalez-Meler, Miquel A
Flower, Charles E
[et al.](#)

Publication Date

2013-07-01

DOI

10.1111/nph.12271

Supplemental Material

<https://escholarship.org/uc/item/6cf6x8tb#supplemental>

Peer reviewed

Research review

Ecosystem-level controls on root-rhizosphere respiration

Author for correspondence:

Miquel A. Gonzalez-Meler

Tel: +1 312 355 3928

Email: mmeler@uic.edu

Received: 29 December 2012

Accepted: 12 March 2013

Francesca Hopkins¹, Miquel A. Gonzalez-Meler², Charles E. Flower²,
Douglas J. Lynch², Claudia Czimczik¹, Jianwu Tang³ and Jens-Arne Subke⁴

¹Department of Earth System Science, University of California, Irvine, CA 92697, USA; ²Ecology and Evolution Group, Department of Biological Sciences, University of Illinois at Chicago, Chicago, IL 60607, USA; ³The Ecosystems Center, Marine Biological Laboratory, 7 MBL St, Woods Hole, MA 02543, USA; ⁴Biological and Environmental Sciences, School of Natural Sciences, University of Stirling, Stirling, FK9 4LA, UK

New Phytologist (2013) **199**: 339–351

doi: 10.1111/nph.12271

Key words: autotrophic, global change, gross primary productivity (GPP), heterotrophic, nitrogen (N), rhizosphere, root respiration, soil respiration.

Summary

Recent advances in the partitioning of autotrophic from heterotrophic respiration processes in soils in conjunction with new high temporal resolution soil respiration data sets offer insights into biotic and environmental controls of respiration. Besides temperature, many emerging controlling factors have not yet been incorporated into ecosystem-scale models. We synthesize recent research that has partitioned soil respiration into its process components to evaluate effects of nitrogen, temperature and photosynthesis on autotrophic flux from soils at the ecosystem level. Despite the widely used temperature dependence of root respiration, gross primary productivity (GPP) can explain most patterns of ecosystem root respiration (and to some extent heterotrophic respiration) at within-season time-scales. Specifically, heterotrophic respiration is influenced by a seasonally variable supply of recent photosynthetic products in the rhizosphere. The contribution of stored root carbon (C) to root respiratory fluxes also varied seasonally, partially decoupling the proportion of photosynthetic C driving root respiration. In order to reflect recent insights, new hierarchical models, which incorporate root respiration as a primary function of GPP and which respond to environmental variables by modifying C allocation belowground, are needed for better prediction of future ecosystem C sequestration.

Introduction

The majority of the 140 Pg of carbon (C) fixed annually by terrestrial gross primary productivity (GPP) passes through soil, with 8–52% respired back to the atmosphere by the rhizosphere (Lambers *et al.*, 2008), and 23–83% of GPP ending up in plant tissue C (DeLucia *et al.*, 2007) which is ultimately returned to the atmosphere as CO₂ via the activity of decomposer organisms. Soil respiration is a primary process governing net C sequestration in terrestrial ecosystems (Valentini *et al.*, 2000); this soil flux is likely to change as a result of the impacts of global change factors such as nitrogen (N) deposition, warming, and rising CO₂ concentrations (Houghton *et al.*, 1998). Because of its importance to the global C cycle, soil respiration has been intensively studied for several decades (Luo & Zhou, 2006), yet the ability to predictively model soil respiration and its components through time and space is still elusive (Chen *et al.*, 2011; Leuzinger & Thomas, 2011).

Spatial and temporal heterogeneity of soil respiration is a major challenge to ecosystem scientists, making it difficult to interpret

effects of climate variables on soil CO₂ efflux (Davidson & Janssens, 2006). This variability stems from the multitude of different sources and pathways for the production of CO₂ throughout the soil profile, each of which is controlled to a varying extent by biotic and abiotic drivers (Janssens *et al.*, 2001; Gonzalez-Meler *et al.*, 2004; Davidson & Janssens, 2006; Fig. 1). A fundamental distinction between soil CO₂ sources in terms of C turnover and temporal dynamics is that of CO₂ derived from the decomposition of organic matter by soil fauna and microbial organisms (fungi, bacteria and protozoans), from that originating from plant roots and rhizospheric organisms (including symbiotic mycorrhizal fungi). These two generic CO₂ sources are commonly referred to as heterotrophic and autotrophic soil CO₂ flux (Högberg *et al.*, 2001), with some debate about how heterotrophic organisms associated with autotrophic C supply should be categorized (Högberg *et al.*, 2006).

Autotrophic contributions represent a substantial component of seasonal soil CO₂ efflux (Gonzalez-Meler & Taneva, 2005), and may even represent the majority of soil respired CO₂ during periods




Fig. 1 Soil respiration divided into conceptual and modeled components. Autotrophic respiration consists of growth respiration (R_G) and maintenance respiration (R_M), which are derived from new photosynthate and plant carbon (C) storage pools. Heterotrophic respiration consists of respiration from litter decomposition (R_L) and soil organic matter decomposition (R_S). It is increasingly being recognized by experimentalists that a significant portion of soil respiration is neither strictly autotrophic, as it passes through microorganisms such as mycorrhizal fungi, nor heterotrophic, as substrates are coming directly from roots (they do not pass as tissue through the litter pool to decomposers). This respiration has been termed 'rhizosphere respiration,' and is thought to include respiration of mycorrhizal fungi (R_{MF}) and respiration from soil decomposer organisms that is 'primed' by additions of labile substrates in root exudates (R_P). Rhizosphere processes are identified in gray as they are not implicitly incorporated into ecosystem-scale models. Colored boxes represent different methods of isolating different conceptual respiration fluxes. Addition of C isotope measurements to measurements of these fluxes can provide information on the source pool and/or turnover time of C and the combination of methods can theoretically resolved any source of soil CO_2 efflux.

of high productivity (Subke *et al.*, 2006; Gomez-Casanovas *et al.*, 2012). As the temporal dynamics of this flux component are linked to aboveground C assimilation and C transport to roots via the plants' phloem, it is not exclusively controlled by soil temperature or moisture (Högberg *et al.*, 2001; Trueman & Gonzalez-Meler, 2005). As the peak growing season generally coincides with the warmest time of year (at least in ecosystems not limited by soil moisture), high apparent responses of autotrophic soil CO_2 efflux with temperature may have been reported. Indeed, much of the observed variation in soil respiration, and its temperature sensitivity in particular, is probably attributable to variation in substrate availability in space and time (Davidson *et al.*, 2006; Subke & Bahn, 2010). Nevertheless, temperature is still used as the primary driver for descriptive and prognostic models of soil respiration (Davidson *et al.*, 2006 and references therein).

The availability of labile C such as from root exudation or fresh litter input sustains microbial populations capable of decomposing slower-turnover soil organic matter (SOM) (Fontaine *et al.*, 2004; Guenet *et al.*, 2012). This leads to soil C priming (or more specifically a 'rhizosphere priming effect' (RPE); Kuzyakov, 2010), which is a natural process in soils. The dynamics of RPEs are tightly linked to autotrophic C supply, but may affect older soil C pools previously assumed to be stable. Current ecosystem models do not include RPEs, and a better understanding of belowground C

dynamics, including roots, rhizosphere and associated organisms, is required.

Most ecosystem-scale C cycle models include the effect of substrate supply on respiration through its relationship to GPP, either indirectly through a fixed ratio of net primary productivity (NPP) to GPP (e.g. the CASA model; Field *et al.*, 1995), or directly through allocation schemes that partition GPP to growth, storage and respiration (e.g. the Biome-BGC model; Thornton & Rosenbloom, 2005). Heterotrophic respiration is determined by litter and SOM pool sizes (which are controlled by GPP at multiple time-scales), and the effect of temperature and soil moisture on decomposition rates. Autotrophic respiration, however, is not always explicitly represented. In models based in units of NPP, autotrophic respiration is an emergent process that represents the remainder of GPP (Gifford, 2003). By contrast, models such as Biome-BGC explicitly represent autotrophic respiration as the sum of growth respiration and maintenance respiration, which in turn depend on plant C pool size and on the magnitude of GPP (Thornton & Rosenbloom, 2005; Fig. 1).

In the hierarchy of biological processes, NPP-based models were designed to represent processes at a higher level than that of physiology (Gifford, 2003); however, physiological processes may be an important part of the response of the C cycle to climate change. In particular, decades-long data sets of ecosystem respiration are increasingly being used to parameterize C cycle sensitivity to climate change (Mahecha *et al.*, 2010). Such data sets will also be an important part of validating the process and climate response in land models (Luo *et al.*, 2012). The incorporation of explicit allocation schemes, storage pools, and substrate supply as drivers of autotrophic respiration in C cycle models will probably help to explain disparities between models and data (Keenan *et al.*, 2012; Richardson *et al.*, 2013). Furthermore, the role of acclimation of soil respiration components to forcing factors is largely ignored in large-scale models (Leuzinger & Thomas, 2011).

Here we review the evidence for effects of N, temperature, and photosynthate supply (including stored C) on root respiration in order to identify their importance as drivers of autotrophic respiration for ecosystem-scale models. We place particular emphasis on photosynthetic controls on root respiration assessed using recent ^{14}C and ^{13}C data, which show patterns of stored C use in root respiration, and elaborate on a continuous ^{13}C whole-ecosystem tracer that gives powerful insights into effects of current photosynthesis on soil autotrophic respiration.

Techniques to measure root respiration at the ecosystem level

Soil respiration is a combination of root respiration, microbial respiration, and possibly root-associated mycorrhizal respiration. In order to determine effects of environmental change and substrate supply on root-rhizosphere respiration alone, the components of soil respiration must be separated, and their dynamics analyzed independently (Taneva & Gonzalez-Meler, 2011). However, partitioning of soil respiration has been notoriously difficult (Hanson *et al.*, 2000; Kuzyakov & Larionova, 2005; Trumbore, 2006), and there is no ideal method to separate soil respiration into

its functional components. Current methods isolate a different set of respiration processes (Fig. 1), having corresponding strengths and weaknesses.

In root exclusion experiments (Fig. 1), a physical barrier prevents roots from entering excluded soils, and this method is widely used to obtain *in situ* measurements of root respiration by subtracting respiration from the root-free plot from respiration from the control. Trenching, soil coring, root in-growth cores, and gap analysis are common root exclusion methods. Trenching requires digging a narrow trench around an open plot (1–10 m²), inserting plastic sheets to block root in-growth, and then refilling trenched area with soil to replicate original conditions (Tang *et al.*, 2005a). Soil coring consists of placing polyvinyl chloride (PVC) pipes (*c.* 10–30 cm diameter) into soils to block the in-growth of roots (Bond-Lamberty *et al.*, 2011) and can be modified through the use of mesh bags that allow mycorrhizas to grow in but exclude roots (41 µm), or exclude both mycorrhizas and roots (1 µm) (Heinemeyer *et al.*, 2007). The gap analysis method subtracts heterotrophic respiration measured in natural ecosystem gaps (no roots) from respiration measured in vegetated areas (with roots) (e.g. Tang & Baldocchi, 2005).

Drawbacks to the root exclusion method center on the assumption that heterotrophic respiration remains unchanged after elimination of root inputs (Ewel *et al.*, 1987; Bowden *et al.*, 1993; Epron *et al.*, 1999; Tang *et al.*, 2005a). This assumption has been challenged (Trueman & Gonzalez-Meler, 2005), as root exclusion methods alter substrate supply to microbes by decreasing labile root inputs to microbial activity and initial increases in substrate supply from dead roots (Subke *et al.*, 2006). Root exclusion zones also block root water uptake, resulting in higher soil water contents compared with 'control' areas, thereby confounding differences in microbial activity between experimental treatments (Heinemeyer *et al.*, 2012).

Root excision measures root respiration directly as excised intact roots are directly picked from soils, and their CO₂ efflux measured on a per unit mass basis (Burton *et al.*, 2008; Fig. 1). Flux rates show considerable variation with this method, with significant impacts of CO₂ concentration in measuring cuvettes, time elapsed between root excision and CO₂ flux measurement, and whether roots were separated under 'wet' or 'dry' conditions (Subke *et al.*, 2006). Cuvettes attached to excavated surface roots without excision may circumvent some of these problems (Chen *et al.*, 2010). Root biomass is needed to calculate autotrophic respiration from soils but the few roots sampled for respiration may not be representative of the entire fine-root community.

Tree girdling can estimate root respiration from soils without disturbance of the physical integrity of roots (Högberg *et al.*, 2001; Fig. 1). Tree girdling terminates the supply of recent photosynthate to roots through the phloem while not disturbing water flow through the xylem (Högberg *et al.*, 2001, 2009; Subke *et al.*, 2011; Levy-Varon *et al.*, 2012; Liu *et al.*, 2012). This technique has problems similar to those of trenching methods; however, root respiration and root-rhizosphere effects could be maintained by carbohydrates stored in roots, and root water uptake is maintained in treatment plots.

Isotopic methods are often noninvasive and require either the use of natural C tracers or the experimental addition of isotopically altered CO₂ (¹³CO₂ or ¹⁴CO₂) to separate root respiration from

microbial respiration. Isotopic labeling experiments include free air CO₂ enrichment (FACE), where CO₂ added to the ecosystem is fossil-derived (Pataki *et al.*, 2003; Taneva *et al.*, 2006), and pulse-chase experiments, where a one-time supply of isotopically distinct CO₂ is tracked to belowground tissues and soil efflux (Brüggemann *et al.*, 2011). Tracer addition techniques measure the oxidation of labeled C by roots but not necessarily total autotrophic flux because contributions of unlabeled stored C to root respiration are not accounted for (Fig. 1). Low-level ¹⁴C pulse labeling studies can potentially circumvent this problem on subannual time-scales by comparing ¹⁴CO₂ values before and after the pulse (Carbone & Trumbore, 2007).

In certain conditions, natural differences in both ¹³C (Hanson *et al.*, 2000; Kuzyakov & Larionova, 2005; Gomez-Casanovas *et al.*, 2012) and ¹⁴C (Borken *et al.*, 2006; Czimczik *et al.*, 2006; Schuur & Trumbore, 2006) can be utilized to partition autotrophic and heterotrophic components. Natural abundance ¹⁴C measurements allow the determination of the mean age of C pools and fluxes (Trumbore, 2000). This technique takes advantage of a spike in atmospheric ¹⁴C content resulting from nuclear weapons testing in the early 1960s and subsequent decline following the banning of testing in 1963 (Levin *et al.*, 2010). The current annual change in atmospheric ¹⁴C values (5‰; Levin *et al.*, 2010) is similar to or greater than the precision at which ¹⁴C measurements can currently be made (2–5‰; Southon & Santos, 2004) and therefore useful to separate current-year C from stored C fueling root respiration.

Some problems are associated with isotope analyses. Analyses of ¹³C are sensitive to post-photosynthetic fractionation of respiratory substrates which may challenge the quantification of C sources and their ages to root respiration (Lynch *et al.*, 2013). In the field, isotopic fractionation during diffusive non-steady-state gas transport may induce errors in quantification of plant respired CO₂ from soils (Risk *et al.*, 2012). For ¹⁴C analyses, cost and sampling size are major drawbacks. Also, the time resolution at natural abundance levels can encompass 1–3 yr (error of ¹⁴C measurement with accelerator mass spectrometry (AMS)). In addition, isotopic techniques may be unable to distinguish between decomposition originating from the turnover of very short-lived roots (heterotrophic) and maintenance respiration from roots (autotrophic), as these two sources are composed of C of similar ages (Fig. 1).

These methods remain imperfect approximations of the continuum of physiological processes that govern soil respiration and the conceptual components represented in process models (Fig. 1). Nevertheless, their application has begun to disentangle the often opposing effects of global change factors on autotrophic and heterotrophic respiration. Future progress will be made by further studies that combine these partitioning techniques (which can potentially identify multiple sources of soil respired CO₂) in global change manipulation experiments to quantify these fluxes and their biotic and abiotic controls.

Nitrogen effects on root respiration

At the tissue level, specific respiration rates are often proportional to N content (Ryan, 1991, 1995; Reich *et al.*, 1996; Ryan *et al.*, 1996; Gonzalez-Meler *et al.*, 2004), as N constitutes most of the

maintenance costs of tissues (Bouma *et al.*, 1994). This is the case for fine roots, as increased N content correlates well with increased specific respiration rates (Burton *et al.*, 2002; Chen *et al.*, 2010; Jia *et al.*, 2010; Wang *et al.*, 2010). Therefore, root N content has been used as a proxy to estimate total fine-root respiration (Jia *et al.*, 2011). However, scaling this relationship to ecosystems is problematic, as fine roots have many-fold differences in N content within a diameter class (Iversen, 2010), affecting the function and respiratory activity of individual roots (Lambers *et al.*, 1996; Pregitzer *et al.*, 1998; Scheurwater *et al.*, 1998; Burton *et al.*, 2000; Chapin *et al.*, 2002), and hence preventing the use of root N to estimate total root respiration in some ecosystems (Vose & Ryan, 2002). Despite these problems, the relationship between N content and respiration of fine roots has become important in C-cycling modeling to estimate autotrophic respiration from soils (Thornton & Rosenbloom, 2005).

Despite the clear relationship between N content and specific respiration rates of roots, the direction and magnitude of the response of autotrophic respiration to increased N availability (e.g. N deposition) are less certain (Smithwick *et al.*, 2013). Increased nutrient availability has been shown to both increase (Ryan *et al.*, 1996; Pregitzer *et al.*, 2000; Gough *et al.*, 2004; Jia *et al.*, 2011) and reduce root respiration (Zogg *et al.*, 1996; Maier & Kress, 2000; Janssens *et al.*, 2010; Burton *et al.*, 2012). These contrasting results are probably attributable to changes in standing crop root biomass which have been documented in CO₂ and N fertilization experiments (Matamala & Schlesinger, 2000; Iversen, 2010; Burton *et al.*, 2012). Therefore, N-driven changes in root biomass may impact autotrophic respiration from soils to a greater extent than intrinsic variations in N content (Burton *et al.*, 2012).

N deposition changes the way in which roots interact with other soil organisms, by reducing C allocation to mycorrhizas (Högberg *et al.*, 2010; Vallack *et al.*, 2012), although changes in mycorrhizal respiration may depend on the quantity of N applied (Hasselquist *et al.*, 2012). Additionally, N deposition may reduce root exudation rates and associated microbial activity (Phillips *et al.*, 2011), which may be the cause of observed changes in heterotrophic respiration rates with added N (Neff *et al.*, 2002; Nowinski *et al.*, 2009). Future research efforts are needed to examine how N-induced changes in C allocation belowground will affect not just autotrophic respiration, but associated heterotrophic processes, including the RPE. Although changes in root biomass (and thus autotrophic flux from soils) in response to global change factors are beginning to be documented, these effects are seldom included in ecosystem models, which largely predict changes in the autotrophic component of soil respiration to passively respond to changes in soil temperature.

Temperature effects on the autotrophic and heterotrophic soil C flux

There is considerable concern that an increase in mean air temperatures globally will erode C stored in soils (Davidson & Janssens, 2006; Hartley & Ineson, 2008; Bond-Lamberty & Thomson, 2010; Hopkins *et al.*, 2012) and significant research effort has been directed at distinguishing between temperature

responses of labile and stable SOM which have different importance to soil C storage (Taneva *et al.*, 2006; Hartley & Ineson, 2008; Karhu *et al.*, 2010; Taneva & Gonzalez-Meler, 2011). Predicting the rate and amount of soil C vulnerable to loss with warming has posed an important challenge for empirical studies and model predictions (Schlesinger & Andrews, 2000; Fang *et al.*, 2005; Reichstein *et al.*, 2005; Davidson & Janssens, 2006; Friedlingstein *et al.*, 2006; Hartley & Ineson, 2008). Many studies rely on the temperature sensitivity (Q_{10}) of bulk soil respiration and its components, yet these estimates are confounded by the temporally and environmentally varying role of the different processes controlling soil flux. Attempts to predict the fate of soil C stocks simply based on changes in temperature alone are clearly unrealistic, as shifts in climate will have profound impacts on vegetation composition and activity. Consequential changes in plant C assimilation, belowground C allocation, root growth, rooting depth, and mycorrhizal associations are all likely to impact soil C inputs and mineralization rates (Fontaine *et al.*, 2007; Milcu *et al.*, 2011), changing soil C stores at long time-scales. To gain a better understanding of the temperature sensitivity of these distinct processes, it is paramount to separately assess the independent temperature responses of the autotrophic and heterotrophic components of soil respiration (Moyano *et al.*, 2008; Gomez-Casanovas *et al.*, 2012; Fig. 1).

In the absence of acclimation, kinetic theory predicts that the metabolic processes governing root and microbial respiration respond positively to warming. However, if we further partition root respiration into growth respiration and maintenance respiration (Ryan, 1990) and add respiratory acclimation, the temperature sensitivity of autotrophic respiration from soils becomes more likely to vary with season. Because root growth may vary across species and seasons, growth respiration is probably driven by phenology and fuelled by the availability of C transported from recent photosynthate or from stored C pools. Maintenance respiration of roots (which includes nutrient uptake costs) is primarily driven by tissue N concentration and influenced by temperature. Therefore, nutrient and energy demands largely determine rates of growth and maintenance respiration, suggesting that root respiration will rapidly acclimate to temperature (Atkin *et al.*, 2008). However, root respiration may appear correlated with temperature in model assessments because peak GPP and plant phenology are often correlated with temperature in most ecosystems (Subke & Bahn, 2010).

Indeed, increasing evidence that photosynthesis influences soil respiration rates (Craine *et al.*, 1999; Högberg *et al.*, 2001; Janssens *et al.*, 2001; Irvine *et al.*, 2005; Tang *et al.*, 2005b; Gomez-Casanovas *et al.*, 2012; Savage *et al.*, 2013) confirms the importance of current photosynthetic substrate for soil respiration. In addition, diel and seasonal effects on substrate supply belowground are a major confounding factor for assessments of the temperature sensitivity of soil respiration, explaining the existence of hysteresis patterns common in soil respiration data (Davidson *et al.*, 2006) and the wide range of belowground autotrophic Q_{10} values reported in the literature (Boone *et al.*, 1998; Lavigne *et al.*, 2003; Tang *et al.*, 2005a; Hartley *et al.*, 2007; Burton *et al.*, 2008). Ecosystem-scale studies are lacking, but a grassland study showed

that GPP and soil moisture interactions were the major drivers for the soil autotrophic flux at multiple time-scales, whereas GPP and temperature interactions explained variations in the heterotrophic component of soil respiration (Gomez-Casanovas *et al.*, 2012). In fact, a recent meta-analysis (using 84 manipulation studies) suggests that soil respiration may be more sensitive to changes in precipitation than to changes in temperature (Wu *et al.*, 2011). Consequently, the apparent empirical relationship between temperature and soil respiration data sets may simply be the result of temperature fluctuations co-occurring with variations in GPP on diel and seasonal time-scales influencing at least the root-rhizosphere component of soil respiration.

Recent results indicate that the RPE leads to an apparent increase in the temperature sensitivity of heterotrophic respiration (Zhu & Cheng, 2011), and the stability of supposedly recalcitrant organic matter depends strongly on how accessible it is to decomposing organisms (Dungait *et al.*, 2012), rather than on changes in temperature *per se*. The confounding phenomena of the RPE implicitly recognize the connection of plant root activity with the activity of soil heterotrophs which may affect SOM mineralization. In fact, photosynthate supply may exert a strong control on both autotrophic and heterotrophic respiration (Bahn *et al.*, 2009; Taneva & Gonzalez-Meler, 2011; Gomez-Casanovas *et al.*, 2012), highlighting the intermediate area between strictly autotrophic and heterotrophic processes, such as photosynthate-derived, substrate-supply-driven respiration by mycorrhizal associates (Drigo *et al.*, 2010). The emerging paradigm for heterotrophic respiration is that substrate supply (GPP) influences not only the component of respiration most closely associated with the rhizosphere (Phillips *et al.*, 2011) but also the overall heterotrophic component of soil respiration after some time lag (Gomez-Casanovas *et al.*, 2012).

For the purpose of prognostic modeling, therefore, Q_{10} and temperature dependences of soil respiration and its flux components are not meaningful proxies, as the magnitude of autotroph-derived soil CO_2 efflux is determined by the amount of C assimilated by plants and subsequent allocation to roots (Subke & Bahn, 2010). In order to model this particular flux component, it would be necessary to provide estimates of the assimilation and allocation patterns of vegetation under globally changed conditions, including overall warmer temperatures, changes in precipitation, higher CO_2 concentrations, and (regionally) altered light conditions. A better process understanding of the availability of photosynthetic products (including reserves) at diurnal and seasonal time-scales (Lloyd & Taylor, 1994; Högberg *et al.*, 2001; Tang *et al.*, 2005b; Davidson *et al.*, 2006; Taneva & Gonzalez-Meler, 2011) as a predictor of autotrophic respiration rates (and accounting for acclimation and adaptation of root respiration to soil thermal changes; Rachmilevitch *et al.*, 2007; Atkin *et al.*, 2008) provides a more meaningful basis for a realistic projection of ecosystem responses to environmental change. Combinations of soil warming and air warming experiments provide an opportunity to tease apart the confounding influence of temperature on photosynthesis and autotrophic respiration from soils. Soil warming would initially (before soil nutrient availability is significantly changed by warming) illustrate the effects of temperature on soil processes without the similar temperature

effect on GPP (other than natural variation), and therefore direct effects of temperature on root respiration could be untangled from natural variation in air temperature influencing GPP. Recent advances have been made using time-scale analysis of high-frequency soil respiration data sets to relate temporal patterns of soil respiration to biophysical drivers (Vargas *et al.*, 2011); however, the most progress will be made by combining multiple measurement techniques (including partitioning techniques reviewed here) and manipulations (soil and air warming combinations) with process-based models (including photosynthesis, C allocation, and plant functional type) to link to the ecosystem scale.

^{14}C and ^{13}C measurements of root respiration to infer contributions of stored C

Forest trees appear to use stored carbohydrates that are several years old and would buffer changes in GPP or belowground C allocation over days or seasons. Combined with manipulative experiments, ^{14}C approaches have the potential to inform studies about the use of stored C in ecosystem metabolism and its role in the resistance of forests to stress, disturbance or responses to climate change. Here, we review ^{14}C and some ^{13}C evidence highlighting the use of stored carbohydrates in forests during the growing season, among growing seasons and over successional stages.

Seasonal variability

Data from temperate deciduous and coniferous forest in the northeastern USA (Borken *et al.*, 2006; S. Trumbore, unpublished) suggest a decline in the mean C age of root respiration ($\Delta\Delta^{14}\text{C}$, the difference in $\Delta^{14}\text{C}$ between a sample and the atmosphere; Trumbore, 2006) between June and September (Fig. 2). In a *Quercus* spp. dominated woodland, Cisneros-Dozal (2005) observed a decline in the $\Delta^{14}\text{C}$ - CO_2 emitted by excised roots ($\Delta^{14}\text{C}_{\text{root respiration}}$) throughout the growing season (May to July, and March to May). Analysis of the ^{14}C isotopic composition of starch pools extracted from live roots at this oak site suggested that stored C contributed 70% (before leaf-out) to < 10% (August)




Fig. 2 The $\Delta\Delta^{14}\text{C}$ of root respiration over the growing season for Howland forest (91% conifer, 8% hardwood (solid triangles)) and Harvard forest (mixed site 86% hardwood, 14% conifer (open squares); hemlock (*Tsuga canadensis*) roots only (open diamonds)). $\Delta\Delta^{14}\text{C}$, which gives an estimate of C age, is calculated as the difference between $\Delta^{14}\text{C}$ of root respiration and the $\Delta^{14}\text{C}$ of the atmosphere in the year of sampling. The error bars are the standard deviation of the mean of replicate samples, or the propagated error of the $\Delta\Delta^{14}\text{C}$ calculation, taking the larger value.




Fig. 3 Fraction of stored carbon (C) from past gross primary productivity (GPP) contributing to root respiration at a free air CO₂ enrichment (FACE) site in a *Liquidambar styraciflua* plantation at Oak Ridge, TN, USA. Storage C was determined using an isotope mixing model taking advantage of the cessation of a long-term ecosystem ¹³C isotopic tracer (1998–2009) that was incorporated during CO₂ fumigation. Results demonstrate that C stored during fumigation was utilized for root respiration in 2010 (storage C that is at least 1 yr old) and in 2011 (storage C that is at least 2 yr old). Data were extracted and redrawn from Lynch *et al.* (2013) with the authors' permission.

of the total root-respired CO₂ (Cisneros-Dozal, 2005). In a temperate *Liquidambar styraciflua* plantation, Lynch *et al.* (2013) found that carbohydrates stored from GPP in a given year can substantially fuel root respiration for at least two subsequent years (Fig. 3). These results indicate that stored carbohydrates can be mobilized under different scenarios, particularly under conditions of low GPP.

Interannual variability

Stored carbohydrates are used by roots to balance energy demands during periods of low GPP (e.g. prior to leaf expansion), in response to stressful conditions (e.g. drought, high vapor deficit and/or high temperature), or when C allocation belowground is limited. Czimczik *et al.* (2006) found a greater use of stored C in a dry and warm year, compared with the following normal year, where root respired C had a $\Delta^{14}\text{C}$ value consistent with the current atmosphere. Schuur & Trumbore (2006) also found a small difference in $\Delta^{14}\text{C}$ (c. 2 yr) in respiration of *Picea mariana* roots between two consecutive sampling years. The age of stored C can be multiple years old (Figs 2–4), so increasing storage reserves during years of high productivity can ameliorate current photosynthetic supply shortages during years when conditions are not favorable.

Stand age Forest productivity declines with stand age, possibly as a result of increases in respiration costs over constant GPP (DeLucia *et al.*, 2007; Goulden *et al.*, 2011), suggesting that the respiratory demand for stored carbohydrates may be higher as forest stands age. There is some evidence that stand age may affect the $\Delta^{14}\text{C}$ value of root respired CO₂. For instance, *P. mariana* (black spruce) trees at least 40 yr old were respiring stored C up to 6 yr old, but young *P. mariana* trees from adjacent sites involved almost no stored carbohydrates during the normal respiration of roots (Czimczik *et al.*, 2006).




Fig. 4 Minimum and maximum $\Delta^{14}\text{C}$ for available measurements at one tropical, two temperate, and two boreal forest sites. Sites are: Harvard Forest (Borken *et al.*, 2006); Thompson (Czimczik *et al.*, 2006); Delta Junction (Schuur & Trumbore, 2006); and Santarém (Trumbore *et al.*, 2006). Mean C age is quantified by $\Delta^{14}\text{C}$ (the difference between $\Delta^{14}\text{C}$ of root respiration and the $\Delta^{14}\text{C}$ of the atmosphere in the year of sampling). The error bars are the standard deviation of the mean of replicate samples.

Seasonal, ontogenic or environmental conditions affect the amount of stored carbohydrates used to support autotrophic respiration from soils. Therefore, the demand for stored carbohydrates will increase as growing season length shortens with latitude. A synthesis of current ¹⁴C data provides some indication of a latitudinal trend in minimum and maximum $\Delta^{14}\text{C}$ (Fig. 4). Specifically, at the tropical site, where there is very little seasonality, there is a small difference between the minimum and maximum $\Delta^{14}\text{C}$ in root respiration values. Maximum $\Delta^{14}\text{C}$ appears to increase with latitude, suggesting a larger role for stored C pools at higher latitudes, which experience greater seasonality and more temperature variations affecting GPP. By contrast, there is no regular trend in minimum $\Delta^{14}\text{C}$, suggesting a period when surplus photosynthate is available, and is used exclusively in root respiration and allocated to storage for use in subsequent years.

Radiocarbon evidence indicates the ability of plants to draw on carbohydrate reserves to support root respiration when energy demand by the root system surpasses the substrate availability supplied by current photosynthesis. An important emerging outcome from these isotope approaches is that stored carbohydrates buffer the variability of current photosynthate supply to roots during ontogeny, phenology or environmental conditions that lead to low GPP, suggesting that root respiration may be limited by substrate availability. Therefore, current photosynthesis may play a more important role in regulating autotrophic respiration from soils than temperature (Street *et al.*, 2011; Gomez-Casanovas *et al.*, 2012; Subke *et al.*, 2012).

Current photosynthate supply effects on root respiration: the Duke FACE case study

Carbon isotope tracers and pulse-chase experiments have been used across a wide range of vegetation types to investigate the way in

which C assimilated by plants is exchanged with the atmosphere and the soil biota (Taneva *et al.*, 2006; Bowling *et al.*, 2008; Subke *et al.*, 2012; Figs 2, 3). The speed of transfer between photosynthesis and soil efflux differs between vegetation types, with belowground transfer reported in the range of hours to days (Högberg *et al.*, 2001; Trueman & Gonzalez-Meler, 2005; Carbone & Trumbore, 2007; Bahn *et al.*, 2009; Mencuccini & Holta, 2010) to months and years for more stable soil pools (Taneva *et al.*, 2006). There are further reported differences between species, probably linked to phloem anatomy, assimilation rates or life history (Dannoura *et al.*, 2011). In addition, whole-plant or ecosystem-level isotopic experiments can help unravel the transfer of assimilated C to heterotrophic microbes in the soil. These experiments have shown that root-associated mycorrhizal fungal species have high uptake rates of plant assimilated C and contribute greatly to soil respiration (Subke *et al.*, 2012). Pulse labeling experiments have enabled isotope-specific model descriptions (Ohlsson, 2011) to estimate patterns of C allocation, C use efficiency (CUE) or mobilization of stored C for root respiration (Bahn *et al.*, 2009; Street *et al.*, 2011; Subke *et al.*, 2012). These experiments indicate that photosynthesis has a substantial effect on root respiration, with stored carbohydrates buffering respiratory demand for photosynthate when assimilation rates are not sufficient (Figs 2–4).

At the Duke FACE experiment, a *Pinus taeda* plantation was exposed to elevated CO₂ starting in 1996 using a constant isotopically depleted source of ¹³CO₂ (see Supporting Information Notes S1; Figs S1–S4). The C turnover rates of leaves and soil pools at the site are relatively slow and in the range of 3–20+ yr (Schlesinger & Lichter, 2001; Matamala *et al.*, 2003; Taneva *et al.*, 2006; Feng *et al.*, 2010). In addition, the continuous isotope label has been used to separate the autotrophic from the heterotrophic components of soil respiration (Andrews *et al.*, 1999; Luo *et al.*, 2001a; Moore *et al.*, 2008; Taneva & Gonzalez-Meler, 2011; see Table S1). This plethora of data along with the extensive ecosystem fluxes and biomass pools measured at the site (DeLucia *et al.*, 1999; Matamala & Schlesinger, 2000; Luo *et al.*, 2001b; Hui & Luo, 2004; Taneva *et al.*, 2006; Table S2; Fig. S5) offer a unique opportunity to investigate the effects of GPP on root respiration. This is particularly true during the first year of the experiment, when the newly labeled treatment C was marginally incorporated into litter and soil pools (Table 1, Fig. 5).

In the first year of fumigation, GPP increased by 27% in forests grown at elevated CO₂ (Table 1). Although trees exposed to elevated CO₂ conditions allocated more C belowground in absolute terms than those exposed to ambient CO₂, trees in both treatments invested roughly 40% of GPP to belowground tissues and root-rhizosphere respiration and maintained similar values for CUE (i.e. NPP-to-GPP ratio; Table 1). As a result, increases in belowground autotrophic respiration in plants grown at elevated CO₂ appeared to be proportional to both increases in GPP and increases in root biomass. These results are consistent with successive observations made at the site, where root respiration had a clear diel and seasonal pattern not solely driven by temperature and moisture variations (Taneva & Gonzalez-Meler, 2011). Further evidence that photosynthesis may exert a larger leverage on root respiration than soil moisture or temperature has

Table 1 Carbon partitioning to above- and belowground components during the first year of fumigation (1997) in a pine forest plantation exposed to ambient and elevated CO₂ at the Duke free air CO₂ enrichment (FACE) study

	Ambient	Elevated	Elevated/ ambient
Gross primary production	1277	1624	1.27
Total aboveground allocation (including coarse roots)	768	941	1.22
Aboveground respiration	335	391	1.17
Aboveground biomass	433	549	1.27
Total belowground allocation	509	683	1.34
Belowground biomass	105	150	1.42
Belowground respiration	403	533	1.32
Carbon use efficiency	0.42	0.43	1.02

Units are in g C m⁻² and are the average of three replicated rings. Data were extracted from Andrews *et al.* (1999), DeLucia *et al.* (1999), Matamala & Schlesinger (2000), Luo *et al.* (2001a,b), Hamilton *et al.* (2002), Matamala *et al.* (2003) and Taneva *et al.* (2006).




Fig. 5 The relationship between daily gross primary productivity (GPP) and daily respiration of labeled new photosynthetic carbon (C) by root-rhizosphere respiration during the first year of fumigation (1997) in a *Pinus taeda* forest plantation exposed to elevated CO₂ at the Duke free air CO₂ enrichment (FACE) study. Each point represents a 24-h mean of respiration and GPP. A k-means clustering analysis was performed and data were separated into clusters, which segregated by seasonal changes in GPP (Supporting Information Table S4). Nongrowing season, closed circles; growing season with low GPP, open circles; growing season, triangles. A third cluster (open squares), imposed ad hoc, revealed anomalies that occurred during the growing season under conditions with high autotrophic respiration from soils and comparatively low GPP, probably because of overcast days during the growing season, and where C storage could have buffered low GPP. The fit of the piece-wise linear model is $R^2 = 0.51$. Soil respiration and isotope data were obtained from Andrews *et al.* (1999), Hui & Luo (2004) and Taneva *et al.* (2006) and GPP from Luo *et al.* (2001b), and calculations were performed as described in Notes S1.

recently been obtained in grasslands (Gomez-Casanovas *et al.*, 2012) and this has also been suggested in other ecosystems (Street *et al.*, 2011; Subke *et al.*, 2012).

The isotope mixing models used here identify the amount of recently assimilated C respired from roots during the first year of

CO₂ fumigation (1997) at Duke FACE (Fig. 5, following Andrews *et al.*, 1999 and Taneva *et al.*, 2006; Table S1). As a result of the contribution of stored C (e.g. Bahn *et al.*, 2009; Gaudinski *et al.*, 2009; Lynch *et al.*, 2013; Figs 2–4), total root respiration is probably underestimated, as storage C would have the pretreatment isotopic signal. Nevertheless, we used these data to further investigate the relationships between photosynthesis and the use of current-year fixed C in root-rhizosphere respiration at the ecosystem level (Fig. 5). A nonhierarchical k-means cluster analysis (Sugar & James, 2003) identified two distinct clusters of respiration rate which was determined by seasonal GPP (Figs 5, S6–S8; Table S4). Thus, during periods of high GPP, root respiration is also high, correlating with high demands for whole-plant water and nutrients (Janssens *et al.*, 2001; Notes S1; Tables S2, S3). These results also suggest that root respiration can respond to shifts in GPP at daily time-scales, indicating a rapid response of root activity to environmental variables, changes in plant demands and/or C availability. Furthermore, a piece-wise linear model indicates that GPP can explain ~51% of the variance in use of current-year C for root respiration (Fig. 5). While GPP approaches zero for several days during the year, an apparent basal maintenance rate of respiration of 0.3 g C m^{-2} represents a minimum substrate contribution to root respiration. Total root respiration was probably supported by storage C during the low GPP periods, as suggested above (Fig. 2; Lynch *et al.*, 2013; Gough *et al.*, 2009). These results suggest that models should consider a hierarchical approach to parameterize root respiration, which responds directly to environmental drivers such as soil temperature and moisture but also responds rapidly to changes in photosynthesis.

Synthesis and goals for future research

Methods to separate *in situ* the autotrophic from the heterotrophic components of soil respiration have biases that have weakened the formulation of clear biotic and abiotic controls on soil fluxes (Fig. 1). Among all the methods currently used, pulse-chase isotope methods and natural abundance ¹³C and ¹⁴C methods show the most promise in unraveling ecosystem controls on soil respiration and its components, particularly when used in combination with other approaches (Table 2). These methods are revealing the importance of GPP and stored C as factors that drive root respiration at the ecosystem level. However, there is still much to be done to fully document patterns of photosynthate use by roots and their variation in time and space through phenology, ontogeny, succession, and climate regimes (Table 2). Comparisons made across successional stages within an ecosystem and between ecosystems or biomes are desperately needed (e.g. comparisons of tropical versus temperate versus boreal forests, grassland types, and managed ecosystems). Separating autotrophic from heterotrophic soil flux in combination with long-term monitoring or manipulation experiments (e.g. flux sites and National Ecological Observatory Network) would be highly beneficial to identifying the role of biotic and abiotic influences on each of the soil components over time under a variety of ecosystems and environmental conditions. A summary of these challenges and potential approaches is given in Table 2.

Current models can simulate GPP relationships with autotrophic soil flux, yet the parameterization of models to capture the apparent physiological temperature response of ecosystems is an emerging challenge for both the modeling and flux communities. Both communities need to adopt a hierarchical approach to advance our process understanding at small scales (Table 2) to increasingly larger temporal and spatial scales and ultimately earth system models (ESMs; Table 2). The predictive ability of ESMs will be improved by explicit representations of root respiration that are primarily driven by GPP and plant C allocation which are currently absent.

Future progress will be made by using the continuous ecosystem CO₂ flux data that exist for a wide range of natural, semi-natural and agricultural ecosystems, across most biomes (Luysaert *et al.*, 2007), to ask questions that either challenge or improve model representations of the temporal controls on photosynthesis and C storage (Table 2). A recent example of this approach identified the importance of endogenous circadian rhythms as a control on GPP (Resco de Dios *et al.*, 2012). Incorporating this process into ESMs could help drive diel photosynthesis patterns without invoking a simple temperature response function. Similarly, the importance of photosynthesis-driven priming processes to heterotrophic SOM turnover should be tested using model–data fusion approaches. Key challenges for future research are the detection of RPE in more heterogeneous ecosystems than those used in manipulative experiments or tightly controlled field and laboratory studies. This will provide boundaries to the temperature sensitivity of belowground C fluxes and hierarchical responses of GPP to air temperature, as well as root respiratory responses to GPP acclimation and soil temperature change (Table 2). Soil versus air warming experiments may offer an opportunity to disentangle the effects of temperature–GPP interactions on soil fluxes (Table 2), but most current warming experiments have been focused on total soil respiration instead of, more importantly, the autotrophic and heterotrophic components.

It is important to note that increases in respiratory demand by belowground tissues can also influence total belowground C allocation. Because plants allocate C to maximize photosynthesis and growth (Thornley, 1969; Thornley & Cannell, 2000), plants will partition GPP into tissues that would minimize negative impacts of limiting resources on growth (DeLucia *et al.*, 2007; Litton *et al.*, 2007; Franklin *et al.*, 2012). Increased allocation of C to belowground tissues in response to nutrient and water demands (Gower *et al.*, 1996; Giardina *et al.*, 2003) potentially increases respiration costs (Odum, 1969) and reduces the proportion of GPP invested in growth (DeLucia *et al.*, 2007). Climate and phenological factors may lead to proportional changes in aboveground and belowground C allocation (and potentially maintenance of root-to-shoot ratios of species; Litton *et al.*, 2007; Gough *et al.*, 2010), and factors such as water availability, temperature, atmospheric CO₂ or nutrient availability have been documented to affect total belowground C allocation and autotrophic and heterotrophic respiration from soils (Haynes & Gower, 1995; King *et al.*, 2002; Norby *et al.*, 2004; Giardina *et al.*, 2005; Ryan & Law, 2005; Trueman & Gonzalez-Meler, 2005; Bryla *et al.*, 2008). C allocation patterns are used in models to predict the growth and C balance of ecosystems under climate change scenarios (Friedlingstein *et al.*, 2006; Fisher

Table 2 Suggested future research needs and potential experimental and modeling approaches for understanding the combined effects of abiotic and biotic drivers of root respiration from soils

Environmental control	Research needs	Experimental and modeling approaches for future research
Nitrogen (N)	Resolve whether changes in N availability result in changes in root respiration mediated by biomass and/or N content. Impact of N availability on carbon (C) allocation to heterotrophic soil organisms (e.g. mycorrhizas), and ultimately on direction of the plant response to N.	Experiments: Combine N fertilization and isotope tracers with biomass collection, and soil respiration fluxes and flux partitioning approaches. Quantify C allocation to mycorrhizas and the magnitude of the rhizosphere priming effect (RPE). Models: Use this quantitative understanding of how N availability affects belowground C allocation/respiration, and root distribution (size, order and depth). Include RPE effects on N mineralization in models.
Temperature	Test root acclimation to temperature in field experiments. Disentangle the relative significance of GPP from temperature in determining root respiration from soils. Understand the effects of temperature on belowground C allocation and nutrient demand that affects root mass and its respiratory flux. Understand the effects of temperature on nutrient availability mediated by decomposition because it will have indirect effects on root mass, plant growth and nutrient demand (seen N).	Experiments: Use multiple temperature treatments to obtain a temperature response function. Combine air and soil warming experiments that will allow for characterization of GPP versus direct temperature effects on root respiration. Document and understand temperature-driven changes in C allocation. Use a hierarchical approach to root respiration where primary (GPP and temperature), secondary (allocation), and tertiary (nutrient availability and heterotrophic responses) effects are identified. Models: Use process-based models to represent hierarchical temperature effects (e.g. temperature effect on autotrophic and heterotrophic metabolic rates, allocation patterns, and GPP) rather than simple empirical relationships (e.g. rely less on Q_{10}). Incorporate process understanding of the availability of photosynthetic products on root respiration, including stored C, on diurnal and seasonal time-scales.
C supply and seasonality	Disentangle phenology and diel patterns from temperature-driven changes in respiration. Document the role of stored C in buffering interannual, seasonal, and diel changes in substrate availability. Quantify C demand and respiratory cost for root turnover. Temporal changes in autotrophic substrate supply in the rhizosphere may be apparent in heterotrophic respiration after a time lag.	Experiments: Use radiocarbon, pulse-chase experiments, biometric approaches and/or phloem sap flux to determine patterns of belowground C allocation and instantaneous C use efficiency. Document the role of stored C in buffering interannual, seasonal, and diel changes in substrate availability. Identify the fraction of fine roots that turns over seasonally using minirhizotrons or isotopes. Determine temporal patterns for C allocation to heterotrophs in the rhizosphere. Models: Include root respiration as a function of current photosynthate or that buffered by stored C when GPP is insufficient. Reconcile the time steps of models with the temporal resolution of empirical data. Incorporate GPP controls on heterotrophic respiration.

et al., 2010; Franklin *et al.*, 2012). However, these models often operate at longer temporal and larger spatial scales than empirical data at which belowground C allocation are gathered (Gower *et al.*, 1996; Norby *et al.*, 2004; Litton *et al.*, 2007; Table 2). Additionally, poor constraints on C turnover in fine roots increase uncertainties in quantifying C allocation belowground and movement of C from plants to soils (Lynch *et al.*, 2013). New promising methods based on diurnal stem diameter change that correlate phloem sap flow with total sugar flux may provide new insights at tree and stand levels at various temporal scales (Hölttä *et al.*, 2006; Mencuccini *et al.*, 2013).

In summary, the typical representation of the temperature sensitivity of root respiration in models contrasts with empirical evidence gathered at the tissue to ecosystem levels that suggests otherwise. For instance, although tissue N content is a good predictor of specific rates of fine-root respiration, ecosystem-level autotrophic soil respiration may better scale with changes in fine-root biomass than root N content (Table 2). In addition to

changes in root biomass, autotrophic soil respiration is further modulated by the respiratory acclimation to changes in temperature (seasonal or resulting from warming). Our findings suggest that while all of these processes act in combination, substrate availability (a product of GPP and storage) is the ultimate driver of the autotrophic soil flux (Table 2). Identifying these factors in a hierarchical way is paramount to advance our predictive understanding of the responses of ecosystems to climate change.

Acknowledgements

We thank the Department of Energy (DOE) Office of Biological and Environmental Research (BER) for funding the workshop 'Scaling Root Processes: Global Impacts,' in which the topics of this paper were developed. Grants from the DOE (ER65188 to M.A.G.-M.; DE-SC0006951 to J.T.) and the US National Science Foundation (DEB-0919276 to M.A.G.-M.; AGS-1005663 and ARC-0902109 to J.T.) supported this research and participation of

D.J.L. in the workshop. We also thank Susan Trumbore for the generous use of radiocarbon data.

References

- Andrews JA, Harrison KG, Matamala R, Schlesinger WH. 1999. Separation of root respiration from total soil respiration using carbon-13 labeling during free-air carbon dioxide enrichment (FACE). *Soil Science Society of America Journal* 63: 1429–1435.
- Atkin OK, Edwards EJ, Loveys BR. 2008. Response of root respiration to changes in temperature and its relevance to global warming. *New Phytologist* 147: 141–154.
- Bahn M, Schmitt M, Siegwolf R, Richter A, Bruggemann N. 2009. Does photosynthesis affect grassland soil-respired CO₂ and its carbon isotope composition on a diurnal timescale? *New Phytologist* 182: 451–460.
- Bond-Lamberty B, Bronson D, Bladyka E, Gower ST. 2011. A comparison of trenched plot techniques for partitioning soil respiration. *Soil Biology & Biochemistry* 43: 2108–2114.
- Bond-Lamberty B, Thomson A. 2010. Temperature-associated increases in the global soil respiration record. *Nature* 464: 579–582.
- Boone RD, Nadelhoffer KJ, Canary JD, Kaye JP. 1998. Roots exert a strong influence on the temperature sensitivity of soil respiration. *Nature* 396: 570–572.
- Borken W, Savage K, Davidson EA, Trumbore SE. 2006. Effects of experimental drought on soil respiration and radiocarbon efflux from a temperate forest soil. *Global Change Biology* 12: 177–193.
- Bouma TJ, Janssen HJA, de Kock MJ, van Leeuwen PH, de Visser R. 1994. Respiratory energy requirements and rate of protein turnover in vivo determined by the use of an inhibitor of protein synthesis and a probe to assess its effect. *Physiologia Plantarum* 92: 585–594.
- Bowden RD, Nadelhoffer KJ, Boone RD, Melillo JM, Garrison JB. 1993. Contributions of aboveground litter, belowground litter, and root respiration to total soil respiration in a temperate mixed hardwood forest. *Canadian Journal of Forest Research* 23: 1402–1407.
- Bowling DR, Pataki DE, Randerson JT. 2008. Carbon isotopes in terrestrial ecosystem pools and CO₂ fluxes. *New Phytologist* 178: 24–40.
- Bruggemann N, Gessler A, Kayler Z, Keel SG, Badeck F, Barthel M, Boeckx P, Buchmann N, Brugnoli E, Esperschütz J *et al.* 2011. Carbon allocation and carbon isotope fluxes in the plant–soil–atmosphere continuum: a review. *Biogeosciences* 8: 3457–3489.
- Bryla DR, Bouma TJ, Eissenstat DM. 2008. Root respiration in citrus acclimates to temperature and slows during drought. *Plant, Cell & Environment* 20: 1411–1420.
- Burton AJ, Jarvey JC, Jarvi MP, Zak DR, Pregitzer KS. 2012. Chronic N deposition alters root respiration–tissue N relationship in northern hardwood forests. *Global Change Biology* 18: 258–266.
- Burton AJ, Melillo JM, Frey SD. 2008. Adjustment of forest ecosystem root respiration as temperature warms. *Journal of Integrative Plant Biology* 50: 1467–1483.
- Burton AJ, Pregitzer KS, Hendrick RL. 2000. Relationships between fine root dynamics and nitrogen availability in Michigan northern hardwood forests. *Oecologia* 125: 389–399.
- Burton AJ, Pregitzer KS, Ruess RW, Hendrik RL, Allen MF. 2002. Root respiration in North American forests: effects of nitrogen concentration and temperature across biomes. *Oecologia* 131: 559–568.
- Carbone MS, Trumbore SE. 2007. Contribution of new photosynthetic assimilates to respiration by perennial grasses and shrubs: residence times and allocation patterns. *New Phytologist* 176: 124–135.
- Chapin FS III, Matson PA, Mooney HA. 2002. *Principles of terrestrial ecosystem ecology*. New York, NY, USA: Springer, 125–127.
- Chen D, Zhou L, Rao X, Lin Y, Fu S. 2010. Effects of root diameter and root nitrogen concentration on in situ root respiration among different seasons and tree species. *Ecological Research* 25: 983–993.
- Chen X, Post WM, Norby RJ, Classen AT. 2011. Modeling soil respiration and variations in source components using a multi-factor global climate change experiment. *Climatic Change* 107: 459–490.
- Cisneros-Dozal LM. 2005. *Using a ¹⁴C release to partition soil respiration sources in a southeastern hardwood forest*. PhD thesis, University of California, Irvine, CA, USA.
- Craine JM, Wedin DA, Chapin FS. 1999. Predominance of ecophysiological controls on soil CO₂ flux in a Minnesota grassland. *Plant and Soil* 207: 77–86.
- Czimczik CI, Trumbore SE, Carbone MS, Winston GC. 2006. Changing sources of soil respiration with time since fire in a boreal forest. *Global Change Biology* 12: 957–971.
- Dannoura M, Maillard P, Fresneau C, Plain C, Berveiller D, Gerant D, Chipeaux C, Bosc A, Ngao J, Damesin C *et al.* 2011. In situ assessment of the velocity of carbon transfer by tracing ¹³C in trunk CO₂ efflux after pulse labeling: variations among tree species and seasons. *New Phytologist* 190: 181–192.
- Davidson EA, Janssens IA. 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440: 165–173.
- Davidson EA, Janssens IA, Luo Y. 2006. On the variability of respiration in terrestrial ecosystems: moving beyond Q₁₀. *Global Change Biology* 12: 154–164.
- DeLucia E, Drake JE, Thomas RB, Gonzalez-Meler MA. 2007. Forest carbon use efficiency: is respiration a constant fraction of gross primary production? *Global Change Biology* 13: 1157–1167.
- DeLucia EH, Hamilton JG, Naidu SL, Thomas RB, Andrews JA, Finzi A, Lavine M, Matamala R, Mohan JE, Hendrey GR *et al.* 1999. Net primary production of a forest ecosystem with experimental CO₂ enrichment. *Science* 284: 1177–1179.
- Drigo B, Pijl AS, Duyts H, Kielak AM, Gamper HA, Houtekamer MJ, Boschker HTS, Bodelier PLE, Whiteley AS, van Veen JA *et al.* 2010. Shifting carbon flow from roots into associated microbial communities in response to elevated atmospheric CO₂. *Proceedings of the National Academy of Sciences, USA* 107: 10938–10942.
- Dungait JAJ, Hopkins DW, Gregory AS, Whitmore AP. 2012. Soil organic matter turnover is governed by accessibility not recalcitrance. *Global Change Biology* 18: 1781–1796.
- Epron D, Farque L, Lucot E, Badot PM. 1999. Soil CO₂ efflux in a beech forest: the contribution of root respiration. *Annals Forest Science* 56: 289–295.
- Ewel KC, Cropper WP, Gholz HL. 1987. Soil carbon dioxide evolution in Florida [USA] slash pine plantations: II. Importance of root respiration. *Canadian Journal of Forest Research* 17: 330–333.
- Fang C, Smith P, Moncrieff JB, Smith JU. 2005. Similar response of labile and resistant soil organic matter pools to changes in temperature. *Nature* 433: 57–59.
- Feng X, Xu Y, Jaffé R, Schlesinger WH, Simpson MJ. 2010. Turnover rates of hydrolysable aliphatic lipids in Duke Forest soils determined by compound specific ¹³C isotopic analysis. *Organic Geochemistry* 6: 573–579.
- Field CB, Randerson JT, Malmström CM. 1995. Global net primary production: combining e colony and remote sensing. *Remote Sensing Environment* 51: 74–88.
- Fisher RA, McDowell N, Purves D, Moorcroft P, Sitch S, Cox P, Huntingford C, Meir P, Woodward FI. 2010. Assessing uncertainties in a second-generation dynamic vegetation model caused by ecological scale limitations. *New Phytologist* 187: 666–681.
- Fontaine S, Bardoux G, Abbadie L, Mariotti A. 2004. Carbon input to soil may decrease soil carbon content. *Ecology Letters* 7: 314–320.
- Fontaine S, Barot S, Barre P, Bdioui N, Mary B, Rumpel C. 2007. Stability of organic carbon in deep soil layers controlled by fresh carbon supply. *Nature* 450: 277–280.
- Franklin O, Johansson J, Dewar RC, Dieckmann U, McMurtrie RE, Brannstrom A, Dybzinski R. 2012. Modeling carbon allocation in trees: a search for principles. *Tree Physiology* 32: 648–666.
- Friedlingstein P, Cox P, Betts R, Bopp L, von Bloh W, Brovkin V, Cadule P, Doney S, Eby M, Fung I *et al.* 2006. Climate-carbon cycle feedback analysis: results from the C4MIP model intercomparison. *Journal of Climate* 19: 3337–3353.
- Gaudinski JB, Torn MS, Riley WJ, Swanston C, Trumbore SE, Joslin JD, Majdi H, Dawson TE, Hanson PJ. 2009. Use of stored carbon reserves in growth of temperate tree roots and leaf buds: analyses using radiocarbon measurements and modelling. *Global Change Biology* 15: 992–1014.
- Giardina CP, Coleman MD, Hancock JE, King JS, Lilleskov EA, Loya WM, Pregitzer KS, Ryan MG, Trettin CC. 2005. The response of belowground carbon allocation in forests to global change. In: Binkley D, Menyailo O, eds. *Tree species effects on soils: implications for global change*. Dordrecht, the Netherlands: Springer in cooperation with NATO Public Diplomacy Division, 119–154.
- Giardina CP, Ryan MG, Binkley D, Fownes JH. 2003. Primary production and carbon allocation in relation to nutrient supply in a tropical experimental forest. *Global Change Biology* 9: 438–450.

- Gifford RM. 2003. Plant respiration in productivity models: conceptualization, representation and issues for global terrestrial carbon-cycle research. *Functional Plant Biology* 30: 171–186.
- Gomez-Casanovas N, Matamala R, Cook DR, Gonzalez-Meler MA. 2012. Net ecosystem exchange modifies the relationship between the autotrophic and heterotrophic components of soil respiration with abiotic factors in prairie grasslands. *Global Change Biology* 18: 2532–2545.
- Gonzalez-Meler MA, Taneva L. 2005. Integrated effects of atmospheric CO₂ concentration on plant and ecosystem respiration. In: Lambers H, Ribas-Carbo M, eds. *Plant respiration*. Dordrecht, the Netherlands: Kluwer-Academic Publishers, 211–259.
- Gonzalez-Meler MA, Taneva L, Trueman RJ. 2004. Plant respiration and elevated atmospheric CO₂ concentration: cellular responses and global significance. *Annals of Botany* 94: 467–656.
- Gough CM, Flower CE, Vogel CS, Curtis PS. 2010. Phenological and temperature controls on the temporal non-structural carbohydrate dynamics of *Populus grandidentata* and *Quercus rubra*. *Forests* 1: 65–81.
- Gough CM, Flower CE, Vogel CS, Dragoni D, Curtis PS. 2009. Whole-ecosystem labile carbon production in a north temperate deciduous forest. *Agricultural and Forest Meteorology* 149: 1531–1540.
- Gough CM, Seiler JR, Maier CA. 2004. Short-term effects of fertilization on loblolly pine (*Pinus taeda*L.) physiology. *Plant, Cell & Environment* 27: 876–886.
- Goulden ML, McMillan AMS, Winston GC, Rocha AV, Mainies KL, Harden JW, Bond-Lamberty BP. 2011. Patterns of NPP, GPP, respiration, and NEP during boreal forest succession. *Global Change Biology* 17: 855–871.
- Gower ST, Pongracic S, Landberg JJ. 1996. A global trend in belowground carbon allocation: can we use the relationship at smaller scales? *Ecology* 77: 1750–1755.
- Guenet B, Juarez S, Bardoux G, Abbadie L, Chenu C. 2012. Evidence that stable C is as vulnerable to priming effect as is more labile C in soil. *Soil Biology & Biochemistry* 52: 43–48.
- Hamilton JG, DeLucia EH, George K, Naidu SL, Finzi AC, Schlesinger WH. 2002. Forest carbon balance under elevated CO₂. *Oecologia* 131: 250–260.
- Hanson PJ, Edwards NT, Garten CT, Andrews JA. 2000. Separating root and soil microbial contributions to soil respiration: a review of methods and observations. *Biogeochemistry* 48: 115–146.
- Hartley IP, Heinemeyer A, Evans SP, Ineson P. 2007. The effect of soil warming on bulk soil vs. rhizosphere respiration. *Global Change Biology* 13: 2654–2667.
- Hartley IP, Ineson P. 2008. Substrate quality and the temperature sensitivity of soil organic matter decomposition. *Soil Biology & Biochemistry* 40: 1567–1574.
- Hasselquist NJ, Metcalfe DB, Högborg P. 2012. Contrasting effects of low and high nitrogen additions on soil CO₂ flux components and ectomycorrhizal fungal sporocarp production in a boreal forest. *Global Change Biology* 18: 3596–3605.
- Haynes BE, Gower ST. 1995. Belowground carbon allocation in unfertilized and fertilized red pine plantations in northern Wisconsin. *Tree Physiology* 15: 317–325.
- Heinemeyer A, Hartley IP, Evans SP, De la Fuente JAC, Ineson P. 2007. Forest soil CO₂ flux: uncovering the contribution and environmental responses of ectomycorrhizas. *Global Change Biology* 13: 1786–1797.
- Heinemeyer A, Wilkinson M, Vargas R, Subke JA, Casella E, Morison JIL, Ineson P. 2012. Exploring the “overflow tap” theory: linking forest soil CO₂ fluxes and individual mycorrhizosphere components to photosynthesis. *Biogeosciences* 9: 79–95.
- Högborg P, Bhupinderpal S, Lofvenius MO, Nordgren A. 2009. Partitioning of soil respiration into its autotrophic and heterotrophic components by means of tree-girdling in old boreal spruce forest. *Forest Ecology and Management* 257: 1764–1767.
- Högborg P, Buchmann N, Read DJ. 2006. Comments on Yakov Kuzyakov’s review ‘Sources of CO₂ efflux from soil and review of partitioning methods’ [Soil Biology & Biochemistry 38, 425–448]. *Soil Biology & Biochemistry* 38: 2997–2998.
- Högborg P, Nordgren A, Buchmann N, Taylor AFS, Ekblad A, Högborg MN, Nyberg G, Ottosson-Lofvenius M, Read DJ. 2001. Large-scale forest girdling shows that current photosynthesis drives soil respiration. *Nature* 411: 789–792.
- Högborg MN, Briones MJI, Keel SG, Metcalfe DB, Campbell C, Midwood AJ, Thornton B, Hurry V, Linder S, Nasholm T *et al.* 2010. Quantification of effects of season and nitrogen supply on tree below-ground carbon transfer to ectomycorrhizal fungi and other soil organisms in a boreal pine forest. *New Phytologist* 187: 485–493.
- Hölttä T, Vesala T, Sevanto S, Perämäki M, Nikinmaa E. 2006. Modeling xylem and phloem water flows in trees according to cohesion theory and Münch pressure flow hypothesis. *Trees* 20: 67–78.
- Hopkins FM, Torn MS, Trumbore SE. 2012. Warming accelerates decomposition of decades-old carbon in forest soils. *Proceedings of the National Academy of Sciences, USA* 109: E1753–E1761.
- Houghton RA, Davidson EA, Woodwell GM. 1998. Missing sinks, feedbacks and understanding the role of terrestrial ecosystems in the global carbon balance. *Global Biogeochemical Cycles* 12: 24–34.
- Hui DF, Luo YQ. 2004. Evaluation of soil CO₂ production and transport in Duke Forest using a process-based modeling approach. *Global Biogeochemical Cycles* 18: GB4029.
- Irvine J, Law BE, Kurpius MR. 2005. Coupling of canopy gas exchange with root and rhizosphere respiration in a semi-arid forest. *Biogeochemistry* 73: 271–282.
- Iversen CM. 2010. Digging deeper: fine-root responses to rising atmospheric CO₂ concentration in forested ecosystems. *New Phytologist* 186: 346–357.
- Janssens IA, Dieleman W, Luysaert S, Subke JA, Reichstein M, Ceulemans R, Ciais P, Dolman AJ, Grace J, Matteucci G *et al.* 2010. Reduction of forest soil respiration in response to nitrogen deposition. *Nature Geoscience* 3: 315–322.
- Janssens IA, Lankreijer H, Matteucci G, Kowalski AS, Buchmann N, Epron D, Pilegaard K, Kutsch W, Longdoz B, Grunwald T *et al.* 2001. Productivity overshadows temperature in determining soil and ecosystem respiration across European forests. *Global Change Biology* 7: 269–278.
- Jia S, Wang Z, Li X, Zhang X, McLaughlin NB. 2011. Effects of nitrogen fertilizer, root branch order and temperature on respiration and tissue N concentration of fine roots in *Larix gmelinii* and *Fraxinus mandshurica*. *Tree Physiology* 31: 718–726.
- Jia SX, Wang ZQ, Li XP, Sun Y, Zhang XP, Liang AZ. 2010. N fertilization affects on soil respiration, microbial biomass and root respiration in *Larix gmelinii* and *Fraxinus mandshurica* plantations in China. *Plant and Soil* 333: 325–326.
- Karhu K, Fritze H, Hamalainen K, Vanhala P, Jungner H, Oinonen M, Sonninen E, Tuomi M, Spetz P, Kitunen V *et al.* 2010. Temperature sensitivity of soil carbon fractions in boreal forest soil. *Ecology* 91: 370–376.
- Keenan TF, Davidson E, Moffat AM, Munger W, Richardson AW. 2012. Using model-data fusion to interpret past trend, and quantify uncertainties in future projections, or terrestrial ecosystem carbon cycling. *Global Change Biology* 18: 2555–2569.
- King JS, Albaugh TJ, Allen HL, Buford M, Strain BR, Dogherty B. 2002. Below-ground carbon input to soil is controlled by nutrient availability and fine root dynamics in loblolly pine. *New Phytologist* 154: 389–398.
- Kuzyakov Y. 2010. Priming effects: interactions between living and dead organic matter. *Soil Biology & Biochemistry* 42: 1363–1371.
- Kuzyakov Y, Larionova AA. 2005. Root and rhizomicrobial respiration: a review of approaches to estimate respiration by autotrophic and heterotrophic organisms in soil. *Journal of Plant Nutrition and Soil Science* 168: 503–520.
- Lambers H, Chapin FS, Pons TL. 2008. *Plant physiological ecology*. New York, NY, USA: Springer.
- Lambers H, Schwurwater I, Atkin OK. 1996. Respiratory patterns in roots in relation to their functioning. In: Waisel Y, Eshel A, Kafkaki K, eds. *Plant roots: the hidden half*. New York, NY, USA: Marcel Dekker, 323–362.
- Lavigne MB, Boutin R, Foster RJ, Goodine G, Bernier PY, Robitaille G. 2003. Soil respiration responses to temperature are controlled more by roots than by decomposition in balsam fir ecosystems. *Canadian Journal of Forest Research* 33: 1744–1753.
- Leuzinger S, Thomas RQ. 2011. How do we improve Earth system models? Integrating Earth system models, ecosystem models, experiments and long-term data. *New Phytologist* 191: 15–18.
- Levin I, Naegler T, Kromer B, Diehl M, Francey RJ, Gomez-Pelaez AJ, Steele LP, Wagenbach D, Weller R, Worthy DE. 2010. Observations and modelling of the global distribution and long-term trend of atmospheric ¹⁴CO₂. *Tellus Series B* 62: 26–46.
- Levy-Varon JH, Schuster WSF, Griffin KL. 2012. The autotrophic contribution to soil respiration in a northern temperate deciduous forest and its response to stand disturbance. *Oecologia* 169: 211–220.
- Litton CM, Raich JW, Ryan MG. 2007. Carbon allocation in forest ecosystems. *Global Change Biology* 13: 2089–2109.

- Liu ZF, Wu JP, Zhou LX, Lin YB, Fu SL. 2012. Tree girdling effect on bacterial substrate utilization pattern depending on stand age and soil microclimate in Eucalyptus plantations. *Applied Soil Ecology* 54: 7–13.
- Lloyd J, Taylor JA. 1994. On the temperature dependence of soil respiration. *Functional Ecology* 8: 315–323.
- Luo Y, Medlyn B, Hui D, Ellsworth D, Reynolds J, Katul G. 2001b. Gross primary productivity in Duke Forest: modeling synthesis of CO₂ experiment and eddy-flux data. *Ecological Applications* 11: 239–252.
- Luo YQ, Randerson JT, Abramowitz G, Bacour C, Blyth E, Carvalhais N, Ciais P, Dalmonech D, Fisher J, Fisher R *et al.* 2012. A framework for benchmarking land models. *Biogeosciences* 9: 3857–3874.
- Luo Y, Wu L, Andrews JA, White L, Matamala R, Schäfer KVR, Schlesinger WH. 2001a. Elevated CO₂ differentiates ecosystem carbon processes: deconvolution analysis of Duke Forest FACE data. *Ecological Monographs* 71: 357–376.
- Luo Y, Zhou X. 2006. *Soil respiration and the environment*. Amsterdam, the Netherlands: Academic Press.
- Luyssaert S, Inglima I, Jung M, Richardson AD, Reichstein M, Papale D, Piao SL, Schulze ED, Wingate L, Matteucci G *et al.* 2007. CO₂-balance of boreal, temperate and tropical forest derived from a global database. *Global Change Biology* 13: 2509–2537.
- Lynch DJ, Matamala R, Iverson CM, Norby RJ, Gonzalez-Meler MA. 2013. Stored carbon partly fuels fine-root respiration but is not used for production of new fine roots. *New Phytologist*. doi: 10.1111/nph.12290.
- Mahecha MD, Reichstein M, Carvalhais N, Lasslop G, Lange H, Seneviratne SI, Vargas R, Ammann C, Arain MA, Cescatti A *et al.* 2010. Global convergence in the temperature sensitivity of respiration at ecosystem level. *Science* 329: 838–840.
- Maier CA, Kress LW. 2000. Soil CO₂ evolution and root respiration in 11 year-old loblolly pine (*Pinus taeda*) plantations as affected by moisture and nutrient availability. *Canadian Journal of Forest Research* 30: 347–359.
- Matamala R, Gonzalez-Meler MA, Jastrow JD, Norby RJ, Schlesinger WH. 2003. Impacts of fine root turnover on forest NPP and soil C sequestration potential. *Science* 302: 1385–1387.
- Matamala R, Schlesinger WH. 2000. Effects of elevated atmospheric CO₂ on fine root production and activity in an intact temperate forest ecosystem. *Global Change Biology* 6: 967–979.
- Mencuccini M, Holtta T. 2010. The significance of phloem transport for the speed with which canopy photosynthesis and belowground respiration are linked. *New Phytologist* 185: 189–203.
- Mencuccini M, Hölttä T, Sevanto S, Nikinmaa E. 2013. Concurrent measurements of change in the bark and xylem diameters of trees reveal a phloem-generated turgor signal. *New Phytologist*. doi: 10.1111/nph.12224.
- Milcu A, Heim A, Ellis RJ, Scheu S, Manning P. 2011. Identification of general patterns of nutrient and labile carbon control on soil carbon dynamics across a successional gradient. *Ecosystems* 14: 710–719.
- Moore DJP, Gonzalez-Meler MA, Taneva L, Phippen JS, Kim H-S, DeLucia EH. 2008. The effect of carbon dioxide enrichment on apparent stem respiration from *Pinus taeda* L. is confounded by high levels of soil carbon dioxide. *Oecologia* 158: 1–10.
- Moyano FE, Kutsch WL, Rebmann C. 2008. Soil respiration fluxes in relation to photosynthetic activity in broad-leaf and needle-leaf forest stands. *Agricultural and Forest Meteorology* 148: 135–143.
- Neff JC, Townsend AR, Gleixner G, Lehman SJ, Turnbull J, Bowman WD. 2002. Variable effects of nitrogen additions on the stability and turnover of soil carbon. *Nature* 419: 915–917.
- Norby RJ, Ledford J, Reilly CD, Miller NE, O'Neill EG. 2004. Fine-root production dominates response of a deciduous forest to atmospheric CO₂ enrichment. *Proceedings of the National Academy of Sciences of the United States of America* 101: 9689–9693.
- Nowinski NS, Trumbore SE, Jimenez G, Fenn ME. 2009. Alteration of belowground carbon dynamics by nitrogen addition in southern California mixed conifer forests. *Journal of Geophysical Research* 114: g02005.
- Odum EP. 1969. The strategy of ecosystem development. *Science* 164: 262–270.
- Ohlsson KEA. 2011. Theoretical model of the abiotic component of soil (CO₂)-C-13 tracer efflux in ¹³C pulse-labeling experiments on plant-soil systems. *Soil Biology & Biochemistry* 43: 675–681.
- Pataki DE, Ellsworth DS, Evans RD, Gonzalez-Meler M, King J, Leavitt SW, Lin GH, Matamala R, Pendall E, Siegwolf R *et al.* 2003. Tracing changes in ecosystem function under elevated carbon dioxide conditions. *BioScience* 53: 805–818.
- Phillips RP, Finzi AC, Bernhardt ES. 2011. Enhanced root exudation induces microbial feedbacks to N cycling in a pine forest under long-term CO₂ fumigation. *Ecology Letters* 14: 187–194.
- Pregitzer KS, King JS, Burton AJ. 2000. Responses of tree fine roots to temperature. *New Phytologist* 147: 105–115.
- Pregitzer KS, Laskowski MJ, Burton AJ, Lessard VC, Zak DR. 1998. Variation in sugar maple root respiration with root diameter and soil depth. *Tree Physiology* 18: 665–670.
- Rachmilevitch S, Xu Y, Gonzalez-Meler MA, Huang B, Lambers H. 2007. Cytochrome and alternative pathway activity in roots of thermal and non-thermal *Agrostis* species in response to high soil temperature. *Physiologia Plantarum* 129: 163–174.
- Reich PB, Oleksyn J, Tjoelker MG. 1996. Needle respiration and nitrogen concentration in Scots Pine populations from a broad latitudinal range: a common garden test with field grown trees. *Functional Ecology* 10: 768–776.
- Reichstein M, Katterer T, Andren O, Ciais P, Schulze E, Cramer W, Papale D, Valentini R. 2005. Temperature sensitivity of decomposition in relation to soil organic matter pools: critique and outlook. *Biogeosciences* 2: 317–321.
- Resco de Dios V, Goulden ML, Ogle K, Richardson AD, Hollinger DY, Davidson EA, Alday JG, Barron-Gafford GA, Carrara A, Kowalski AS *et al.* 2012. Endogenous circadian regulation of carbon dioxide exchange in terrestrial ecosystems. *Global Change Biology* 18: 1956–1970.
- Richardson AD, Carbone MS, Keenan TF, Czimczik CI, Hollinger DY, Murakami P, Schaberg PG, Xu X. 2013. Seasonal dynamics and age of stemwood nonstructural carbohydrates in temperate forest trees. *New Phytologist* 197: 850–861.
- Risk D, Nickerson N, Phillips CL, Kellman L, Moroni M. 2012. Drought alters respired delta(CO₂)-C-13 from autotrophic, but not heterotrophic soil respiration. *Soil Biology & Biochemistry* 50: 26–32.
- Ryan MG. 1990. Growth and maintenance respiration in stems of *Pinus contorta* and *Picea engelmannii*. *Canadian Journal of Forest Research* 20: 48–57.
- Ryan MG. 1991. Effects of climate change on plant respiration. *Ecological Applications* 1: 157–167.
- Ryan MG. 1995. Foliar maintenance respiration of subalpine and boreal trees and shrubs in relation to nitrogen content. *Plant, Cell & Environment* 18: 765–772.
- Ryan MG, Hubbard RM, Pongracic S, Raison RJ, McMurtire RE. 1996. Foliage, fine-root, woody-tissue and stand respiration in *Pinus radiata* in relation to nitrogen status. *Tree Physiology* 16: 333–343.
- Ryan MG, Law BE. 2005. Interpreting, measuring, and modeling soil respiration. *Biogeochemistry* 73: 3–27.
- Savage K, Davidson EA, Tang J. 2013. Diel patterns of autotrophic and heterotrophic respiration among phenological stages. *Global Change Biology* in press.
- Scheurwater I, Cornelissen C, Dictus F, Welschen R, Lambers H. 1998. Why do fast- and slow-growing grass species differ so little in their rate of root respiration, considering the large differences of growth and ion uptake? *Plant, Cell & Environment* 21: 995–1005.
- Schlesinger WH, Andrews JA. 2000. Soil respiration and the global carbon cycle. *Biogeochemistry* 48: 7–20.
- Schlesinger WH, Lichter J. 2001. Limited carbon storage in soil and litter of experimental forest plots under increased atmospheric CO₂. *Nature* 411: 466–469.
- Schuur EAG, Trumbore SE. 2006. Partitioning sources of soil respiration in boreal black spruce forest using radiocarbon. *Global Change Biology* 12: 165–176.
- Smithwick EAH, Eissenstat DM, Lovett GM, Bowden RD, Rustad LE, Driscoll CT. 2013. Root stress and nitrogen deposition: consequences and research priorities. *New Phytologist* 197: 712–719.
- Southon J, Santos G. 2004. Ion source development at KCCAMS, University of California, Irvine. *Radiocarbon* 46: 33–39.
- Street LE, Subke JA, Sommerkorn M, Heinemeyer A, Williams M. 2011. Turnover of recently assimilated carbon in arctic bryophytes. *Oecologia* 167: 325–337.
- Subke JA, Bahn M. 2010. On the 'temperature sensitivity' of soil respiration: can we use the immeasurable to predict the unknown? *Soil Biology & Biochemistry* 42: 1653–1656.

- Subke JA, Heinemeyer A, Vallack HW, Leronni V, Baxter R, Ineson P. 2012. Fast assimilate turnover revealed by in situ CO₂-C-13 pulse-labelling in subarctic tundra. *Polar Biology* 35: 1209–1219.
- Subke JA, Inglema I, Cotrufo MF. 2006. Trends and methodological impacts in soil CO₂ efflux partitioning: a meta-analytical review. *Global Change Biology* 12: 921–943.
- Subke JA, Voke NR, Leronni V, Garnett MH, Ineson P. 2011. Dynamics and pathways of autotrophic and heterotrophic soil CO₂ efflux revealed by forest girdling. *Journal of Ecology* 99: 186–193.
- Sugar CA, James GM. 2003. Finding the number of clusters in a dataset: an information-theoretic approach. *Journal of the American Statistical Association* 98: 750–763.
- Taneva L, Gonzalez-Meler MA. 2011. Distinct patterns in the diurnal and seasonal variability in four components of soil respiration in a temperate forest under free-air CO₂ enrichment. *Biogeosciences* 8: 3077–3092.
- Taneva L, Phippen JS, Schlesinger WH, Gonzalez-Meler MA. 2006. The turnover of carbon pools contributing to soil CO₂ and soil respiration in a temperate forest exposed to elevated CO₂ concentration. *Global Change Biology* 12: 983–994.
- Tang J, Baldocchi DD. 2005. Spatial-temporal variation in soil respiration in an oak-grass savanna ecosystem in California and its partitioning into autotrophic and heterotrophic components. *Biogeochemistry* 73: 183–207.
- Tang J, Baldocchi DD, Xu L. 2005b. Tree photosynthesis modulates soil respiration on a diurnal time scale. *Global Change Biology* 11: 1298–1304.
- Tang J, Misson L, Gershenson A, Cheng WX, Goldstein AH. 2005a. Continuous measurements of soil respiration with and without roots in a ponderosa pine plantation in the Sierra Nevada Mountains. *Agricultural and Forest Meteorology* 132: 212–227.
- Thornley JM. 1969. A model to describe the partitioning of photosynthate during vegetative plant growth. *Annals of Botany* 33: 419–430.
- Thornley JHM, Cannell MGR. 2000. Modeling the components of plant respiration: representation and realism. *Annals of Botany* 85: 55–67.
- Thornton PE, Rosenbloom NA. 2005. Ecosystem model spin-up: estimating steady state conditions in a coupled terrestrial carbon and nitrogen cycle model. *Ecological Modelling* 189: 25–48.
- Trueman RJ, Gonzalez-Meler MA. 2005. Accelerated belowground C cycling in a managed agriforest ecosystem exposed to elevated carbon dioxide concentrations. *Global Change Biology* 11: 1258–1271.
- Trumbore S. 2006. Carbon respired by terrestrial ecosystems - recent progress and challenges. *Global Change Biology* 12: 141–153.
- Trumbore S, Da Costa ES, Nepstad DC, De Camargo PB, Martinelli L, Ray D, Restom T, Silver W. 2006. Dynamics of fine root carbon in Amazonian tropical ecosystems and the contribution of roots to soil respiration. *Global Change Biology* 12: 217–229.
- Trumbore SE. 2000. Age of soil organic matter and soil respiration: radiocarbon constraints on belowground C dynamics. *Ecological Applications* 10: 399–411.
- Valentini R, Matteucci G, Doman AJ, Schulze ED, Reimann C, Moors EJ, Granier A, Gross P, Jensen NO, Pilegaard K *et al.* 2000. Respiration as the main determinant of carbon balance in European forests. *Nature* 404: 861–863.
- Vallack HW, Leronni V, Metcalfe DB, Höglberg P, Ineson P, Subke JA. 2012. Application of nitrogen fertilizer to a boreal pine forest has a negative impact on the respiration of ectomycorrhizal hyphae. *Plant and Soil* 352: 405–417.
- Vargas R, Carbone MS, Reichstein M, Baldocchi DD. 2011. Frontiers and challenges in soil respiration research: from measurements to model-data integration. *Biogeochemistry* 102: 1–13.
- Vose JM, Ryan MG. 2002. Seasonal respiration of foliage, fine roots, and woody tissues in relation to growth, tissue N, and photosynthesis. *Global Change Biology* 8: 182–193.
- Wang W, Peng S, Fang J. 2010. Root respiration and its relation to nutrient contents in soil and root and EVI among 8 ecosystems, northern China. *Plant and Soil* 333: 391–401.
- Wu Z, Dijkstra P, Koch GW, Peñuelas J, Hungate BA. 2011. Responses of terrestrial ecosystems to temperature and precipitation change: a meta-analysis of experimental manipulation. *Global Change Biology* 17: 927–942.
- Zhu B, Cheng W. 2011. Rhizosphere priming effect increases the temperature sensitivity of soil organic matter decomposition. *Global Change Biology* 17: 2172–2183.
- Zogg GP, Zak DR, Burton AJ, Pregitzer KS. 1996. Fine root respiration in northern hardwood forests in relation to temperature and nitrogen availability. *Tree Physiology* 16: 719–725.

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 Increasing CO₂ by 200 μl l⁻¹ lowers the δ¹³C of the CO₂ in the elevated FACE plots to -20‰.

Fig. S2 Isotopic composition of fumigation CO₂ at the Duke FACE site.

Fig. S3 Pool size isotope correction for estimating fluxes from pool residence times that was applied to live and dead roots of different diameter classes and soil CO₂.

Fig. S4 Model for correcting incorporation of post-treatment C in root decomposition.

Fig. S5 Soil respiration for the elevated plots as simulated by Hui and Luo (2004).

Fig. S6 Cluster profile plots from the k-means analysis demonstrate that GPP is primarily responsible for the clustering, and root respiration rate is not.

Fig. S7 Daily GPP versus the day of year partitioned into two clusters using k-means cluster analysis.

Fig. S8 Daily current-year C utilized for root respiration versus the day of year.

Table S1 Isotope end-members for calculations of pretreatment C (A, ambient) and post-treatment C (E, elevated) at the Duke FACE site

Table S2 Averaged monthly values of the proportion of current photosynthate (% new C) respired by soils that was not originated in decomposition of leaf and root litter pools

Table S3 Partitioning of GPP into ecosystem components and fluxes at the Duke FACE site

Table S4 Summary statistics for k-means clustering analysis

Notes S1 Contribution of current photosynthate to root-rhizosphere respiration.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.