

# UC San Diego

## Recent Work

### Title

Distribution Theory for the Studentized Mean for Long, Short, and Negative Memory Time Series

### Permalink

<https://escholarship.org/uc/item/35c7r55c>

### Authors

McElroy, Tucker S.  
Politis, Dimitris N.

### Publication Date

2012-05-01

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/3.0/>

# Distribution Theory for the Studentized Mean for Long, Short, and Negative Memory Time Series

Tucker McElroy  
Statistical Research Division  
U.S. Census Bureau  
4600 Silver Hill Road  
Washington, D.C. 20233-9100  
tucker.s.mcelroy@census.gov

Dimitris N. Politis\*  
Department of Mathematics  
University of California at San Diego  
La Jolla, CA 92093-0112  
dpolitis@ucsd.edu

## Abstract

We consider the problem of estimating the variance of the partial sums of a stationary time series that has either long memory, short memory, negative/intermediate memory, or is the first-difference of such a process. The rate of growth of this variance depends crucially on the type of memory, and we present results on the behavior of tapered sums of sample autocovariances in this context when the bandwidth vanishes asymptotically. We also present asymptotic results for the case that the bandwidth is a fixed proportion of sample size, extending known results to the case of flat-top tapers. We adopt the fixed proportion bandwidth perspective in our empirical section, presenting two methods for estimating the limiting critical values – both the subsampling method and a plug-in approach. Extensive simulation studies compare the size and power of both approaches as applied to hypothesis testing for the mean. Both methods perform well – although the subsampling method appears to be better sized – and provide a viable framework for conducting inference for the mean. In summary, we supply a unified asymptotic theory that covers all different types of memory under a single umbrella.

**Keywords.** Kernel, Lag-windows, Overdifferencing, Spectral estimation, Subsampling, Tapers, Unit-root problem.

**Disclaimer** This report is released to inform interested parties of research and to encourage discussion. The views expressed on statistical issues are those of the authors and not necessarily those of the U.S. Census Bureau.

---

\*Research partially supported by NSF grants DMS-07-06732 and DMS-10-07513.

# 1 Introduction

Consider a sample  $Y = \{Y_1, Y_2, \dots, Y_n\}$  from a strictly stationary time series with mean  $EY_t = \mu$ , autocovariance function (acf)  $\gamma_h = Cov(Y_t, Y_{t+h})$ , and integrable spectral density function  $f(w) = \sum_h \gamma_h e^{-ihw}$ . We are interested in studying the distribution of the studentized sample mean where the normalization involves the summation of sample autocovariances weighted by an arbitrary taper, and when the stochastic process exhibits either short or long memory or even when the process is overdifferenced. The latter case is especially tricky – and not well-studied in the literature – because in this case the studentization is achieved by dividing with a quantity that tends to zero. The objective in studentizing/self-normalizing the mean is the generation of a pivotal asymptotic distribution that can serve as the basis for the construction of confidence intervals and hypothesis tests for the unknown mean  $\mu$ .

In the case that the autocovariances  $\gamma_h$  are (absolutely) summable with  $\sum_h \gamma_h > 0$ , it is well-known – under regularity conditions – that the sample mean  $\bar{Y} = n^{-1} \sum_{t=1}^n Y_t$  is asymptotically normal with variance  $f(0) = \sum_h \gamma_h$ , i.e.,  $\sqrt{n}(\bar{Y} - \mu) \xrightarrow{\mathcal{L}} N(0, f(0))$ . A consistent estimate of  $f(0)$  is given by

$$W_{\Lambda, M} = \sum_{|h| \leq M} \Lambda_M(h) \tilde{\gamma}_h, \quad (1)$$

where  $\Lambda_M$  is an arbitrary taper (described in Section 4), and  $\tilde{\gamma}_h$  are the sample autocovariance estimators defined by

$$\tilde{\gamma}_k = \frac{1}{n} \sum_{t=1}^{n-|k|} (Y_{t+|k|} - \bar{Y})(Y_t - \bar{Y}) \quad \text{for } |k| < n. \quad (2)$$

As usual,  $M = M(n)$  is a bandwidth parameter tending to  $\infty$  as  $n \rightarrow \infty$  but in such a way that  $M \ll n$ . Define the *bandwidth-fraction* to be  $b = b(n) = M(n)/n$ . The literature on such taper-based, “lag-window” spectral estimators is extensive, going back over fifty years; see e.g. Hannan (1970), Brillinger (1981), Priestley (1981), Rosenblatt (1985), Brockwell and Davis (1991), and the references therein. Also see Grenander and Rosenblatt (1957), Blackman and Tukey (1959), and Percival and Walden (1993). Much is already known about the classical case where  $f(0)$  is bounded above and bounded away from zero, but also about the long memory case where  $f(0)$  is infinite – see, e.g., Beran (1994), Robinson (1994), and Palma (2007). Other recent literature includes Sun (2004) and Robinson (2005). However, little is yet known in the case that  $f(0) = 0$ , although this possibility was brought to the forefront early on by Rosenblatt (1961). We attempt to remedy this situation in the paper at hand. The case  $f(0) = 0$  will be referred to henceforth as the *superefficient (SE) case*, since it implies  $\bar{Y} = \mu + o_P(n^{-1/2})$ , i.e., superefficient estimation of the mean.

The SE case has received little attention in the literature despite the fact that it is of some importance in applied econometrics. We illustrate this through a brief discussion of the unit root problem that arises from over-differencing. Although most economic time series exhibit obvious

trends, much debate rages over whether processes are stationary or  $I(1)$ ; witness the extensive literature on unit-root testing, starting with Dickey and Fuller (1979, 1981). Hamilton (1994) gives an overview; also see Phillips and Perron (1988) and Parker, Paparoditis, and Politis (2006) and the references therein. It is commonly felt that an economic time series is rarely  $I(2)$ , and yet many such  $I(2)$  models are selected by automatic model identification software, such as X-12-ARIMA and TRAMO-SEATS (both of which often select the Box-Jenkins airline model, which is  $I(2)$ ); see the discussion in Findley, Monsell, Bell, Otto, and Chen (1998), and Maravall and Caparello (2004).

Over-differencing relates to over-specification of the order of differencing, i.e., modeling a process as  $I(1)$  when it is stationary, or as  $I(2)$  when it is only  $I(1)$ . Naturally, estimation of parameters (such as the mean and other regression effects, but also maximum likelihood estimation of ARMA parameters) is performed on the differenced series, where the nonstationarity has been removed. But if the differencing order has been over-specified, then there will be over-differencing; this results in the spectrum of the differenced data being zero at frequency zero. For example, if  $\{Y_t\}$  is stationary with spectral density  $f$  but is viewed as  $I(1)$ , then the spectrum of the data's first difference is  $|1 - e^{-i\lambda}|^2 f(\lambda)$ . We mention in passing that similar issues exist, at least in theory, for seasonal frequencies, i.e., taking a seasonal difference when the seasonal component of the time series is actually stationary. In this case, the zero in the spectrum takes place at the seasonal frequencies corresponding to the angular portions of the zeroes of the seasonal differencing operator. Such zeroes offer no impediment to the estimation of the sample mean, but may generate other problems in model estimation; we do not pursue this point further here.

One approach to this problem is to do a pre-test of a possible unit root before differencing the data (Dickey and Fuller, 1979). Also see the treatment in Tanaka (1990, 1996). However, given that Type II errors will occur some of the time (the power of the Dickey-Fuller statistic is explored in many articles, including Lopez (1997)), we advocate in this paper the use of robust studentized sample mean estimates, where the robustness is with respect to the three basic cases for  $f(0)$ : infinite (long memory), finite and positive (short memory), or zero (the SE case, distinguished in what follows as either negative memory or differential memory). To that end, we study the finite-sample and asymptotic distributional properties of studentized sample means, where the normalization is of the form (1) and the stochastic process satisfies some very general conditions. Note that  $W_{\Lambda, M}$  must mirror the properties of the variance of the sample mean under the three scenarios: for long memory, it must diverge at the appropriate rate; for short memory it should converge to the same constant; and for negative/differential memory it should tend to zero at the same rate. Our work determines what conditions on a taper are needed in order to ensure such a robustness against different alternative memory scenarios.

We first provide a common context in econometrics for the problems at hand, which is intended as a motivation for our work (Section 2). Our main goal is to understand the joint distribu-

tional properties of  $\bar{Y}$  and  $W_{\Lambda, M}$  for various stochastic processes and various tapers, distinguishing between the case that the bandwidth-fraction  $b$  is vanishing and the case that it is a constant proportion. In the former case, we obtain central limit theorems for the sample mean, while the variance estimate tends in probability to a constant when appropriately normalized (Section 4). But in the latter case, the variance estimate tends to a random limit (Section 5). In Section 3 we give precise definitions of the four types of memory (LM, SM, NM, and DM), and also discuss some basic properties. Given the limit results, it is not obvious how to proceed with testing for the mean. This is addressed in Section 6 through two proposed methods, one for the vanishing bandwidth-fraction case and one for the fixed bandwidth-fraction case. These procedures are extensively evaluated through size and power simulations, covering many stochastic processes, tapers, and bandwidth-fractions. Section 7 summarizes our findings, and proofs are gathered in an Appendix.

## 2 A Context for Anti-Persistence in Applied Econometrics

A primal modeling assumption for many economic time series of interest is that they arise from sampling an  $I(d)$  process, where  $d$  is integer. One can associate a  $d$ th degree polynomial (in time) to the process as well, as this only affects the differenced process by a mean shift. This older paradigm, discussed in Box and Jenkins (1976), has in recent decades been altered to accommodate long memory. Whereas previously the differenced process was modeled via some short memory device, such as an ARMA model, now more complex covariance structures can be entertained – see Palma (2007) for an overview.

So we might now conceive that the underlying process is  $I(D)$  plus a degree  $d$  polynomial, where  $D \in [d - 1/2, d + 1/2)$ . That is, suppose the observed data process  $\{X_t\}$  satisfies  $X_t = p(t) + Z_t$ , where  $p$  is a degree  $d$  polynomial and  $\{Z_t\}$  is  $I(D)$ . Then

$$Y_t = (1 - B)^d X_t = d! p_d + (1 - B)^d Z_t,$$

where  $B$  is the backshift operator and  $p_d$  is the leading coefficient of the polynomial  $p$ . Now  $\{(1 - B)^d Z_t\}$  is a mean zero  $I(D - d)$  process; because  $D - d \in [-1/2, 1/2)$ , the process  $\{Y_t\}$  is stationary with mean  $\mu = d! p_d$ .

If  $D - d = 0$ , then  $\{Y_t\}$  is short memory – this corresponds to the classical case alluded to above. If  $D - d > 0$ , we have a stationary long memory process. Note that if we differenced only  $b$  times instead – for integer  $b < d$  – so that  $D - b \geq 1/2$ , then  $\{Y_t\}$  would be non-stationary, and this might be detected by inspection of the sample autocorrelations. Then  $b$  could be incremented through higher integers until  $D - b$  falls into the range corresponding to stationarity. One could also proceed more rigorously by using unit root tests.

Over-differencing, whereby  $D - b < -1/2$ , would be harder to guard against. This is because in

practice both the appropriately differenced series and the over-differenced series often have sample autocorrelations consistent with a stationary hypothesis. One way this could happen is when we believe  $b$  differencings are needed to reduce the data to stationarity, but really the polynomial  $p$  is order  $b$  and the process  $\{Z_t\}$  only requires  $d = b - 1$  differencings. It may not be readily apparent that this is the case, rather than  $\{Z_t\}$  requiring the full  $b$  differences instead. So differencing  $b$  times produces a constant plus an over-differenced series. Again, unit root testing techniques can be employed to guard against this error, but there is some evidence that they have lower power in the presence of long memory (Diebold and Rudebusch, 1991). When  $b$  is correctly identified as  $b = d$ , then the differenced series is integrated with order between  $-1/2$  and  $1/2$ , implying that super-efficiency (SE) has an equal *a priori* chance of arising as long range dependence.

This discussion, which applies to many time series being examined for the purposes of forecasting, signal extraction, or other analysis, indicates how long memory, short memory, negative memory, and differential memory may arise. When the order of differencing  $b$  is selected correctly (i.e., it equals  $b$ ), then  $D \in (d, d + 1/2)$  implies that  $\{Y_t\}$  will have (stationary) long memory; if  $D = d$  then  $\{Y_t\}$  will have short memory. But if  $D \in [d - 1/2, d)$ , then  $\{Y_t\}$  will have negative memory (this is part of the SE case). Finally, if the data is over-differenced by mistakenly taking  $b > d$ , then  $(1 - B)^b X_t = (1 - B)^b Z_t$  is  $I(D - b)$  with mean zero, and  $D - b < -1/2$ . This case corresponds to differential memory (the other part of the SE case), and the sample mean has a radically different behavior. If the underlying polynomial actually has order  $b$  rather than  $d$ , then the over-differenced data will have a non-zero mean effect.

So in any of these scenarios, the differenced data  $\{Y_t\}$  is stationary with mean  $\mu$  and has one of four types of memory. The natural estimate for  $\mu$  is the sample mean  $\bar{Y}$ , essentially being an ordinary least squares estimate. A generalized least squares estimate could be contrived, but knowledge of the autocovariances would be required, and this is not possible without modeling or extensive estimation. The practitioner may desire to know  $\mu$ , because it is important for forecasting or filtering the original series  $\{X_t\}$ ; forecast functions will have different formulas when  $\mu = 0$ , so it is also of interest to test this hypothesis. If we estimate  $\mu$  via  $\bar{Y}$ , we must also assess the statistical uncertainty in order to know whether it is safe to assume  $\mu = 0$  in our applications. It is adequate to measure the second-order structure of  $\bar{Y}$  through the variance  $V_n$  of  $S_n = n\bar{Y}$  (where  $n$  is the sample size of the differenced data); knowledge of the sampling distribution of  $(S_n - n\mu)/\sqrt{V_n}$  will permit inference for  $\mu$ . As the sampling distribution is typically not available, the asymptotic distribution is utilized instead. The key to this is to estimate  $V_n$  in such a way that the studentized statistic still converges, no matter what type of memory pertains. This is the topic of the remainder of the paper.

### 3 Types of Memory

We now discuss in detail the different memory scenarios. Throughout this paper we use the notation  $A_n \sim B_n$  to denote  $A_n/B_n \rightarrow 1$  as  $n \rightarrow \infty$ . By Short Memory (SM), we refer to the condition that the autocovariances – recall that  $\gamma_h = Cov(Y_t, Y_{t+h})$  – are absolutely summable and their sum  $f(0)$  is a nonzero constant. By Long Memory (LM), we mean that the autocovariances are not summable, and the partial sums of them, denoted by  $W_n$ , satisfy

$$W_n = \sum_{|k| \leq n} \gamma_k = L(n)n^\beta. \quad (3)$$

In (3)  $L$  is slowly varying at infinity (Ebrechts, Klüppelberg, and Mikosch, 1997), with a limit that can be zero,  $C$ , or infinity, where  $C$  is a positive constant. Most importantly,  $\beta > 0$  (and is less than 1). In the SM case (3) applies with  $\beta = 0$  and  $L$  tending to  $C = \sum_k \gamma_k = f(0)$ . The case that  $\beta = 0$  but  $L$  tends to infinity is also LM (e.g., say  $\gamma_k = k^{-1}$  for  $k \geq 1$ ).

We will denote by Negative Memory (NM) the case of an absolutely summable autocovariance sequence such that (3) holds, but with  $\beta < 0$  (though the case that  $\beta = 0$  and  $L$  tends to zero is also considered to be NM). Some authors have used the term “intermediate memory” for this concept (Brockwell and Davis, 1991). Our nomenclature is due to the negative memory exponent, and also the result that most of the autocovariances are negative in this case (see Remark 2 below); the same conditions on  $L$  apply here. When the autocovariances are zero past a certain threshold, we obtain an example of Differential Memory (DM). For example, consider an  $MA(q)$  data process that is over-differenced; the resulting autocovariances are identically zero for lags exceeding  $q$ , and  $\sum_{|k| < n} \gamma_k = 0$  for  $n > q + 1$ . These definitions encompass ARFIMA models (Hosking, 1981), FEXP models (Beran (1993, 1994)), and fractional Gaussian Noise models, as well as the case of over-differenced processes. Some authors prefer to parametrize memory in terms of the rate of explosion of  $f$  or  $1/f$  at frequency zero, but it is more convenient for us to work in the time domain; see Palma (2007) for an overview.

To distinguish between the LM, SM, NM, and DM memory cases, the key determinant in the limit theorems for  $S_n = n\bar{Y}$  is the rate of growth of  $V_n = Var(S_n)$ , which in turn is related to  $\sum_{|k| \leq n} \gamma_k$ . In the rest of this paper we study  $S_n/\sqrt{\widehat{V}_n}$ , where  $\widehat{V}_n$  is some estimate of the variance  $V_n$  such that  $\sqrt{\widehat{V}_n} = O_P(S_n)$ . Recall that  $W_n = \sum_{|k| \leq n} \gamma_k$  by definition, which in turn has asymptotic behavior given by (3) by assumption; then we have the following identity:

$$V_n = \sum_{k=0}^{n-1} W_k \quad (4)$$

that is proved using summation by parts. Now for LM proceses,  $W_n$  diverges, whereas in the SM case  $W_n$  tends to a nonzero constant. The superefficient case (SE) where  $f(0) = 0$  is characterized

by  $W_n$  tending to zero; however, we distinguish the case that  $W_n$  is summable (DM case) versus the case that it is not (NM case).

**Definition 1** Define the four types of memory as follows:

- (i) LM:  $W_n$  has asymptotics given by (3) with either  $\beta \in (0, 1)$  or  $\beta = 0$  and  $L$  is tending to infinity.
- (ii) SM:  $W_n \rightarrow C$  with  $C > 0$ .
- (iii) NM:  $W_n$  has asymptotics given by (3) with either  $\beta < 0$  or  $\beta = 0$  and  $L$  is tending to zero. When  $\beta = -1$ , suppose that  $W_n$  is not summable.
- (iv) DM:  $W_n$  is summable.

The condition (3) on  $W_n$  in case (i) is denoted  $LM(\beta)$ , and the same condition on  $W_n$  in case (iii) is denoted  $NM(\beta)$ .

**Remark 1** Note that cases (i), (ii), and (iii) are mutually exclusive, since case (ii) essentially corresponds to  $\beta = 0$  in (3) with  $L$  tending to a nonzero constant. When  $\beta = -1$  in (3), it is possible for  $W_n$  to be either summable or not summable; the former case belongs to case (iv). If  $\beta < -1$  in (3), then necessarily this is case (iv). Proposition 1 below shows that most of the autocovariances are negative in case (iii), and this justifies the nomenclature of “negative memory.” In case (iv)  $\{Y_t\}$  can be represented as the difference of another stationary process (Proposition 2), and hence the appellation “differential memory.” The case of an over-differenced series, as discussed in Section 1, is always included in case (iv). Both NM and DM have  $f(0) = 0$ , so both are part of the SE case.

**Remark 2** These four cases are mutually exclusive but do not cover all possibilities, since we can consider a function  $L$  in (3) that is not slowly-varying (e.g.,  $L(n) = 1 + \sin(n)$ ). However, the four cases *are* exhaustive for processes satisfying (3) with  $L$  slowly-varying and  $\beta < 1$ .

The following representation for slowly-varying functions (Theorem A3.3 of Embrechts, Klüppelberg, and Mikosch, 1997) will be used below: we can write  $L$  in the form

$$L(n) = c(n) \exp \int_z^n \eta(u)/u \, du, \tag{5}$$

with  $c(n) \rightarrow c$  a positive constant,  $z$  some fixed positive constant, and  $\eta(u)$  tending to zero as  $u \rightarrow \infty$ . Note that in (5) we can take the variable argument of  $L$  to be a continuous variable  $x$ . The following result gives the behavior of  $V_n$  in each of the four cases described in Definition 1, and also discusses the implied asymptotic behavior of  $\gamma_k$ .



**Proposition 1** With  $W_n$  and  $V_n$  given by (3) and (4) respectively,

$$V_n \sim \frac{1}{\beta+1} nW_n = \frac{L(n)n^{\beta+1}}{\beta+1} \quad (6)$$

for cases (i) and (ii), and for  $\beta > -1$  in case (iii). When  $\beta = -1$  we have  $V_n/(nW_n) \rightarrow \infty$  (so in a sense (6) holds true). In case (ii), we let  $\beta = 0$  in the formula and  $L(n) \sim f(0)$ . In case (iv),  $V_n \rightarrow \sum_{k \geq 0} W_k$ . If  $\beta \neq 0$  in cases (i) and (iii), we also have

$$\gamma_n \sim \frac{\beta}{2} L(n)n^{\beta-1}. \quad (7)$$

In cases (i) and (iii) with  $\beta = 0$  we have  $\gamma_n \sim -.5 \int_{n-1}^n [\eta(u)/u] du = o(n^{-1})$ .

This result shows that  $V_n \rightarrow \infty$  in cases (i), (ii), and (iii), as long as  $\beta > -1$ . This will facilitate a fairly standard limit theorem for  $\bar{Y}$  under some additional conditions. Case (iv) produces a very different sort of limit theorem; these results are discussed in Section 3 below.

**Remark 3** Note that  $L(n)$  must be non-negative for all  $n$  larger than some  $n_0$ , say; this follows from (6) and the fact that  $V_n > 0$  for all  $n$ . Hence, for large  $n$  all the  $\gamma_n$  are *negative* in the NM case and *positive* in the LM case by (7). Essentially, NM is due to heavy negative correlation and LM to heavy positive correlation; this justifies the name “negative memory” for the NM case.

**Example 1** Let  $\beta > 0$  with  $\gamma_h = h^{-\beta}$  for  $h \geq 1$  and  $\gamma_0$  chosen suitably large to guarantee the sequence is positive definite; then this is the acf of a LM process. If we (temporally) difference the process, then the resulting acf is  $2\gamma_h - \gamma_{h+1} - \gamma_{h-1}$ , which equals

$$2h^{-\beta} - (h+1)^{-\beta} - (h-1)^{-\beta} = -\beta(\beta+1)h^{-(\beta+2)} + o(1)$$

when  $h \geq 2$ . Although this appears at first to be NM (comparing to (7)), in fact it can be shown that  $W_n$  is summable so that the differenced process is DM (see Proposition 2 below).

**Example 2** Let  $\beta < 0$  and  $\gamma_h = -h^{\beta-1}$  for  $h > 0$ , and  $\gamma_0 = 2 \sum_{h \geq 1} h^{\beta-1}$ . Then the discrete fourier transform of  $\{\gamma_h\}$  is  $2 \sum_{h \geq 1} h^{\beta-1} (1 - \cos \lambda h)$  for  $\lambda \in [-\pi, \pi]$ , which is always non-negative. Hence  $\{\gamma_h\}$  is the acf of a time series process; by (7) it seems to have the form of a NM acf, but we must check the summability of  $W_n$ . Direct calculation shows that

$$W_n = 2 \sum_{h > n} h^{\beta-1} = O(n^\beta).$$

Thus if  $\beta \geq -1$  the process is NM, but otherwise is DM.

In case (ii) there is no result of the form (7), but in case (iv) the autocovariances have a particular structure if we suppose in addition that  $\{Y_t\}$  is purely nondeterministic. This is discussed in the proposition below, whose result is similar to Theorem 8.6 of Bradley (2007).

**Proposition 2** Suppose that  $W_k$  is summable and  $\{Y_t\}$  is purely nondeterministic with mean  $\mu$  and acf  $\{\gamma_h\}$ . Then there exists a strictly stationary process  $\{Z_t\}$  with autocovariance sequence  $r_k$  such that  $Y_t = Z_t - Z_{t-1} + \mu$  and  $\gamma_k = 2r_k - r_{k+1} - r_{k-1}$ . This decomposition is not unique. Also  $W_k = 2(r_k - r_{k+1})$  and  $r_0 = \sum_{k \geq 0} W_k/2$ . Conversely, given any stationary process  $\{Z_t\}$  with acf  $r_k$  tending to zero, the differenced process  $Z_t - Z_{t-1}$  is DM.

We note that the process  $\{Z_t\}$  might be LM, SM or NM, and might even be DM.

**Example 3** Let  $\{Z_t\}$  be iid and let  $Y_t = Z_t - Z_{t-1}$ . Then clearly  $\{Y_t\}$  is DM with autocorrelation of  $-.5$  at lag one and zero at higher lags. If the innovation variance is unity, then  $W_0 = 2$  and  $W_k = 0$  for  $k \geq 1$ ; this is clearly a summable sequence.

**Example 4** A more interesting example is given as follows: let  $Y_t = \sum_{j=0}^{\infty} \psi_j \epsilon_{t-j}$  where  $\{\epsilon_t\}$  is white noise and  $\psi_j = (-1)^j/j^p$  for  $p > 0$  and  $j \geq 1$ , with  $\psi_0 = -\sum_{j=1}^{\infty} \psi_j$  (which clearly exists by the alternating series test). Hence  $\sum_j \psi_j = 0$ , which implies that  $\sum_k \gamma_k = 0$ . Notice that the variance of  $Y_t$  only exists if  $p > 1/2$ , since

$$\sum_{j=0}^{\infty} \psi_j^2 = \psi_0^2 + \sum_{j=1}^{\infty} j^{-2p} = F^2(1/2, p) + F(1, 2p),$$

where  $F(x, s)$  is the periodic-zeta function given by  $F(x, s) = \sum_{n \geq 1} e^{2\pi i x n} n^{-s}$ . The variance of  $\sum_{j=0}^n \psi_j \epsilon_{t-j}$  is  $\sum_{j=1}^n j^{-2p} + \psi_0^2 \sim n^{1-2p}$ , which diverges unless  $p > 1/2$ ; we assume this henceforth, as we are not concerned with infinite variance time series in this paper. As in the proof of Proposition 2, define

$$\theta_j = \sum_{k=0}^j \psi_k = - \sum_{k=j+1}^{\infty} \frac{(-1)^k}{k^p}.$$

The sum of any two consecutive terms of this series, up to a minus sign, can be written as  $k^{-p} - (k+1)^{-p} = ((1+1/k)^p - 1)/(k+1)^{-p}$ , which by Taylor series expansion about zero yields an approximation of  $p k^{-1}(k+1)^{-p}$  plus terms that decay at order  $k^{-2-p}$ . Thus asymptotically, the sum of consecutive terms in  $\theta_j$  is  $p k^{-p-1}$ , plus other terms that decay even faster. So such a sequence is summable, and we find that  $\theta_j = O(j^{-p})$  as  $j \rightarrow \infty$ . Since  $p > 1/2$ , this sequence is square summable. This implies that the time series  $Z_t = \sum_{j \geq 0} \theta_j \epsilon_{t-j}$  is well-defined, i.e., is finite almost surely, since it has finite variance. Clearly  $Y_t = Z_t - Z_{t-1}$ , and the other assertions of Proposition 2 apply; in particular,  $\{Y_t\}$  is DM.

**Example 5** Suppose that an observed time series  $\{X_t\}$  is an ARFIMA  $(0, D, 0)$ , so that  $(1 - B)^D X_t = \epsilon_t$  is white noise, where  $D \in [0, 1]$ . If  $D = 1$  this is just a random walk, and if  $D \in [.5, 1)$  the process is said to have nonstationary long memory. If  $D < .5$  the process is stationary, but with long memory if  $D > 0$ . Of course  $D = 0$  corresponds to short memory (white noise). If the observed process is differenced once to produce  $Y_t = X_t - X_{t-1}$ , it is easy to see that the result is stationary

with memory parameter  $\beta = 2D - 2$ . That is, if  $D = 1$  we obtain short memory (white noise); if  $D \in [.5, 1)$  we obtain a negative memory process of parameter  $\beta \in [-1, 0)$ ; if  $D < .5$  then we obtain a process with differential memory. The borderline case  $D = .5$  is interesting: we don't get a differential memory process, since the original process is nonstationary – instead we get a negative memory process with  $\beta = -1$  and nonsummable  $W_k$  sequence.

## 4 Limit Theory for the Case of Vanishing Bandwidth-Fraction

In the case that  $b(n) \rightarrow 0$  as  $n \rightarrow \infty$ , we can treat the asymptotics of  $S_n$  and  $W_{\Lambda, M}$  separately (recall that  $W_{\Lambda, M}$  was defined in (1)), because the variance estimate, when appropriately normalized, always converges to a constant. Let us then consider the partial sums first; it is necessary to impose some additional assumptions. Typical assumptions for limit theorems involve either moment and mixing conditions, or linearity of the process involved. Limit theorems have also been derived under the assumption that the given process is a direct function of an underlying Gaussian process. It turns out (see discussion below) that mixing assumptions are not helpful when (3) holds. Instead, one can make fairly strong assumptions on the higher order cumulants of the time series. The  $k$ th order cumulant of  $\{Y_t\}$  is defined by  $c_k(u_1, u_2, \dots, u_{k-1}) = \text{cum}(Y_{t+u_1}, Y_{t+u_2}, \dots, Y_{t+u_{k-1}}, Y_t)$  for any  $t$  and integers  $u_1, \dots, u_{k-1}$ , where  $k \geq 1$  (Taniguchi and Kakizawa, 2000). Letting  $u$  denote the  $k - 1$ -vector of indices, we write  $c_k(u)$  for short. Also let  $\|\cdot\|$  denote the sup-norm of a vector, so that  $\sum_{\|u\| < n} c_k(u)$  is a short-hand for summing the cumulant over all indices such that  $|u_j| < n$  for each  $j$ . We also require the concept of Hermite rank (Taqqu, 1975): if  $g \in \mathbb{L}^2(\mathbb{R}, e^{-x^2/2})$ , then it can be expanded in terms of the Hermite polynomials  $H_k$ , with coefficients  $\langle g, H_k \rangle$  (the bracket denotes the inner product of the Hilbert space) for  $k \geq 0$ . The Hermite rank is the index of the first nonzero coefficient. Given these notations, the assumptions on our process  $\{Y_t\}$  that we will consider are given below:

- **Process P1.**  $\{Y_t\}$  is a linear process:  $Y_t = \sum_j \psi_j \epsilon_{t-j}$  with  $\{\psi_j\}$  square summable and  $\{\epsilon_t\}$  iid with finite variance. Also,  $\{Y_t\}$  is either (i) LM( $\beta$ ) with  $\beta \in [0, 1)$ ; (ii) SM; or (iii) NM( $\beta$ ) with  $\beta \in [-1, 0]$ .
- **Process P2.**  $Y_t = g(X_t)$  for each  $t$ , where  $g$  is a function in  $\mathbb{L}^2(\mathbb{R}, e^{-x^2/2})$  of Hermite rank  $\tau$ , and  $\{X_t\}$  is a Gaussian process with autocovariance  $r_k$ . Assume that  $\{X_t\}$  is either (i) LM( $\beta$ ) with  $\beta \in [0, 1)$ ; (ii) SM; or (iii) is NM( $\beta$ ) with  $\beta \in (-1, 0)$ . In case (i) we also assume that  $(1 - \beta)\tau < 1$ , but in cases (ii) and (iii) there is no restriction on  $\tau$ .
- **Process P3.**  $\{Y_t\}$  is a process that is either (i) LM( $\beta$ ) with  $\beta \in [0, 1)$ ; (ii) SM; or (iii) NM( $\beta$ ) with  $\beta \in (-1, 0]$ . Also assume the  $k$ th order cumulants exist and are summable over its  $k$  indices, for all  $k \geq 1$ . Moreover, when  $\beta < 0$  we also assume that  $\sum_{\|u\| < n} c_k(u) = O(n^\beta)$ .

**Remark 4** Yet another type of assumption would require  $2 + \delta$  moments (for some  $\delta > 0$ ) and a mixing assumption. It is known that the strong mixing condition of Rosenblatt (1956)<sup>1</sup> is not satisfied by long memory Gaussian processes – Gaussian processes are strongly mixing iff their spectrum is bounded away from zero and infinity (Kolmogorov and Rozanov, 1960). If the process is not Gaussian, it is conceivable that a strong mixing condition could be satisfied, although no examples of this have been published. The weak dependence condition of Doukhan and Louhichi (1999) might be used instead, because in Bardet, Doukhan, Land, and Ragache (2008) it is shown to be compatible with long range dependence. Along with a Lindeberg condition on higher moments of the partial sum, a central limit theorem can be established utilizing the Bernstein blocks method described in Rosenblatt (1956) and Doukhan and Louhichi (1999). Unfortunately, such a result is not compatible with (3) when  $\beta \neq 0$ : examination of Theorem 1 of Rosenblatt (1984) indicates that the rate of convergence in this type of CLT must be  $k_n V_{p_n}$ , where  $k_n$  is the number of big blocks and  $p_n$  is the size of the big blocks, and moreover  $n \sim k_n p_n$ . This is not a paradox: in order to prove a CLT, fairly strong assumptions must be placed upon the mixing coefficients – either they must be strong mixing, or must have rapid decay rate if they are weak dependence coefficients, etc. – which then actually precludes the possibility of (3). For this reason, we don't formulate limit theorems under mixing conditions in this paper.

Each of the above assumptions P1, P2, and P3 provide sufficient conditions for a limit theorem for  $S_n$ , as shown below. Note that these cover only cases of LM, SM, and NM; the DM case must be handled separately in what follows. The  $\beta = -1$  case is only handled under P1. It is clear that P1 and P2 cover distinct cases, because in the latter the process can be nonlinear. The restriction on  $\tau$  in the LM case of P2 is for convenience – limit results are also available for the case that  $(1 - \beta)\tau > 1$ , though the boundary case  $(1 - \beta)\tau = 1$  cannot be handled without additional knowledge about the acf. Note that in the case of P3 with a Gaussian process, all the cumulant assumptions are automatically satisfied. In any of these scenarios, the assumptions are typically unverifiable, or difficult to verify from the data; these should be viewed as working assumptions.

The following result covers the cases of LM, SM, and NM under the three different conditions P1, P2, and P3. We also establish a result for the DM process, restricting to purely non-deterministic processes so that Proposition 2 can be applied.

**Theorem 1** *Suppose that  $\{Y_t\}$  is a strictly stationary process with finite variance that satisfies (3). If the process satisfies one of P1, P2, or P3, then  $\frac{S_n - n\mu}{\sqrt{V_n}} \xrightarrow{\mathcal{L}} B$  as  $n \rightarrow \infty$ , where  $B$  is an absolutely continuous random variable that is standard normal except when  $\tau > 1$  in the LM case of P2. If  $\{Y_t\}$  is purely non-deterministic with summable  $W_k$ , and if the process  $\{Z_t\}$  of Proposition 2 is*

---

<sup>1</sup>A stationary process  $\{Y_t\}$  is strongly mixing (Rosenblatt, 1956) if  $\alpha_k \rightarrow 0$  as  $k \rightarrow \infty$  where  $\alpha_k = \sup_{A \in \mathcal{F}_{-\infty}^0, B \in \mathcal{F}_k^\infty} |\mathbb{P}(A \cap B) - \mathbb{P}(A)\mathbb{P}(B)|$ , and  $\mathcal{F}_j^m$  is the  $\sigma$ -algebra generated by  $\{Y_k, j \leq k \leq m\}$ .

either linear or strong mixing, then  $\frac{S_n - n\mu}{\sqrt{V_n}} \xrightarrow{\mathcal{L}} (Z_* - Z_0)/\sqrt{2r_0}$  as  $n \rightarrow \infty$ , where  $Z_*$  is a random variable equal in distribution to  $Z_0$ , but independent of it.

To utilize Theorem 1 it is important to know  $V_n$ , or to estimate it. In view of (3), the taper-based estimate given by  $W_{\Lambda, M}$  in (1) is a nonparametric estimate, and can be related to estimation of  $V_n$  via (4) and (6). For example, with an SM process  $\widehat{V}_n = nW_{\Lambda, M}$  is often used as an estimator.

The asymptotic behavior of  $W_{\Lambda, M}$  depends on the type of taper as well as the type of memory of the process. The tapers that we consider are very general:  $\Lambda_M$  is a piecewise smooth (i.e., piecewise differentiable), even function on the integers such that  $\Lambda_M(h) = 0$  for  $|h| \geq M$ . Letting  $U_M$  denote the maximum value of  $\Lambda_M(h)$  for all  $h$ , we suppose that  $U_M$  does not grow too fast as  $M \rightarrow \infty$ . Classical tapers are bounded, in which case  $U_M$  can be taken constant. The triangular (Bartlett) kernel, the trapezoidal (Politis and Romano, 1995), and the more general flat-top kernels (Politis 2001, 2005) all satisfy these conditions.

Our next main result is that under some conditions the estimator  $W_{\Lambda, M}$  is asymptotic to the deterministic sequence  $\widetilde{W}_M$  as  $M \rightarrow \infty$  (recall that  $M/n \rightarrow 0$ , though), which is defined via

$$\widetilde{W}_M = \sum_h \Lambda_M(h) \left(1 - \frac{|h|}{n}\right) \gamma_h. \quad (8)$$

We require the following condition, which parallels Assumption A of Andrews (1991), to establish our result:

**Assumption B:**  $\mathbb{E}|Y_t^4| < \infty$  and the fourth order cumulant  $c_4$  is absolutely summable.

The above assumption is compatible with the process conditions P1, P2, and P3 for the LM, SM, and NM cases. Assumption B is also compatible with the DM case. As discussed in Andrews (1991), linear processes with absolutely summable coefficients and finite fourth moments satisfy Assumption B, even if the process is LM. As Lemma 1 of Andrews (1991) shows, Assumption B is also implied by a strong mixing plus moments condition. Of course, a long memory Gaussian process trivially satisfies Assumption B, because its fourth order cumulants are zero.

**Proposition 3** *Suppose that  $\{Y_t\}$  is a strictly stationary process satisfying Assumption B, which is either LM, SM, NM, or DM. Also suppose that  $b(n) + 1/M(n) \rightarrow 0$  as  $n \rightarrow \infty$ . Then  $W_{\Lambda, M} = \widetilde{W}_M(1 + o_P(1))$ .*

Next we consider the asymptotics of the deterministic sequence  $\widetilde{W}_M$  (8), which depends upon the memory assumptions. Together with Proposition 3, this will show that  $W_{\Lambda, M}$  is asymptotic to a constant times  $W_M$ . This constant will be denoted by  $\zeta$ , and depends upon  $\beta$  and the taper used. We consider tapers of the form  $\Lambda_M(h) = \Lambda(h/M)$  for a fixed function  $\Lambda(x)$ . We assume  $\Lambda$  is an even, piecewise smooth function, that is real-analytic on every such interval; by  $\dot{\Lambda}_+(x)$  for  $x \geq 0$ , we denote the derivative from the right (and  $\Lambda_+^{(j)}(x)$  for higher order derivatives). An example is

a “flat-top” taper where there exists an interval  $[0, c]$  (with  $c > 0$ ) for which  $\Lambda$  equals unity – see Politis (2001). Then define  $\zeta = 1$  when  $\Lambda$  is the truncation taper (i.e.,  $\Lambda = 1_{[-1,1]}$ ), and otherwise

$$\zeta = - \int_0^1 \dot{\Lambda}_+(x) x^\beta dx. \quad (9)$$

Then we have the following result.

**Theorem 2** *Let  $\Lambda(x)$  be an even, piecewise differentiable function supported on  $[-1, 1]$ , with  $\Lambda_M(h) = \Lambda(h/M)$ . Suppose that  $\{Y_t\}$  is a strictly stationary process satisfying Assumption B, which is either LM, SM, NM, or DM. Also suppose that  $b(n) + 1/M(n) \rightarrow 0$  as  $n \rightarrow \infty$ . Then if the process is LM, SM, or NM with  $\beta \in (-1, 1)$ ,*

$$\frac{\widetilde{W}_M}{W_M} \rightarrow \zeta \text{ as } M \rightarrow \infty. \quad (10)$$

*If the process is DM, then*

$$\widetilde{W}_M \sim -2r_0 \dot{\Lambda}_+(0) M^{-1} (1_{\{c=0\}} + o(1)).$$

In the SM case recall that  $C \sim \sum_k \gamma_k$ , so that (9) with  $\beta = 0$  yields  $\zeta = \Lambda(0)$ ; this equals unity as long as  $\Lambda(0) = 1$ , which is commonly assumed. Thus  $W_{\Lambda, M} \sim C\Lambda(0)$  and  $V_n \sim nC$  can be estimated by  $nW_{\Lambda, M}$  when  $\Lambda(0) = 1$ . In the DM case for non-flat-top kernels, the overall error is controlled by the first derivative  $\dot{\Lambda}_+(0)$ . This is because  $c = 0$ ; but when  $c > 0$ , the rate of decay is even faster, and is hard to describe in a general result. For example, with the Bartlett taper  $\dot{\Lambda}_+(0) = -1$ , and  $\widetilde{W}_M \sim 2r_0 M^{-1}$ . The quantity  $\zeta$  is asymptotic to one plus the relative bias  $(\widetilde{W}_M - W_M)/W_M$ , measuring the asymptotic discrepancy between our variance estimate  $W_{\Lambda, M}$  and the sequence  $W_M$ . We refer to  $\zeta$  as the *quotient bias* hereafter. Note that  $\zeta$  is well-defined, since the derivative of  $\Lambda$  exists almost everywhere.

**Remark 5** It follows from Proposition 3, Theorem 2, and Proposition 1 – when their respective conditions are satisfied – that in probability

$$\begin{aligned} MW_{\Lambda, M} &\sim (\beta + 1)\zeta V_M \\ MW_{\Lambda, M} &\sim -2r_0 \dot{\Lambda}_+(0) (1_{\{c=0\}} + o(1)) \end{aligned} \quad (11)$$

for the cases of LM/SM/NM and DM respectively. Although  $V_M = Var(S_M)$  is of some interest, we really need to obtain a quantity asymptotic to  $V_n$ . Unfortunately, since  $b(n) = M(n)/n \rightarrow 0$ , we have  $V_M/V_n \rightarrow 0$  except in the DM case. So we cannot normalize  $S_n$  by  $\sqrt{MW_{\Lambda, M}}$  in the LM/SM/NM case, since the normalization rate is not correct (also there is the matter of the unknown  $(\beta + 1)\zeta$  factor). In the DM case, if we use a non-flat-top taper, we can indeed utilize  $S_n (-MW_{\Lambda, M}/\dot{\Lambda}_+(0))^{-1/2}$  since  $2r_0 = \sum_{k \geq 0} W_k = V_\infty$ ; by Theorem 1, this studentized statistic converges to  $Z_* - Z_0$ .

Now if  $\beta > 0$ , then  $\zeta$  can be rewritten as  $\beta \int_0^1 \Lambda(x)x^{\beta-1} dx$  using integration by parts piecewise. In this case, we proceed to calculate  $\zeta$  for some commonly used tapers. The simplest flat-top taper is the trapezoidal taper of Politis and Romano (1995) given by

$$\Lambda^{T,c}(x) = \begin{cases} 1 & \text{if } |x| \leq c \\ \frac{|x|-1}{c-1} & \text{if } c < |x| \leq 1 \\ 0 & \text{else} \end{cases}$$

with  $c \in (0, 1]$ . Then it follows that

$$\zeta^{T,c} = - \int_0^1 \dot{\Lambda}^{T,c}(x)x^\beta dx = \frac{1 - c^{\beta+1}}{(1-c)(1+\beta)}.$$

Also in the DM case, it can be shown rather easily that  $M\widetilde{W}_M \sim \frac{2}{1-c}(r_{[cM]} - r_M)$ , which tends to zero at a rate that depends upon the autocovariance sequence  $\{r_k\}$  and the truncation  $c$ . The triangular (Bartlett's) taper is obtained as the limiting case of  $\Lambda^{T,c}$  as  $c \rightarrow 0$ ; in this case,  $\zeta^{Bar} = 1/(\beta + 1)$ . Interestingly, the factor  $(1 + \beta)\zeta$  appearing in (11) is then unity for any  $\beta \in (-1, 1)$ . Also since  $\dot{\Lambda}_+(0) = -1$  for the Bartlett, we have  $MW_{\Lambda,M} \sim V_M$  for all cases, including DM.

The asymptotic quotient bias  $\zeta$  is also easily computed for the Parzen taper, given by

$$\Lambda^{Par}(x) = \begin{cases} 1 - 6|x|^2 + 6|x|^3 & \text{if } |x| \leq 1/2 \\ 2(1 - |x|)^2 & \text{if } 1/2 < |x| \leq 1 \\ 0 & \text{if } |x| > 1 \end{cases}.$$

Then  $\zeta^{Par} = (2 - (1/2)^\beta)(3/(\beta + 3) - 6/(\beta + 2) + 3/(\beta + 1))$ .

## 5 Limit Theory for the Case of Fixed Bandwidth-Fraction

The vanishing bandwidth-fraction results of Theorem 2 and Remark 5 indicate a difficulty with using tapers when LM or NM is present, because it is difficult to capture the correct rate for all types of memory. In fact, there is an asymptotic distortion equal to the quotient bias  $\zeta$  that depends on the unknown  $\beta$  parameter. In addition, there is the presence of the slowly-varying function  $L$ . These problems can be resolved – except in the DM case – by using a fixed bandwidth-fraction approach, which is described in this section.

As in Kiefer and Vogelsang (2005), let the bandwidth  $M$  be proportional to sample size  $n$ , i.e.,  $M = bn$  with  $b \in (0, 1]$ . We stress that in this section  $b$  is a fixed number, and does not grow with  $n$  as in the previous Section. Let  $\widehat{S}_i = \sum_{t=1}^i (Y_t - \bar{Y})$  (so that  $\widehat{S}_n = 0$ ). A derivative of  $\Lambda$  from the left is denoted  $\dot{\Lambda}_-$ , whereas the second derivative is  $\ddot{\Lambda}$ . The greatest integer function is denoted by

[·]. We consider tapers from the following family:

$$\{\Lambda \text{ is even and equals unity for } |x| \leq c, c \in [0, 1]. \text{ Furthermore, } \Lambda \text{ is supported on } [-1, 1], \\ \text{is continuous, and is twice continuously differentiable on } (c, 1) \cup (-1, -c).\} \quad (12)$$

This assumption is slightly less restrictive than the conditions of Theorem 2, because we only require two continuous derivatives. This family of tapers includes the family of “flat-top” kernels of Politis (2005) where  $c > 0$ , as well as the Bartlett kernel (letting  $c = 0$  and a linear decay of  $\Lambda$ ), and other kernels considered in Kiefer and Vogelsang (2005). The following result was proved in McElroy and Politis (2009), and is restated here for convenience.

**Proposition 4** (McElroy and Politis, 2009) *Let  $\Lambda$  be a kernel from family (12), and let the bandwidth be  $M = bn$ . Let  $b \in (0, 1]$  be a constant bandwidth-fraction. Then*

$$\begin{aligned} nW_{\Lambda, M} &= \sum_{i, j=1}^n \widehat{S}_i \widehat{S}_j \left( 2\Lambda \left( \frac{i-j}{M} \right) - \Lambda \left( \frac{i-j+1}{M} \right) - \Lambda \left( \frac{i-j-1}{M} \right) \right) \\ &= -\frac{2}{bn} \sum_{i=1}^{n-[c bn]} \widehat{S}_i \widehat{S}_{i+[c bn]} \left( \dot{\Lambda}_+(c) + \frac{1}{2bn} \ddot{\Lambda}(c) + O(n^{-2}) \right) \\ &\quad - \frac{1}{b^2 n^2} \sum_{[c bn] < |i-j| < [bn]} \widehat{S}_i \widehat{S}_j \left( \ddot{\Lambda} \left( \frac{|i-j|}{bn} \right) + O(n^{-1}) \right) + \frac{2}{bn} \sum_{i=1}^{n-[bn]} \widehat{S}_i \widehat{S}_{i+[bn]} \left( \dot{\Lambda}_-(1) + O(n^{-1}) \right). \end{aligned}$$

**Remark 6** In case the taper is continuously differentiable at  $c$ ,  $\dot{\Lambda}_+(c) = 0$  and the second derivative becomes dominant in the first term, which can then be recombined with the second term to yield

$$-\frac{1}{b^2 n^2} \sum_{[c bn] \leq |i-j| < [bn]} \widehat{S}_i \widehat{S}_j \left( \ddot{\Lambda} \left( \frac{|i-j|}{bn} \right) + O(n^{-1}) \right).$$

Likewise, if there is no kink at  $|x| = 1$ , then  $\dot{\Lambda}_-(1) = 0$  and the third term vanishes completely.

In order to apply this result, we need functional limit theorems for the partial sums, since  $\widehat{S}_i = S_i - i/n S_n$ . For the LM, SM, and NM cases such limit theorems can be proved which extend Theorem 1 under more restrictive conditions; the DM case is treated separately.

Functional limit theorems are often formulated in the Skorohod space, denoted  $D[0, 1]$  – see Karatzas and Shreve (1991). Because it is more convenient to prove tightness in  $C[0, 1]$ , the space of continuous functions, we will construct a linearly-interpolated process for the partial sums, and prove its convergence to Fractional Brownian Motion (FBM), which is defined in Samorodnitsky and Taqqu (1994). This provides a result of independent interest (Marinucci and Robinson (2000) work in  $D[0, 1]$ ), and also facilitates our main application, given in Theorem 4 below, under fairly simple conditions. One additional stricture, which seems unavoidable, is the requirement for higher moments in the NM case; this is because the sample paths are less smooth in the NM case than



the LM case, so that the tightness criterion is satisfied only by requiring higher moments. (The same problem affects results in the Skorohod space, as shown in Marinucci and Robinson (2000).)

So we consider the step function sum process  $S_n(t) = S_{[nt]}$ , and its linear interpolant  $\xi_n(t) = S_{[nt]} + (nt - [nt])Y_{[nt]+1}$ . The step function process is mean-centered at  $[nt]\mu$ , while the latter is mean-centered by  $nt\mu$ ; both will be normalized by the sequence  $\sqrt{V_n}$ . It is immediate that  $S_n \in D[0, 1]$  and  $\xi \in C[0, 1]$ .

**Theorem 3** *Let  $\kappa = 2 \wedge [2/(1 + \beta)]$  and suppose that  $\{Y_t\}$  is a strictly stationary process satisfying (3), and with moments of order  $\kappa + \delta$  for some  $\delta > 0$ . Moreover, suppose that  $\mathbb{E}[|S_n - n\mu|^{\kappa+\delta}] = O(V_n^{(\kappa+\delta)/2})$ . Furthermore, assume that the process satisfies one of P1, P2, or P3, with  $\beta \in (-1, 1)$ , and that in the case of a P2 process with  $\beta > 0$ , the Hermite rank is unity. Then as  $n \rightarrow \infty$*

$$V_n^{-1/2} (\xi_{[n\cdot]} - n\mu) \xrightarrow{\mathcal{L}} B \quad (13)$$

*in the sense that the corresponding probability measures on  $C[0, 1]$  converge weakly.  $B$  is a FBM process of parameter  $\beta$ .*

Letting  $\widehat{S}_n(t) = S_n(t) - \frac{[tn]}{n} S_n$  and  $\widehat{\xi}_n(t) = \xi_n(t) - \frac{[tn]}{n} \xi_n(1)$ , it follows from Theorem 3 that  $\widehat{S}_{[rn]}/\sqrt{V_n}$  converges weakly to the process  $\widetilde{B}(r) = B(r) - rB(1)$ , which is a Fractional Brownian Bridge (FBB). Then putting Proposition 4 and Theorem 3 together – with the fact that  $\widehat{S}_n$  and  $\widehat{\xi}_n$  are equivalent stochastic processes – yields the following result.

**Theorem 4** *Let  $\Lambda$  be a kernel from family (12), and let the bandwidth  $M = bn$ . Let  $b \in (0, 1]$  be a constant bandwidth-fraction. Let  $\kappa = 2 \wedge [2/(1 + \beta)]$  and suppose that  $\{Y_t\}$  is a strictly stationary process satisfying (3), and with moments of order  $\kappa + \delta$  for some  $\delta > 0$ . Moreover, suppose that  $\mathbb{E}[|S_n - n\mu|^{\kappa+\delta}] = O(V_n^{(\kappa+\delta)/2})$ . Furthermore, assume that the process satisfies one of P1, P2, or P3, with  $\beta \in (-1, 1)$ , and that in the case of a P2 process with  $\beta > 0$ , the Hermite rank is unity. Then as  $n \rightarrow \infty$*

$$\frac{S_n - n\mu}{\sqrt{nW_{\Lambda, M}}} \xrightarrow{\mathcal{L}} \frac{B(1)}{\sqrt{Q(b)}}, \quad (14)$$

*where (recall that  $c$  is defined in (12))*

$$Q(b) = -\frac{2}{b} \dot{\Lambda}_+(c) \int_0^{1-cb} \widetilde{B}(r) \widetilde{B}(r+cb) dr - \frac{1}{b^2} \int_{cb < |r-s| < b} \widetilde{B}(r) \widetilde{B}(s) \ddot{\Lambda} \left( \frac{|r-s|}{b} \right) dr ds \quad (15)$$

$$+ \frac{2}{b} \dot{\Lambda}_-(1) \int_0^{1-b} \widetilde{B}(r) \widetilde{B}(r+b) dr.$$

The case of a DM process is treated separately below, because the requisite conditions are fairly different.

**Theorem 5** *Let  $\Lambda$  be a kernel from family (12), and let the bandwidth  $M = bn$ . Let  $b \in (0, 1]$  be a constant bandwidth-fraction. Suppose that  $\{Y_t\}$  is a strictly stationary purely nondeterministic*

process with finite variance and summable  $W_k$ . Also suppose that the process  $\{Z_t\}$  of Proposition 2 is either linear or strong mixing. Let  $\tilde{C}(r) = (rZ_* + (1-r)Z_0)/\sqrt{r_0}$ ; here  $Z_*$  is a random variable equal in distribution to  $Z_0$ , but independent of it, and recall that  $r_0$  is the variance of the  $\{Z_t\}$  process. Then as  $n \rightarrow \infty$

$$\frac{S_n - n\mu}{\sqrt{nW_{\Lambda,M}}} \xrightarrow{\mathcal{L}} \frac{\tilde{C}(1) - \tilde{C}(0)}{\sqrt{P(b)}}, \quad (16)$$

where  $P(b)$  is defined as the sum of  $-2b^{-1}\dot{\Lambda}_+(0)1_{\{c=0\}}$  and  $Q(b)$  given by

$$\begin{aligned} Q(b) = & -\frac{2}{b}\dot{\Lambda}_+(c) \int_0^{1-cb} \tilde{C}(r)\tilde{C}(r+cb) dr - \frac{1}{b^2} \int_{cb < |r-s| < b} \tilde{C}(r)\tilde{C}(s)\tilde{\Lambda}\left(\frac{|r-s|}{b}\right) dr ds \\ & + \frac{2}{b}\dot{\Lambda}_-(1) \int_0^{1-b} \tilde{C}(r)\tilde{C}(r+b) dr. \end{aligned}$$

**Remark 7** The first result (14) was derived in a preliminary calculation in McElroy and Politis (2009), and the distribution (15) has been tabulated. The joint distribution of  $B(1)$  and  $Q(b)$  was also explored in McElroy and Politis (2011) through the device of the joint Fourier-Laplace Transform. The DM case (16) is novel here. Note that the first term of  $P(b)$  in the  $c = 0$  case, i.e.,  $-2b^{-1}\dot{\Lambda}_+(0)$  agrees with the first term in the expansion of  $\tilde{W}_M$  in the DM case of Theorem 2, since in the vanishing bandwidth-fraction case we have  $nW_{\Lambda,M} \sim -2r_0\frac{n}{M}\dot{\Lambda}_+(0)$  plus higher order terms.

**Remark 8** The expression for  $P(b)$  in the DM case really reduces to a quadratic in  $Z_*/\sqrt{r_0}$  and  $Z_0/\sqrt{r_0}$ , the coefficients of which can be calculated in terms of the taper's derivatives. However, knowing these quantities is not helpful towards understanding the distribution of  $P(b)$ , since the distribution of  $Z_*$  is unknown.

Unlike the special case studied by Kiefer, Vogelsang, and Bunzel (2000), the numerator  $B(1)$  of (14) is not independent of the denominator  $Q(b)$  if  $\beta \neq 0$ . To elaborate, Kiefer et al. (2000) considered the case  $b = 1$  and  $c = 0$ , the kernel is the Bartlett, and  $\beta = 0$  (although later work by Kiefer and Vogelsang (2002, 2005) generalizes to  $b < 1$ ). Then  $Q(1) = 2 \int_0^1 \tilde{B}^2(r) dr$ , and the authors note that  $B(1)$  is independent of  $\tilde{B}(r)$ . As shown in McElroy and Politis (2011), this is true for other kernels as long as  $\beta = 0$ ; however, if  $\beta \neq 0$ , then  $B(1)$  and  $Q(b)$  are dependent. Fortunately, it is a simple matter to determine the limiting distribution numerically for any given value of  $\beta$ , and any choice of taper and bandwidth fraction  $b$ .

## 6 Applications and Numerical Studies

The preceding two sections give two different perspectives on the asymptotic behavior of taper-normalized sample means. If we normalize  $S_n - n\mu$  by  $\sqrt{nW_{\Lambda,M}}$  or by  $\sqrt{MW_{\Lambda,M}}$ , the studentized statistic does not converge – except in the SM and DM cases, respectively for the two normalizations

– when adopting the vanishing bandwidth-fraction perspective (see Remark 5). However,  $(S_n - n\mu)/\sqrt{nW_{\Lambda,M}}$  converges to a nondegenerate distribution by Theorem 4 when adopting the fixed bandwidth-fraction approach. Since bandwidth choice is up to the practitioner, it appears that the fixed bandwidth-fraction viewpoint might be preferable in our attempt towards a unified treatment of inference for the mean that is valid in all kinds of scenarios.

However, the limit distribution of the studentized sample mean will generally depend on  $\beta$ . Either one must estimate the nuisance parameters – including  $\beta$  – or a nonparametric technique such as the bootstrap or subsampling (Politis, Romano, and Wolf (1999)) must be utilized to get the limit quantiles. The parametric bootstrap is not feasible here (since no model is specified for the data in our context) and the block bootstrap tends to perform badly when autocorrelation dies gradually (Lahiri, 2003). However, given that the limit distribution has been tabulated for some values of  $\beta \in (-1, 1)$  and some popular tapers (cf. McElroy and Politis, 2009), one can utilize a plug-in estimator of  $\beta$  instead; this is similar in spirit to the approach advocated in Robinson (2005).

These two techniques are described in more detail below, along with statistical justification. A new estimator of  $\beta$ , based on the rate estimation ideas of McElroy and Politis (2007), is also discussed. Then in the following subsection, both methods are applied to the study of size and power for testing the null hypothesis that  $\mu = 0$ . We look at Gaussian processes exhibiting LM, SM, NM, or DM, at a variety of sample sizes and choices of taper.

## 6.1 Subsampling Methodology for Obtaining Critical Values

Firstly, we consider the subsampling method applied to the statistic  $S_n/\sqrt{nW_{\Lambda,M}}$ , or equivalently,  $\bar{Y}/\sqrt{W_{\Lambda,M}/n}$ . Letting  $M = bn$  as usual with  $b$  fixed and constant, under the assumptions of Theorem 4 we obtain the nondegenerate limit distribution  $B(1)/\sqrt{Q(b)}$  for the LM/SM/NM case, and  $[\tilde{C}(1) - \tilde{C}(0)]/\sqrt{P(b)}$  in the DM case. The subsampling distribution estimator (sde) will be consistent for this limit distribution under a mixing condition, such as weak dependence (see discussion in Section 4). In Jach, McElroy, and Politis (2011) the weak dependence condition is shown to be sufficient to establish consistency of subsampling distribution estimators for studentized statistics; Ango-Nze et al. (2003) was an earlier work on subsampling under weak dependence.

As for the sde itself, we first select a subsampling blocksize  $an$ , where  $a$  is the subsampling-fraction; as usual, this is assumed to be vanishing, i.e.,  $a = a(n) \rightarrow 0$ , though  $an \rightarrow \infty$ . Then  $n - an + 1 = (1 - a)n + 1$  contiguous overlapping blocks of the time series are constructed, and the statistics  $\bar{Y}$  and  $W_{\Lambda,bn}$  are evaluated on each block. This means a sample mean over  $an$  random variables, and the corresponding tapered variance estimate based on this subsample, so that the bandwidth is actually  $abn$  rather than  $bn$ . As a practical matter, unless  $b$  and  $a$  are taken fairly large, the bandwidth  $abn$  becomes unmanageably small. The subsampled statistics can be collected

into a set:

$$\left\{ \frac{\bar{Y}_{an,i} - \bar{Y}_n}{\sqrt{W_{\Lambda,abn,i}/(an)}} \text{ for } i = 1, \dots, n - an + 1. \right\}$$

Now taking the order statistics on this collection produces the quantiles of the sde. Further details of the construction can be found in Politis, Romano, and Wolf (1999), but we here sketch the remaining theoretical details. First we note a general result that follows from Theorem 2 and Remark 5: if  $M/n \rightarrow 0$  as  $n \rightarrow \infty$ , then

$$\frac{V_n/n^2}{\overline{W}_M/M} \rightarrow 0. \quad (17)$$

This is true for the LM/SM/NM case, since by Proposition 1 the limit is  $O([M/n]^{1-\beta})$ . The convergence in (17) is also true for the DM case if a non-flat-top taper (i.e.,  $c = 0$ ) is used, since the above limit will be asymptotic to  $-M/\dot{\Lambda}_+(0)n$ . (The result is not guaranteed for flat-top tapers; this will depend on the rate that  $M/n \rightarrow 0$ , versus the rate of decay of  $\{r_k\}$ .)

Therefore, the probability that the  $i$ th subsample statistic exceeds a given  $x$  is

$$\mathbb{P} \left[ \sqrt{an} \frac{\bar{Y}_{an,i} - \bar{Y}_n}{\sqrt{W_{\Lambda,abn,i}}} > x \right] = \mathbb{P} \left[ \frac{S_{an} - an\mu}{\sqrt{anW_{\Lambda,abn}}} - b^{-1/2} \frac{S_n - n\mu}{\sqrt{V_n}} \sqrt{\frac{V_n/n^2}{W_{\Lambda,abn}/(abn)}} > x \right].$$

Using Theorems 1, 4, and 5, as well as Proposition 3 and (17), the second term in the probability will tend to zero. Hence (in the LM/SM/NM case)

$$\mathbb{P} \left[ \frac{S_{an} - an\mu}{\sqrt{anW_{\Lambda,abn}}} > x \right] \rightarrow \mathbb{P} \left[ \frac{B(1)}{\sqrt{Q(b)}} > x \right]$$

as  $n \rightarrow \infty$ , noting that  $an \rightarrow \infty$  by assumption (and  $b$  is fixed). In the DM case the limit is  $[\tilde{C}(1) - \tilde{C}(0)]/\sqrt{P(b)}$ . Now, as long as the  $i$ th subsample statistic is approximately independent to the  $j$ th one when  $|i - j|$  is large – which is implied by the weak dependence condition – the sde is consistent for the target limit distribution, and subsampling is valid. Note that we assume the fixed bandwidth-fraction condition for this result, but end up utilizing some of the vanishing bandwidth-fraction results, since the actual bandwidth-fraction for the subsampled tapered variance estimate is the vanishing quantity  $ab$ .

## 6.2 Plug-in Methodology for Obtaining Critical Values

Alternatively, one can use a plug-in approach as in McElroy and Politis (2011). Adopting the fixed bandwidth-fraction asymptotics (so assume  $b$  is constant throughout), and assuming that  $\beta \in (-1, 1)$  (so that the DM case is explicitly excluded), we proceed to estimate the quantiles of the limit distribution  $B(1)/\sqrt{Q(b)}$  via first estimating  $\beta$  from the data, and then utilizing  $x_\alpha(\hat{\beta})$ , where  $x_\alpha(\beta)$  is the upper right  $\alpha$  quantile of  $B(1)/\sqrt{Q(b)}$ . That is,

$$\mathbb{P} \left[ \frac{B(1)}{\sqrt{Q(b)}} > x_\alpha(\hat{\beta}) \right] = \alpha.$$

These quantiles have been tabulated for  $\beta \in \mathbb{B} = \{-.8, -.6, -.4, -.2, 0, .2, .4, .6, .8\}$ , three values of  $\alpha$ , several commonly used tapers, and all values of  $b$  (via regression) – see the tables in McElroy and Politis (2009). Since the distribution is continuous in  $\beta$ , any consistent estimate  $\widehat{\beta}$  can be utilized. Then one finds the member of  $\mathbb{B}$  closest to the given  $\widehat{\beta}$  – call this  $\widetilde{\beta}$  – and utilizes  $x_\alpha(\widetilde{\beta})$ . This will be called the empirical plug-in method. Clearly, a finer mesh of simulation values for  $\mathbb{B}$  would improve the procedure, but we may yet expect to obtain results superior to just using  $\beta = 0$  in ignorance of the true memory. This latter approach, which essentially assumes that only short memory is present, will be referred to as the default plug-in method, and will be utilized as a benchmark for the empirical plug-in method.

Now many nonparametric estimators of  $\beta$  can be utilized. Here we consider the simple estimator proposed in McElroy and Politis (2011), namely:

$$\widehat{\beta} = \frac{\log W_{\Lambda, bn}}{\log n}. \quad (18)$$

This is consistent for  $\beta$  under the assumptions common to this paper.

**Proposition 5** *Assume that  $b \in (0, 1]$  is fixed, as well as the hypotheses of Theorem 4. Then  $\widehat{\beta}$  defined by (18) converges in probability to  $\beta$ .*

This estimator is similar in spirit to the tail index estimator of Meerschaert and Scheffler (1998), since it is based upon a convergence rate. The performance of  $\widehat{\beta}$  can be poor in finite sample when long memory is present, but it is quite versatile and simple to implement. Note that it is easy to show that  $\widehat{\beta} \xrightarrow{P} -1$  in the DM case, but since  $x_\alpha(-1)$  does *not* correspond to the quantile of  $[\widetilde{C}(1) - \widetilde{C}(0)]/\sqrt{P(b)}$ , we must exclude the DM case by assumption to avoid an inconsistent procedure.

### 6.3 Size and Power of Methods

We next evaluate the two methodologies – subsampling and empirical plug-in – through simulations. We adopt the perspective of testing a null hypothesis of  $\mu = \mu_0$ , since it is easier to evaluate the finite-sample properties through size and power, as opposed to the confidence interval perspective. In the simulations we take  $\mu_0 = 0$ . For each method, we consider values of  $\mu$  between 0 and 1, with  $\mu = 0$  corresponding to the null hypothesis. We compute the statistic  $S_n/\sqrt{n\overline{W_{\Lambda, bn}}}$  for a few different values of  $b$ , and five different tapers. The critical values of the limit distribution can be approximated using either of the two methods described above (though the plug-in method cannot be used with DM processes, and we cannot justify the use of flat-top tapers in the DM case). Then we record the proportion of times that the statistic exceeds these critical values, using a two-sided test. Note that when  $\mu > 0$ , this assessment is interpreted as empirical power, but when  $\mu = 0$  we obtain the empirical size.

The size of the plug-in approach has been partially addressed in McElroy and Politis (2009, 2011). However, the subsampling method’s performance has not been previously studied, so we provide some additional material regarding its size. Tables 1 through 10 display size results for various Type I error rates, for a two-sided test with  $\alpha = .10, .05, .01$ . The sampling fraction for the subsampling method was selected at values  $a = .2, .1, .04$ . Also we have the five tapers – Bartlett, Trapezoid with  $c = .25$ , Trapezoid with  $c = .5$ , Parzen, and Daniell – with three choices of bandwidth fraction  $b = .1, .5, 1$ . The sample size is varied from  $n = 250, 500, 1000$ . Results are for the empirical coverage, and so the target for the columns are the values  $.90, .95, .99$ .

For data generation processes we focus on the Gaussian distribution, and consider simple white noise for the SM case (since serially correlated processes have been considered in previous literature, it suffices to take the simple white noise case); we consider four NM processes with  $\beta = -.2, -.4, -.6, -.8$ , where the autocovariance function is determined from Example 2 of Section 2. Also there are four LM processes, with  $\beta = .2, .4, .6, .8$  and autocovariance function given in Example 1 of Section 2. Finally, our tenth process is DM, generated by the first difference of a white noise process. Informally, we will refer to this via  $\beta = -1$ , by an abuse of notation. We generated 1000 simulations of each specification.

The power surfaces are organized a bit differently. We focus on one sample size  $n = 250$  and one Type I error rate  $\alpha = .05$  (for a two-sided test for the mean, so we use the upper one-sided critical value at  $.975$  from the subsampling distributions and the tabulated values). Restricting to this  $\alpha$  value gives the general behavior, and a reasonable sense of the power can be gleaned from the  $n = 250$  case – higher sample sizes tend to shift the contours upwards (not dramatically for high  $\beta$ ), but the overall shape is the same. We consider the same tapers (Bartlett, Trapezoid (.25), Trapezoid (.5), Parzen, and Daniell), but with bandwidth fractions  $b = .2, .5, 1$ . The sampling fractions are  $a = .04, .12, .2$ . These choices are convenient, as it is always guaranteed that  $abn$  is an integer. The range of  $\mu$  was chosen so as to capture the main qualitative features of the power surface across all data processes:  $\mu \in \{j/20\}_{j=0}^{19}$  proved to yield power close to 50 percent for the long memory processes, while being small enough to allow visual discrimination of cases.

The six methods are placed in each figure as sub-panels. Moving from top left to bottom right, the first three methods correspond to subsampling with various sampling fractions. Then we have the empirical plug-in method, followed by the default plug-in (which uses  $\beta = 0$  critical values throughout) method. Note that these methods are greatly flawed when the true process is DM, but we present the results anyways for thoroughness. The final panel is an omniscient plug-in, based on knowing the true value of  $\beta$  (so it is not a practicable method, but is helpful for understanding power). The discrepancy in power between this and the empirical plug-in method (middle right panels) is mainly due to error in our estimator of  $\beta$ .

Now we discuss these numerical results. The size results for the SM case are fairly standard, being adequate at sample size  $n = 500$  and greater; the higher bandwidth fractions gave slightly

better results. At  $n = 1000$  all the tapers were roughly comparable in performance. For NM the coverage improved for higher values of  $\beta$ ; the flat-top tapers and other tapers perform similarly. The results for the LM case were much worse, with poor coverage at  $\beta = .2$ ; higher bandwidths seemed to improve performance slightly. At  $\beta = .4$  the results are best with the  $b = .5$  bandwidth, but for  $\beta = .6, .8$  the coverage deteriorates. Finally, for the DM case it seems that a smaller bandwidth is preferable. The Bartlett taper performs badly, but the Trapezoidal (.5) and Truncated tapers are adequate (except for  $b = .1$ ). In summary, if there is DM, NM, or SM, then one can use a flat-top (with higher value of  $c$ ) or other taper with a low or middle bandwidth fraction, such as  $b = .1$  or  $b = .5$ , and obtain adequate coverage for larger samples. If LM is present, a larger bandwidth is preferable.

If a statistic rejects too often under the null hypothesis – as is seen to happen in the tables for the subsampling methods – then it is liable to have higher power than otherwise. The fact that all methods tend to be mis-sized is evident in the surface plots in Figures 1 through 15 by examination of the  $\mu = 0$  cross-sectional curve towards the right side of the surface. But as  $\mu$  increases, the DM and NM processes generate high power relatively quickly, giving a mesa shape to the surfaces. For SM and weaker LM, the rise to full power is slower. An ironic feature is that when  $\mu$  is quite low, the power for strong LM is better than for weaker LM, essentially due to the methods being over-sized. This is seen in the “ruffle” feature of the curves along the  $\beta = .8$  cross-section.<sup>2</sup> Since power approaches 50 percent for all processes as  $\mu$  increases to unity, there is the question of how this is meaningful relative to the variation in the process. All were constructed with  $\gamma_0 = 1$  (an alternative way to normalize is to set the innovation variances equal, by in each case dividing through by the square root of the integral of the log spectrum), so the coefficient of variation is  $1/\mu$  for all processes. Primarily, we view these figures as a way to contrast the power of methods and tapers.

For a given taper, there is moreover the issue of selecting the best possible bandwidth fraction. If there is only a minor discrepancy between the true mean and the null mean, then higher power will be associated with smaller (upper one-sided) critical values. Leaving aside the issue of size, we can maximize power in the worst possible scenario (i.e., where the true mean  $\mu$  is approximately equal to the null mean) by finding  $b$  such that the quantile is minimized. Table 11 provides such bandwidth fractions for each given taper, as a function of the true  $\beta$ , for a variety of  $\alpha$  values (.10, .05, .025, and .01). For NM and SM a small bandwidth is best, but larger bandwidths – between .02 and .28 – provide better asymptotic power. This result also tends to parallel the size results, in that large bandwidths should be avoided in the NM and SM cases, while being more appropriate for the LM case.

---

<sup>2</sup>Some authors prefer to investigate what the power would be were these statistics to be adjusted to be correctly sized; however, in practice such a procedure is impossible to implement on real data. We have chosen to display the power that would occur were a practitioner to utilize any of the methods on real data.

## 7 Discussion

This paper sets out a thorough study of self-normalized mean estimation when long memory or negative memory is present. The main statistic of interest is the sample mean of a stationary time series (appropriately differenced beforehand), normalized by a tapered sum of sample autocovariances. The behavior of autocovariances changes greatly depending on whether a time series has long range dependence, anti-persistence, or short memory. This in turn has a large impact on the convergence rates of sample mean and tapered autocovariances. We provide a unified treatment of the various types of memory, including the important Super-Efficient (SE) case, wherein the partial sums are  $o_P(\sqrt{n})$ . The SE scenario is important, since it can easily arise from over-differencing of a time series suspected of having trend nonstationarity.

Several novel results on the memory of a time series are presented, which – together with examples – furnish some intuition for the qualitative behavior. The properties of Differential Memory (DM) processes are elucidated, and shown to be distinct from the behavior of Negative Memory (NM) processes – together, the DM and NM cases partition the important SE case. But our main interest is in the asymptotics of sample mean and tapered autocovariances, and we treat these topics through several theorems. For the asymptotic results we consider both the vanishing bandwidth-fraction case (a more classical approach, going back to Parzen (1957)) and the fixed bandwidth-fraction case (a more recent approach espoused in Kiefer, Vogelsang, and Bunzel (2000)). We both summarize known results, and prove new ones, examining three broad classes of data process that exhibit the various types of memory described herein.

In order to make use of the asymptotic results, it is still necessary to get the critical values of the limiting distributions, which in the fixed bandwidth-fraction case are functionals of the Fractional Brownian Bridge. We propose two methodologies: subsampling, which avoids explicit estimation of the memory parameter  $\beta$  but requires selection of a sampling fraction  $\alpha$ ; and the plug-in approach, which requires an estimate of  $\beta$  and a look-up table of critical values (computed ahead of time via simulation) for the limiting distributions. These methods are compared through extensive finite-sample size and power simulations, which are succinctly summarized here.<sup>3</sup> While power tends to deteriorate with greater memory, Type II error is comparatively quite small with anti-persistent processes. There are size problems with the plug-in method, whereas in contrast the subsampling method tends to have superior coverage (i.e., empirical size is closer to the nominal level).

One outstanding problem in the literature on this topic is the question of bandwidth selection. If one seeks an optimal bandwidth, it must be selected to minimize a pertinent criterion. Recent work by Sun, Phillips, and Jin (2008) provides an attractive approach based on examining both type I and type II errors. However, their paper is focused on the SM case; any attempt to mimic their procedure makes the bandwidth dependent on  $\beta$ , which then must be estimated. Also, higher-

---

<sup>3</sup>All code and results are available from the first author.



level assumptions are typically required for such an analysis, which is at war with the fairly generic assumptions of this paper.

What is really needed in practice is a prescription for applied statisticians. Choice of taper, bandwidth, and subsampling fraction are all to be determined by the user, and the optimal combination of such depends on the unknown data process. Moreover, any such optimality is typically derived using asymptotic criteria, whose impact on finite samples is less clear; also see Table 11. However, some of the lessons from our simulations can be repeated here: critical values vary more widely with respect to  $\beta$  when  $b$  is low; flat-top tapers and smooth tapers (e.g., the Parzen taper) perform adequately in terms of size and power, except in the LM case; size and power can depend substantially on the subsampling fraction. Note that the subsampling fraction might be selected using the technique of Bickel and Sakov (2008), as described in Jach, McElroy, and Politis (2011). Or one might utilize the plug-in approach described in Section 6, which requires simulation of quantiles ahead of time.

For any of the methods discussed, neither computer programming nor computation time is burdensome (unless quantiles are being simulated). Therefore, one can generate results for a multitude of tapers and bandwidths, and include all outcomes as alternative explanations of the data. If it is paramount to produce a single confidence interval or  $p$ -value, the practitioner might just retain the narrowest interval, as this corresponds to the lowest type II error (apart from approximation error). However, if for some reason the practitioner is unable to produce a spectrum of results, we recommend utilizing a small bandwidth (say  $b$  close to .02) when it is judged that only NM or SM is present, but a bandwidth fraction closer to .1 or .2 if LM is suspected. To obtain decent performance over a range of data processes, either the Parzen or a trapezoidal taper is recommended.

In summary, we provide a viable framework for conducting inference for the mean, supplying a unified asymptotic theory that covers all different types of memory under a single umbrella. This framework is robust against different memory specifications, obviating the need to do extensive modeling. Future work should examine the bandwidth selection problem, using these theoretical results as groundwork.

**Acknowledgements:** We thank the associate editor and two anonymous referees for helpful comments.

## Appendix

**Proof of Proposition 1.** First consider cases (i), (ii), and (iii) with  $\beta \in (-1, 1)$ . By Bingham, Goldie, and Teugels (1987), a slowly varying function satisfies, for any  $\delta > 0$ ,  $|L(k)/L(n)| \leq \tilde{C}(|k/n|^\delta + |n/k|^\delta)$  for  $1 \leq k \leq n$  and some constant  $\tilde{C} > 0$ . Then by the Dominated Convergence

Theorem

$$\frac{V_n}{nW_n} = n^{-1} \sum_{k=0}^{n-1} \frac{W_k}{W_n} = n^{-1} \sum_{k=0}^{n-1} \frac{L(k)}{L(n)} (k/n)^\beta \rightarrow \int_0^1 x^\beta dx = \frac{1}{\beta+1}.$$

In case (iii) when  $\beta = -1$ , we have  $V_n/(nW_n) \sim \sum_{h < n} h^{-1}L(h)/L(n) \rightarrow \infty$  by Theorem A3.6 of Embrechts, Klüppelberg, and Mikosch (1997). Case (iv) is immediate from its definition. For (7), in cases (i) and (iii) with  $\beta \in [-1, 1) \setminus \{0\}$  we have

$$\gamma_k = \frac{1}{2} (W_k - W_{k-1}) = \frac{1}{2} \left[ (L(k) - L(k-1))k^\beta \right] + \frac{1}{2} \left[ L(k-1)(k^\beta - (k-1)^\beta) \right].$$

Dividing through by  $L(k)k^{\beta-1}/2$  yields

$$2 \frac{\gamma_k}{L(k)k^{\beta-1}} = [(1 - L(k-1)/L(k))k] + \left[ \frac{kL(k-1)}{L(k)} (1 - (1 - 1/k)^\beta) \right].$$

The second expression on the right hand side is asymptotic to  $\beta$  via Taylor series. The first expression on the right hand side is asymptotic to  $-kg(k)$ , where  $g(x) = \int_{x-1}^x \eta(u)/u du$  by expanding the exponential term in the slowly-varying function  $L$ . But  $kg(k) = o(1)$  as  $k \rightarrow \infty$ . So when  $\beta \neq 0$ , the acf is asymptotic to  $\beta L(k)k^{\beta-1}/2$ . But when  $\beta = 0$  we have instead

$$\frac{\gamma_k}{L(k)} = \frac{1}{2} (1 - L(k-1)/L(k)) \sim -g_k/2$$

by the previous analysis, which is  $o(k^{-1})$ .  $\square$

**Proof of Proposition 2.** Without loss of generality, suppose that  $\mu = 0$ . By the Wold decomposition (see Brockwell and Davis, 1991) there exists an uncorrelated sequence  $\{\epsilon_t\}$  and square summable coefficients  $\{\psi_j\}$  such that  $Y_t = \sum_{j \geq 0} \psi_j \epsilon_{t-j}$ . Let  $\sigma^2 = \text{Var}(\epsilon_t)$  be equal to one for simplicity. We have  $0 = \lim_{k \rightarrow \infty} W_k = \sum_h \gamma_h = (\sum_j \psi_j)^2 \sigma^2$ , so  $\sum_j \psi_j = 0$ . It follows that  $1 - z$  should divide  $\Psi(z) = \sum_{j \geq 0} \psi_j z^j$ , though we must establish that  $\Theta(z) = \Psi(z)/(1 - z)$  converges. Extending  $\psi_j$  to be zero if  $j < 0$ , we define  $\theta_j = \sum_{k=-\infty}^j \psi_k$  for any integer  $j$ . Note that  $\theta_j \rightarrow 0$  as  $j \rightarrow \infty$  and is zero if  $j < 0$ . We will show that the  $\theta_j$  sequence is square summable, so that  $\sum_j \theta_j \epsilon_{t-j}$  is finite with probability one; then this will define  $Z_t$ , from which  $Y_t = Z_t - Z_{t-1}$  follows at once. Note that  $\theta_j = \sum_{k \geq 0} \psi_{j-k}$ ; let  $\theta_{j,m} = \sum_{k=0}^m \psi_{j-k}$ . Then  $\sum_j \theta_{j,m}^2 = \sum_{i,k=0}^m \gamma_{i-k} = \sum_{k=0}^m W_k$  for each  $m$ . Then by Fatou's Lemma

$$\sum_j \theta_j^2 \leq \liminf_{m \rightarrow \infty} \sum_j \theta_{j,m}^2 = \sum_{k \geq 0} W_k < \infty.$$

This establishes the existence of a series  $\{Z_t\}$  with the required properties. However, the series given by  $\sum_j \theta_j \epsilon_{t-j} + A$ , where  $A$  is uncorrelated with the series  $\{\epsilon_t\}$ , also has the requisite properties, as the temporal difference will be the same as that of  $Z_t - Z_{t-1}$ . Although an additional term of  $\text{Var}(A)$  is added on to the acf, this will cancel out in  $\gamma_k = 2r_k - r_{k+1} - r_{k-1}$ . So the series  $\{Z_t\}$  cannot be determined uniquely.

Next, we note that the formula for  $W_k$  follows from summing the  $\gamma_k$  via (3); the formula for  $r_0$  is obtained by summing the formula for  $W_k$ , and using the property of telescoping sums. The assertions of the converse are now immediate.  $\square$

**Proof of Theorem 1.** From Proposition 1 we can utilize (6). Also without loss of generality we can take  $\mu = 0$  in the proof. Let us first consider that the process satisfies P1, so that  $Y_t = \sum_j \psi_j \epsilon_{t-j}$  for an *iid* sequence  $\{\epsilon_t\}$ , and by assumption  $\gamma_0 \propto \sum_j \psi_j^2$  exists. So Theorem 5.2.3 of Taniguchi and Kakizawa (2000) (which is due to Ibragimov and Linnik (1971, p. 359)) gives the result. Note that by Hosoya (1996), we could relax the independence assumption on  $\{\epsilon_t\}$  to a type of mixing condition when  $\beta = 0$ .

Under P2 first consider the LM case. Since  $\beta > 0$ , we have  $r_k \sim \frac{\beta}{2} L(k) k^{\beta-1}$  by Proposition 1. Then we apply Theorem 5.2.2 of Taniguchi and Kakizawa (2000) (which is due to Taqqu, 1975) when  $(1 - \beta)\tau < 1$ , obtaining a non-central limit theorem if  $\tau > 1$ . In this case  $\{Y_t\}$  is  $LM(1 - (1 - \beta)\tau)$ . The distribution of  $B$  is described in the above references, and is non-Gaussian if  $\tau > 1$ . Note that the LM behavior of  $\{Y_t\}$  is inherited from that of  $\{X_t\}$ , as shown by Theorem 5.2.1 of Taniguchi and Kakizawa (2000). If the  $\{X_t\}$  process is SM, then a central limit theorem holds by applying Breuer and Major (1983).

Under P2 and NM, write  $g(x) = \sum_{q \geq 0} H_q(x) J_q / q!$ , with  $H_q$  the  $q$ th Hermite polynomial and  $J_q = \mathbb{E}[g(Z) H_q(Z)]$  where  $Z$  is standard normal. Then we have  $\gamma_k = \mathbb{E}[Y_t Y_{t+k}] = \sum_q r_k^q J_q^2 / q!$  (see 5.2.10 of Taniguchi and Kakizawa (2000)). Since  $\{Y_t\}$  is  $NM(\beta)$ , we have  $0 = \sum_k \gamma_k = \sum_q J_q^2 \sum_k r_k^q / q!$ . The sum of autocovariances  $W_n$  is related to the  $r_k$  sequence via

$$W_n = \sum_{q \geq 0} \frac{J_q^2}{q!} \sum_{|k| \leq n} r_k^q. \quad (\text{A.1})$$

Now  $W_n$  satisfies (3) by assumption, and moreover  $\sum_{|k| \leq n} k^{(\beta-1)q} \sim O(n^{(\beta-1)q+1})$ ; hence the limit of  $\sum_{|k| \leq n} r_k^q W_n^{-1}$  exists, and will be denoted by  $D_q$ . Also  $D_q = O(n^{(\beta-1)(q-1)})$ , which is zero when  $q > 1$ . Then dividing (A.1) by  $W_n$  and applying the Dominated Convergence Theorem yields  $1 = J_1^2 D_1$ . Thus

$$\sum_{t=1}^n Y_t = J_1 \sum_{t=1}^n X_t + \sum_{q > 1} \frac{J_q}{q!} \sum_{t=1}^n H_q(X_t)$$

and  $n^{-1} \text{Var}(\sum_{t=1}^n H_q(X_t)) = o(W_n)$  if  $q > 1$  by equation 5.2.9 of Taniguchi and Kakizawa (2000) and Theorem A3.6 of Embrechts et al. (1997). Hence

$$V_n^{-1/2} \sum_{t=1}^n Y_t = o_P(1) + J_1 \sum_{t=1}^n X_t / \sqrt{V_n}. \quad (\text{A.2})$$

This latter term is Gaussian with asymptotic variance  $J_1^2 D_1 = 1$ , i.e., the partial sum is asymptotic to the first (Gaussian) term of the Hermite expansion in the NM case, the higher order terms being negligible.

Next, consider P3. The  $k$ -fold cumulant of  $S_n$  is

$$\text{cum}(S_n, \dots, S_n) = \sum_{t_k=1}^n \sum_{t_1=1}^n \cdots \sum_{t_{k-1}=1}^n c_k(t_1 - t_k, t_2 - t_k, \dots, t_{k-1} - t_k),$$

which is  $O(n \cdot n^{\beta \wedge 0})$  when  $k > 2$ . Letting  $\kappa_{k,n}$  denote the  $k$ th order cumulant of  $S_n V_n^{-1/2}$ , we obtain  $\kappa_{k,n} = O(n^{(\beta+1) \wedge 1} \cdot V_n^{-k/2})$ , which tends to zero as  $n \rightarrow \infty$  if  $k > 2$ . Of course  $\kappa_{1,n} = 0$  and  $\kappa_{2,n} = 1$ , and the limit theorem is proved by expanding the characteristic function of  $S_n V_n^{-1/2}$  as in equation (2.19) of Hall (1992).

Finally, we turn to the DM case. Applying Proposition 2, we have  $S_n = Z_n - Z_0$ . If  $\{Z_t\}$  is strongly mixing, let  $\tilde{Z}_n$  be equal in distribution to  $Z_n$  for every  $n$ , but independent of  $Z_0$ . Then

$$\left| \mathbb{E} \exp\{i\nu(Z_n - Z_0)\} - \mathbb{E} \exp\{i\nu(\tilde{Z}_n - Z_0)\} \right| \leq 16 \alpha_n$$

by Lemma B.0.6 of Politis, Romano, and Wolf (1999), which is due to Ibragimov (1962). Now  $\phi_{\tilde{Z}_n - Z_0}(\nu) = \phi_Z(\nu) \cdot \phi_Z(-\nu)$  where  $\phi$  denotes the characteristic function, and  $Z$  has the common distribution of the  $\{Z_t\}$  process.

If  $\{Z_t\}$  is linear, then  $\{\epsilon_t\}$  are *iid* in the representation  $Z_t = \sum_{j \geq 0} \theta_j \epsilon_{t-j}$ , with the convention that  $\theta_j = 0$  if  $j < 0$ . Then

$$S_n = \sum_j (\theta_{n-j} - \theta_{-j}) \epsilon_j = \sum_{j=1}^n \theta_{n-j} \epsilon_j + \sum_{j \leq 0} (\theta_{n-j} - \theta_{-j}) \epsilon_j.$$

These two terms are independent, and the first is equal in distribution to  $\sum_{j=0}^{n-1} \theta_j \epsilon_j$ , which tends in probability (and thus weakly) to  $\sum_{j \geq 0} \theta_j \epsilon_j$  by Theorem 22.6 of Billingsley (1995). This limit is equal in distribution to  $Z$ , so  $\sum_{j=1}^n \theta_{n-j} \epsilon_j \xrightarrow{\mathcal{L}} Z$ . For the second term, we have  $\sum_{j \leq 0} \theta_{n-j} \epsilon_j \xrightarrow{P} 0$ , since its variance is equal to  $\sum_{j \geq n} \theta_j^2 \sigma^2$ . What's left is  $-\sum_{j \leq 0} \theta_{-j} \epsilon_j$ , which is equal to  $-Z$  in distribution. Since the two terms are independent, we have that  $S_n$  converges weakly to the difference of two independent random variables, each with the distribution of  $Z$ .  $\square$

**Proof of Proposition 3.** The expression  $\tilde{\gamma}_h - \gamma_h$  is unchanged if we replace  $Y_t$  by  $Y_t - \mu$ , so without loss of generality suppose that  $\mu = 0$ . We begin by decomposing  $W_{\Lambda, M}$ :

$$\begin{aligned} W_{\Lambda, M} &= \tilde{W}_M + E_M^{(1)} + E_M^{(2)} \\ E_M^{(1)} &= \sum_h \Lambda_M(h) (\tilde{\gamma}_h - \bar{\gamma}_h) \\ E_M^{(2)} &= \sum_h \Lambda_M(h) (\bar{\gamma}_h - \gamma_h), \end{aligned} \tag{A.3}$$

where  $\bar{\gamma}_k = \frac{1}{n} \sum_{t=1}^{n-|k|} (Y_t - \mu)(Y_{t+k} - \mu)$  for  $k \geq 0$ , and recall  $\tilde{\gamma}_k$  is given by (2). The terms  $E_M^{(1)}$  and  $E_M^{(2)}$  are stochastic errors, whereas  $\tilde{W}_M$  is deterministic and serves as an approximation to

$W_M$ . We will show that the error terms  $E_M^{(1)}$  and  $E_M^{(2)}$  of (A.3) satisfy  $E_M^{(1)} = O_P(U_M M V_n n^{-2})$  and  $E_M^{(2)} = O_P(U_M M n^{-\eta} K(n))$  as  $n \rightarrow \infty$ , where  $K$  is slowly-varying and  $\eta = 1/2$  if  $\beta < 1/2$  and  $\eta = 1 - \beta$  if  $\beta \geq 1/2$  (in the LM case); then the statement of the proposition follows. For each  $h \geq 0$

$$\begin{aligned}\tilde{\gamma}_h &= \gamma_h + (\bar{\gamma}_h - \gamma_h) + (\tilde{\gamma}_h - \bar{\gamma}_h) \\ &= \left(1 - \frac{h}{n}\right) \gamma_h + \frac{1}{n} \sum_{t=1}^{n-h} (Y_t Y_{t+h} - \gamma_h) + \bar{Y} (\bar{Y} - \bar{Y}_{1:n-h} - \bar{Y}_{h+1:n}),\end{aligned}$$

where  $\bar{Y}_{1:n-h} = \sum_{t=1}^{n-h} Y_t/n$  and  $\bar{Y}_{h+1:n} = \sum_{t=h+1}^n Y_t/n$ . Let  $c_n = V_n n^{-2}$ . Then  $\bar{Y} = O_P(n^{-1} V_n^{-1/2})$ , so the third term above is  $O_P(c_n)$  uniformly in  $h$  (this is true since  $|h| \leq M = o(n)$ ). Now  $c_n = n^{\beta-1} L(n)$  in the LM and NM cases, with  $\beta \in [-1, 1)$ , but  $c_n = n^{-1}$  for the SM case and  $c_n = n^{-2}$  for the DM case. Using the  $U_M$  bound on  $\Lambda_M(h)$ , we obtain  $E_M^{(1)} = O_P(U_M M c_n)$  as claimed. For the error term  $E_M^{(2)}$ , we compute

$$\begin{aligned}\text{Var}(E_M^{(2)}) &\leq n^{-2} \sum_{h,k} \Lambda_M(h) \Lambda_M(k) \sum_{t,s=1}^n \text{Cov}(Y_t Y_{t+h}, Y_s Y_{s+k}) \\ &\leq n^{-2} \sum_{h,k} \Lambda_M(h) \Lambda_M(k) \sum_{t,s=1}^n (\text{cum}(Y_t, Y_{t+h}, Y_s, Y_{s+k}) + \gamma_{t-s} \gamma_{t-s+h-k} + \gamma_{t-s-k} \gamma_{t+h-s}) \\ &\leq n^{-1} \sum_{h,k} \Lambda_M(h) \Lambda_M(k) \sum_{|l| \leq n} (1 - |l|/n) (\text{cum}(Y_0, Y_h, Y_l, Y_{l+k}) + \gamma_l \gamma_{l-h+k} + \gamma_{l+k} \gamma_{h-l}).\end{aligned}$$

The inequality used here only concerns  $|h|$  terms, which is negligible with respect to  $n$ . The sum of the cumulant function is bounded using Assumption B, resulting in an overall bounds of  $U_M^2/n$ . For the sum over  $\gamma_l \gamma_{l-h+k}$ , we can use the Cauchy-Schwarz inequality to obtain the bound  $O(M^2 U_M^2 n^{\xi-1} K(n))$ , where  $K$  is a slowly-varying function and  $n^\xi K(n)$  represents the order of  $\sum_{|l| \leq n} \gamma_l^2$ ; using Proposition 1 we have  $\xi = 0$  if  $\beta < 1/2$  (as this results in a square summable sequence) or  $\xi = 2\beta - 1$  if  $\beta > 1/2$ . When  $\beta = 1/2$  and  $L(n) \equiv 1$ , we take  $\xi = 0$  and  $K(n) = \log n$ . The analysis of the sum over  $\gamma_{l+k} \gamma_{h-l}$  yields a similar order. Taking square roots, we learn that  $E_M^{(2)} = O_P(M U_M n^{-\eta} K^{1/2}(n))$  as asserted.  $\square$

**Proof of Theorem 2.** First note that we can remove the term  $(1 - |h|/n)$  since  $M/n \rightarrow 0$ . The case of the truncation filter is trivial. Case (iv) is treated differently, so we first consider cases (i), (ii), and (iii). We proceed to break the sum over  $h$  up according to the intervals of smoothness for  $\Lambda$ . The first such interval is  $[-c, c]$ , which corresponds to the flat-top interval; if there is no flat-top region, then  $c = 0$ . In general, consider an interval  $(r, s]$  such that the restriction of  $\Lambda$  is continuously differentiable there. Then

$$\sum_{[rM] < |h| \leq [sM]} \Lambda\left(\frac{h}{M}\right) \gamma_h = \Lambda\left(\frac{[sM]}{M}\right) W_{[sM]} - \Lambda\left(\frac{[rM]}{M}\right) W_{[rM]} + \sum_{h=[rM]}^{[sM]-1} \left[ \Lambda\left(\frac{h}{M}\right) - \Lambda\left(\frac{h+1}{M}\right) \right] W_h \quad (\text{A.4})$$

via summation by parts. The first two terms on the right hand side will cancel with other like terms for the other intervals, leaving the last term on the right hand side. The terms in the square brackets consist of values of  $\Lambda$  restricted to  $(r, s]$ , excepting only the first term  $h = [rM]$ ; however, since  $\Lambda([rM]/M) - \Lambda(([rM] + 1)/M) \cong -M^{-1}\dot{\Lambda}_+(r)$  by continuity, the first term's analysis is the same as the others. In general,  $\Lambda(h/M) - \Lambda(h + 1/M) = -\dot{\Lambda}(h/M)M^{-1} + O(M^{-2})$ . The case of  $r > 0$  for an interval  $(r, s]$  has a different analysis from the  $r = 0$  case, which we consider later. So long as  $r > 0$ , we have  $h \rightarrow \infty$  as  $M \rightarrow \infty$  in the above summation. Thus in case (ii),  $W_h \rightarrow W_\infty$  and this convergence occurs uniformly in  $h \in (rM, sM)$  as  $M \rightarrow \infty$ . Using the boundedness of  $\dot{\Lambda}(x)$  and the limit of a Cesaro sum, we obtain a limit of  $-W_\infty \int_r^s \dot{\Lambda}(x) dx$ . The argument can be extended to cases (i) and (iii) as follows:

$$\frac{\sum_{h=[rM]}^{[sM]-1} [\Lambda(\frac{h}{M}) - \Lambda(\frac{h+1}{M})] W_h}{CM^\beta L(M)} = \sum_{h=[rM]}^{[sM]-1} \left\{ \Lambda\left(\frac{h}{M}\right) - \Lambda\left(\frac{h+1}{M}\right) \right\} (h/M)^\beta \left[ W_h h^{-\beta} / CL(M) \right],$$

and the expression in square brackets equals one plus error tending to zero as  $M \rightarrow \infty$ , uniformly in  $h \in ([rM], [sM])$ . This is because

$$\left| \frac{W_h}{Ch^\beta L(M)} - 1 \right| \leq \left| \frac{W_h}{Ch^\beta L(h)} - 1 \right| + \left| \frac{W_h}{Ch^\beta L(h)} \left( \frac{L(h)}{L(M)} - 1 \right) \right|.$$

The first term tends to zero uniformly in  $h$  as  $M \rightarrow \infty$ . For the second, we have  $L(h)/L(M) \rightarrow 1$  uniformly in  $h$  as  $M \rightarrow \infty$  as well, which is seen by using the representation (5). Note that these arguments hold for  $\beta = 0$ . Using the boundedness of  $x^\beta$  for  $x \in (r, s]$  and  $r > 0$ , we obtain a limit of  $-\int_r^s \dot{\Lambda}(x)x^\beta dx$  as  $M \rightarrow \infty$ . This argument works for any  $\beta \in (-1, 1)$ .

The first interval must be treated differently – unless it is flat-top, i.e.,  $c > 0$ , in which case it is trivially given by  $W_{[cM]}$ , which cancels with boundary term in the next intervals. More generally, the first interval has the form  $\sum_{0 \leq |h| \leq [sM]} \Lambda(h/M)\gamma_h$ , which tends to  $\Lambda(0)W_\infty$  in case (ii). Otherwise in cases (i) and (iii) we have

$$\sum_{|h|=0}^{[sM]} \Lambda\left(\frac{h}{M}\right)\gamma_h = \Lambda(0)W_{[sM]} + 2 \sum_{h=1}^{[sM]} \left[ \Lambda\left(\frac{h}{M}\right) - \Lambda(0) \right] \gamma_h.$$

Note that  $\beta > -1$  by assumption. The expression in brackets can be expanded in the Taylor series  $\sum_{j \geq 1} \Lambda^{(j)}(0)h^j M^{-j}/j!$ . Noting that  $\sum_{h=1}^M h^j \gamma_h$  is divergent and asymptotic to  $\beta M^j W_M / 2(\beta + j)$  (proved by L'Hopital's rule and (7)) for any  $j \geq 1$ , we can interchange summations to obtain

$$W_{[sM]} + \beta W_{[sM]} \sum_{j \geq 1} \Lambda^{(j)}(0) \frac{s^j}{\beta + j} \sim W_M \left( s^\beta + \beta \int_0^s (\Lambda(x) - 1)x^{\beta-1} dx \right)$$

after some algebra. Now piecing all intervals together, taking into account cancelations and integration by parts, we arrive at (10).

Now we turn to case (iv). Letting the partition  $0 = s_0 < s_1 < \dots < s_T < s_{T+1} = 1$  denote the points of non-differentiability in  $\Lambda$ , we can write

$$\sum_{|h|=0}^{[sM]} \Lambda\left(\frac{h}{M}\right) \gamma_h = \sum_{j=0}^T \sum_{h=[s_j M]}^{[s_{j+1} M]-1} \left[ \Lambda\left(\frac{h}{M}\right) - \Lambda\left(\frac{h+1}{M}\right) \right] W_h.$$

The first term is identically zero for a flat-top kernel; otherwise when  $c = 0$ , we can use a Taylor series expansion at  $h/M$  again. Since  $\{W_h\}$  is a summable sequence, we can apply the Dominated Convergence Theorem and obtain  $2r_0$  times the Taylor series at zero, as stated in the theorem. As for the other intervals, note that  $\sum_{h=[s_j M]}^{[s_{j+1} M]-1} W_h = 2(r_{[s_j M]} - r_{[s_{j+1} M]})$  is tending to zero as  $M \rightarrow \infty$ , since  $j \geq 1$ . Thus these other terms decay even faster than the first term. Hence for flat-top tapers, the rate of convergence is  $o(M^{-1})$ .  $\square$

**Proof of Theorem 3.** The proof proceeds by first showing that the finite-dimensional distributions of  $\xi_n$  converge to those of FBM. Then we show tightness. We will now show that  $S_n$  and  $\xi_n$  are asymptotically equivalent processes, i.e., any linear combination over any set of times of their difference tends to zero in probability. Because  $\xi_n(t) - S_n(t) = (nt - [nt])Y_{[nt]+1}$ , clearly for every  $\epsilon > 0$  and any collection of times  $t_1, \dots, t_k$  and constants  $\alpha_1, \dots, \alpha_k$  (for any  $k \geq 1$ ),

$$\mathbb{P} \left[ \left| \sum_{j=1}^k \alpha_j (\xi_n(t_j) - S_n(t_j)) V_n^{-1/2} \right| > \epsilon \right] \rightarrow 0$$

as  $n \rightarrow \infty$ . This follows from (6). Hence it suffices to show that the finite-dimensional distributions of  $S_n$  converge to those of FBM. We may as well assume  $\mu = 0$  henceforth. We proceed to show this for the three cases of P1, P2, and P3 in turn.

For the linear case P1, we note that historically Davydov (1970) and Gorodetskii (1977) provide a proof of the result requiring higher moments. Marinucci and Robinson (2000) relax the requirement to  $2 + \delta$  moments for some  $\delta > 0$ . We will adapt the argument used in Theorem 5.2.3 of Taniguchi and Kakizawa (2000). The linear representation allows us to write  $Y_t = \sum_j \epsilon_j \psi_{t-j}$  as in the proof of Theorem 1. We have  $\sum_{j=1}^m \alpha_j S_{[r_j n]} = \sum_j \epsilon_j b_{j,n}$ , with  $b_{j,n} = \sum_{i=1}^m \alpha_i \sum_{t=1}^{[r_i n]} \psi_{t-j}$ . Since  $m < \infty$ , the same types of bounds used in the proof of Theorem 5.2.3 of Taniguchi and Kakizawa (2000) still apply. Hence  $\sum_{j=1}^m \alpha_j S_{[r_j n]}$  is asymptotically standard normal when normalized by the square root of  $\sum_j b_{j,n}^2 \sigma^2$ , which by algebra equals the variance of  $\sum_{j=1}^m \alpha_j S_{[r_j n]}$ . Expanding this expression yields

$$\begin{aligned} & \sum_{i_1, i_2=1}^m \alpha_{i_1} \alpha_{i_2} \sum_{t_1=1}^{[r_{i_1} n]} \sum_{t_2=1}^{[r_{i_2} n]} \gamma_{t_1-t_2} = \sum_{i_1=1}^m \alpha_{i_1}^2 \sum_{t_1, t_2=1}^{[r_{i_1} n]} \gamma_{t_1-t_2} + 2 \sum_{i_1 < i_2} \alpha_{i_1} \alpha_{i_2} \sum_{t_1=1}^{[r_{i_1} n]} \sum_{t_2=1}^{[r_{i_2} n]} \gamma_{t_1-t_2} \\ & = \sum_{i_1=1}^m \alpha_{i_1}^2 V_{[r_{i_1} n]} + \sum_{i_1 < i_2} \alpha_{i_1} \alpha_{i_2} \left( V_{[r_{i_1} n]} + V_{[r_{i_2} n]} - V_{[(r_{i_2} - r_{i_1})n]} \right). \end{aligned} \tag{A.5}$$

This final breakdown of the variance is true for any variance of a sum, and is generic. Dividing (A.5) through by  $V_n$  gives the asymptotic expression

$$\sim \sum_{i_1=1}^m \alpha_{i_1}^2 r_{i_1}^{\beta+1} + \sum_{i_1 < i_2} \alpha_{i_1} \alpha_{i_2} \left( r_{i_1}^{\beta+1} + r_{i_2}^{\beta+1} - (r_{i_2} - r_{i_1})^{\beta+1} \right),$$

which is the same as the variance of  $\sum_{i=1}^m \alpha_i B(r_i)$ . In other words the limit of  $V_n^{-1/2} \sum_{j=1}^m \alpha_j S_{[r_j n]}$  has the same distribution as  $\sum_{i=1}^m \alpha_i B(r_i)$ .

The P2 case is handled as follows. If LM and  $\beta > 0$ , we apply Theorem 5.2.2 of Taniguchi and Kakizawa (2000); note that we only obtain a FBM limit if  $\tau = 1$ . In the SM case a FBM limit is also obtained by the same arguments (but things are easier since all powers of the acf are summable). For the NM case, as in the proof of Theorem 1 we have (A.2), and we can substitute  $[nr]$  for  $n$  in the summation. Clearly  $\sum_{t=1}^{[nr]} X_t / \sqrt{V_n}$  is Gaussian, and its asymptotic variance is  $D_1 r^{\beta+1}$ . It is easy to see that this has the finite-dimensional distribution of FBM using (A.5).

For the P3 case the  $k$ th order cumulant of  $\sum_{i=1}^m \alpha_i S_{[r_i n]}$  can be expanded into a  $k$ -fold sum

$$\sum_{i_1=1}^m \sum_{i_2=1}^m \cdots \sum_{i_k=1}^m \alpha_{i_1} \alpha_{i_2} \cdots \alpha_{i_k} \text{cum}(Z_{i_1}, Z_{i_2}, \dots, Z_{i_k}),$$

where  $Z_{i_j} = \sum_{t=[r_{i_j} n]+1}^{[r_{i_{j+1}} n]} Y_t$ . The same overall bounds can then be obtained as in the proof of Theorem 1, showing that all higher order cumulants with  $k > 2$  of the normalized sum will tend to zero. Again (A.5) shows that the variance converges to the variance of FBM, and so the result follows.

Finally, we establish tightness. The criterion we use is given by Problem 4.11 of Karatzas and Shreve (1991), which is appropriate for  $C[0, 1]$ . Letting  $\gamma = (\kappa + \delta)/2$ , and taking any times  $s < t$  and any  $n$ ,

$$\begin{aligned} & \mathbb{E} \left[ \left| (\xi_n(t) - nt\mu) - (\xi_n(s) - ns\mu) \right|^{2\gamma} V_n^{-\gamma} \right] \\ &= V_n^{-\gamma} \mathbb{E} \left[ \left| \sum_{j=[ns]+1}^{[nt]} Y_j + (nt - [nt])Y_{[nt]+1} - (ns - [ns])Y_{[ns]+1} - n\mu(t-s) \right|^{2\gamma} \right] \\ &\sim V_n^{-\gamma} \mathbb{E} \left[ \left| S_{[nt]-[ns]} - ([nt] - [ns])\mu \right|^{2\gamma} \right] \\ &= V_n^{-\gamma} O(V_{[n(t-s)]}^\gamma) = O((t-s)^{(\beta+1)\gamma}) \end{aligned}$$

by Proposition 1. Because  $\gamma > 1/(\beta + 1)$ , tightness is assured. This completes the proof.  $\square$

**Proof of Theorem 4.** We wish to apply Theorem 3, which is in terms of  $\xi_n(t)$ , to the result of Proposition 4, which is in terms of  $S_n(t)$ . We can do this because  $\widehat{\xi}_n(t) = \widehat{S}_n(t) + (nt - [nt])Y_{[nt]+1}$ , so that  $(\widehat{\xi}_n(t) - nt\mu)/\sqrt{V_n}$  is asymptotically equivalent to  $(\widehat{S}_n(t) - [nt]\mu)/\sqrt{V_n}$ . Then all linear or quadratic integral expressions of such processes will also be asymptotically the same. So note



that  $\widehat{S}_i = S_{[rn]} - rS_n$  with  $r = i/n$ , and recognize the summations in the expression for  $W_{\Lambda, M}$  in Proposition 4 as Riemann sums; the result now follows at from Theorem 3.  $\square$

**Proof of Theorem 5.** As in the proof of Theorem 4, it suffices to work with  $S_n(t)$  rather than  $\xi_n(t)$ . We have  $\widehat{S}_i = Z_i - \frac{i}{n}Z_n - (1 - \frac{i}{n})Z_0$ , and hence without loss of generality suppose that  $Z_i$  is mean zero. Using the abbreviation  $\Delta_n(i-j) = 2\Lambda_n(i-j) - \Lambda_n(i-j+1) - \Lambda_n(i-j-1)$ ,

$$\begin{aligned} nW_{\Lambda, M} &= \sum_{i,j=1}^n \widehat{S}_i \widehat{S}_j \Delta_n(i-j) \\ &= \sum_{i,j=1}^n Z_i Z_j \Delta_n(i-j) - 2Z_n \sum_{i,j=1}^n \frac{i}{n} Z_j \Delta_n(i-j) - 2Z_0 \sum_{i,j=1}^n \left(1 - \frac{i}{n}\right) Z_j \Delta_n(i-j) \\ &\quad + 2Z_0 Z_n \sum_{i,j=1}^n \left(1 - \frac{i}{n}\right) \frac{j}{n} \Delta_n(i-j) + Z_n^2 \sum_{i,j=1}^n \frac{ij}{n^2} \Delta_n(i-j) + Z_0^2 \sum_{i,j=1}^n \left(1 - \frac{i}{n}\right) \left(1 - \frac{j}{n}\right) \Delta_n(i-j). \end{aligned} \tag{A.6}$$

Using the Taylor series approximations for  $\Delta_n(h)$  derived in the proof of Proposition 4, we can compute each of the six terms above. Each of these is of the form (up to multiplication by other random variables)  $\sum_{i,j=1}^n X_i X_j \Delta_n(i-j)$  for some random variables  $X_i$ , which is approximately

$$\begin{aligned} &- 2b^{-2} \frac{1}{n} \sum_{h=[cbn]+1}^{[bn]-1} \ddot{\Lambda} \left( \frac{h}{bn} \right) \frac{1}{n} \sum_{i=1}^{n-h} X_i X_{i+h} \\ &- 2b^{-1} \dot{\Lambda}_+(c) \frac{1}{n} \sum_{i=1}^{n-[cbn]} X_i X_{i+[cbn]} \\ &+ 2b^{-1} \dot{\Lambda}_-(1) \frac{1}{n} \sum_{i=1}^{n-[bn]} X_i X_{i+[bn]}, \end{aligned}$$

up to  $o_P(1)$  terms. Now in the first term in (A.6), we have  $X_i = Z_i$  and  $\frac{1}{n} \sum_{i=1}^{n-h} Z_i Z_{i+h} \xrightarrow{P} (1-h/n)r_h$ . Now if  $c = 0$  we obtain  $\frac{1}{n} \sum_{i=1}^{n-[cbn]} W_i W_{i+[cbn]} \xrightarrow{P} r_0$ ; all other portions of the first term tend to zero in probability, whether or not  $c = 0$ . For the second and third terms of (A.6), we have  $X_j = Z_j$  and  $X_i$  deterministic; then these terms tends to zero since  $Z_j$  is mean zero. For the fourth term of (A.6), the limit is  $2Z_0 Z_*$  times

$$-b^{-2} \int_{cb < |r-s| < b} (1-r)s \ddot{\Lambda} \left( \frac{|r-s|}{b} \right) dr ds - 2b^{-1} \dot{\Lambda}_+(c) \int_0^{1-cb} (1-r)(r+cb) dr + 2b^{-1} \dot{\Lambda}_-(1) \int_0^{1-b} (1-r)(r+b) dr.$$

The fifth term of (A.6) is asymptotic to  $Z_*^2$  times

$$-b^{-2} \int_{cb < |r-s| < b} rs \ddot{\Lambda} \left( \frac{|r-s|}{b} \right) dr ds - 2b^{-1} \dot{\Lambda}_+(c) \int_0^{1-cb} r(r+cb) dr + 2b^{-1} \dot{\Lambda}_-(1) \int_0^{1-b} r(r+b) dr,$$

while the sixth term of (A.6) tends to  $Z_0^2$  times

$$\begin{aligned} & -b^{-2} \int_{cb < |r-s| < b} (1-r)(1-s) \ddot{\Lambda}\left(\frac{|r-s|}{b}\right) dr ds - 2b^{-1} \dot{\Lambda}_+(c) \int_0^{1-cb} (1-r)(1-r-cb) dr \\ & + 2b^{-1} \dot{\Lambda}_-(1) \int_0^{1-b} (1-r)(1-r-b) dr. \end{aligned}$$

Combining the fourth, fifth, and sixth terms together yields  $Q(b)$  with  $\tilde{C}(r) = Z_0 + r(Z_* - Z_0)$ , which interpolates between  $Z_0$  and  $Z_*$ . This establishes the convergence of  $nW_{\Lambda, M}$ ; for the joint convergence, note that we can rework the same proof for  $\alpha(S_n - n\mu) + \gamma nW_{\Lambda, M}$  for any  $\alpha$  and  $\gamma$ , since  $S_n - n\mu = Z_n - Z_0$ . Hence up to terms  $o_P(1)$ ,  $\alpha(S_n - n\mu) + \gamma nW_{\Lambda, M}$  can be expressed as a bilinear combination of  $Z_0$  and  $Z_n$ , which converges to  $\alpha(Z_* - Z_0) + \gamma P(b)$  after gathering like terms.  $\square$

**Proof of Proposition 5.** By Theorem 4 we know that  $Q_n := nW_{\Lambda, bn}/V_n \xrightarrow{\mathcal{L}} Q(b)$ , and by Proposition 1 we have  $V_n/n \sim n^\beta CL(n)/(\beta+1)$  in the LM/SM/NM case. Thus it follows that (18) satisfies

$$\hat{\beta} \sim \beta + \frac{\log Q_n}{\log n} + \frac{\log(CL(n)/(\beta+1))}{\log n}.$$

Since  $Q_n = O_P(1)$  and  $\log L(n)/\log n \rightarrow 0$  for slowly-varying functions  $L$ , the estimator will be consistent. Note that if the process is DM, we obtain

$$\hat{\beta} = -1 + \frac{\log Q_n}{\log n} + \frac{\log V_n}{\log n}$$

since  $Q_n = nW_{\Lambda, bn}/V_n$ . Now  $V_n \rightarrow 2r_0$  in the DM case, so  $\hat{\beta} \xrightarrow{P} -1$ .  $\square$

## References

- [1] Andrews, D. (1991) Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica* **59**, 817–858.
- [2] Ango-Nze, P., Dupoirion, S., and Rios, R. (2003) Subsampling under weak dependence conditions. Technical report from CREST, *DT2003 – 42*.
- [3] Bardet, J., Doukhan, P., Lang, G., and Ragache, N. (2008) Dependent Lindeberg Central Limit Theorem and some applications. *Probability and Statistics* **12**, 154–172.
- [4] Beran, J. (1993) Fitting long-memory models by generalized linear regression. *Biometrika*, **80**, 817–822.
- [5] Beran, J. (1994) *Statistics for Long Memory Processes*. New York: Chapman and Hall.
- [6] Billingsley, P. (1995) *Probability and Measure*. New York: John Wiley and Sons.

- [7] Bingham, N., Goldie, C., and Teugels, J. (1987) *Regular Variation*. Cambridge: Cambridge University Press.
- [8] Blackman, R.B. and Tukey, J.W. (1959). *The Measurement of Power Spectra from the Point of View of Communications Engineering*. New York: Dover.
- [9] Box, G. and Jenkins, G. (1976) *Time Series Analysis, Forecasting and Control*. San Francisco: Holden-Day.
- [10] Bradley, R. (2007) *Introduction to Strong Mixing Conditions*, Volume I. Heber City, Utah: Kendrick Press.
- [11] Breuer, P. and Major, P. (1983) Central limit theorems for non-linear functionals of Gaussian fields. *Journal of Multivariate Analysis* **13**, 425–441.
- [12] Brillinger, D.R. (1981) *Time Series: Data Analysis and Theory*. New York: Holden-Day.
- [13] Brockwell, P. and Davis, R. (1991) *Time Series: Theory and Methods, 2nd Ed.* New York: Springer.
- [14] Davydov, Y. (1970) The invariance principle for stationary processes. *Theory of Probability and its Applications* **14**, 487–498.
- [15] Dickey, D. and Fuller, W. (1979) Distribution of the estimators for autoregressive time series with a unit root. *Journal of the American Statistical Association* **74**, 427–431.
- [16] Dickey, D. and Fuller, W. (1981) Likelihood ratio statistics for autoregressive time series with a unit root. *Econometrica* **49**, 1057–1072.
- [17] Diebold, F. and Rudebusch, G. (1991) On the power of the Dickey-Fuller tests against fractional alternatives. *Economics Letters* **35**, 155–160.
- [18] Doukhan, P. and Louhichi, S. (1999) A new weak dependence and applications to moment inequalities. *Stochastic Processes and Their Applications* **84**, 313–343.
- [19] Embrechts, P., Klüppelberg, C., and Mikosch, T. (1997). *Modeling Extremal Events for Insurance and Finance*. Berlin: Springer-Verlag.
- [20] Findley, D. F., Monsell, B. C., Bell, W. R., Otto, M. C. and Chen, B. C. (1998) New capabilities and methods of the X-12-ARIMA seasonal adjustment program. *Journal of Business and Economic Statistics* **16**, 127–177 (with discussion).
- [21] Gorodetskii, V. (1977) On convergence to semi-stable Gaussian processes. *Theory Prob. Appl.* **22**, 498–508.

- [22] Grenander, U. and Rosenblatt, M. (1957) *Statistical Analysis of Stationary Time Series*. New York: Wiley.
- [23] Hall, P. (1992) *The Bootstrap and Edgeworth Expansion*. Springer-Verlag: New York.
- [24] Hamilton, J.D. (1994) *Time Series Analysis*. Princeton: Princeton University Press.
- [25] Hannan, E.J. (1970) *Multiple Time Series*. New York: John Wiley.
- [26] Hosking, J.R.M. (1981) Fractional differencing. *Biometrika*, **68**, 165-176.
- [27] Hosoya, Y. (1996) Some limit theorems on stationary processes with long-range dependence. *Athens conference on applied probability and time series* **2**, 233–245. eds. P.M. Robinson and R. Rosenblatt, Springer.
- [28] Ibragimov, I. (1962) Some limit theorems for stationary processes. *Theory of Probability and its Applications* **7**, 349–382.
- [29] Ibragimov, I. and Linnik, Y. (1971) *Independent and Stationary Sequences of Random Variables*. Wolters-Noordhof Publishing Groningen.
- [30] Jach, A., McElroy, T., and Politis, D. (2011) Subsampling Inference for the Mean of Heavy-tailed Long Memory Time Series. *Journal of Time Series Analysis* **33**, 96–111.
- [31] Kiefer, N., Vogelsang, T. and Bunzel, H. (2000) Simple robust testing of regression hypotheses. *Econometrica* **68**, 695–714.
- [32] Kiefer, N. and Vogelsang, T. (2002) Heteroscedastic-autocorrelation robust standard errors using the Bartlett kernel without truncation. *Econometrica* **70**, 2093–2095.
- [33] Kiefer, N. and Vogelsang, T. (2005) A new asymptotic theory for heteroskedasticity-autocorrelation robust tests. *Econometric Theory* **21**, 1130–1164.
- [34] Kolmogorov, A. and Rozanov, J. (1960) On a strong mixing condition for a stationary Gaussian process. *Theory of Probability and its Applications* **5**, 204–208.
- [35] Lahiri, S. (2003) *Resampling Methods for Dependent Data*. New York: Springer-Verlag.
- [36] Lopez, J. (1997) The power of the ADF test. *Economics Letters* **57**, 5–10.
- [37] Maravall, A. and Caporello, G. (2004) Program TSW: Revised Reference Manual. *Working Paper 2004, Research Department, Bank of Spain*. <http://www.bde.es>
- [38] Marinucci, D. and Robinson, P. (2000) Weak convergence of multivariate fractional processes. *Stochastic Processes and their Applications* **86**, 103–120.

- [39] McElroy, T. and Politis, D. (2007) Computer-intensive rate estimation, diverging statistics, and scanning. *Annals of Statistics* **35**, 1827–1848.
- [40] McElroy, T. and Politis, D. (2009) Fixed-b asymptotics for the studentized mean from time series with short, long or negative memory. UC San Diego: Department of Economics, UCSD.
- [41] McElroy, T. and Politis, D. (2011) Fixed-b asymptotics for the studentized mean for long and negative memory time series. *Econometric Theory* **28**, 1–11.
- [42] Palma, W. (2007) *Long-Memory Time Series*. New York: John Wiley and Sons.
- [43] Parker, C., Paparoditis, E., and Politis, D. (2006) Unit root testing via the stationary bootstrap. *Journal of Econometrics* **133**, 601–638.
- [44] Parzen, E. (1957) On consistent estimates of the spectrum of a stationary time series. *Annals of Mathematical Statistics* **28**, 329–348.
- [45] Percival, D.B. and Walden, A.T.(1993). *Spectral analysis for physical applications. Multitaper and conventional univariate techniques*. Cambridge: Cambridge University Press.
- [46] Phillips, P. and Perron, P. (1998) Testing for a unit root in time series regression. *Biometrika* **75**, 335–346.
- [47] Politis, D. (2001) On nonparametric function estimation with infinite-order flat-top kernels, in *Probability and Statistical Models with applications*, Ch. Charalambides et al. (Eds.), Chapman and Hall/CRC: Boca Raton, pp/ 469–483.
- [48] Politis, D. (2005) Higher-order accurate, positive semi-definite estimation of large-sample covariance and spectral density matrices, UCSD Dept. of Economics Discussion Paper 2005-03.
- [49] Politis, D. and Romano, J. (1995) Bias-corrected nonparametric spectral estimation. *Journal of Time Series Analysis* **16**, 67–103.
- [50] Politis, D., Romano, J., and Wolf, M. (1999) *Subsampling*. New York: Springer-Verlag.
- [51] Priestley, M.B. (1981) *Spectral Analysis and Time Series*. New York: Academic Press.
- [52] Robinson, P. (1994) Semiparametric analysis of long memory time series. *Annals of Statistics* **22**, 515–539.
- [53] Robinson, P. (2005) Robust covariance matrix estimation: HAC estimates with long memory/antipersistence correction. *Econometric Theory* **21**, 171–180.
- [54] Rosenblatt, M. (1956) A central limit theorem and a strong mixing condition. *Proceedings of the National Academy of Sciences* **42**, 43–47.

- [55] Rosenblatt, M. (1961) Independence and dependence. *Proc. Fourth Berkeley Symp. on Math. Statist. and Prob., Vol.2*, University of California Press, 431–443.
- [56] Rosenblatt, M. (1984) Stochastic processes with short-range and long-range dependence. In *Statistics: An Appraisal, Proceedings of the 50th Anniversary Conference* (David, H.A. and David, H.T., Eds.), The Iowa State University Press, Ames, IA, pp. 509–520.
- [57] Rosenblatt, M. (1985) *Stationary Sequences and Random Fields*. Boston: Birkhäuser.
- [58] Samorodnitsky, G. and Taqqu, M. (1994) *Stable Non-Gaussian Random Processes*. New York City, New York: Chapman & Hall.
- [59] Sun, Y. (2004) A convergent t-statistic in spurious regressions. *Econometric Theory* **20**, 943–962.
- [60] Tanaka, K. (1990) Testing for a moving average unit root. *Econometric Theory* **6**, 433–444.
- [61] Tanaka, K. (1996) *Time Series Analysis: Nonstationary and Noninvertible Distribution Theory*. New York City, New York: John Wiley and Sons.
- [62] Taniguchi, M. and Kakizawa, Y. (2000) *Asymptotic Theory of Statistical Inference for Time Series*. New York City, New York: Springer-Verlag.
- [63] Taqqu, M. (1975) Weak convergence to fractional Brownian motion and to the Rosenblatt process. *Z. Wahr. Verw. Geb.* **31**, 287–302.

Differential Memory					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	82.3 88.9 95.5	75.9 85.5 94.5	75.3 85.3 94.7	58.5 65.3 75.4	58.9 67.6 81.7
	86.2 91.7 97.6	40.1 48.7 61.9	69.5 82.7 93.8	45.0 54.6 67.3	41.0 50.3 63.9
	100 100 100	100 100 100	89.4 92.9 95.3	100 100 100	100 100 100
$b = .5$	81.8 88.4 95.0	78.8 89.3 98.1	78.5 90.0 98.1	66.2 74.4 86.7	74.1 85.2 95.6
	88.1 93.4 97.9	87.9 95.5 99.0	81.0 92.8 97.7	62.3 74.7 90.8	84.8 97.0 99.3
	99.6 100 100	55.4 63.9 74.3	60.1 68.9 80.8	92.2 97.2 99.7	67.4 77.4 89.4
$b = 1$	82.8 88.9 94.3	83.8 91.4 97.8	75.3 85.9 94.8	72.0 82.3 92.4	77.9 86.4 95.6
	88.2 94.6 98.2	87.6 96.0 99.2	81.5 91.1 97.3	81.3 92.6 98.8	86.9 96.9 99.4
	93.9 98.0 99.8	91.7 97.3 99.4	79.2 89.2 96.6	86.9 97.1 99.6	93.5 98.9 100
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	82.1 88.8 95.6	78.0 88.9 97.5	73.7 85.0 95.4	54.4 64.0 76.3	64.6 73.7 86.1
	86.7 92.7 98.0	79.4 90.7 98.5	79.8 89.6 97.4	50.6 58.5 71.6	56.0 68.2 86.8
	91.4 96.7 99.7	43.0 52.6 77.4	69.5 85.2 97.2	40.3 49.3 63.5	44.8 52.7 73.8
$b = .5$	80.2 87.4 94.3	78.5 88.6 98.1	78.8 89.4 98.0	67.3 75.7 87.6	74.2 84.4 95.3
	86.1 91.6 97.6	89.4 97.3 99.0	88.9 95.8 99.0	67.0 79.8 94.9	86.1 95.4 99.2
	91.1 96.9 100	96.6 98.4 99.7	91.9 96.9 99.2	66.8 79.1 97.9	91.2 97.0 99.4
$b = 1$	81.4 88.2 95.4	80.3 89.2 97.6	79.5 88.6 96.9	71.2 80.0 90.2	78.1 84.1 93.8
	87.2 93.1 98.7	88.5 95.4 98.8	81.5 89.9 97.5	84.1 93.7 99.2	82.8 93.2 99.6
	92.0 97.8 99.7	88.7 95.8 98.9	80.5 89.4 98.0	88.6 98.2 99.9	90.9 99.1 100
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	80.9 88.0 96.5	76.9 88.0 97.4	75.4 86.1 96.8	56.8 63.8 76.9	65.4 76.2 88.8
	86.1 92.1 97.8	82.7 92.9 98.7	76.8 90.2 97.8	51.2 58.6 71.7	62.8 74.4 92.8
	89.9 95.2 99.1	82.2 93.0 98.3	76.3 89.4 97.7	35.5 42.8 58.4	48.1 59.7 85.8
$b = .5$	81.9 87.6 95.0	75.3 86.1 97.2	77.5 87.8 98.1	67.0 73.8 83.6	70.3 79.3 93.3
	86.4 91.5 97.8	86.6 95.7 99.2	89.1 97.0 99.5	71.7 81.6 97.1	81.7 94.2 99.4
	90.0 95.5 99.8	96.3 98.2 99.6	93.9 98.1 99.7	69.3 84.4 99.3	94.0 98.5 99.5
$b = 1$	82.2 88.1 94.6	79.9 88.4 97.4	78.4 89.0 96.9	68.9 76.1 88.0	78.4 84.1 91.4
	87.1 93.1 97.9	85.8 94.4 99.6	81.2 90.9 98.4	80.1 91.8 99.4	83.2 92.3 99.3
	89.5 95.2 99.1	88.7 96.9 99.0	79.3 90.8 98.3	88.7 98.8 100	91.5 99.3 100

Table 1: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Differential Memory.

Negative Memory: $\beta = -.8$					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	86.6 90.7 95.4	81.3 88.5 93.9	81.5 87.8 93.3	84.4 89.7 94.3	83.6 89.4 94.4
	88.9 93.7 97.6	76.6 83.9 90.4	81.7 89.0 95.5	83.3 89.9 95.7	82.0 87.8 93.1
	100 100 100	99.9 100 100	99.8 100 100	100 100 100	100 100 100
$b = .5$	84.5 90.6 95.0	85.0 90.2 95.8	84.5 91.1 96.5	81.7 87.6 92.2	83.4 89.0 94.5
	90.1 95.2 97.6	87.8 93.7 98.0	88.0 94.1 97.4	85.4 91.3 95.9	86.6 93.5 96.7
	99.4 100 100	93.8 96.9 99.4	89.6 94.1 97.3	99.2 99.8 100	95.8 98.8 100
$b = 1$	84.1 88.6 94.0	84.0 90.4 96.7	83.0 89.5 96.0	81.7 88.2 95.4	85.9 92.5 97.3
	90.1 94.1 97.1	88.2 94.7 97.8	88.9 94.8 98.1	89.4 94.5 98.2	88.0 94.5 98.2
	94.0 97.4 99.6	89.3 93.7 98.7	91.1 94.9 98.7	90.2 95.1 98.3	90.5 95.2 98.9
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	84.7 90.9 96.6	82.0 87.4 92.0	80.0 85.8 91.7	80.9 86.9 93.8	78.5 84.2 91.2
	90.7 95.0 98.1	83.2 89.1 94.8	85.2 89.8 95.6	86.7 91.1 96.8	85.0 90.9 95.6
	95.7 98.4 99.6	85.2 90.6 96.5	88.4 92.2 97.7	89.7 93.5 98.2	86.2 92.3 96.8
$b = .5$	85.4 90.7 95.5	83.3 88.6 96.1	82.7 89.5 96.5	81.6 86.4 91.9	82.4 89.4 95.9
	91.4 94.8 98.0	87.4 93.3 98.5	86.9 92.9 98.0	85.5 91.2 96.9	88.2 93.0 97.6
	93.5 97.1 99.6	89.0 94.2 98.9	87.9 93.0 98.3	88.8 93.9 97.8	89.5 93.9 98.1
$b = 1$	86.8 92.4 96.6	86.8 93.0 97.4	83.7 91.1 97.4	82.0 88.5 94.4	84.1 90.1 97.0
	89.6 94.5 98.0	89.3 93.8 98.4	89.0 94.0 98.8	88.3 93.2 97.1	88.0 94.3 98.4
	90.9 96.3 99.5	89.9 93.4 98.6	89.4 95.0 98.8	90.2 95.0 99.5	89.7 94.0 98.4
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	86.1 90.7 96.1	84.0 89.3 95.0	80.5 87.2 93.4	82.3 89.3 94.0	81.4 86.4 92.3
	89.1 94.4 98.4	85.7 91.0 95.8	86.9 92.7 98.0	87.1 91.9 97.8	86.7 92.5 96.4
	92.4 97.4 99.2	86.9 92.8 97.8	87.9 93.3 98.5	86.3 91.2 97.6	87.6 93.2 98.2
$b = .5$	84.4 89.5 95.6	82.1 89.9 96.9	83.2 88.9 97.5	83.6 89.0 94.0	83.5 89.7 95.6
	87.9 93.6 97.9	88.7 94.0 98.4	89.2 94.2 98.2	86.7 91.5 97.4	87.6 93.1 98.5
	92.0 96.1 99.6	88.5 92.7 98.5	90.6 95.6 98.5	90.5 94.0 98.6	89.4 94.1 98.8
$b = 1$	85.3 90.5 95.1	85.2 93.0 98.5	84.6 90.7 97.9	85.0 92.0 96.9	84.9 91.6 97.3
	91.4 95.5 98.5	90.2 94.9 99.3	88.2 93.9 98.9	86.9 92.6 97.3	91.1 95.7 98.9
	92.5 97.3 99.7	88.8 94.3 98.8	91.1 95.4 99.3	88.0 94.8 99.1	90.8 95.6 99.4

Table 2: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Negative Memory with  $\beta = -.8$ .



Negative Memory: $\beta = -.6$					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	84.0 88.2 94.1	81.9 86.3 91.7	78.1 84.7 90.5	82.0 87.6 92.4	82.8 88.0 92.6
	89.9 94.8 97.6	81.2 87.0 92.9	84.2 90.3 95.4	85.3 92.3 96.5	84.5 90.5 94.5
	99.8 100 100	99.9 100 100	99.6 100 100	100 100 100	99.9 99.9 100
$b = .5$	85.3 89.8 94.0	84.8 89.0 94.9	81.8 89.6 96.5	82.4 87.6 91.8	82.2 88.0 94.3
	88.3 92.9 96.6	87.6 92.4 96.7	87.4 92.3 96.4	85.4 91.2 95.6	85.8 91.6 96.5
	98.4 99.1 99.5	94.0 97.0 98.8	90.9 93.4 97.4	97.4 99.2 100	95.7 98.2 99.5
$b = 1$	85.9 90.8 93.2	84.5 90.0 96.0	83.7 90.7 95.8	84.1 88.4 94.4	85.7 90.4 96.6
	89.0 93.3 96.6	87.5 93.3 97.3	89.1 93.5 97.6	86.3 92.0 96.1	88.7 94.4 99.0
	92.0 96.5 99.2	91.6 95.7 98.4	89.7 94.6 98.8	89.6 94.3 97.8	90.5 94.6 98.7
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	84.7 88.8 94.4	80.3 86.7 92.8	82.8 88.2 92.6	84.7 89.8 95.3	84.2 88.8 95.3
	89.5 94.7 98.5	85.7 92.2 95.9	85.2 90.4 95.4	87.1 92.4 96.9	86.7 92.2 96.5
	92.9 96.0 98.8	86.2 91.8 97.3	88.9 94.0 98.6	89.9 95.6 98.6	88.1 92.5 97.4
$b = .5$	82.8 87.7 92.5	81.9 89.5 96.0	82.3 88.8 96.2	79.7 84.8 91.1	81.0 88.0 94.4
	87.6 91.9 96.7	86.5 91.9 97.3	86.6 92.6 98.3	87.5 92.7 96.8	87.8 92.4 97.9
	91.8 96.2 98.9	89.5 94.4 98.4	88.8 93.5 98.1	88.7 93.8 97.2	89.6 94.5 98.9
$b = 1$	86.9 91.0 95.2	83.6 90.7 97.3	85.3 91.2 97.7	79.4 87.0 94.0	85.2 90.7 97.1
	89.0 93.5 97.2	89.1 94.4 98.5	88.4 93.1 98.6	87.1 91.9 97.1	86.5 92.6 98.3
	92.6 96.5 99.2	90.0 94.9 98.8	87.3 93.4 98.4	89.3 94.0 98.6	89.7 95.2 99.0
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	83.6 89.1 93.7	82.1 86.5 91.5	83.3 88.1 92.8	83.7 89.7 94.4	81.6 87.6 93.7
	88.3 93.2 97.4	85.5 91.1 96.5	86.6 92.7 97.1	87.2 93.0 96.8	84.0 90.2 95.3
	93.2 96.4 99.1	90.5 95.3 98.7	88.3 92.6 97.8	90.3 94.0 97.9	89.0 93.2 98.2
$b = .5$	82.9 89.4 94.5	83.3 89.5 96.7	83.7 89.9 97.3	78.8 84.8 92.4	82.8 88.0 94.5
	90.3 93.9 97.9	85.8 92.5 97.9	88.0 93.4 98.0	86.5 91.8 96.8	87.0 92.3 97.5
	90.6 95.1 99.1	89.7 93.4 98.7	88.3 93.6 98.8	89.0 93.9 98.3	86.9 93.0 98.5
$b = 1$	86.0 91.0 96.0	85.5 91.0 96.0	86.0 92.4 97.4	84.5 89.8 96.0	84.9 91.3 97.1
	88.1 93.4 97.2	87.4 93.3 98.0	87.2 91.8 98.4	85.2 91.7 97.1	87.5 94.1 98.7
	91.8 95.9 99.1	88.2 94.3 98.7	88.7 93.3 99.2	88.3 93.5 98.5	89.5 95.5 99.5

Table 3: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Negative Memory with  $\beta = -.6$ .

Negative Memory: $\beta = -.4$					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	85.2 89.0 93.2	82.2 86.5 92.0	82.2 86.6 90.7	83.7 88.7 92.6	79.7 84.9 90.8
	87.5 92.0 96.0	84.3 89.1 93.8	83.9 88.5 93.6	86.6 90.8 94.4	83.0 88.7 93.3
	99.5 99.9 100	99.3 99.9 100	98.8 99.6 99.9	99.6 100 100	99.5 100 100
$b = .5$	79.2 83.2 88.9	80.8 86.8 93.4	82.2 87.8 95.5	81.3 85.8 90.2	80.7 86.0 92.8
	84.9 90.5 95.2	84.9 91.2 96.5	85.0 92.6 96.8	85.1 90.1 94.5	87.2 92.9 97.2
	96.9 98.7 99.7	92.5 96.0 98.6	90.3 94.1 97.7	96.7 98.6 99.6	94.1 96.9 99.3
$b = 1$	80.7 87.5 91.8	85.2 91.0 97.3	83.8 89.6 95.8	83.2 88.1 92.8	86.4 91.6 97.0
	89.0 94.0 97.4	90.1 95.1 98.3	87.6 94.6 98.2	86.8 92.2 96.6	86.0 92.0 97.0
	90.1 95.1 98.3	90.5 94.8 98.1	90.4 94.5 98.5	90.3 94.9 98.0	89.9 94.4 98.3
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	81.2 86.8 92.4	80.0 85.3 90.5	82.0 87.1 92.1	83.5 88.0 92.8	82.3 88.1 93.0
	87.0 90.4 95.8	85.0 89.4 94.7	86.6 91.0 95.0	85.9 91.2 95.3	87.2 91.5 95.2
	90.9 95.8 99.0	86.6 91.4 96.2	89.7 94.2 98.6	89.0 93.9 97.6	88.6 93.2 97.6
$b = .5$	82.4 87.3 92.2	81.6 88.9 95.7	81.1 88.9 95.6	81.5 85.8 89.6	81.7 87.7 94.3
	87.2 92.1 96.2	87.7 93.3 97.8	86.9 92.6 98.2	85.9 91.1 96.4	85.8 92.0 97.1
	90.8 96.0 98.5	87.3 94.2 98.1	88.9 94.4 98.2	88.6 93.5 97.6	88.7 94.1 98.3
$b = 1$	84.1 87.1 91.5	84.0 90.5 96.6	83.2 89.8 96.9	82.3 87.7 93.4	82.9 90.1 96.8
	86.2 91.2 96.4	88.3 94.0 98.1	86.1 92.1 97.8	87.5 92.5 96.6	88.1 93.5 98.4
	91.2 95.1 98.5	88.4 93.6 99.0	88.9 94.7 98.9	89.1 94.7 99.3	88.9 94.1 98.7
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	84.9 88.7 92.7	82.9 87.8 92.1	82.0 87.2 91.1	83.3 88.0 92.9	83.3 89.1 92.7
	88.5 92.3 95.7	84.2 90.2 95.2	87.2 92.2 96.2	85.9 90.3 95.7	86.4 90.8 95.4
	88.4 93.4 97.6	89.5 93.8 97.6	89.3 93.9 97.5	87.7 93.0 98.1	88.3 93.0 96.7
$b = .5$	82.6 87.1 92.6	83.0 87.8 94.9	83.6 88.2 94.5	80.9 85.7 90.5	81.0 86.8 93.1
	88.7 92.8 96.2	84.6 90.9 98.4	86.8 93.1 98.2	85.5 92.2 95.9	86.5 91.2 97.1
	90.0 94.0 97.9	89.4 95.0 99.2	86.8 93.8 99.3	89.3 95.0 98.4	89.2 94.3 98.0
$b = 1$	82.6 88.6 93.1	84.3 90.6 97.3	86.4 92.2 98.4	81.0 86.8 93.2	86.1 91.3 97.1
	86.6 91.6 96.3	87.1 93.6 98.4	86.7 93.5 98.9	86.5 91.9 96.2	86.5 92.8 98.1
	88.0 94.1 98.3	88.1 93.7 98.6	87.6 94.4 99.4	88.9 93.8 98.4	90.4 94.7 99.1

Table 4: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Negative Memory with  $\beta = -.4$ .

Negative Memory: $\beta = -.2$					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	79.0 83.8 88.6	79.3 84.1 88.1	80.1 83.7 88.1	79.8 85.4 90.9	78.6 84.2 89.7
	84.9 89.7 93.7	83.2 87.7 92.2	83.8 88.3 92.8	84.0 89.6 93.8	82.8 88.9 92.9
	97.6 99.3 99.8	97.1 99.0 99.9	96.4 98.4 99.5	97.8 99.1 99.6	96.6 99.2 99.8
$b = .5$	80.1 85.3 90.3	80.4 86.5 92.6	80.7 87.2 94.7	80.6 84.6 89.0	79.6 84.7 91.1
	85.5 92.3 94.8	84.6 90.2 96.7	84.7 91.3 96.1	86.2 90.9 95.2	84.8 90.0 95.2
	94.2 97.1 99.7	91.3 94.4 98.5	90.3 93.9 97.8	93.3 96.5 99.3	91.6 94.7 98.4
$b = 1$	80.2 86.0 91.0	82.7 88.6 95.4	81.3 87.8 94.8	81.0 85.8 91.2	83.5 90.1 96.1
	85.6 90.4 94.4	88.0 93.6 97.5	86.6 93.5 97.3	87.3 91.6 95.7	87.2 93.8 98.0
	87.2 93.8 97.7	89.2 93.9 98.7	87.7 93.3 98.2	87.6 92.5 97.3	88.6 94.8 98.4
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	80.4 85.8 90.0	80.9 85.4 90.1	79.8 84.6 89.0	80.8 85.1 90.9	80.5 84.3 90.3
	87.1 91.8 95.3	86.8 90.0 93.8	85.3 90.7 95.4	85.6 89.3 94.3	85.5 89.3 93.1
	90.7 95.4 98.5	86.5 92.0 96.8	89.8 94.9 98.5	86.4 91.6 97.1	90.3 93.9 97.6
$b = .5$	79.1 84.6 88.9	81.3 87.6 94.3	82.0 89.7 95.2	81.3 85.8 91.4	81.0 86.8 92.6
	88.2 91.6 95.7	86.3 91.7 96.6	87.2 92.9 98.1	84.3 89.0 94.3	84.4 89.5 95.2
	89.9 94.7 98.2	90.3 94.3 98.0	86.7 93.1 98.6	87.6 92.5 96.8	88.2 93.9 98.3
$b = 1$	81.4 86.4 91.2	82.6 88.8 96.6	84.8 90.6 96.7	80.9 86.6 92.2	83.3 89.7 95.7
	84.5 89.9 95.1	87.8 92.6 97.7	86.7 92.2 98.0	87.9 93.0 96.5	88.0 93.0 98.5
	89.8 94.8 98.5	89.8 94.7 98.9	87.4 93.1 98.7	88.6 93.9 98.2	90.5 94.9 99.4
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	81.2 86.2 90.9	78.9 83.8 88.7	79.9 85.7 90.4	80.1 85.5 90.1	81.4 86.2 91.0
	86.9 89.5 94.6	85.6 90.2 95.1	86.4 90.3 94.8	84.6 89.8 94.3	83.8 89.9 95.5
	89.4 94.3 97.9	88.6 93.3 97.2	89.1 94.3 97.7	90.6 94.6 97.8	88.4 92.5 97.3
$b = .5$	80.7 85.1 89.7	82.7 88.2 94.8	81.5 87.0 95.7	79.2 84.1 90.3	80.6 87.5 93.5
	85.4 90.5 96.0	86.7 92.1 97.9	84.4 91.3 98.3	84.7 90.8 95.3	85.7 89.7 97.0
	87.0 91.9 97.9	86.6 92.3 97.6	87.9 94.1 98.9	88.0 92.8 97.7	87.7 93.1 98.6
$b = 1$	82.9 87.2 90.7	85.3 91.4 97.0	83.3 89.1 96.9	83.0 88.7 94.1	83.1 89.3 96.1
	85.8 91.2 95.6	86.5 92.2 98.2	87.5 92.8 98.3	86.7 92.4 97.1	86.5 91.6 97.5
	89.7 94.4 98.2	88.4 93.4 98.5	87.8 93.2 98.6	87.2 93.0 97.5	87.9 93.3 98.4

Table 5: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Negative Memory with  $\beta = -.2$ .

Short Memory					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	77.8 82.0 86.2	77.3 82.1 85.7	77.8 82.8 87.6	78.6 83.7 88.4	77.0 82.6 85.7
	84.1 89.9 93.1	80.7 86.3 90.6	81.8 88.4 93.0	85.7 89.9 93.3	85.8 90.0 93.4
	91.6 95.8 98.0	91.5 95.5 98.6	92.2 95.9 98.7	91.4 96.4 98.9	92.6 95.7 98.8
$b = .5$	79.4 84.7 88.0	79.3 85.4 91.2	81.5 86.4 93.8	80.0 84.1 87.5	77.3 83.2 89.6
	84.3 89.9 93.7	84.6 89.4 94.9	85.2 92.4 96.9	85.0 90.2 93.6	83.3 89.3 94.9
	89.6 94.7 98.3	89.7 94.1 97.8	89.0 92.8 97.2	91.2 95.3 98.4	90.1 93.6 98.2
$b = 1$	79.3 83.4 87.9	81.3 88.6 94.9	82.7 89.9 95.8	79.5 84.6 88.9	83.0 88.4 94.2
	81.4 87.9 92.4	87.1 92.9 97.2	85.5 91.9 96.9	84.0 90.7 94.7	84.2 91.8 97.8
	87.4 92.0 97.2	88.9 93.3 97.7	88.7 92.9 98.1	89.5 94.2 98.2	89.0 93.9 97.8
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	77.7 83.9 87.5	77.4 81.3 85.0	81.7 86.7 90.6	78.7 82.5 87.0	77.6 82.8 86.7
	84.6 88.8 92.1	84.5 89.4 94.4	85.4 90.3 94.1	84.8 88.9 93.3	82.9 88.1 92.6
	88.2 92.9 97.2	87.7 92.4 96.7	87.4 93.1 97.7	86.9 92.1 97.0	88.3 93.6 97.3
$b = .5$	80.6 84.9 89.1	81.1 85.2 91.3	81.7 88.7 94.8	75.7 80.4 86.3	78.7 83.9 90.7
	85.1 89.4 93.9	86.5 91.6 97.7	85.6 91.9 97.2	84.0 88.9 93.4	83.7 89.7 96.5
	86.9 91.6 96.8	88.2 93.8 98.7	87.0 92.8 98.4	87.0 92.3 97.2	90.0 93.4 98.3
$b = 1$	78.5 84.6 88.8	83.2 89.1 95.3	83.6 89.5 95.5	81.2 85.6 90.1	82.8 88.4 94.5
	85.3 91.1 95.6	85.3 90.6 97.0	88.3 93.2 97.7	86.7 90.6 95.6	87.2 93.3 97.6
	86.7 92.5 97.3	87.2 93.1 98.6	87.8 94.4 98.4	86.5 92.5 98.0	87.4 93.9 99.0
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	77.5 82.3 86.3	77.9 82.2 86.9	80.1 84.2 87.8	78.9 82.9 87.8	78.7 82.4 87.1
	84.4 89.1 93.6	84.2 89.5 92.9	84.9 89.3 93.7	84.8 89.0 93.2	84.6 89.1 93.0
	86.4 93.1 97.2	87.8 92.5 97.7	88.6 93.9 97.5	88.5 93.1 96.8	87.2 91.9 97.5
$b = .5$	76.6 81.6 86.2	79.6 85.3 92.5	80.5 87.0 94.0	77.5 83.6 88.0	80.0 84.7 91.1
	83.4 88.1 93.9	84.6 90.7 97.5	85.8 91.9 97.8	83.1 88.8 93.6	86.1 90.1 95.9
	88.0 93.0 96.7	89.2 93.4 97.7	88.2 92.4 98.4	88.6 93.7 97.7	86.7 91.3 98.0
$b = 1$	77.2 82.3 88.2	82.9 89.4 96.2	80.8 87.7 95.9	76.0 82.2 87.7	81.2 87.3 94.0
	85.3 90.2 93.9	88.4 93.2 99.1	85.1 91.8 97.7	88.3 92.6 96.3	85.2 91.9 97.6
	87.9 92.4 96.9	87.0 93.3 98.1	89.6 93.7 99.0	88.5 92.6 98.2	88.7 93.6 98.6

Table 6: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Short Memory.

Long Memory: $\beta = .2$					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	78.2 82.1 84.7	81.3 85.6 88.5	75.0 78.4 82.4	80.0 82.6 85.9	80.2 83.7 86.6
	89.5 92.4 94.3	90.1 93.1 94.6	87.7 91.8 94.1	91.0 94.2 95.6	89.7 92.5 94.4
	80.8 87.6 94.7	81.4 88.6 95.6	84.0 90.3 95.5	75.7 84.5 93.6	84.3 90.0 95.6
$b = .5$	79.2 82.6 85.3	78.6 83.4 88.7	81.1 85.7 91.7	77.9 81.3 84.4	78.3 82.8 87.2
	85.6 90.9 94.1	83.5 89.2 93.3	84.5 90.2 95.5	86.2 90.6 95.0	83.3 89.3 94.3
	93.8 96.2 98.3	94.4 97.1 99.2	95.2 97.2 99.1	92.7 96.1 98.5	95.8 97.6 99.1
$b = 1$	77.5 81.8 86.3	81.1 87.4 92.7	81.4 87.4 94.8	76.8 81.6 86.8	80.3 86.5 91.4
	87.4 92.8 95.6	85.0 91.6 97.0	88.8 94.1 98.1	83.7 90.2 94.4	84.7 90.7 96.3
	91.1 94.8 98.3	91.8 95.2 98.8	89.7 94.0 97.8	91.8 96.0 99.0	90.1 94.7 98.7
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	78.2 82.9 85.8	77.5 81.9 85.1	75.8 79.2 83.3	78.2 81.7 85.4	77.1 82.4 85.8
	87.7 91.6 94.0	85.1 88.8 92.6	87.3 91.4 94.6	89.7 91.5 94.9	87.6 90.9 93.8
	94.3 97.3 98.9	93.4 96.0 98.4	94.4 96.9 98.7	95.1 97.5 98.8	95.4 97.5 98.9
$b = .5$	77.6 81.1 85.2	77.1 81.8 87.7	77.9 84.6 92.4	77.8 82.6 86.1	79.8 84.9 89.3
	85.5 90.7 94.6	84.3 89.6 95.7	84.0 90.3 95.9	85.4 90.0 93.9	83.8 89.3 94.7
	92.3 95.8 98.3	91.5 94.1 98.3	89.2 94.8 99.4	89.9 94.1 98.1	89.0 93.6 98.3
$b = 1$	79.3 84.1 87.4	79.3 86.7 93.1	81.5 87.9 95.1	77.5 83.0 88.4	79.6 85.9 92.9
	88.4 92.6 96.0	86.7 92.5 97.5	88.3 92.3 97.9	84.9 88.9 93.5	86.7 90.7 96.5
	91.5 94.9 97.7	89.3 94.5 98.8	92.3 95.4 98.8	89.4 94.7 98.6	89.3 94.7 97.6
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	79.3 82.5 86.1	78.1 82.1 84.8	78.3 82.6 84.7	78.5 82.4 86.5	78.0 82.1 84.6
	85.8 90.4 93.8	85.2 89.8 93.8	86.6 90.4 94.9	88.4 91.0 93.6	85.4 89.9 93.1
	93.1 95.8 98.1	93.1 96.1 98.3	92.6 95.6 98.0	93.8 97.0 99.0	92.6 96.1 98.8
$b = .5$	76.1 81.2 85.0	78.2 83.6 89.0	77.2 83.8 92.7	76.2 79.7 83.1	77.7 82.3 87.1
	84.5 89.8 93.2	83.7 90.0 94.7	85.5 91.9 96.7	84.0 89.1 92.9	84.8 89.7 95.1
	91.3 95.3 98.8	87.3 92.5 98.2	90.2 95.1 98.4	90.6 94.2 98.2	90.5 94.6 98.0
$b = 1$	77.9 83.0 87.8	82.8 88.0 94.5	81.1 89.1 96.1	78.7 83.1 87.9	80.4 86.0 92.8
	84.4 90.1 94.7	86.8 93.0 97.5	88.1 92.8 98.5	86.1 91.3 94.9	84.0 90.7 96.8
	91.7 94.7 98.3	90.2 94.2 98.6	91.4 95.7 99.5	88.9 93.8 97.6	90.7 94.2 98.4

Table 7: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Long Memory with  $\beta = .2$ .

Long Memory: $\beta = .4$					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	76.7 80.6 83.4	74.3 78.0 80.3	76.1 79.2 82.1	75.9 78.9 81.7	76.0 79.5 83.0
	87.4 90.7 94.1	87.5 92.0 94.4	85.7 89.6 92.0	87.0 90.9 93.2	86.8 90.1 93.4
	73.4 80.8 88.1	76.1 83.9 91.6	81.6 87.1 93.5	69.8 77.5 86.8	76.0 83.7 91.7
$b = .5$	72.6 76.8 81.7	76.7 80.8 85.9	79.2 84.9 90.0	74.6 79.1 81.8	74.4 78.4 84.4
	83.6 88.2 92.3	81.9 88.0 93.6	83.3 89.6 95.1	84.8 90.0 92.6	82.8 88.9 94.0
	90.0 95.0 98.0	93.3 96.5 98.4	94.4 96.7 98.7	90.3 94.5 97.4	91.0 95.0 97.3
$b = 1$	73.3 79.2 83.8	76.7 83.6 91.7	78.3 85.8 93.0	76.5 81.3 86.3	76.4 82.8 89.2
	83.2 88.5 92.1	84.6 90.4 96.2	84.2 91.6 97.6	82.7 88.5 93.1	84.0 89.6 94.8
	91.2 95.7 98.5	89.0 94.4 98.0	91.1 95.4 98.6	90.7 95.5 99.2	89.5 93.8 97.8
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	75.8 80.7 83.7	74.7 77.8 80.5	74.3 78.6 81.9	78.2 81.5 83.8	76.9 80.0 83.4
	86.8 90.3 93.4	86.0 89.4 92.1	85.4 89.3 92.6	86.9 89.8 94.2	84.9 88.5 91.8
	92.2 95.7 98.0	93.4 96.3 98.7	93.6 96.6 98.7	94.1 96.4 98.8	95.0 97.7 98.9
$b = .5$	75.0 79.3 83.0	75.2 80.7 85.9	76.6 81.8 89.7	73.3 78.3 81.7	76.1 79.6 85.2
	83.6 87.9 91.9	83.2 87.8 94.1	83.4 89.1 95.7	81.8 86.9 91.7	85.1 91.0 94.7
	89.2 93.9 98.3	90.3 93.5 98.3	89.4 93.1 97.5	89.2 94.1 97.6	87.5 93.5 97.3
$b = 1$	73.9 78.8 83.8	79.5 85.2 93.6	79.0 85.6 94.7	75.6 80.0 85.8	79.1 84.0 90.5
	83.1 88.2 93.3	83.8 90.4 96.4	84.4 90.7 96.4	82.8 88.2 92.6	82.5 88.5 94.6
	90.7 94.3 98.1	87.8 93.0 98.1	90.8 94.4 98.7	90.2 94.3 98.7	89.3 94.5 98.4
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	75.5 78.7 82.0	74.6 78.1 80.8	71.5 76.1 80.6	74.9 78.2 82.5	74.6 78.6 82.4
	87.6 90.2 93.0	85.1 89.1 92.0	84.2 87.8 91.9	85.1 89.0 91.7	84.9 88.8 91.8
	91.8 95.7 98.2	91.1 94.0 97.7	89.3 93.9 97.6	92.9 96.5 99.0	90.7 94.0 97.0
$b = .5$	73.7 77.8 81.3	75.5 80.2 85.4	75.3 82.3 89.9	74.9 80.2 84.3	73.3 79.5 85.9
	83.5 89.4 93.6	82.4 88.8 93.8	83.8 89.8 96.0	83.4 87.8 92.0	84.1 89.8 94.7
	90.4 94.7 97.8	88.5 94.2 97.9	88.3 92.7 97.4	86.7 91.4 96.1	88.4 93.0 97.4
$b = 1$	76.0 80.9 85.0	79.0 85.0 93.1	80.4 86.8 94.3	75.1 82.0 87.3	77.9 83.9 91.0
	81.8 86.7 91.1	85.0 91.2 97.7	84.6 91.4 97.0	82.2 87.0 92.6	87.3 92.2 96.5
	89.3 93.6 97.9	88.2 93.1 98.5	88.8 94.4 98.5	90.1 94.4 98.2	88.8 93.9 98.3

Table 8: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Long Memory with  $\beta = .4$ .

Long Memory: $\beta = .6$					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	70.9 74.7 77.4	71.6 76.6 79.3	70.5 74.6 77.5	71.3 74.6 78.1	70.9 74.9 76.7
	82.3 86.2 89.1	84.0 88.5 91.7	81.5 86.8 90.5	81.8 85.8 88.9	84.9 88.5 91.7
	61.2 70.5 81.4	65.3 73.2 83.4	73.3 80.3 87.9	59.0 67.6 80.2	63.7 72.0 81.0
$b = .5$	71.6 76.6 80.5	70.8 78.0 84.1	72.0 78.3 84.9	67.7 72.6 75.1	71.3 76.3 82.5
	79.4 84.8 89.9	79.5 84.7 90.9	79.9 86.9 93.8	80.4 85.9 89.3	80.2 86.8 92.2
	86.7 90.4 95.1	89.5 93.4 96.6	88.9 93.5 97.0	86.9 91.0 95.1	87.1 90.9 95.3
$b = 1$	71.2 75.9 80.2	73.9 80.8 90.8	74.8 82.4 90.9	69.7 75.5 80.8	75.9 81.4 88.3
	82.0 87.7 92.0	81.3 88.9 94.0	82.8 90.0 95.3	80.6 85.8 91.8	83.2 89.3 95.0
	86.8 91.1 96.4	89.6 94.0 98.1	88.2 93.8 97.7	88.3 93.0 97.9	87.9 92.4 97.4
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	70.6 74.3 77.7	67.3 71.7 74.9	67.8 71.5 74.6	69.4 73.5 76.8	70.4 74.8 78.0
	79.4 83.6 88.7	83.1 86.8 90.2	81.5 85.9 89.1	81.3 84.8 88.1	79.1 84.6 88.2
	88.3 92.5 97.0	88.5 92.6 97.6	89.8 94.0 97.5	91.1 94.3 97.1	89.0 93.2 97.0
$b = .5$	70.2 74.2 78.2	73.8 79.5 84.6	73.0 80.2 87.8	68.9 73.2 76.7	72.4 76.6 81.6
	81.6 87.2 90.9	77.2 84.6 92.6	79.2 86.5 93.9	79.3 84.2 88.8	81.7 88.2 92.7
	85.9 92.2 96.9	86.2 91.3 97.3	84.8 91.3 97.6	87.2 91.8 97.4	85.7 90.3 96.3
$b = 1$	71.5 77.1 81.6	75.6 82.4 92.9	77.1 84.2 93.0	68.7 75.7 81.9	74.2 81.0 88.2
	77.6 83.5 88.9	84.0 89.4 96.8	82.3 88.9 96.5	82.7 86.9 92.8	80.0 86.0 93.9
	87.2 91.5 96.5	87.7 92.8 98.2	88.2 94.6 98.5	87.8 93.0 97.6	89.2 94.4 99.0
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	68.6 73.0 76.4	69.8 73.8 76.2	67.8 71.8 75.2	67.0 70.6 74.1	68.1 71.5 74.8
	79.4 83.9 88.2	79.7 84.5 88.7	77.7 82.5 87.6	81.2 85.1 88.9	81.1 86.9 90.5
	87.4 91.2 95.8	88.0 92.2 96.3	86.6 93.0 97.3	88.2 92.6 96.0	88.3 93.3 96.3
$b = .5$	71.9 77.1 80.8	71.3 77.3 83.0	70.8 76.9 85.4	68.5 74.2 77.6	72.2 77.6 83.5
	79.5 84.7 90.0	80.9 86.5 93.2	80.4 87.7 94.7	77.2 82.8 87.8	79.3 85.7 92.8
	85.1 90.9 96.0	86.8 91.5 97.7	87.2 94.0 98.2	84.5 90.3 95.7	86.7 91.5 97.4
$b = 1$	69.5 74.9 79.6	73.8 81.3 91.4	74.4 82.2 92.3	72.4 78.8 83.2	72.8 80.3 87.9
	80.3 85.6 93.1	83.5 89.9 96.5	83.1 90.9 97.5	79.2 84.8 92.6	81.4 89.0 96.1
	86.4 92.5 97.7	86.7 92.7 97.8	86.4 92.8 98.4	85.1 91.6 96.9	86.3 91.5 97.2

Table 9: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Long Memory with  $\beta = .6$ .

Long Memory: $\beta = .8$					
Sample Size	Tapers				
n= 250	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	59.8 63.0 65.8	59.2 62.4 64.7	60.6 64.2 68.3	58.9 63.3 65.9	60.4 63.9 66.5
	72.8 78.4 81.6	75.8 80.5 83.9	76.0 81.2 84.9	73.3 78.6 83.4	73.9 78.9 82.4
	49.3 58.7 68.3	50.0 59.6 69.9	56.0 64.1 73.7	43.6 50.9 61.5	49.8 56.9 68.4
$b = .5$	61.1 66.8 70.9	61.8 68.3 76.2	65.0 71.6 81.2	58.8 63.8 67.5	62.6 67.0 72.7
	73.9 81.2 86.5	70.6 78.6 86.8	76.4 84.0 92.4	72.0 78.7 84.5	74.2 81.2 86.5
	74.8 81.9 89.0	77.6 84.5 91.6	82.1 87.3 92.8	72.4 78.9 87.4	79.4 85.2 91.8
$b = 1$	62.3 67.2 72.6	68.1 76.8 85.0	66.8 75.9 85.9	64.2 70.5 76.6	65.4 72.0 83.0
	74.9 80.6 86.4	75.9 84.4 92.8	79.0 87.0 94.4	75.5 82.3 89.3	76.1 84.9 92.0
	82.1 88.6 93.9	85.0 91.1 97.7	85.4 91.9 97.3	80.8 87.3 94.2	83.7 89.9 96.5
n = 500	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	59.0 62.9 66.0	59.6 63.7 66.8	59.5 63.9 66.0	55.6 59.0 62.4	57.2 61.2 63.6
	69.0 74.1 79.1	71.1 75.9 80.9	69.4 74.7 79.4	72.3 76.7 81.8	71.8 75.7 81.8
	79.3 84.9 92.3	81.0 85.9 91.7	81.2 86.2 92.2	81.4 87.0 92.1	80.9 86.2 92.8
$b = .5$	59.5 64.2 68.8	64.4 70.8 78.1	63.0 70.9 81.1	60.1 64.1 68.6	62.0 68.0 74.7
	69.2 75.5 83.7	74.1 81.9 89.9	72.2 80.3 91.3	72.9 79.8 85.9	73.5 80.9 88.7
	82.8 88.3 94.5	78.4 86.3 95.3	82.5 89.1 96.6	81.7 87.5 95.1	81.9 87.7 96.0
$b = 1$	63.3 69.9 75.0	68.6 77.6 87.3	63.2 73.5 86.1	62.8 70.0 75.4	66.7 74.7 84.0
	75.1 79.7 88.0	78.3 85.0 94.0	77.6 84.3 94.7	73.6 80.2 86.8	76.3 85.2 93.8
	79.9 87.1 94.3	84.0 90.4 97.6	83.9 90.2 96.8	80.3 88.3 95.5	82.0 89.2 96.3
n = 1000	Bartlett	Trapezoid (.25)	Trapezoid (.5)	Parzen	Daniell
$b = .1$	56.6 60.3 63.7	55.6 60.3 63.2	54.8 60.4 64.6	55.4 58.4 61.2	57.7 61.1 64.9
	68.5 74.6 80.4	70.4 75.5 79.7	67.6 74.1 80.5	65.5 71.9 77.7	68.1 73.3 78.6
	79.3 85.7 92.1	80.2 86.4 91.0	78.8 83.1 89.9	80.6 85.3 91.8	80.2 86.5 92.4
$b = .5$	63.1 67.4 70.7	63.1 69.2 77.3	67.1 74.8 83.9	61.1 65.2 68.8	62.4 68.8 76.9
	69.5 76.3 83.6	75.9 82.1 89.5	75.5 83.5 91.9	71.4 78.4 83.8	69.7 78.5 86.2
	83.8 88.7 95.1	79.2 86.9 95.8	80.8 87.7 95.4	81.5 87.6 93.4	77.3 85.8 94.2
$b = 1$	61.3 66.7 71.6	69.0 76.7 87.3	66.4 75.5 87.3	61.0 67.7 74.5	67.9 74.7 83.2
	74.0 80.0 86.2	78.1 86.1 94.8	74.4 81.9 93.1	74.4 81.9 89.1	76.1 82.6 91.8
	79.8 86.4 93.8	80.4 87.4 96.3	82.8 89.8 98.5	78.6 86.3 94.4	83.5 89.9 96.4

Table 10: Empirical size for two-sided test with Type I error rate  $\alpha = .10, .05, .01$  for the left, middle, and right hand entries respectively, in each cell. In each cell, the first row is for sampling fraction  $a = .2$ , the second row for  $a = .1$ , and the third row for  $a = .04$ . Various sample sizes and taper bandwidths  $b$  are considered. The data process is Long Memory with  $\beta = .8$ .



Optimal Bandwidth Fraction									
	Beta								
	- .8	- .6	- .4	- .2	0	.2	.4	.6	.8
Bartlett									
.90	.02	.02	.02	.02	.02	.10	.14	.14	.18
.95	.02	.02	.02	.02	.02	.06	.10	.14	.18
.975	.02	.02	.02	.02	.02	.06	.10	.14	.14
.99	.02	.02	.02	.02	.02	.04	.10	.14	.14
Trapezoid (.25)									
.90	.02	.02	.02	.02	.02	.06	.12	.14	.14
.95	.02	.02	.02	.02	.02	.06	.10	.12	.14
.975	.02	.02	.02	.02	.02	.04	.08	.14	.14
.99	.02	.02	.02	.02	.02	.04	.08	.08	.10
Trapezoid (.50)									
.90	.02	.02	.02	.02	.02	.04	.10	.12	.16
.95	.02	.02	.02	.02	.02	.04	.10	.10	.12
.975	.02	.02	.02	.02	.02	.04	.06	.10	.12
.99	.02	.02	.02	.02	.02	.02	.04	.06	.06
Parzen									
.90	.02	.02	.02	.02	.02	.12	.20	.22	.28
.95	.02	.02	.02	.02	.02	.06	.16	.20	.20
.975	.02	.02	.02	.02	.02	.06	.10	.22	.20
.99	.02	.02	.02	.02	.02	.06	.08	.14	.20
Daniell									
.90	.02	.02	.02	.02	.02	.06	.10	.16	.18
.95	.02	.02	.02	.02	.02	.06	.10	.16	.14
.975	.02	.02	.02	.02	.02	.06	.10	.08	.14
.99	.02	.02	.02	.02	.02	.04	.06	.08	.10

Table 11: Optimal bandwidth fractions  $b$  as a function of taper and memory parameter  $\beta$ . Each row corresponds to either the (upper one-sided) .90, .95, .975, or .99 quantile.




Figure 1: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Bartlett taper with bandwidth fraction  $b = .2$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 2: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Bartlett taper with bandwidth fraction  $b = .5$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 3: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Bartlett taper with bandwidth fraction  $b = 1$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 4: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6, .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Trapezoidal (.25) taper with bandwidth fraction  $b = .2$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 5: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6, .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Trapezoidal (.25) taper with bandwidth fraction  $b = .2$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 6: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Trapezoidal (.25) taper with bandwidth fraction  $b = 1$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 7: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Trapezoidal (.5) taper with bandwidth fraction  $b = .2$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .






Figure 8: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Trapezoidal (.5) taper with bandwidth fraction  $b = .5$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 9: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Trapezoidal (.5) taper with bandwidth fraction  $b = .5$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 10: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6, .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Parzen taper with bandwidth fraction  $b = .2$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 11: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6, .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Parzen taper with bandwidth fraction  $b = .5$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 12: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Parzen taper with bandwidth fraction  $b = 1$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 13: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6., .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Daniell taper with bandwidth fraction  $b = .2$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 14: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6, .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Daniell taper with bandwidth fraction  $b = .5$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .




Figure 15: Power surfaces by Memory parameter  $\beta \in \{-1, -.8, -.6, -.4, -.2, 0, .2, .4, .6, .8\}$  and true mean  $\mu \in [0, 1)$ . These power surfaces are for the Daniell taper with bandwidth fraction  $b = 1$ . Panel (a) corresponds to subsampling with sampling fraction  $a = .04$ ; panel (b) corresponds to subsampling with sampling fraction  $a = .12$ ; panel (c) corresponds to subsampling with sampling fraction  $a = .2$ . Panel (d) corresponds to using the plug-in estimate of  $\beta$ ; panel (e) corresponds to using the plug-in method with  $\beta = 0$ ; panel (f) corresponds to using the true (unknown) value of  $\beta$ .