

AN ELEMENTARY PROOF OF THE RECONSTRUCTION CONJECTURE

First Author^{*1}, Some Second Author², and Some Third Author³

¹*Department of Inconsequential Studies, Solatido College, North Kentucky, U.S.A.
fa@solatido.edu*

^{2,3}*School of Hard Knocks, University of Western Nowhere, Somewhere, Australia
ssa@uwn.edu.au, sta@uwn.edu.au*

Submitted: Jan 1, 2020; Accepted: Jan 2, 2020; Published: TBD

© The authors. Released under the CC BY license (International 4.0).

Abstract. The reconstruction conjecture states that the multiset of vertex-deleted subgraphs of a graph determines the graph, provided it has at least 3 vertices. This problem was independently introduced by Stanisław Ulam (1960) and Paul Kelly (1957). In this paper, we prove the conjecture by elementary methods. It is only necessary to integrate the Lenkle potential of the Broglington manifold over the quantum supervacillatory measure in order to reduce the set of possible counterexamples to a small number (less than a trillion). A simple computer program that implements Pipletti's classification theorem for torsion-free Aramaic groups with symplectic socles can then finish the remaining cases.

Keywords. Broglington manifolds, permutations

Mathematics Subject Classifications. 05C88, 05C89

1. Introduction

The reconstruction conjecture states that the multiset of unlabeled vertex-deleted subgraphs of a graph determines the graph, provided it has at least three vertices. This problem was independently introduced by Ulam [Ula60] and Kelly [Kel57]. The reconstruction conjecture is widely studied [Bol90, FGH72, HHRT07, KSU10, SW18] and is very interesting because it is. See [BH06] for more about the reconstruction conjecture.

Definition 1.1. A graph is *fabulous* if *rest of definition here*.

Theorem 1.2. *All planar graphs are fabulous.*

Proof. Suppose on the contrary that some planar graph is not fabulous. Then we have a contradiction. □

*Supported by NASA grant ABC123.

2. Broglington Manifolds

This section describes background information about Broglington Manifolds.

Lemma 2.1. *Broglington manifolds are abundant.*

Proof. A proof is given here. □

3. Proof of Theorem 1.2

In this section we complete the proof of Theorem 1.2.

Proof of Theorem 1.2. Let G be a graph. We have

$$\begin{aligned} |X| &= a + b + c \\ &= \alpha\beta\gamma. \end{aligned} \tag{3.1}$$

This completes the proof of Theorem 1.2. □

Figure 3.1: Here is an informative figure.

Acknowledgements

Thanks to Professor Qwerty for suggesting the proof of Lemma 2.1 and introducing us to the work of [Bjö95] as well as [KAH⁺13].

Some of the computation of the present paper have been done using the SageMath software [The18]. You can find our code over here [DG18].

References

- [BH06] J. A. Bondy and R. Hemminger. Graph reconstruction-a survey. *Journal of Graph Theory*, 1:227 – 268, 10 2006. doi:10.1002/jgt.3190010306.
- [Bjö95] Anders Björner. Topological methods. In *Handbook of combinatorics, Vol. 1, 2*, pages 1819–1872. Elsevier Sci. B. V., Amsterdam, 1995.
- [Bol90] Béla Bollobás. Almost every graph has reconstruction number three. *Journal of Graph Theory*, 14:1 – 4, 03 1990. doi:10.1002/jgt.3190140102.
- [DG18] Joseph Doolittle and Bennet Goeckner. A counterexample to Stanley’s k-fold acyclic boolean interval decomposition conjecture., 2018. doi:10.5281/zenodo.5604926.

- [FGH72] J. Fisher, R. L. Graham, and F. Harary. A simpler counterexample to the reconstruction conjecture for denumerable graphs. *Journal of Combinatorial Theory Series B - JCTB*, 12:203–204, 04 1972. doi:10.1016/0095-8956(72)90026-3.
- [HHRT07] Edith Hemaspaandra, Lane Hemaspaandra, Stanislaw Radziszowski, and Rahul Tripathi. Complexity results in graph reconstruction. *Discrete Applied Mathematics*, 155:103–118, 01 2007. doi:10.1016/j.dam.2006.04.038.
- [KAH⁺13] Veselin Kostov, Daniel Allan, Nikolaus Hartman, Scott Guzewich, and Justin Rogers. “Winter is coming”. 2013. arXiv:1304.0445.
- [Kel57] Paul Kelly. A congruence theorem for trees. *Pacific J. Math*, 7, 03 1957. doi:10.2140/pjm.1957.7.961.
- [KSU10] Masashi Kiyomi, Toshiki Saitoh, and Ryuhei Uehara. Reconstruction of interval graphs. *Theor. Comput. Sci.*, 411:3859–3866, 10 2010. doi:10.1016/j.tcs.2010.07.006.
- [SW18] Hannah Spinoza and Douglas West. Reconstruction from the deck of k -vertex induced subgraphs. *Journal of Graph Theory*, 90, 10 2018. doi:10.1002/jgt.22409.
- [The18] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 8.4)*, 2018. URL: <https://www.sagemath.org>.
- [Ula60] S.M. Ulam. *A Collection of Mathematical Problems*. Interscience tracts in pure and applied mathematics. Interscience Publishers, 1960. URL: https://books.google.ca/books?id=u_kHAAAAMAAJ.