

Lawrence Berkeley National Laboratory

Recent Work

Title

TARGET FRAGMENT ANGULAR DISTRIBUTIONS FOR THE INTERACTION OF 25.2 GeV 12c WITH 197Au

Permalink

<https://escholarship.org/uc/item/005149g1>

Authors

Morita, Y.
Loveland, W.
Seaborg, G.T.

Publication Date

1983-06-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED
LAWRENCE
BERKELEY LABORATORY

OCT 19 1983

LIBRARY AND
DOCUMENTS SECTION

To be published in Physical Review C


TARGET FRAGMENT ANGULAR DISTRIBUTIONS FOR THE
INTERACTION OF 25.2 GeV ^{12}C WITH ^{197}Au

Y. Morita, W. Loveland, and G.T. Seaborg

June 1983

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.*



LBL-16271
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

TARGET FRAGMENT ANGULAR DISTRIBUTIONS FOR THE
INTERACTION OF 25.2 GeV ^{12}C WITH ^{197}Au .

Y. Morita, W. Loveland* and G.T. Seaborg

Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

The target fragment angular distributions for the interaction of 25.2 GeV ^{12}C with ^{197}Au were measured. The data are generally in agreement with predictions based on the two step model and thick target-thick catcher recoil data.

NUCLEAR REACTIONS Au (^{12}C , x), $E_{^{12}\text{C}} = 25.2$ GeV,
 $\sigma(\theta)$, comparison of data with two step model,
firestreak model

PACS Numbers 25.70 Np

One of the least well understood aspects of relativistic nucleus-nucleus collisions is the momentum transferred to the target nucleus during the collision. There have been several studies of this problem with heavy target nuclei utilizing radiochemical techniques^{1,2,3,4,5} and physical measurements.^{6,7,8} From these studies we know that the target fragments may be divided into three general classes, fission fragments (produced with low to intermediate momenta), spallation products (whose momenta increase with increasing mass loss from the target) and the light fragments ($A < 60$) whose momenta in high energy heavy ion reactions is the greatest of all the fragments. Heckman⁹ has shown the relative magnitudes of the momenta of the light fragments and fission fragments can be understood in terms of a simple kinematic model that assumes the reaction takes place in two steps, a fast initial interaction between the projectile and the target and a second slow deexcitation step. Some of the assumptions of this model have been verified by Kaufman et al.⁷ However no current theoretical model of relativistic heavy ion collisions correctly predicts the magnitudes of the target fragment momenta.^{3,4}

In this brief report, we shall present additional information about the kinematic properties, i.e., the angular distributions, of these fragments and the class of fragments not studied by physical techniques, the spallation products. In particular, we shall report an extension of the previous measurements¹⁰ of target fragment angular distributions in the reaction of 3 and 12 GeV ^{12}C with ^{197}Au and ^{238}U to the case of 25.2 GeV ^{12}C interacting with ^{197}Au .

The experiment was performed at the LBL Bevalac using a 2.1 GeV/u ^{12}C beam. The target fragment angular distributions were measured utilizing exactly the same techniques, target thicknesses, etc. that have been described previously.¹⁰ The total beam fluence was $\sim 10^{12}$ particles over a period of 10 hours.

The measured fragment angular distributions are shown in Figure 1. In qualitative agreement with previous² thick target-thick catcher recoil studies, one observes very forward-peaked distributions for nuclides with $145 \leq A \leq 171$ (thought to be deep spallation products), a slightly forward-peaked distribution for a possible fission product (^{97}Ru) and a nearby isotropic distribution for the light fragment ($^{44\text{m}}\text{Sc}$). In all distributions the first point (23°) is considerably elevated with relation to the other three points in the distribution.

It is of interest to compare the measured fragment angular distributions to those predicted by the two step model of nuclear reactions and relevant recoil data.^{2,11} Porile et al.¹² have shown that the fragment angular distributions can be expressed in terms of the two step model recoil parameters η_{11} and b/a as

$$F_L(\theta_L) = \frac{1 + (b/a) \cos^2 [\theta_L + \sin^{-1} (\eta_{11} \sin \theta_L)]}{1 + b/3a} \times \frac{[\eta_{11} \cos \theta_L + (1 - \eta_{11}^2 \sin^2 \theta_L)^{1/2}]^2}{(1 - \eta_{11}^2 \sin^2 \theta_L)^{1/2}} \quad (1)$$

where $F_L(\theta_L)$ is the laboratory differential cross section at angle θ_L . Taking η_{11} from Ref. 2 and 11 (and assuming $b/a = 0$), equation (1) was used to calculate the distributions shown in Figure 1. Apart from the point at 23° , the predicted and measured distributions are in general agreement, indicating some general validity for the use of the two step model to describe average fragment properties.

It is also of interest to compare the predictions of a current theoretical model of relativistic heavy ion collisions, the nuclear firestreak model, with the measured distributions. To do this we used the firestreak model as implemented by McGaughey¹³ to predict the momenta of the emerging primary targets from the $^{12}\text{C} - ^{197}\text{Au}$ interaction. (In making this calculation, we used realistic nucleon-nucleon scattering cross sections. The model predicted a total reaction cross section of 2931 mb, in good agreement with the "soft spheres"¹⁴ estimate of 2825 mb and the Bradt-Peters^{15,16} estimate of 3169 mb.) The final fragment velocity and angular distribution was calculated from the primary distribution using the DFF code as modified by McGaughey and Morrissey.¹³ This code allows deexcitation of primary fragments by particle emission and fission and allows one to calculate the effects of these processes upon the fragment velocities and direction of motion. The resulting distributions for $39 \leq A \leq 49$ and $92 \leq A \leq 102$ are shown in Figure 1. The distributions predicted by the firestreak model and the two step model have qualitatively different shapes, especially for the light fragments where the firestreak model predicts a distribution peaking at intermediate angles. The data, unfortunately,

do not permit any definitive statements about the relative merits of the two sets of predicted distributions.

All of the distributions do show an elevated point at the most forward angle which, if taken at face value, would imply a shape of the deep spallation distributions that varies as $\sin^{-n}(\theta_L)$ where $n \approx 2 - 2.5$. It is important to note that at the same time the distributions described in this paper were measured, measurements were made of the fission fragment angular distributions from the interaction of 25.2 GeV ^{12}C with ^{238}U . These measurements showed isotropic fragment distributions, indicating no systematic effects in the experiment or data analysis leading to artificially elevated differential cross sections at forward angles. In all candor, we must state that we do not understand the process leading to these elevated differential cross sections at the most forward angles but we are convinced that because we do not observe such effects in the other systems we have studied using these techniques they are not artifacts.

This work was supported in part by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U. S. Department of Energy under contract DE-AC03-76SF00098.


References

*Present address; Department of Chemistry, Oregon State University,
Corvallis, Oregon 97331

1. S.B. Kaufman, E.P. Steinberg, and B.D. Wilkins, Phys. Rev. Lett. 41 1359 (1978).
2. S.B. Kaufman, E.P. Steinberg, B.D. Wilkins, and D.J. Henderson, Phys. Rev. C22, 1897 (1980).
3. W. Loveland, D.J. Morrissey, K. Aleklett, G.T. Seaborg, S.B. Kaufman, E.P. Steinberg, B.D. Wilkins, J.B. Cumming, P.E. Hausstein, and H.C. Hseuh, Phys. Rev. C23, 253 (1981).
4. W. Loveland, C. Luo, P.L. McGaughey, D.J. Morrissey, and G.T. Seaborg, Phys. Rev. C24 464 (1981).
5. G.D. Cole and N.T. Porile, Phys. Rev. C25, 244 (1982).
6. W.G. Meyer, H.H. Gutbrod, Ch. Lukner, and A. Sandoval, Phys. Rev. C22, 179 (1980).
7. S.B. Kaufman, M.S. Freedman, D.J. Henderson, E.P. Steinberg, B.D. Wilkins, A. Baden, H.H. Gutbrod, M.R. Maier, J. Peter, H.G. Ritter, H. Stelzer, A.I. Warwick, H.H. Wieman, and F. Weik, Phys. Rev. C26, 2694 (1982).
8. A.E. Warwick, H.H. Wieman, H.H. Gutbrod, M.R. Maier, J. Peter, H.G. Ritter, H. Stelzer, F. Weik, M. Freedman, D.J. Henderson, S.B. Kaufman, E.P. Steinberg, and B.D. Wilkins, Phys. Rev. C27, 1083 (1983).
9. H.H. Heckman, in Dynamics of Heavy-Ion Collisions, N. Cindro, R.A. Ricci, and W. Greiner, eds. (North-Holland, 1981) p. 339.
10. Y. Morita, W. Loveland, P.L. McGaughey, and G.T. Seaborg, Phys. Rev. C26, 511 (1982).
11. Li Wenxin (private communication, 1983). The recoil parameters measured in this work are in general agreement with the data of Ref. 2 but include values for the specific products studied in this work which were not discussed in reference 2.
12. N.T. Porile, S. Pandian, H. Klonk, C.R. Rudy and E.P. Steinberg, Phys. Rev. C19, 1832 (1979).
13. P.L. McGaughey, Lawrence Berkeley Laboratory Report LBL-15325 (November, 1982).
14. P.J. Karol, Phys. Rev. C11, 1203 (1975).
15. H.L. Bradt and B. Peters, Phys. Rev. 75, 1779 (1949).
16. H.H. Heckman, D.E. Greiner, P.J. Lindstrom, and H. Shue, Phys. Rev. C17, 1735 (1978).

Figure Captions

Figure 1. Fragment angular distributions (in the laboratory system) for the interaction of 25.2 GeV ^{12}C with ^{197}Au . Solid lines indicate predictions of the two step model, while dashed lines represent firestreak model predictions.



This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720