

# UC Santa Cruz

## UC Santa Cruz Previously Published Works

### Title

Multiple conserved regulatory domains promote Fezf2 expression in the developing cerebral cortex

### Permalink

<https://escholarship.org/uc/item/00h1h3cz>

### Journal

Neural Development, 9(1)

### ISSN

1749-8104

### Authors

Eckler, Matthew J

Larkin, Kathryn A

McKenna, William L

et al.

### Publication Date

2014-12-01


### DOI

10.1186/1749-8104-9-6

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



## Multiple conserved regulatory domains promote *Fezf2* expression in the developing cerebral cortex

Eckler *et al.*

RESEARCH ARTICLE

Open Access

# Multiple conserved regulatory domains promote *Fezf2* expression in the developing cerebral cortex

Matthew J Eckler<sup>1</sup>, Kathryn A Larkin<sup>1†</sup>, William L McKenna<sup>1†</sup>, Sol Katzman<sup>2</sup>, Chao Guo<sup>1</sup>, Robin Roque<sup>1</sup>, Axel Visel<sup>3,4</sup>, John L R Rubenstein<sup>5</sup> and Bin Chen<sup>1\*</sup>

## Abstract

**Background:** The genetic programs required for development of the cerebral cortex are under intense investigation. However, non-coding DNA elements that control the expression of developmentally important genes remain poorly defined. Here we investigate the regulation of *Fezf2*, a transcription factor that is necessary for the generation of deep-layer cortical projection neurons.

**Results:** Using a combination of chromatin immunoprecipitation followed by high throughput sequencing (ChIP-seq) we mapped the binding of four deep-layer-enriched transcription factors previously shown to be important for cortical development. Building upon this we characterized the activity of three regulatory regions around the *Fezf2* locus at multiple stages throughout corticogenesis. We identified a promoter that was sufficient for expression in the cerebral cortex, and enhancers that drove reporter gene expression in distinct forebrain domains, including progenitor cells and cortical projection neurons.

**Conclusions:** These results provide insight into the regulatory logic controlling *Fezf2* expression and further the understanding of how multiple non-coding regulatory domains can collaborate to control gene expression *in vivo*.

**Keywords:** *Fezf2*, Enhancer, Promoter, Cerebral cortex, Gene regulation, Transcription

## Background

Current estimates suggest that greater than half of evolutionarily conserved sequences in the human genome do not correspond to protein coding regions [1-4]. Transcriptional enhancers, non-coding DNA sequences that promote gene expression in a spatial and temporal manner, constitute a substantial portion of these regions [5]. Genome-wide association studies (GWAS) have identified sequence variations within putative enhancers that are associated with a wide range of neurological disorders such as autism, epilepsy, and schizophrenia [6-10]. However, the function of these DNA elements during normal brain development remains poorly understood [11].

The cerebral cortex contains six layers of projection neurons that are generated in a stereotyped manner such

that deep-layer neurons, layers 5 and 6 (L5 and L6), are born before upper-layer neurons [12]. It has previously been shown that the transcription factor *Fezf2* is necessary and sufficient for the generation of deep-layer subcerebral projection neurons (SCPNs) [13-16]. Collectively, these studies suggest that *Fezf2* functions high in a transcriptional hierarchy that regulates SCPN development and fate specification [14,17-19]. Despite the essential role of *Fezf2* in neural development, little is known about how its transcription is precisely regulated.

To investigate the regulatory mechanisms controlling *Fezf2* transcription, we utilized chromatin immunoprecipitation combined with high throughput DNA sequencing (ChIP-seq) to identify a transcription factor-binding signature around the *Fezf2* locus. Guided by our ChIP-seq data, we mapped a promoter that was sufficient for reporter gene expression in the cerebral cortex throughout cortical development, and investigated the activity of two putative *Fezf2* enhancers. We demonstrate that a

\* Correspondence: bchen@ucsc.edu

†Equal contributors

<sup>1</sup>Department of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, CA, USA

Full list of author information is available at the end of the article

downstream enhancer, 434, is sufficient to drive expression in cortical progenitor cells while a putative upstream enhancer, 1316, is strictly active in deep-layer neurons within the cortex. Taken together, these results provide insight into the developmental programs that promote *Fezf2* expression during development of the cerebral cortex.

## Results

### *Fezf2* is dynamically expressed during development of the cerebral cortex

Towards understanding how *Fezf2* transcription is regulated, we first examined its expression during cortical neurogenesis (Figure 1A-C'). Consistent with previous reports [20,21], at E12.5 *Fezf2* expression was detected in ventricular zone (VZ) progenitors and deep-layer cortical neurons (Figure 1A-A'). At E15.5 *Fezf2* expression began to decline in L6, but remained high in L5 (Figure 1B-B'). Expression at P0 was similar to E15.5 (Figure 1C-C'). Thus, at late stages of cortical development, expression of *Fezf2* in the cerebral cortex was restricted to three distinct domains: higher expression in L5 neurons, and lower expression in L6 neurons and progenitors (Figure 1C-C'). This dynamic spatial and temporal expression pattern suggests that *Fezf2* transcription is under the control of a complex regulatory program.

### Identification of putative regulatory regions around the *Fezf2* locus


Recent work has identified several transcription factors that are essential for cortical neuron fate specification and differentiation. Among these, SOX5 and TBR1 appear to directly regulate *Fezf2* transcription [22-24]. Using chromatin isolated from E15.5 mouse cortices, we mapped the binding of transcription factors expressed in L5 and L6 that were previously shown to be important for cortical development. These included FOXG1, NFIB, SOX5, and TBR1 [22-33]. We found that FOXG1, SOX5, and TBR1 bound to the evolutionarily conserved

region immediately upstream of *Fezf2* (Figure 2B), suggesting that this region may contain the *Fezf2* promoter. Interestingly, all of these factors bound to a previously identified enhancer downstream of *Fezf2*, enhancer 434 (also known as E4) (Figure 2B). In addition to the upstream region and enhancer 434, FOXG1 and TBR1 bound a region in the neighboring *Cadps* gene, near the 5' end of *Fezf2* (Figure 2C). Taken together, our ChIP-seq results indicate that multiple transcription factors that are expressed in deep-layer neurons and are critical for cortical development bind at or near the *Fezf2* locus.


To determine the significance of our ChIP-seq data we searched the VISTA Enhancer Browser (enhancer.lbl.gov) for sequences in and near *Fezf2* that may regulate its expression [34]. This approach uncovered several enhancers around *Fezf2*. These included an enhancer at the 5' side of *Fezf2*, enhancer 1316, and the previously studied enhancer at the 3' side, enhancer 434 (Figure 2A). Our ChIP-seq data demonstrated binding within or near these enhancers, suggesting they are likely important for regulating *Fezf2* transcription within the cerebral cortex (Figure 2B,C).

### The 2.7 kb promoter is active across all *Fezf2* expression domains

Previous work identified the 2.7 kb region upstream of and including the *Fezf2* start codon as sufficient to drive forebrain expression before the onset of cortical neurogenesis [35]. This region was bound by multiple transcription factors in our ChIP-seq dataset (Figure 2B). We generated stable transgenic lines expressing the LacZ reporter under the control of this 2.7 kb sequence to study its activity throughout cortical development. When the earliest-born cortical projection neurons were being generated at E11.5, we observed LacZ activity throughout the cortex that mimicked endogenous *Fezf2* expression (compare Figure 3A with Figure 1A). We observed  $\beta$ -gal expression in SOX2<sup>+</sup> cortical progenitor cells and in TBR1<sup>+</sup> and Tuj1<sup>+</sup> postmitotic neurons (Figure 3B-D).



**Figure 1** *Fezf2* is dynamically expressed during cortical development. *In situ* hybridization at E12.5 (A-A'), E15.5 (B-B'), and P0 (C-C') for *Fezf2* reveals its expression pattern during development. Ctx, cortex; LGE, lateral ganglionic eminence; LV, lateral ventricle; MGE, medial ganglionic eminence; PP, preplate; Str, striatum; VZ, ventricular zone. Scale bars: (A) 100  $\mu$ m, (A') 10  $\mu$ m, (B) 250  $\mu$ m, (B') 25  $\mu$ m, (C) 500  $\mu$ m, (C') 50  $\mu$ m.




We next examined 2.7 kb promoter activity at P0. Whole mount X-Gal staining revealed expression throughout the cerebral cortex in a high anterior-medial to low posterior-lateral pattern (Figure 3E). However, unlike endogenous *Fezf2*, which is expressed in the VZ and in deep cortical layers (L5 and L6), the 2.7 kb-LacZ transgene was active in all cortical layers (compare Figure 3F with Figure 1C). LacZ was expressed in SOX2<sup>+</sup> progenitor cells in the VZ and in the CTIP2<sup>+</sup> or SATB2<sup>+</sup> projection neurons (Figure 3G-I). LacZ expression at P12 was similar to that observed at P0 (Additional file 1). Taken together, these results indicate that the 2.7 kb promoter is active

throughout cortical development and is sufficient to drive transcription across all endogenous *Fezf2* expression domains (progenitor cells, L5 and L6 neurons). However, its expression in upper cortical layers indicates that additional *cis*-elements are required to properly coordinate *Fezf2* expression.

#### Enhancer 434 activity is strongest in cortical progenitor cells


Previous work has shown that enhancer 434 is highly conserved across vertebrate species and is necessary for *Fezf2* expression in the cortex [36]. Deletion of this



**Figure 3 Promoter activity of the 2.7 kb fragment during cortical development.** (A) X-Gal staining at E11.5 revealed reporter activity in the cerebral cortex, similar to endogenous *Fezf2* expression. (B-D) Immunohistochemistry with  $\beta$ -Gal antibody combined with SOX2 (B), Tuj1 (C), and TBR1 (D) antibodies. (E) Whole mount X-Gal staining of P0 brains demonstrating expression in the forebrain. (F) X-Gal staining of P0 brain sections. (G-I') Immunohistochemistry using  $\beta$ -Gal antibody combined with antibodies for SOX2 (G, G'), CTIP2 (H, H') and SATB2 (I, I'). Arrows represent co-expression of LacZ and the indicated markers. Arrowheads indicate a lack of co-expression. PP, preplate. Scale bars: (A) 200  $\mu$ m, (D) 50  $\mu$ m, (E) 250  $\mu$ m, (F) 500  $\mu$ m, (I) 100  $\mu$ m, (I') 50  $\mu$ m.

enhancer from either a bacterial artificial chromosome (BAC) containing a *Fezf2-EGFP* reporter or from its endogenous locus resulted in a loss of EGFP or *Fezf2* expression in the cerebral cortex [36]. However, it remains unknown whether enhancer 434 alone is sufficient to promote *Fezf2* expression in subcerebral projection neurons.

We generated stable transgenic lines expressing LacZ under the control of enhancer 434 and the hsp68 minimal promoter [37]. At E11.5,  $\beta$ -gal was expressed in SOX2<sup>+</sup> and PAX6<sup>+</sup> cortical progenitor cells (Figure 4A-B), including the metaphase cells that were labeled by PHH3 antibody at the ventricular surface (Figure 4C).



In addition, Tuj1<sup>+</sup> neurons located in the preplate expressed  $\beta$ -gal (Figure 4D). Whole mount staining of P0 brains revealed strong LacZ activity throughout multiple brain regions (Figure 4E). However, the majority of enhancer 434 activity was in the VZ of the cortex and basal ganglia (Figure 4F). Immunohistochemistry with antibodies for LacZ and the radial glial cell (RGC) marker BLBP demonstrated significant co-localization (Figure 4G). Additionally, some LacZ<sup>+</sup> cells in the VZ expressed the RGC marker PAX6 (Figure 4H). Some cells in the cortical plate expressed LacZ, however a minority of these LacZ<sup>+</sup> cells expressed the projection neuron markers SOX5 and SATB2 (Additional file 2). Thus, at P0 enhancer 434 activity is strongest in cortical progenitor cells. This strong LacZ signal observed in the VZ is consistent with the activity of enhancer 434 reported by the VISTA Enhancer Browser (enhancer.lbl.gov).

In adult mice, LacZ expression was observed in the cerebral cortex, white matter, hippocampus, and striatum (Figure 4I-L). Most of the LacZ<sup>+</sup> cells exhibited typical astrocyte-like morphology and expressed the astrocyte marker GFAP (Figure 4K, L). A minority of LacZ<sup>+</sup> cells in the cerebral cortex expressed the cortical projection neuron marker SATB2 (Figure 4J'). This shift in enhancer 434 activity suggests that it may have divergent functions during development *versus* in adult animals. Taken together, these data demonstrate that during cortical development enhancer 434 activity is strongest in progenitor cells, one of the endogenous expression domains of *Fezf2*.

#### Enhancer 1316 is active in deep-layer projection neurons in the cerebral cortex

We next investigated the activity of enhancer 1316. MultiZ alignment of this region showed strong vertebrate conservation, especially within the middle 500 bp (Additional file 3). We generated stable transgenic lines expressing LacZ under control of enhancer 1316 and the hsp68 minimal promoter. At E11.5,  $\beta$ -gal was present throughout the newly generated preplate (Figure 5A), in Tuj1<sup>+</sup> neurons (Figure 5D). LacZ<sup>+</sup> cells did not express the progenitor cell markers PAX6 or SOX2 (Figure 5B, C).

Whole mount staining at P0 revealed LacZ activity in deep layers of the cerebral cortex (Figure 5E). LacZ was expressed in the subplate and in some deep-layer neurons, but not in cortical progenitors or astrocytes (Figure 5F). Some LacZ<sup>+</sup> cells expressed CTIP2 and SATB2 (Figure 5H-I'). None of the LacZ<sup>+</sup> cells expressed SOX2 (Figure 5G, G'). Analysis at P12 revealed a similar LacZ expression pattern to that observed at P0 (Additional file 4). Taken together, these results indicate that within the cerebral cortex, enhancer 1316 is strictly active in deep-layer neurons.


#### Discussion

We set out to understand how expression of *Fezf2* is spatially and temporally controlled during development of the cerebral cortex. Previous work demonstrates that *Fezf2* transcription levels are critically important for specification of cortical projection neuron fates [13-17,22-24,31]. Our ChIP-seq data indicate that multiple transcription factors that are expressed in deep-layer neurons bind to evolutionarily conserved sequences around the *Fezf2* locus. The large number of transcription factors that bound enhancer 434 underscores its developmental importance. Recent deletion analysis demonstrated that this enhancer is critical for subcerebral projection neuron identity and connectivity [36]. In the current study, the majority of enhancer 434 activity during cortical neurogenesis was in progenitor cells, suggesting that this element also functions to promote *Fezf2* expression in the VZ, one of the endogenous *Fezf2* expression domains (Figure 6).

An alternative interpretation of this result is that different experimental strategies may account for the somewhat distinct conclusions regarding the activity of enhancer 434. In the study by Shim et al. [36], enhancer 434 was deleted from either a bacterial artificial chromosome containing a *Fezf2-EGFP* reporter or from its endogenous genomic locus. In both cases, deletion of enhancer 434 resulted in a loss of GFP or *Fezf2* expression, respectively, and failed extension of subcerebral projections. In the present study, we assayed the activity of enhancer 434 in isolation and found that it drove LacZ activity strongest in progenitor cells with lower levels of reporter activity in postmitotic neurons. These data indicate that when taken out of its endogenous locus, enhancer 434 is sufficient to drive reporter expression in the progenitor cells. It is important to note however that both the Shim et al. study [36] and the data presented here indicate a critical role for enhancer 434 in the regulation of *Fezf2* transcription.

Although the present study identifies conserved regulatory elements around the *Fezf2* locus, functional enhancer-promoter interactions cannot be established based upon transgenic reporter assays alone. Specifically, whether enhancer 1316 functionally interacts with the *Fezf2* promoter to regulate transcription remains unknown. Recently published high throughput chromosome conformation capture (Hi-C) suggests an interaction between enhancer 1316 and the *Fezf2* promoter [38]. However, this enhancer does not appear to be essential for *Fezf2* expression during development. Transgenic mice containing a modified BAC (RP23-141E17) that express EGFP or CRE from the *Fezf2* open reading frame have been generated. Although enhancer 1316 is not contained within this BAC, GFP, or CRE expression in these transgenic mice recapitulates endogenous *Fezf2* gene expression [36,39,40], suggesting that this enhancer is not essential for *Fezf2* expression during






**Figure 5** Enhancer 1316 is active in deep-layer cortical neurons. (A-D) Immunohistochemical analysis of E11.5 brain sections using antibodies for  $\beta$ -Gal (A),  $\beta$ -Gal and PAX6 (B),  $\beta$ -Gal and SOX2 (C), and  $\beta$ -Gal and Tuj1 (D). (E, F) X-Gal staining of P0 whole mount brains (E) and brain sections (F). (G-I) Immunohistochemistry using  $\beta$ -Gal antibody combined with antibodies for SOX2 (G), CTIP2 (H), and SATB2 (I). Arrows represent co-expression of LacZ and the indicated markers. Arrowheads indicate a lack of co-expression. Scale bars: (A) 200  $\mu$ m, (D) 50  $\mu$ m, (E) 250  $\mu$ m, (F) 500  $\mu$ m, (I) 100  $\mu$ m, (I') 25  $\mu$ m.

development. Thus, it is possible that additional enhancer (s) present on the RP23-141E17 BAC may function redundantly with enhancer 1316 to promote *Fezf2* expression in deep layer neurons.

Multiple *cis*-regulatory sequences around *Fezf2* exhibited distinct yet overlapping activities during cortical development. These results are similar to the regulation of the *eve* gene locus in *Drosophila* where multiple, partially

redundant enhancers control expression in individual stripes along the anterior-posterior axis of the embryo [41,42]. Ultimately, combined activity of these enhancers is required to drive the precise spatio-temporal expression of *eve*. Similar regulatory paradigms have also been uncovered in mammals through analysis of the interferon-beta [43] or *Fgf8* [44] loci. In our study, neither of the identified enhancers nor the 2.7 kb promoter, when assayed in



**Figure 6** A model for the regulation of *Fezf2* transcription in the cortex. The 2.7 kb promoter was sufficient to drive transcription across all endogenous *Fezf2* expression domains throughout development of the cerebral cortex. Enhancer 434 exhibited strongest activity in cortical progenitor cells and in contrast enhancer 1316 was strictly active in deep-layer postmitotic neurons.

isolation, were sufficient to drive LacZ reporter expression in an identical pattern to that of endogenous *Fezf2* expression. This suggests the presence of an additional enhancer (enhancers) that promotes high levels of expression specifically in L5 neurons. Moreover, the strong activity of the 2.7 kb promoter in upper cortical layers indicates that repressor sequences are necessary to restrict its activity to cortical progenitor cells and deep-layer neurons. Identification of these regulatory sequences combined with an increased understanding of the chromatin environment surrounding the *Fezf2* locus should help to shed light on the mechanisms that precisely regulate its transcription.

We recently reported that in addition to deep-layer cortical neurons, *Fezf2* is expressed in progenitor cells throughout cortical neurogenesis [39]. Phenotypic analysis of *Fezf2*<sup>-/-</sup> mice indicates that it is required for both the patterning and maintenance of forebrain progenitors [45], and the fate specification of deep-layer cortical projection neurons [13-16]. In *Fezf2*<sup>-/-</sup> mutant cortices, subcerebral projection neurons switch their identity to become callosal or corticothalamic projection neurons. However, it remains unclear whether *Fezf2* functions within cortical progenitor cells, postmitotic neurons or a combination of both in order to properly specify the fate of deep-layer cortical projection neurons. Identifying distinct promoter and enhancer sequences that regulate *Fezf2* transcription may help provide answers to this question.

## Conclusions

Given recent progress identifying putative mammalian enhancers [37], it is critical that we understand the spatial and temporal activities of these sequences in the regulation of individual genes, in order to appreciate the deleterious effects of mutations in non-coding sequences identified through GWAS. Focusing on *Fezf2*, a key regulator of cortical development and deep-layer projection neuron fate, our study suggests that multiple enhancer and promoter activities can coordinate gene expression during brain development. Recent work indicates that *Fezf2* is expressed in neocortical progenitors throughout development. Although mice lacking *Fezf2* exhibit developmental defects largely within deep-layer neurons (L5

and L6), its role in cortical progenitors *versus* postmitotic neurons remains unclear. Identifying the regulatory mechanisms that control *Fezf2* transcription will enable the functional dissection of its role within these distinct cellular populations. We propose that the combined activities of the 2.7 kb promoter, enhancer 434, and possibly enhancer 1316 constitute a *cis*-regulatory module that controls *Fezf2* in distinct domains of the cerebral cortex, ultimately leading to its dynamic expression during cortical development (Figure 6).

## Methods

### Animals

All experiments in this study were carried out in accordance with protocols approved by the IACUC at the University of California, Santa Cruz and were performed in accordance with institutional and federal guidelines. The day of vaginal plug detection was designated as embryonic day 0.5 (E0.5). The day of birth was designated as postnatal day 0 (P0). Transgenic mice were generated using standard protocols [46]. To generate stable transgenic lines, founders were outcrossed to CD1 wild-type females for a minimum of three generations before analysis. Number of founder lines generated: 2.7 kb-*LacZ* (2), enhancer 434-*LacZ* (3), enhancer 1316-*LacZ* (4).

### Plasmids

Generation of the 2.7 kb-*lacZ* plasmid was previously described [35]. For enhancer 434-*lacZ*, the genomic region was obtained from mouse genomic DNA by PCR using primers: ATCGCTCGAGCAGGCTGTAGGATGG GCAGCAGGAGTTTC and ATCGAAGCTTGTAACA AGTCAGGTGAGCAGGCGGTA. The product was then cloned into the hsp68-*lacZ* expression vector using Gateway cloning according to the manufacturer's protocol [11]. Enhancer 1316-*LacZ* was generated from human genomic DNA using primers AAACCACACAGCTGGTTTCC and TTTCCCGATAGATCGTCAGC and cloned into the hsp68-*lacZ* expression vector [37].

### Histology

Immunostaining and *in situ* hybridization were performed as previously described [47]. The *Fezf2* cRNA probe

corresponds to nucleotides 644 to 1,374 of mouse *Fezf2* (GenBank: NC\_000080). LacZ staining was performed according to standard protocols. Primary antibodies used: Chicken anti  $\beta$ -gal (Abcam, 1:500), Rabbit anti BLBP (Millipore, 1:500), Rat anti CTIP2 (Abcam, 1:1,000), Guinea Pig anti GFAP (Advanced Immuno Chemical, 1:100), Rabbit anti PAX6 (Covance, 1:100), PHH3 (Cell Signaling, 1:100), Rabbit anti SATB2 (Abcam, 1:1,000), Goat anti SOX2 (Santa Cruz Biotech, 1:500), Rabbit anti SOX5 (Abcam, 1:500), Rabbit anti TBR1 (Abcam, 1:1,000), Mouse anti Tuj1 (Covance, 1:1,000). Primary antibodies were detected using AlexaFluor-conjugated secondary antibodies (Invitrogen, 1:1,000). DNA was visualized with DAPI (1:50,000).

### Microscopy

Bright field and epifluorescence images were captured on an Olympus BX51 microscope using a Q Imaging Retiga EXj camera or a Keyence BZ-9000 microscope. Confocal images were captured on a Leica TCS SP5 confocal microscope. Images were processed using Adobe Photoshop CS5 to adjust brightness and contrast.

### ChIP-Seq

Chromatin immunoprecipitation was performed as previously described [24]. Antibodies used were Rabbit anti FOXG1 (Cell Signaling), Rabbit anti NFIB (Active Motif), Rabbit anti SOX5 (Abcam), and Rabbit anti TBR1 (Abcam). Sequencing libraries were generated using the Illumina TruSeq kit according to the manufacturer's protocol. Sequencing was performed on an Illumina HiSeq 2000 at the UCSC Genome Technology Center. Input DNA was sequenced as control. Sequencing reads were mapped to the mouse genome (mm9) using the Bowtie mapping algorithm. Non-overlapping reads and PCR duplicates were removed. Peaks were called using the MACS algorithm [48].

### Additional files

**Additional file 1: 2.7 kb promoter activity at P12. (A)** The activity of the 2.7 kb promoter at P12 was similar to that observed at P0, with expression throughout the cortex and the strongest activity in upper layers. **(B-C)** LacZ was expressed in some CTIP2<sup>+</sup> and TBR1<sup>+</sup> cells in deep layers. **(D, D')** However, the highest density of LacZ<sup>+</sup> cells was in upper layers cells that expressed SATB2. Panels B'-D' show amplified areas boxed in panels B-D, respectively. Arrows represent co-expression of LacZ and the indicated markers. Arrowheads indicate a lack of co-expression. Ctx, cortex; LV, lateral ventricle; Str, striatum. Scale bars: (a) 500  $\mu$ m, (d) 100  $\mu$ m, (d') 50  $\mu$ m.

**Additional file 2: Enhancer 434 activity at P0. (A-B')** LacZ expression was observed in a few post-mitotic neurons expressing SOX5 **(A, A')** or SATB2 **(B, B')**. A' and B' show amplified areas boxed in panels A and B. Arrows represent co-expression of LacZ and the indicated markers. Arrowheads indicate a lack of co-expression. Scale bars: (B) 100  $\mu$ m, (B') 50  $\mu$ m.

**Additional file 3: Vertebrate conservation of enhancer 1316.** The UCSC genome browser was used to perform Multiz alignment of enhancer 1316 from 30 vertebrates. Strong conservation was most

evident within the middle 500 bp (approximately) of this region. Basewise conservation is represented as greyscale darkness with higher conservation corresponding to darker values. Gap Annotation: single line, no bases in the aligned species; double line, aligning species has one or more un-alignable bases in the gap region; red coloring, aligning species has Ns in the gap region. Genomic Breaks: green square brackets, enclose shorter alignments consisting of DNA from one genomic context in the aligned species nested inside a larger chain of alignments from a different genomic context.

**Additional file 4: Activity of enhancer 1316 at P12. (A-D')** Expression at P12 mirrored P0. **(A)** X-Gal staining. **(B-D')** Immunohistochemistry analysis of the brain sections. Some LacZ positive cells in deep layers expressed CTIP2 **(B-B')**, SATB2 **(C-C')**, and TBR1 **(D-D')**. B'-D' show enlargement of areas boxed in panels B-D. Ctx, cortex; LV, lateral ventricle; Str, striatum. Scale bars: (A) 500  $\mu$ m, (D) 100  $\mu$ m, (D') 20  $\mu$ m.

### Abbreviations

BAC: Bacterial artificial chromosome;  $\beta$ -gal:  $\beta$ -galactosidase; ChIP-seq: Chromatin co-immunoprecipitation and high throughput DNA sequencing; E: Embryonic day; GWAS: Genome-wide association study; L5: Layer 5; L6: Layer 6; P: Postnatal day; PCR: Polymerase chain reaction; RGC: Radial glial cell; SCPN: Subcerebral projection neurons; VZ: Ventricular zone.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

MJE: conception and design, data collection and analysis, manuscript writing, and final approval of the manuscript. KAL: data collection and analysis, final approval of the manuscript. WLM: data collection and analysis, manuscript writing, and final approval of the manuscript. CG: conception and design, data collection, manuscript writing, and final approval of the manuscript. RR: data collection and final approval of the manuscript. SK: analyzing sequencing data and final approval of the manuscript. AV: generating the enhancer 1316-LacZ reporter construct, manuscript editing, and final approval of the manuscript. JLRR: generating the enhancer 1316-LacZ reporter construct, manuscript editing, and final approval of the manuscript. BC: conception and design, data analysis, manuscript writing, and final approval of the manuscript. All authors read and approved the final manuscript.

### Acknowledgments

We are grateful to Masahiko Hibi (Nagoya University) for providing the 2.7 kb-LacZ reporter construct and for valuable insight during the completion of these experiments. We thank Armen Shamaman and the UCSC Transgenic Facility for generating transgenic mice. We are grateful to the UCSC Genome Technology Center for performing DNA sequencing. Ben Abrams and the UCSC Microscopy center provided support with image acquisition. We thank Drs. Lindsay Hinck, David Feldheim, Jennifer Betancourt, and Muriel Kmet for critical review of the manuscript. Members of the Chen lab and Drs. Lindsay Hinck, David Feldheim, and Susan McConnell provided valuable discussion during the completion of these experiments. This work was funded by grants from California Institute of Regenerative Medicine (RN1-00530-01) to BC, the NIMH (R01-MH094589) to BC, the NHGRI (R01HG003988) to AV, the NINDS (R01NS062859A) to AV and the NINDS (NS34661) to JLRR.

### Author details

<sup>1</sup>Department of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, CA, USA. <sup>2</sup>Center for Biomolecular Science and Engineering, University of California, Santa Cruz, CA, USA. <sup>3</sup>Genomics Division, MS 84-171, Lawrence Berkeley National Laboratory, Berkeley, CA, USA. <sup>4</sup>U.S. Department of Energy Joint Genome Institute, Walnut Creek, CA, USA. <sup>5</sup>Nina Ireland Laboratory of Developmental Neurobiology, Department of Psychiatry, University of California, San Francisco, CA, USA.

Received: 3 December 2013 Accepted: 19 February 2014

Published: 12 March 2014

## References

1. Mouse Genome Sequencing Consortium, Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, Agarwala R, Ainscough R, Alexandersson M, An P, Antonarakis SE, Attwood J, Baertsch R, Bailey J, Barlow K, Beck S, Berry E, Birren B, Bloom T, Bork P, Botcherby M, Bray N, Brent MR, Brown DG, Brown SD, Bult C, Burton J, Butler J, Campbell RD, et al: **Initial sequencing and comparative analysis of the mouse genome.** *Nature* 2002, **420**:520–562.
2. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, Clawson H, Spieth J, Hillier LW, Richards S, Weinstock GM, Wilson RK, Gibbs RA, Kent WJ, Miller W, Haussler D: **Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes.** *Genome Res* 2005, **15**:1034–1050.
3. Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, Yandell M, Evans CA, Holt RA, Gocayne JD, Amanatides P, Ballew RM, Huson DH, Wortman JR, Zhang Q, Kodira CD, Zheng XH, Chen L, Skupski M, Subramaniam G, Thomas PD, Zhang J, Gabor Miklos GL, Nelson C, Broder S, Clark AG, Nadeau J, McKusick VA, Zinder N, et al: **The sequence of the human genome.** *Science* 2001, **291**:1304–1351.
4. Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K, Doyle M, FitzHugh W, Funke R, Gage D, Harris K, Heaford A, Howland J, Kann L, Lehoczky J, LeVine R, McEwan P, McKernan K, Meldrim J, Mesirov JP, Miranda C, Morris W, Naylor J, Raymond C, Rosetti M, Santos R, Sheridan A, Sougnez C, et al: **Initial sequencing and analysis of the human genome.** *Nature* 2001, **409**:860–921.
5. Williamson I, Hill RE, Bickmore WA: **Enhancers: from developmental genetics to the genetics of common human disease.** *Dev Cell* 2011, **21**:17–19.
6. Genomes Project C: **A map of human genome variation from population-scale sequencing.** *Nature* 2010, **467**:1061–1073.
7. International Schizophrenia Consortium: **Rare chromosomal deletions and duplications increase risk of schizophrenia.** *Nature* 2008, **455**:237–241.
8. Hamilton SP, Woo JM, Carlson EJ, Ghanem N, Ekker M, Rubenstein JL: **Analysis of four DLX homeobox genes in autistic probands.** *BMC Genetics* 2005, **6**:52.
9. Sebat J, Lakshmi B, Malhotra D, Troge J, Lese-Martin C, Walsh T, Yamrom B, Yoon S, Krasnitz A, Kendall J, Leotta A, Pai D, Zhang R, Lee YH, Hicks J, Spence SJ, Lee AT, Puura K, Lehtimäki T, Ledbetter D, Gregersen PK, Bregman J, Sutcliffe JS, Jobanputra V, Chung W, Warburton D, King MC, Skuse D, Geschwind DH, Gilliam TC, et al: **Strong association of de novo copy number mutations with autism.** *Science* 2007, **316**:445–449.
10. Wray GA: **The evolutionary significance of cis-regulatory mutations.** *Nat Rev Genet* 2007, **8**:206–216.
11. Visel A, Blow MJ, Li Z, Zhang T, Akiyama JA, Holt A, Plajzer-Frick I, Shoukry M, Wright C, Chen F, Afzal V, Ren B, Rubin EM, Pennacchio LA: **ChIP-seq accurately predicts tissue-specific activity of enhancers.** *Nature* 2009, **457**:854–858.
12. Kwan KY, Sestan N, Anton ES: **Transcriptional co-regulation of neuronal migration and laminar identity in the neocortex.** *Development* 2012, **139**:1535–1546.
13. Chen B, Schaevitz LR, McConnell SK: **Fez1 regulates the differentiation and axon targeting of layer 5 subcortical projection neurons in cerebral cortex.** *Proc Natl Acad Sci U S A* 2005, **102**:17184–17189.
14. Chen B, Wang SS, Hattox AM, Rayburn H, Nelson SB, McConnell SK: **The Fezf2-Ctip2 genetic pathway regulates the fate choice of subcortical projection neurons in the developing cerebral cortex.** *Proc Natl Acad Sci U S A* 2008, **105**:11382–11387.
15. Chen JG, Rasin MR, Kwan KY, Sestan N: **Zfp312 is required for subcortical axonal projections and dendritic morphology of deep-layer pyramidal neurons of the cerebral cortex.** *Proc Natl Acad Sci U S A* 2005, **102**:17792–17797.
16. Molyneaux BJ, Arlotta P, Hirata T, Hibi M, Macklis JD: **Fez1 is required for the birth and specification of corticospinal motor neurons.** *Neuron* 2005, **47**:817–831.
17. Rouaux C, Arlotta P: **Fezf2 directs the differentiation of corticofugal neurons from striatal progenitors in vivo.** *Nat Neurosci* 2010, **13**:1345–1347.
18. De la Rossa A, Bellone C, Golding B, Vitali I, Moss J, Toni N, Luscher C, Jabaudon D: **In vivo reprogramming of circuit connectivity in postmitotic neocortical neurons.** *Nat Neurosci* 2013, **16**:193–200.
19. Rouaux C, Arlotta P: **Direct lineage reprogramming of post-mitotic callosal neurons into corticofugal neurons in vivo.** *Nat Cell Biol* 2012, **15**:214–221.
20. Hirata T, Suda Y, Nakao K, Narimatsu M, Hirano T, Hibi M: **Zinc finger gene fez-like functions in the formation of subplate neurons and thalamocortical axons.** *Dev Dyn* 2004, **230**:546–556.
21. Inoue K, Terashima T, Nishikawa T, Takumi T: **Fez1 is layer-specifically expressed in the adult mouse neocortex.** *Eur J Neurosci* 2004, **20**:2909–2916.
22. Han W, Kwan KY, Shim S, Lam MM, Shin Y, Xu X, Zhu Y, Li M, Sestan N: **TBR1 directly represses Fezf2 to control the laminar origin and development of the corticospinal tract.** *Proc Natl Acad Sci U S A* 2011, **108**:3041–3046.
23. Kwan KY, Lam MM, Krsnik Z, Kawasawa YI, Lefebvre V, Sestan N: **SOX5 postmitotically regulates migration, postmigratory differentiation, and projections of subplate and deep-layer neocortical neurons.** *Proc Natl Acad Sci U S A* 2008, **105**:16021–16026.
24. McKenna WL, Betancourt J, Larkin KA, Abrams B, Guo C, Rubenstein JL, Chen B: **Tbr1 and Fezf2 regulate alternate corticofugal neuronal identities during neocortical development.** *J Neurosci* 2011, **31**:549–564.
25. Alamo EA, Chirivella L, Dautzenberg M, Dobrova G, Farinas I, Grosschedl R, McConnell SK: **Satb2 regulates callosal projection neuron identity in the developing cerebral cortex.** *Neuron* 2008, **57**:364–377.
26. Azim E, Jabaudon D, Fame RM, Macklis JD: **SOX6 controls dorsal progenitor identity and interneuron diversity during neocortical development.** *Nat Neurosci* 2009, **12**:1238–1247.
27. Baranek C, Ditttrich M, Parthasarathy S, Bonnon CG, Britanova O, Lanshakov D, Boukhtouche F, Sommer JE, Colmenares C, Tarabykin V, Atanasiuk S: **Protooncogene Ski cooperates with the chromatin-remodeling factor Satb2 in specifying callosal neurons.** *Proc Natl Acad Sci U S A* 2012, **109**:3546–3551.
28. Batista-Brito R, Rossignol E, Hjerling-Leffler J, Denaxa M, Wegner M, Lefebvre V, Pachnis V, Fishell G: **The cell-intrinsic requirement of Sox6 for cortical interneuron development.** *Neuron* 2009, **63**:466–481.
29. Britanova O, de Juan RC, Cheung A, Kwan KY, Schwark M, Gyorgy A, Vogel T, Akopov S, Mitkovski M, Agoston D, Sestan N, Molnar Z, Tarabykin V: **Satb2 is a postmitotic determinant for upper-layer neuron specification in the neocortex.** *Neuron* 2008, **57**:378–392.
30. Hanashima C, Li SC, Shen L, Lai E, Fishell G: **Foxg1 suppresses early cortical cell fate.** *Science* 2004, **303**:56–59.
31. Lai T, Jabaudon D, Molyneaux BJ, Azim E, Arlotta P, Menezes JR, Macklis JD: **SOX5 controls the sequential generation of distinct corticofugal neuron subtypes.** *Neuron* 2008, **57**:232–247.
32. Miyoshi G, Fishell G: **Dynamic FoxG1 expression coordinates the integration of multipolar pyramidal neuron precursors into the cortical plate.** *Neuron* 2012, **74**:1045–1058.
33. Piper M, Moldrich RX, Lindwall C, Little E, Barry G, Mason S, Sunn N, Kurniawan ND, Gronostajski RM, Richards LJ: **Multiple non-cell-autonomous defects underlie neocortical callosal dysgenesis in Nfib-deficient mice.** *Neural Dev* 2009, **4**:43.
34. Visel A, Minovitsky S, Dubchak I, Pennacchio LA: **VISTA Enhancer Browser—a database of tissue-specific human enhancers.** *Nucleic Acids Res* 2007, **35**:D88–D92.
35. Hirata T, Nakazawa M, Muraoka O, Nakayama R, Suda Y, Hibi M: **Zinc-finger genes Fez and Fez-like function in the establishment of diencephalon subdivisions.** *Development* 2006, **133**:3993–4004.
36. Shim S, Kwan KY, Li M, Lefebvre V, Sestan N: **Cis-regulatory control of corticospinal system development and evolution.** *Nature* 2012, **486**:74–79.
37. Visel A, Taher L, Girgis H, May D, Golonzhka O, Hoch RV, McKinsey GL, Pattabiraman K, Silberberg SN, Blow MJ, Hansen DV, Nord AS, Akiyama JA, Holt A, Hosseini R, Phouanavong S, Plajzer-Frick I, Shoukry M, Afzal M, Kaplan T, Kriegstein AR, Rubin EM, Ovcharenko I, Pennacchio LA, Rubenstein JL: **A high-resolution enhancer atlas of the developing telencephalon.** *Cell* 2013, **152**:895–908.
38. Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B: **Topological domains in mammalian genomes identified by analysis of chromatin interactions.** *Nature* 2012, **485**:376–380.
39. Guo C, Eckler MJ, McKenna WL, McKinsey GL, Rubenstein JL, Chen B: **Fezf2 expression identifies a multipotent progenitor for neocortical projection neurons, astrocytes, and oligodendrocytes.** *Neuron* 2013, **80**:1167–1174.
40. Gong S, Zheng C, Doughty ML, Losos K, Didkovsky N, Schambra UB, Nowak NJ, Joyner A, Leblanc G, Hatten ME, Heintz N: **A gene expression atlas of the central nervous system based on bacterial artificial chromosomes.** *Nature* 2003, **425**:917–925.

41. Andrioli LP, Vasisht V, Theodosopoulou E, Oberstein A, Small S: **Anterior repression of a Drosophila stripe enhancer requires three position-specific mechanisms.** *Development* 2002, **129**:4931–4940.
42. Perry MW, Boettiger AN, Levine M: **Multiple enhancers ensure precision of gap gene-expression patterns in the Drosophila embryo.** *Proc Natl Acad Sci U S A* 2011, **108**:13570–13575.
43. Thanos D, Maniatis T: **Virus induction of human IFN beta gene expression requires the assembly of an enhanceosome.** *Cell* 1995, **83**:1091–1100.
44. Marinic M, Aktas T, Ruf S, Spitz F: **An integrated holo-enhancer unit defines tissue and gene specificity of the Fgf8 regulatory landscape.** *Dev Cell* 2013, **24**:530–542.
45. Shimizu T, Nakazawa M, Kani S, Bae YK, Shimizu T, Kageyama R, Hibi M: **Zinc finger genes Fezf1 and Fezf2 control neuronal differentiation by repressing Hes5 expression in the forebrain.** *Development* 2010, **137**:1875–1885.
46. Nagy A: *Manipulating the mouse embryo: a laboratory manual*. 3rd edition. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 2003.
47. Eckler MJ, McKenna WL, Taghvaei S, McConnell SK, Chen B: **Fezf1 and Fezf2 are required for olfactory development and sensory neuron identity.** *J Comp Neurol* 2011, **519**:1829–1846.
48. Zhang Y, Liu T, Meyer CA, Eickhout J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, Liu XS: **Model-based analysis of CHIP-Seq (MACS).** *Genome Biol* 2008, **9**:R137.

doi:10.1186/1749-8104-9-6

**Cite this article as:** Eckler *et al.*: Multiple conserved regulatory domains promote *Fezf2* expression in the developing cerebral cortex. *Neural Development* 2014 **9**:6.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

