

Lawrence Berkeley National Laboratory

Recent Work

Title

AN EXAMPLE OF Ku_4 DECAY IN EMULSION

Permalink

<https://escholarship.org/uc/item/01q154c9>

Authors

Greiner, Douglas E.

Osborne, W.Z.

Barkas, Walter H.

Publication Date

1964-07-23

University of California

**Ernest O. Lawrence
Radiation Laboratory**

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

AN EXAMPLE OF K_{μ^4} DECAY IN EMULSION

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

AN EXAMPLE OF $K_{\mu 4}$ DECAY IN EMULSION

Douglas E. Greiner, W. Z. Osborne and Walter H. Barkas

July 23, 1964

AN EXAMPLE OF $K_{\mu 4}$ DECAY IN EMULSION

Douglas E. Greiner, W. Z. Osborne and Walter H. Barkas

Lawrence Radiation Laboratory
University of California
Berkeley, California

July 23, 1964

We have accumulated ≈ 4000 examples of three-prong K^+ decay by scanning emulsion exposed to a stopping K^+ beam from the Bevatron. The direction of each prong was measured with a digitized-coordinate microscope and an IBM 7094 was used to calculate the deviation from coplanarity of the prongs. The events which violated coplanarity were then examined in greater detail.

This process isolated an event which we believe represents the decay mode $K^+ \rightarrow \pi^+ \pi^- \mu^+ \nu_{\mu}$. A projection drawing of it is shown in Fig. 1. The K^+ decayed at rest as well as could be determined. Each prong was traced to its stopping point, and all follow-throughs were verified by an independent observer. The range of each prong was measured with our digitized-coordinate microscope, and the associated energies and momenta were computed from the range-energy relation with an IBM 7094. The program used includes the correction of ranges to standard emulsion density. The observations and measurements made are listed in Table I.

Table I. Data For $K_{\mu 4}$ event.

Prong	Direction Cosines			Range (mm)	Terminal Behavior	Particle	Momentum (MeV/c)
	x	y	z				
1	0.369 ± 0.038	-0.737 ± 0.041	-0.567 ± 0.060	0.630	one-prong star	π^-	36.78 ± 0.37
2	-0.694 ± 0.014	0.101 ± 0.011	-0.713 ± 0.014	9.952	$\mu^+ \rightarrow e^+$ decay	μ^+	69.13 ± 0.75
3	-0.511 ± 0.022	0.282 ± 0.019	0.812 ± 0.016	4.581	$\pi^+ \rightarrow \mu^+$ decay	π^+	65.85 ± 0.62

The resultant momentum of the three charged products is 70.1 ± 2.4 MeV/c and the missing energy is 68.94 ± 0.56 MeV. The measurements are therefore consistent with a four-body decay wherein the neutral product has vanishing rest mass. Careful examination of track 2 has revealed no sudden changes of grain density or deflections which could be attributed to $\pi \rightarrow \mu$ decay in flight. We therefore have no reason to suspect that the event is not a genuine example of a new decay mode, $K_{\mu 4}$.

We have, however, eliminated the possibility of two conceivable alternative interpretations. One of these consists of an ordinary τ decay with π - μ decay at a point so near the K-decay vertex as to be unobservable. The experimental value of the quantity $[(M_K - E_1 - E_3)^2 - (\vec{P}_1 + \vec{P}_3)^2]^{1/2}$ is $191.13 \pm .68$ MeV. This is clearly inconsistent with the pion rest mass. The second alternative consists of a radiative τ decay ($K^+ \rightarrow \pi^+ \pi^+ \pi^- \gamma$) with subsequent π - μ decay in flight. The measured directions of tracks 1, 2, and 3 together with the measured momenta of particles 1 and 3 are sufficient to determine completely the assumed decay. The maximum length of track 2 obtainable from this scheme is 7.08mm. This is inconsistent with the measured value of 9.95mm.

Although some events remain to be analyzed, we have already seen two examples of the known^{1,2} $K^+ \rightarrow \pi^+ \pi^- e^+ \nu_e$ decay mode. We should expect to see three such decays in our sample from the K_{e4}^+/τ^+ branching ratio³ of $(7.8 \pm 1.6) \times 10^{-4}$. We estimate that our efficiency for finding the $K_{\mu 4}$ mode is almost 100% so the $K_{\mu 4}^+/\tau^+$ branching ratio is $\approx 1/4000$, based on the one event.

¹ E. L. Koller, S. Taylor, T. Huetter, and P. Stamer, Phys. Rev. Letters 9, 328 (1962).

² R. W. Birge, R. P. Ely, G. Gidal, G. E. Kalmus, A. Kernan, W. M. Powell, V. Camerini, W. F. Fry, J. Gaidos, R. H. March, and S. Natali, Phys. Rev. Letters 11, 35 (1963).

³ R. W. Birge, R. P. Ely, G. Gidal, G. E. Kalmus, A. Kernan, W. M. Powell, V. Camerini, W. F. Fry, J. Gaidos, D. Murphree, and C. T. Murphy, University of California Lawrence Radiation Laboratory Report UCRL-11549 (unpublished).

In all $K_{\ell 4}^+$ decays observed to date, the rule $\Delta S = \Delta Q$ is satisfied. However, one of us (W. Z. O.) has pointed out that these decays do not necessarily provide a good test of the rule if the $T = 0$ dipion resonance, σ , postulated by Brown and Singer⁴ does in fact exist. The effect that this resonance would have upon various distributions in $K_{\ell 4}$ decay has been considered by Brown and Faier.⁵ It is at least conceivable that a large enhancement of the $\Delta S = \Delta Q$ modes with respect to the $\Delta S = -\Delta Q$ modes could ensue. It therefore seems hazardous to base far-reaching conclusions upon the observation of even a large excess of the $\Delta S = \Delta Q$ modes before resolution of the presently uncertain^{3,4,6,7,8} experimental situation relevant to the σ -resonance.

We wish to acknowledge the excellent work of our scanning and measuring staff. Mrs. Marilyn Mollin found the event, Mr. Robert Trankle performed the original follow-throughs, and the digitized-coordinate microscope measurements were carried out by Mr. James Webber and Mr. Peter Lindstrom.

⁴ L. M. Brown and P. Singer, Phys. Rev. 133, 812 (1964).

⁵ L. M. Brown and H. Faier, Phys. Rev. Letters 12, 514 (1964).

⁶ F. S. Crawford, Jr., R. A. Grossman, L. J. Lloyd, L. R. Price, and E. C. Fowler, Phys. Rev. Letters 11, 564 (1963).

⁷ R. Del Fabbro, M. De Pretis, R. Jones, G. Marini, A. Odian, G. Stoppini, and L. Tau, Phys. Rev. Letters 12, 674 (1964).

⁸ G. E. Kalmus, A. Kernan, R. T. Pu, W. M. Powell and R. Dowd, Phys. Rev. Letters 13, 99 (1964).



FIGURE CAPTIONS

Fig. 1. Projection drawing of $K_{\mu 4}$ event.

