

UC Berkeley

UC Berkeley Previously Published Works

Title

Measurement of the direct CP asymmetry in $b \rightarrow s$ gamma Decays.

Permalink

<https://escholarship.org/uc/item/01z2k9d8>

Journal

Physical review letters, 93(2)

ISSN

0031-9007

Authors

Aubert, B
Barate, R
Boutigny, D
[et al.](#)

Publication Date

2004-07-01

DOI

10.1103/physrevlett.93.021804

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Measurement of the Direct CP Asymmetry in $b \rightarrow s\gamma$ Decays

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ C. LeClerc,⁵ G. Lynch,⁵ A. M. Merchant,⁵ L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ W. A. Wenzel,⁵ K. Ford,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ S. E. Morgan,⁶ A. T. Watson,⁶ M. Fritsch,⁷ K. Goetzen,⁷ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ M. Steinke,⁷ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸ M. P. Kelly,⁸ T. E. Latham,⁸ F. F. Wilson,⁸ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ N. S. Knecht,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ A. Khan,¹⁰ P. Kyberd,¹⁰ L. Teodorescu,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹² C. Buchanan,¹³ B. L. Hartfiel,¹³ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵ E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ B. Dahmes,¹⁶ S. L. Levy,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. A. Heusch,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ R. E. Schmitz,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ P. Spradlin,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸ A. Dvoretzki,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸ S. Jayatilake,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰ S. Chen,²⁰ W. T. Ford,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. G. Smith,²⁰ J. Zhang,²⁰ L. Zhang,²⁰ A. Chen,²¹ J. L. Harton,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ Q. L. Zeng,²¹ D. Altenburg,²² T. Brandt,²² J. Brose,²² T. Colberg,²² M. Dickopp,²² E. Feltresi,²² A. Hauke,²² H. M. Lacker,²² E. Maly,²² R. Müller-Pfefferkorn,²² R. Nogowski,²² S. Otto,²² A. Petzold,²² J. Schubert,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² J. E. Sundermann,²² D. Bernard,²³ G. R. Bonneaud,²³ F. Brochard,²³ P. Grenier,²³ S. Schrenk,²³ Ch. Thiebaux,²³ G. Vasileiadis,²³ M. Verderi,²³ D. J. Bard,²⁴ P. J. Clark,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ Y. Xie,²⁴ M. Andreotti,²⁵ V. Azzolini,²⁵ D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,²⁵ G. Cibinetto,²⁵ E. Luppi,²⁵ M. Negrini,²⁵ L. Piemontese,²⁵ A. Sarti,²⁵ E. Treadwell,²⁶ R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ M. Piccolo,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Capra,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ S. Bailey,²⁹ G. Brandenburg,²⁹ M. Morii,²⁹ E. Won,²⁹ R. S. Dubitzky,³⁰ U. Langenegger,³⁰ W. Bhimji,³¹ D. A. Bowerman,³¹ P. D. Dauncey,³¹ U. Egede,³¹ J. R. Gaillard,³¹ G. W. Morton,³¹ J. A. Nash,³¹ G. P. Taylor,³¹ G. J. Grenier,³² U. Mallik,³² J. Cochran,³³ H. B. Crawley,³³ J. Lamsa,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ J. Yi,³³ M. Davier,³⁴ G. Grosdidier,³⁴ A. Höcker,³⁴ S. Laplace,³⁴ F. Le Diberder,³⁴ V. Lepeltier,³⁴ A. M. Lutz,³⁴ T. C. Petersen,³⁴ S. Plaszczynski,³⁴ M. H. Schune,³⁴ L. Tantot,³⁴ G. Wormser,³⁴ C. H. Cheng,³⁵ D. J. Lange,³⁵ M. C. Simani,³⁵ D. M. Wright,³⁵ A. J. Bevan,³⁶ J. P. Coleman,³⁶ J. R. Fry,³⁶ E. Gabathuler,³⁶ R. Gamet,³⁶ R. J. Parry,³⁶ D. J. Payne,³⁶ R. J. Sloane,³⁶ C. Touramanis,³⁶ J. J. Back,³⁷ C. M. Cormack,³⁷ P. F. Harrison,^{37,*} G. B. Mohanty,³⁷ C. L. Brown,³⁸ G. Cowan,³⁸ R. L. Flack,³⁸ H. U. Flaecher,³⁸ M. G. Green,³⁸ C. E. Marker,³⁸ T. R. McMahon,³⁸ S. Ricciardi,³⁸ F. Salvatore,³⁸ G. Vaitsas,³⁸ M. A. Winter,³⁸ D. Brown,³⁹ C. L. Davis,³⁹ J. Allison,⁴⁰ N. R. Barlow,⁴⁰ R. J. Barlow,⁴⁰ P. A. Hart,⁴⁰ M. C. Hodgkinson,⁴⁰ G. D. Lafferty,⁴⁰ A. J. Lyon,⁴⁰ J. C. Williams,⁴⁰ A. Farbin,⁴¹ W. D. Hulsbergen,⁴¹ A. Jawahery,⁴¹ D. Kovalskyi,⁴¹ C. K. Lae,⁴¹ V. Lillard,⁴¹ D. A. Roberts,⁴¹ G. Blaylock,⁴² C. Dallapiccola,⁴² K. T. Flood,⁴² S. S. Hertzbach,⁴² R. Kofler,⁴² V. B. Koptchev,⁴² T. B. Moore,⁴² S. Saremi,⁴² H. Staengle,⁴² S. Willocq,⁴² R. Cowan,⁴³ G. Sciolla,⁴³ F. Taylor,⁴³ R. K. Yamamoto,⁴³ D. J. J. Mangeol,⁴⁴ P. M. Patel,⁴⁴ S. H. Robertson,⁴⁴ A. Lazzaro,⁴⁵ F. Palombo,⁴⁵ J. M. Bauer,⁴⁶ L. Cremaldi,⁴⁶ V. Eschenburg,⁴⁶ R. Godang,⁴⁶ R. Kroeger,⁴⁶ J. Reidy,⁴⁶ D. A. Sanders,⁴⁶ D. J. Summers,⁴⁶ H. W. Zhao,⁴⁶ S. Brunet,⁴⁷ D. Côté,⁴⁷ P. Taras,⁴⁷ H. Nicholson,⁴⁸ N. Cavallo,⁴⁹ F. Fabozzi,^{49,†} C. Gatto,⁴⁹ L. Lista,⁴⁹ D. Monorchio,⁴⁹ P. Paolucci,⁴⁹ D. Piccolo,⁴⁹ C. Sciacca,⁴⁹ M. Baak,⁵⁰ H. Bulten,⁵⁰ G. Raven,⁵⁰ L. Wilden,⁵⁰ C. P. Jessop,⁵¹ J. M. LoSecco,⁵¹ T. A. Gabriel,⁵² T. Allmendinger,⁵³ B. Brau,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵

M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ G. Tiozzo,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Ocariz,⁵⁶ M. Pivk,⁵⁶ L. Roos,⁵⁶ S. T'Jampens,⁵⁶ G. Therin,⁵⁶ P. F. Manfredi,⁵⁷ V. Re,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ F. Anulli,^{27,59} M. Biasini,⁵⁹ I. M. Peruzzi,^{27,59} M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ M. Bondioli,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ V. Del Gamba,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ F. Martinez-Vidal,^{60,‡} M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ F. Sandrelli,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ K. Paick,⁶¹ D. E. Wagoner,⁶¹ N. Danielson,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² V. Miftakov,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² F. Bellini,⁶³ G. Cavoto,^{62,63} R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ M. Pierini,⁶³ G. Piredda,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ S. Christ,⁶⁴ G. Wagner,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ N. I. Geddes,⁶⁵ G. P. Gopal,⁶⁵ E. O. Olaiya,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ P.-F. Giraud,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Langer,⁶⁶ M. Legendre,⁶⁶ G. W. London,⁶⁶ B. Mayer,⁶⁶ G. Schott,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ F. X. Yumiceva,⁶⁷ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ G. De Nardo,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ E. E. Elsen,⁶⁸ S. Fan,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ C. Hast,⁶⁸ T. Hryn'ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ S. Petrak,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ G. Simi,⁶⁸ A. Snyder,⁶⁸ A. Soha,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ J. Va'vra,⁶⁸ S. R. Wagner,⁶⁸ M. Weaver,⁶⁸ A. J. R. Weinstein,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ T. I. Meyer,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ M. Saleem,⁷⁰ F. R. Wappler,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² H. Kim,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ C. Borean,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ P. Poropat,^{75,§} L. Vitale,⁷⁵ G. Vuagnin,⁷⁵ R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ H. R. Band,⁷⁸ S. Dasu,⁷⁸ M. Datta,⁷⁸ A. M. Eichenbaum,⁷⁸ M. Graham,⁷⁸ J. J. Hollar,⁷⁸ J. R. Johnson,⁷⁸ P. E. Kutter,⁷⁸ H. Li,⁷⁸ R. Liu,⁷⁸ F. Di Lodovico,⁷⁸ A. Mihalyi,⁷⁸ A. K. Mohapatra,⁷⁸ Y. Pan,⁷⁸ R. Prepost,⁷⁸ A. E. Rubin,⁷⁸ S. J. Sekula,⁷⁸ P. Tan,⁷⁸ J. H. von Wimmersperg-Toeller,⁷⁸ J. Wu,⁷⁸ S. L. Wu,⁷⁸ Z. Yu,⁷⁸ and H. Neal⁷⁹

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at Riverside, Riverside, California 92521, USA

¹⁵University of California at San Diego, La Jolla, California 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁸California Institute of Technology, Pasadena, California 91125, USA

¹⁹University of Cincinnati, Cincinnati, Ohio 45221, USA

²⁰University of Colorado, Boulder, Colorado 80309, USA

²¹Colorado State University, Fort Collins, Colorado 80523, USA

²²Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²³Ecole Polytechnique, LLR, F-91128 Palaiseau, France

- ²⁴University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
- ²⁵Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- ²⁶Florida A&M University, Tallahassee, Florida 32307, USA
- ²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ²⁹Harvard University, Cambridge, Massachusetts 02138, USA
- ³⁰Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
- ³¹Imperial College London, London, SW7 2AZ, United Kingdom
- ³²University of Iowa, Iowa City, Iowa 52242, USA
- ³³Iowa State University, Ames, Iowa 50011-3160, USA
- ³⁴Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
- ³⁵Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- ³⁶University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³⁷Queen Mary, University of London, E1 4NS, United Kingdom
- ³⁸University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ³⁹University of Louisville, Louisville, Kentucky 40292, USA
- ⁴⁰University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴¹University of Maryland, College Park, Maryland 20742, USA
- ⁴²University of Massachusetts, Amherst, Massachusetts 01003, USA
- ⁴³Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- ⁴⁴McGill University, Montréal, Quebec, Canada H3A 2T8
- ⁴⁵Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁶University of Mississippi, University, Mississippi 38677, USA
- ⁴⁷Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Quebec, Canada H3C 3J7
- ⁴⁸Mount Holyoke College, South Hadley, Massachusetts 01075, USA
- ⁴⁹Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁵⁰NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵¹University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵²Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA
- ⁵³The Ohio State University, Columbus, Ohio 43210, USA
- ⁵⁴University of Oregon, Eugene, Oregon 97403, USA
- ⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵⁶Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France
- ⁵⁷Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
- ⁵⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁵⁹Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
- ⁶⁰Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- ⁶¹Prairie View A&M University, Prairie View, Texas 77446, USA
- ⁶²Princeton University, Princeton, New Jersey 08544, USA
- ⁶³Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶⁴Universität Rostock, D-18051 Rostock, Germany
- ⁶⁵Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶⁶DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- ⁶⁷University of South Carolina, Columbia, South Carolina 29208, USA
- ⁶⁸Stanford Linear Accelerator Center, Stanford, California 94309, USA
- ⁶⁹Stanford University, Stanford, California 94305-4060, USA
- ⁷⁰State Univ. of New York, Albany, New York 12222, USA
- ⁷¹University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁷²University of Texas at Austin, Austin, Texas 78712, USA
- ⁷³University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁷⁴Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁵Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷⁶Vanderbilt University, Nashville, Tennessee 37235, USA
- ⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- ⁷⁸University of Wisconsin, Madison, Wisconsin 53706, USA
- ⁷⁹Yale University, New Haven, Connecticut 06511, USA
- (Received 23 March 2004; published 9 July 2004)

We describe a measurement of the direct CP asymmetry between inclusive $b \rightarrow s\gamma$ and $\bar{b} \rightarrow \bar{s}\gamma$ decays. This asymmetry is expected to be less than 0.01 in the standard model, but could be enhanced up to about 0.10 by new physics contributions. We use a sample of 89×10^6 $B\bar{B}$ pairs recorded with the

BABAR detector at SLAC PEP-II, from which we reconstruct a set of 12 exclusive $b \rightarrow s\gamma$ final states containing one charged or neutral kaon and one to three pions. We measure an asymmetry of $A_{CP}(b \rightarrow s\gamma) = 0.025 \pm 0.050(\text{stat}) \pm 0.015(\text{syst})$, corresponding to an allowed range of $-0.06 < A_{CP}(b \rightarrow s\gamma) < +0.11$ at 90% confidence level.

DOI: 10.1103/PhysRevLett.93.021804

PACS numbers: 13.20.He, 11.30.Er, 12.15.Mm

The inclusive decay $b \rightarrow s\gamma$ is a flavor-changing neutral current process described by a radiative penguin loop diagram. The world average branching fraction is $(3.3 \pm 0.4) \times 10^{-4}$ [1] in good agreement with recent theoretical predictions [2]. Earlier experimental values of the branching fraction have been used to constrain new physics beyond the standard model [3]. A measurement of the direct CP asymmetry between $b \rightarrow s\gamma$ and $\bar{b} \rightarrow \bar{s}\gamma$ decays provides an independent and significant test of these predictions. In the standard model the dominant loop contribution contains a top quark, with other contributions being suppressed by Cabibbo-Kobayashi-Maskawa (CKM) factors and the Glashow-Iliopoulos-Maiani mechanism. The lack of interference between comparable amplitude contributions leads to a rather small predicted asymmetry [4]:

$$A_{CP}^{TH} = \frac{\Gamma(b \rightarrow s\gamma) - \Gamma(\bar{b} \rightarrow \bar{s}\gamma)}{\Gamma(b \rightarrow s\gamma) + \Gamma(\bar{b} \rightarrow \bar{s}\gamma)} = 0.0044_{-0.0014}^{+0.0024} \quad (1)$$

which has little sensitivity to the photon energy cutoff or to the distribution of hadronic final states. The dominant errors are due to the uncertainty of the charm quark mass and the choice of the perturbative scale. The inclusion of contributions to the loop beyond the standard model can increase the predicted asymmetry up to about 0.10 [4].

There is a previous measurement of direct CP asymmetry [5] in a sum of $b \rightarrow s\gamma$ and $b \rightarrow d\gamma$ decays. In the standard model, the total of the $b \rightarrow s\gamma$ and $b \rightarrow d\gamma$ asymmetries is exactly zero in the U-spin symmetry limit, $m_d = m_s$, as a consequence of CKM unitarity [6]. The measurement in Ref. [5] gives $-0.27 < 0.965 \times A_{CP}(b \rightarrow s\gamma) + 0.02 \times A_{CP}(b \rightarrow d\gamma) < 0.10$.

We use a sample of $(88.9 \pm 1.0) \times 10^6 \bar{B}\bar{B}$ pairs collected at the $Y(4S)$ resonance with the *BABAR* detector at the SLAC PEP-II asymmetric e^+e^- collider. A detailed description of the detector can be found elsewhere [7]. For this analysis the most important detector elements are the 40-layer drift chamber, situated in a 1.5 T solenoidal magnetic field, which measures charged particle momenta, the CsI(Tl) electromagnetic calorimeter, which measures the energies of the photons, and the detector of internally reflected Cherenkov light (DIRC), which is used to identify charged kaons.

We reconstruct $b \rightarrow s\gamma$ decays as the sum of 12 exclusive final states:

$$\begin{aligned} B^- &\rightarrow K^- \pi^0 \gamma, K^- \pi^+ \pi^- \gamma, K^- \pi^0 \pi^0 \gamma, K^- \pi^+ \pi^- \pi^0 \gamma, \\ \bar{B}^0 &\rightarrow K^- \pi^+ \gamma, K^- \pi^+ \pi^0 \gamma, K^- \pi^+ \pi^0 \pi^0 \gamma, K^- \pi^+ \pi^- \pi^+ \gamma, \\ B^- &\rightarrow K_S^0 \pi^- \gamma, K_S^0 \pi^- \pi^0 \gamma, K_S^0 \pi^- \pi^0 \pi^0 \gamma, K_S^0 \pi^- \pi^+ \pi^- \gamma, \end{aligned}$$

and measure the yield asymmetry with respect to their charge conjugate decays $\bar{b} \rightarrow \bar{s}\gamma$. The identification of charged kaons removes $b \rightarrow d\gamma$ decays. We do not use \bar{B}^0 decays to final states with K_S^0 to determine the direct CP asymmetry, since these are not flavor specific, but we study them to understand systematic effects.

The high energy photon is detected from an isolated energy cluster in the calorimeter, with shape consistent with a single photon, and energy $E_\gamma^* > 1.8$ GeV in the e^+e^- center-of-mass frame. A veto is applied to the high energy photons that combined with another photon form either a π^0 within the mass range 117–150 MeV/ c^2 or an η within the mass range 524–566 MeV/ c^2 .

Neutral kaons are reconstructed as $K_S^0 \rightarrow \pi^+ \pi^-$ candidates with an invariant mass within 9 MeV/ c^2 of the nominal mass [1], and a transverse flight distance > 2 mm from the primary event vertex. Charged kaons are tracks identified as kaons from information in the DIRC. The remaining tracks are considered to be charged pions. Both charged and neutral kaons are required to have a laboratory momentum > 0.7 GeV/ c . Above this threshold the rate for charged pions to be misidentified as kaons is $< 2.0\%$.

Neutral pions are reconstructed from pairs of photons with energies > 30 MeV. A π^0 mass cut is applied between 117 and 150 MeV/ c^2 . Charged and neutral pions are required to have laboratory momenta $> 0.5, 0.3,$ or 0.2 GeV/ c for states with 1, 2, or 3 pions, respectively, to reject combinatoric background.

The mass of the hadronic system, X_s , formed from the kaon and pions is required to be between 0.6 GeV/ c^2 and 2.3 GeV/ c^2 , corresponding to a photon energy threshold $E_\gamma > 2.14$ GeV in the B rest frame.

The signal Monte Carlo sample is generated according to Ref. [8], which predicts that $(83 \pm 5)\%$ of the $b \rightarrow s\gamma$ spectrum is above our photon energy threshold. The hadronic mass range below 1.1 GeV/ c^2 is assigned to the exclusive final state $B \rightarrow K^* \gamma$ and modeled with the known branching fraction and K^* mass distribution [1]. Above 1.1 GeV/ c^2 we use JETSET [9] to hadronize the system of the strange and spectator quarks. Within the selected hadronic mass range, the 12 final states constitute 48% of the total rate. If we were to include the \bar{B}^0 decays to K_S^0 and equate the decays to K_L^0 with those to K_S^0 , this would increase to 73% of the total rate. As a part of our analysis, we check the dependence of the asymmetry on the hadronic mass and final state.

Most of the background in this analysis arises from continuum production of a high energy photon, either by initial state radiation, or from the decays of π^0 and η

mesons. We remove 86% of these backgrounds by selections on the angle between the thrust axis of the B meson candidate and the thrust axis of all the other particles of the event, $|\cos\theta_T^*| < 0.80$, and the angle between the B candidate and the beam axis, $|\cos\theta_B^*| < 0.80$, both defined in the e^+e^- center-of-mass system. We then use a neural network to combine information from a set of event shape variables, including a set of energy flow cones. This halves the continuum background compared to our initial selection.

In 12% of the signal events, we can identify an electron or muon from the decay of the other B [10]. This is a very effective signature for removing continuum background, so the remaining background in this sample comes mostly from other B decays. We present separately our results for the sample of events which are lepton tagged.

Exclusive $b \rightarrow s\gamma$ decays are characterized by two kinematic variables: the beam-energy substituted mass, $m_{ES} = \sqrt{(\sqrt{s}/2)^2 - p_B^{*2}}$, and the energy difference between the B candidate and the beam energy, $\Delta E = E_B^* - (\sqrt{s}/2)$, where E_B^* and p_B^* are the energy and momentum of the B candidate in the e^+e^- center-of-mass frame, and \sqrt{s} is the total center-of-mass energy. We require candidates to have $|\Delta E| < 0.10$ GeV, and remove multiple candidates in each event by selecting the one with the smallest value of $|\Delta E|$. This technique is $> 90\%$ efficient when the true $b \rightarrow s\gamma$ decay is among the reconstructed candidates. We then fit the m_{ES} distribution between 5.22 and 5.29 GeV/ c^2 to extract the signal yield. When calculating m_{ES} , the value of p_B^* is corrected for the tail of the high energy photon response function by scaling the measured E_γ^* to the value that would give $\Delta E = 0$, the value expected for true signal.

In order to fit the m_{ES} distribution in data, we need to understand the different components of the signal and background events. We have identified the following four contributions as shown in Fig. 1. The signal events are described by a crystal ball function [11] with a resolution $\sigma(m_{ES}) = 2.2$ MeV/ c^2 . The continuum background is described by an ARGUS shape [12], which is cross checked by a fit to a sample of 9.6 fb^{-1} of data taken 40 MeV/ c^2 below the $Y(4S)$ resonance. We use a $B\bar{B}$ Monte Carlo sample to model the background from B decays other than $b \rightarrow s\gamma$, which is significant for X_s masses above 1.9 GeV/ c^2 . This background is described by the sum of an ARGUS shape and a peaking component which is modeled by the signal shape.

The last background component is cross feed from incorrectly reconstructed $b \rightarrow s\gamma$ events. This is modeled by the signal Monte Carlo sample, where we identify events reconstructed in the wrong final state. Cross feed occurs when the true $b \rightarrow s\gamma$ decay is not among the reconstructed candidates, or in a multiple candidate event when the wrong candidate is chosen. The shape of the cross feed is described by the sum of an ARGUS shape




FIG. 1. Monte Carlo simulations of the four contributions to the beam-energy substituted mass distribution of events selected as $b \rightarrow s\gamma$, with the corresponding fits: (a) signal, (b) continuum, (c) $B\bar{B}$ decays, and (d) cross feed. The plots are normalized to the luminosity of our data sample.

and a peaking signal shape. We regard cross feed as a background to be subtracted.

We fit the data m_{ES} distributions separately for each flavor. For the total sample, the fit function is parametrized by two ARGUS shapes and a crystal ball function. One ARGUS shape is fixed to be as the continuum ARGUS shape, while the other one is free to represent the sum of the nonpeaking $B\bar{B}$ and cross-feed backgrounds. The crystal ball function fits the combination of the peaking components. For the lepton-tagged sample, we use only one free ARGUS shape and a crystal ball function. In all cases we use an unbinned maximum likelihood fit. The fitting technique has been validated with a large sample of Monte Carlo simulated events. In Fig. 2 we present the final fits to the m_{ES} distributions for




FIG. 2. Fits to the beam-energy substituted mass distributions in data events for: (a) all $b \rightarrow s\gamma$, (b) all $\bar{b} \rightarrow \bar{s}\gamma$, (c) lepton-tagged $b \rightarrow s\gamma$, and (d) lepton-tagged $\bar{b} \rightarrow \bar{s}\gamma$ decays. Contributions are shown from peaking crystal ball (dot-dashed), fixed continuum ARGUS shape (dotted) and free $B\bar{B}$ and cross-feed ARGUS shape (dashed).

$b \rightarrow s\gamma$ and $\bar{b} \rightarrow \bar{s}\gamma$ events. The lower plots are for the lepton-tagged sample. All the fits have χ^2 per degree-of-freedom close to 1, if we make a fit to a binned distribution as shown in Fig. 2. The sum of events in the b and \bar{b} peaks is 1644 ± 72 , of which 201 ± 18 are lepton tagged. To get the true signal yields these have to be corrected for the predicted yield of peaking $B\bar{B}$ and cross-feed backgrounds from Monte Carlo samples (see Fig. 1), which is 88 ± 27 , where 10 ± 8 are lepton tagged.

The direct CP asymmetry is calculated from:

$$A_{CP} = \frac{1}{\langle D \rangle} \left[\frac{(n - \bar{n})}{(n + \bar{n})} - \frac{\Delta D}{2} \right] - A_{CP}^{\text{DET}}, \quad (2)$$

where n and \bar{n} are the numbers of observed $b \rightarrow s\gamma$ and $\bar{b} \rightarrow \bar{s}\gamma$ events after the peaking background is subtracted, $\Delta D = 2(\bar{w} - w)$ is the difference in the wrong flavor fraction between b and \bar{b} decays, and $\langle D \rangle = 1 - (w + \bar{w})$ is the dilution factor from the average wrong flavor fraction. A_{CP}^{DET} is the flavor asymmetry of the detector. We find $\Delta D = 0.001 \pm 0.002$ and $\langle D \rangle = 0.989 \pm 0.001$ from Monte Carlo samples. The small wrong flavor fraction is due to charged pions misidentified as charged kaons.

We need to correct the measured value of A_{CP} for the flavor asymmetry of the detector A_{CP}^{DET} . While it is known that the kaon-nucleon cross sections are asymmetric at low momenta, there are few accurate measurements [1]. This means that our Monte Carlo sample is not expected to model correctly the asymmetries due to the interactions of kaons with the inner part of the detector. The kaon identification efficiency of the DIRC for reconstructed tracks is measured with a control sample of kaons from D^* decays. Averaging over the kaon spectrum in $b \rightarrow s\gamma$ events we obtain a small asymmetry of -0.002 ± 0.001 from particle identification. We measure the overall detector asymmetry of the data events in our m_{ES} and ΔE sidebands, increasing the statistics by removing the neural network cut. Most of these events are from the continuum, where we do not expect any physics mechanism to generate a flavor asymmetry. We observe a significant asymmetry for kaon momenta below 1 GeV/ c . The asymmetry as a function of the kaon momentum is applied to the signal Monte Carlo to determine what shift should be applied to the data. This gives an overall flavor-asymmetry correction $A_{CP}^{\text{DET}} = -0.014 \pm 0.015$.

Table I presents the measured signal yields and corrected CP asymmetries. The lepton-tagged results are consistent with the results for the total sample. We divide the total sample into four bins in X_s mass, and observe no significant mass dependence of the asymmetry. The first bin corresponds to the $K^*(892)$ resonance, for which the world average asymmetry from studies of exclusive $B \rightarrow K^*\gamma$ decays is $A_{CP}(B \rightarrow K^*\gamma) = -0.01 \pm 0.07$ [1]. Our result is consistent with this average.

TABLE I. Signal yields and CP asymmetries for total and only lepton-tagged event samples. The total sample is also divided up into four bins in X_s mass in GeV/ c^2 , and into three types of decay modes. The errors on n and \bar{n} are statistical only, while for A_{CP} we quote the additional systematic error from the detector asymmetry.

Sample	n	\bar{n}	A_{CP}
Total sample	787 ± 54	769 ± 54	$0.025 \pm 0.050 \pm 0.015$
Lepton tagged	91 ± 14	100 ± 13	$-0.04 \pm 0.10 \pm 0.02$
$M_{X_s} = 0.6-1.1$	378 ± 32	396 ± 33	$0.003 \pm 0.059 \pm 0.015$
$M_{X_s} = 1.1-1.5$	162 ± 22	136 ± 23	$0.11 \pm 0.11 \pm 0.02$
$M_{X_s} = 1.5-1.9$	139 ± 19	124 ± 21	$0.07 \pm 0.11 \pm 0.03$
$M_{X_s} = 1.9-2.3$	101 ± 29	67 ± 36	$0.23 \pm 0.30 \pm 0.04$
B^0	455 ± 36	447 ± 38	$0.015 \pm 0.059 \pm 0.014$
$B^\pm \rightarrow K^\pm$	229 ± 31	148 ± 30	$0.22 \pm 0.12 \pm 0.02$
$B^\pm \rightarrow K_S^0$	100 ± 24	166 ± 25	$-0.20 \pm 0.14 \pm 0.03$

We divide our total sample into three types of decay mode: $B^0(\bar{B}^0) \rightarrow K^\pm$, $B^\pm \rightarrow K^\pm$, and $B^\pm \rightarrow K_S^0$. We observe a discrepancy of 2.3σ between the two B^\pm categories which we regard as a statistical fluctuation, since it is not correlated with a specific final state or hadronic mass bin. The combination of the B^\pm samples is consistent with a null asymmetry, as is the B^0 sample.

The dominant systematic error in our measurement is the uncertainty of 0.015 in the flavor asymmetry of the detection efficiency for charged and neutral kaons. For the lepton-tagged sample we add an additional systematic uncertainty of 0.010 to account for a possible charge asymmetry in the lepton-tagging efficiency. This is derived from studies of control samples [10].

We have tested the effect of possible flavor asymmetries in the peaking cross-feed and $B\bar{B}$ backgrounds by varying them within the current experimental bounds (90% C.L.). We added a 0.10 asymmetry to the cross-feed events, and a 0.02 asymmetry to the peaking background from $B\bar{B}$ decays, which comes primarily from $B \rightarrow D^{(*)}\rho$ decays. The change in our measured asymmetry due to these changes in the cross feed and $B\bar{B}$ flavor asymmetries is 0.004, which gives a negligible contribution to the error.

We have checked that the parameters of the ARGUS shapes and crystal ball functions are the same for both flavors within 1σ , so the detector asymmetry is simply an overall normalization difference between the two samples. We have also checked that the neural net distributions for signal and continuum background are flavor symmetric.

Our estimates of the cross-feed background and the detector asymmetry correction, A_{CP}^{DET} , depend on the mix of final states in our signal Monte Carlo sample. We check these, also using information from \bar{B}^0 decays to final states with K_S^0 , by varying the ratios of final states with K^+ or K_S^0 , and π^0 to π^+ measured in our data by $\pm 3\sigma$. Note that the measured ratios are consistent with our

signal Monte Carlo. Changing the ratios has no significant effect on the cross feed or the detector asymmetry correction.

Our final result for the direct CP asymmetry in $b \rightarrow s\gamma$ is $A_{CP} = 0.025 \pm 0.050 \pm 0.015$ for the total sample, and $A_{CP} = -0.04 \pm 0.10 \pm 0.02$ for the lepton-tagged sample. The total sample provides the best constraint, $-0.06 < A_{CP} < +0.11$ at 90% confidence level. This result begins to restrict the range of allowed new physics contributions in the flavor-changing penguin diagram.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.A.), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

*Now at Department of Physics, University of Warwick, Coventry, United Kingdom.

[†]Also with Università della Basilicata, Potenza, Italy.

[‡]Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain.

[§]Deceased.

- [1] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [2] P. Gambino and M. Misiak, Nucl. Phys. **B611**, 338 (2001); A. J. Buras, A. Czarnecki, M. Misiak, and J. Urban, Nucl. Phys. **B631**, 219 (2002).
- [3] J. L. Hewett and J. D. Wells, Phys. Rev. D **55**, 5549 (1997); T. Hurth, Rev. Mod. Phys. **75**, 1159 (2003).
- [4] A. L. Kagan and M. Neubert, Phys. Rev. D **58**, 094012 (1998); T. Hurth, E. Lunghi and W. Porod, hep-ph/0312260 [Nucl. Phys. B (to be published)].
- [5] CLEO Collaboration, T. E. Coan *et al.*, Phys. Rev. Lett. **86**, 5661 (2001).
- [6] J. M. Soares, Nucl. Phys. **367**, 575 (1991).
- [7] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [8] A. L. Kagan and M. Neubert, Eur. Phys. J. C **7**, 5 (1999).
- [9] Torbjörn Sjöstrand, *PYTHIA 5.7 and JETSET 7.4: Physics and Manual*, Lund University, hep-ph/9508391.
- [10] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 201802 (2002).
- [11] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **92**, 111801 (2004).
- [12] ARGUS Collaboration, H. Albrecht *et al.*, Z. Phys. C **48**, 543 (1990).