

UC Berkeley

UC Berkeley Previously Published Works

Title

The MAP4 Kinase SIK1 Ensures Robust Extracellular ROS Burst and Antibacterial Immunity in Plants

Permalink

<https://escholarship.org/uc/item/02g5q2m3>

Journal

Cell Host & Microbe, 24(3)

ISSN

1931-3128

Authors

Zhang, Meixiang

Chiang, Yi-Hsuan

Toruno, Tania Y

et al.

Publication Date

2018-09-01

DOI

10.1016/j.chom.2018.08.007

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Association of germline variants in telomere maintenance genes (*POT1*, *TERF2IP*, *ACD*, and *TERT*) with spitzoid morphology in familial melanoma: A multi-center case series



Alisa M. Goldstein, PhD,^a Richard Qin, BS,^a Emily Y. Chu, MD, PhD,^b David E. Elder, MBChB,^c Daniela Massi, MD, PhD,^d David J. Adams, PhD,^e Paul W. Harms, MD, PhD,^f Carla Daniela Robles-Espinoza, PhD,^{e,g} Julia A. Newton-Bishop, MD, PhD,^h D. Timothy Bishop, PhD,^h Mark Harland, PhD,^h Elizabeth A. Holland, BSc,^{i,j} Anne E. Cust, PhD,^{i,k,l} Helen Schmid, MPH,^{i,j,l} Graham J. Mann, MBBS, PhD,^{j,l,m} Susana Puig, MD, PhD,^{n,o} Miriam Potrony, PhD,^{o,p} Lluçcia Alos, MD, PhD,^q Eduardo Nagore, MD, PhD,^{r,s} David Millán-Esteban, PhD,^{r,s} Nicholas K. Hayward, PhD,^t Natasa Broit, PhD,^t Jane M. Palmer, RN,^t Vaishnavi Nathan, PhD,^t Elizabeth G. Berry, MD,^u Esteban Astiazaran-Symonds, MD,^a Xiaohong R. Yang, PhD,^a Margaret A. Tucker, MD,^a Maria Teresa Landi, MD, PhD,^a Ruth M. Pfeiffer, PhD,^a and Michael R. Sargen, MD^{a,v}

Background: Spitzoid morphology in familial melanoma has been associated with germline variants in *POT1*, a telomere maintenance gene (TMG), suggesting a link between telomere biology and spitzoid differentiation.

From the Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, Maryland^a; Department of Dermatology, Hospital of the University of Pennsylvania, Philadelphia, Pennsylvania^b; Department of Pathology and Laboratory Medicine, Hospital of the University of Pennsylvania, Philadelphia, Pennsylvania^c; Section of Pathological Anatomy, Department of Health Sciences, University of Florence, Florence, Italy^d; Experimental Cancer Genetics, The Wellcome Trust Sanger Institute, Hinxton, England^e; Department of Pathology, University of Michigan, Ann Arbor, Michigan^f; Laboratorio Internacional de Investigación sobre el Genoma Humano, Universidad Nacional Autónoma de México, Campus Juriquilla, Santiago de Querétaro, Qro, Mexico^g; Division of Haematology and Immunology, Institute of Medical Research at St James's, University of Leeds, Leeds, England^h; The Daffodil Centre, The University of Sydney, a joint venture with Cancer Council, NSW, Sydney, Australiaⁱ; Centre for Cancer Research, Westmead Institute for Medical Research, The University of Sydney, Sydney, Australia^j; Sydney School of Public Health, The University of Sydney, Sydney, NSW, Australia^k; Melanoma Institute Australia, The University of Sydney, Sydney, NSW, Australia^l; John Curtin School of Medical Research, Australian National University, Canberra, ACT, Australia^m; Melanoma Unit, Department of Dermatology, Hospital Clínic de Barcelona, IDIBAPS, Barcelona University, Barcelona, Spainⁿ; Centre of Biomedical Research on Rare Diseases (CIBERER), ISCIII, Barcelona, Spain^o; Melanoma Unit, Biochemistry and Molecular Genetics Department, Hospital Clínic de Barcelona, IDIBAPS, Barcelona University, Barcelona, Spain^p; Pathology Department, Hospital Clínic de Barcelona, Universitat de Barcelona, Barcelona, Spain^q; Department of Dermatology, Fundación Instituto Valenciano de Oncología, València, Spain^r; School of Medicine, Universidad Católica de València San

Vicente Mártir, València, Spain^s; QIMR Berghofer Medical Research Institute, Herston, QLD, Australia^t; Department of Dermatology, Oregon Health and Science University, Portland, Oregon^u; and Laboratory of Pathology, Center for Cancer Research, National Cancer Institute, Bethesda, Maryland^v

Funding sources: This work was supported by the Intramural Research Program of the Division of Cancer Epidemiology and Genetics, National Cancer Institute, and National Institutes of Health. This research was also supported by the National Health and Medical Research Council of Australia (Centre of Research Excellence in Melanoma 1135285, and Investigator Grant 2008454). The research at the Melanoma Unit in Hospital Clínic Barcelona is partially financed by Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), of the Instituto de Salud Carlos III, Spain, cofinanced by European Development Regional Fund "A way to achieve Europe" ERDF; Spanish Fondo de Investigaciones Sanitarias grants PI15/00716, PI15/00956, and PI18/00419, of the Instituto de Salud Carlos III, Spain, co-financed by European Development Regional Fund "A way to achieve Europe" ERDF; AGAUR 2017_SGR_1134 of the Catalan Government, Spain; European Commission under the sixth Framework Programme, Contract No. LSHC-CT-2006-018702 (GenoMEL), by the European Commission under the seventh Framework Programme, Diagnostics, and the European Commission under the HORIZON2020 Framework Programme, iTobos and Qualitop; The National Cancer Institute (NCI) of the U.S. National Institutes of Health (NIH) (CA83115); a grant from "Fundació La Marató de TV3" 201331-30, Catalonia, Spain; and a grant from "Fundación Científica de la Asociación Española Contra el Cáncer" GCB15152978SOEN, Spain, and CERCA Programme/Generalitat de Catalunya. Part of the work was carried out at the Esther

Objective: To assess if familial melanoma cases associated with germline variants in TMG (*POT1*, *ACD*, *TERF2IP*, and *TERT*) commonly exhibit spitzoid morphology.

Methods: In this case series, melanomas were classified as having spitzoid morphology if at least 3 of 4 dermatopathologists reported this finding in $\geq 25\%$ of tumor cells. Logistic regression was used to calculate odds ratios (OR) of spitzoid morphology compared to familial melanomas from unmatched noncarriers that were previously reviewed by a National Cancer Institute dermatopathologist.

Results: Spitzoid morphology was observed in 77% (23 of 30), 75% (3 of 4), 50% (2 of 4), and 50% (1 of 2) of melanomas from individuals with germline variants in *POT1*, *TERF2IP*, *ACD*, and *TERT*, respectively. Compared to noncarriers ($n = 139$ melanomas), *POT1* carriers (OR = 225.1, 95% confidence interval: 51.7-980.5; $P < .001$) and individuals with *TERF2IP*, *ACD*, and *TERT* variants (OR = 82.4, 95% confidence interval: 21.3-494.6; $P < .001$) had increased odds of spitzoid morphology.

Limitations: Findings may not be generalizable to nonfamilial melanoma cases.

Conclusion: Spitzoid morphology in familial melanoma could suggest germline alteration of TMG. (JAAD Int 2023;11:43-51.)

Key words: *ACD*; familial melanoma; melanoma; *POT1*; spitzoid melanoma; spitz melanoma; *TERF2IP*; *TERT*.

INTRODUCTION

Approximately 10% of cutaneous melanoma patients have a family history of the disease (familial melanoma).¹ Germline pathogenic variants (GPVs) in melanoma susceptibility genes *CDKN2A*, *CDK4*, and *BAP1* are identified in up to 40%, 0.7%, and 1.0% of familial melanoma cases, respectively.^{1,2} More recently, GPV in telomere maintenance genes (TMG) *POT1*, *TERF2IP*, *ACD*, and *TERT* have been observed in up to 6% of melanoma-prone families.³⁻⁷ *POT1* GPV

CAPSULE SUMMARY

- Germline pathogenic variants in telomere maintenance genes *POT1*, *TERF2IP*, *ACD*, and *TERT* predispose to melanoma.
- Spitzoid morphology is common in familial melanoma cases from individuals with *POT1*, *TERF2IP*, *ACD*, and *TERT* germline variants. Dermatologists and dermatopathologists should be aware of these associations when ordering familial melanoma gene testing panels.

also predispose to lymphoid and myeloid malignancies, angiosarcoma, and glioma.^{8,9}

BAP1 GPV carriers commonly develop spitzoid melanocytic neoplasms, characterized by large epithelioid melanocytes with abundant cytoplasm. The biological mechanisms underlying this genotype-histotype association are unknown.^{10,11} Spitzoid subtype melanomas with complete spitzoid morphology (100% of tumor) and cutaneous melanomas with focal spitzoid morphology ($\geq 25\%$ of tumor) have also been

Koplowitz Center, Barcelona. The work at Fundació Instituto Valenciano de Oncología was partially funded by the Instituto de Salud Carlos III (PI19/00667), the Generalitat Valenciana (Prometeo 21/067), and the EADV (PPRC,-2018-36). The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

IRB approval status: Familial melanoma studies submitting whole slide images to the National Cancer Institute had local institutional review board or ethics review committee approval. Clinicaltrials.gov (or equivalent) listing (if applicable): NCT00556829, NCT00040352.

Prior presentation: Data from this study was presented at the Melanoma Genetics Consortium (GenoMEL) annual meeting in Inverness, United Kingdom (October 22-24, 2022).

Accepted for publication January 19, 2023.

Correspondence and reprint requests to: Michael R. Sargen, MD, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, 9609 Medical Center Dr, Room 6E-542, Rockville, MD 20850. E-mail: michael.sargen@nih.gov.

2666-3287

Published by Elsevier Inc. on behalf of the American Academy of Dermatology, Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.1016/j.jdin.2023.01.013>

Abbreviations used:

CI:	confidence interval
GPV:	germline pathogenic variant
OR:	odds ratio
TMG:	telomere maintenance gene

observed in *POT1* GPV carriers, suggesting dysfunctional telomere maintenance as a possible mechanism for spitzoid histology.¹² Here, we further investigate this hypothesis by evaluating the pathology of familial melanoma cases from individuals with germline variants in TMG.

METHODS

Whole slide images of cutaneous melanomas from individuals with any germline variant in a TMG (*POT1*, *TERF2IP*, *ACD*, and *TERT*) and at least one first, second, or third degree relative with melanoma, were requested during the virtual Melanoma Genetics Consortium (GenoMEL) annual meeting on July 27, 2020. Images were provided to the National Cancer Institute by research groups conducting familial melanoma studies in the United States (1 group), Europe (3 groups), and Australia (2 groups). Collaborators submitted all melanoma cases they had access to from variant carriers, which were independently reviewed by 4 dermatopathologists (MRS, DEE, EYC, and DM) with expertise in melanocytic lesions, who were blinded to clinical information and the specific gene altered. Each dermatopathologist reported histologic subtype and the percentage of tumor cells exhibiting spitzoid morphology, defined as large epithelioid or spindled melanocytes with a cytoplasm to nuclear ratio of greater than or equal to 1. To be consistent with our previous study on this topic, melanomas were classified as spitzoid if at least 3 of 4 dermatopathologists reported spitzoid morphology in 25% or more of the tumor cells.¹²

Logistic regression was used to estimate odds ratios (ORs) and 95% confidence intervals (CIs) for spitzoid morphology and subtype in variant carriers compared to unmatched noncarriers (United States, Europe). All noncarriers had whole exome sequencing performed to exclude GPV in melanoma predisposition genes (*CDKN2A*, *CDK4*, *BAP1*, *POT1*, *TERT*, *ACD*, and *TERF2IP*). Noncarrier cases (all cutaneous melanomas) were previously reviewed by a single dermatopathologist (MRS),¹² and were from individuals with at least one first, second, or third degree relative with melanoma (same as variant carriers). CIs accommodated within person and within family correlations by using a generalized estimating equation approach with a working

independence covariance matrix. Analyses were performed in R Studio and SAS 9.4 (SAS Institute Inc).

Familial melanoma studies submitting whole slide images to the National Cancer Institute had local institutional review board or ethics review committee approval.

RESULTS

Four dermatopathologists reviewed the histology of melanomas from individuals with germline variants in *POT1* ($n = 30$ melanomas, 17 individuals), *TERF2IP* ($n = 4$ melanomas, 3 individuals), *ACD* ($n = 4$ melanomas, 2 individuals), and *TERT* ($n = 2$ melanomas, 2 individuals). Clinical information on variant carriers (28 women [70%], 24 from Europe [60%]) and noncarriers (70 women [50.4%], 81 from Europe [58%]) is available in Table I. Similar proportions of melanomas in the 2 groups were invasive (variant carriers vs noncarriers: 87.5% vs 79.1%, two-sided Fisher's exact $P = 0.36$), ulcerated (variant carriers vs non-carriers: 15.0% vs 19.1%, two-sided Fisher's exact $P = 0.38$), and had a mitotic rate of ≥ 1 mitosis per square millimeter (variant carriers vs non-carriers: 42.9% vs 33.6% for invasive melanomas, two-sided Fisher's exact $P = 0.42$).

Spitzoid morphology involving at least 25% of the tumor was observed by at least 3 dermatopathologists in 77% (23 of 30), 75% (3 of 4), 50% (2 of 4), and 50% (1 of 2) of melanomas from individuals with germline variants in *POT1*, *TERF2IP*, *ACD*, and *TERT*, respectively. Furthermore, 13% (4 of 30) of melanomas from *POT1* carriers and 50% (1 of 2) from *TERT* carriers were completely spitzoid, and therefore, were classified as spitzoid subtype melanomas. The median age at diagnosis for a melanoma with spitzoid morphology was 51.5 years (range: 20-77 years) for the variant carrier group (all genes including *POT1*) and 52 years (range: 35-77 years) for *POT1* variant carriers (Table I).

Compared to noncarriers, individuals with germline *POT1* variants had a 10.5-fold increased odds (95% CI: 2.1-51.7; $P = .004$) of developing a spitzoid subtype melanoma (complete spitzoid morphology) and a 225.1-fold increased odds (95% CI: 51.7-980.5; $P < .001$) of having a melanoma with spitzoid morphology involving at least 25% of the tumor. Restricting the analysis to cases from individuals with likely pathogenic or pathogenic variants in *POT1* based on the American College of Medical Genetics and Genomics and Association for Molecular Pathology guidelines for the interpretation of sequence variants also yielded similar results (OR [spitzoid subtype] = 12.1, 95% CI: 2.2-65.7, $P = .004$; OR [spitzoid morphology] = 205.5, 95% CI: 8.7-1089.9, $P < .001$). *BAP1* germline status was

Table I. Clinical and histologic characteristics of familial melanoma cases

Characteristic	Noncarrier [¶]	Germline variants in telomere maintenance genes			
		<i>ACD</i> [‡]	<i>TERF2IP</i> [‡]	<i>TERT</i> [‡]	<i>POT1</i> [‡]
No. of families	68	2	3	2	16
No. of individuals	132	2	3	2	17
No. of cases (%)	139 (100)	4 (100)	4 (100)	2 (100)	30 (100)
Country, No. (%)					
United States	58 (41.7)	0	0	0	1 (3.3)
Europe [†]	81 (58.3)	0	2 (50)	1 (50)	21 (70.0)
Australia	0	4 (100)	2 (50)	1 (50)	8 (26.7)
Female sex, No. (%)	70 (50.4)	4 (100)	1 (25)	1 (50)	22 (73.3)
Median (range) age at diagnosis, y	49 (15-91)	55 (50-56)	41 (20-59)	29.5 (28-31)	52 (35-77)
Body site, No. (%)					
Head and neck	18 (12.9)	0	0	0	1 (3.3)
Trunk	56 (40.3)	2 (50)	1 (25)	2 (100)	13 (43.3)
Upper extremities	26 (18.7)	1 (25)	3 (75)	0	5 (16.7)
Lower extremities	31 (22.3)	1 (25)	0	0	11 (36.7)
Unknown	8 (5.8)	0	0	0	0
Subtype, No. (%) [‡]					
Superficial spreading melanoma	104 (74.8)	4 (100)	3 (75)	1 (50)	21 (70)
Nodular melanoma	12 (8.6)	0	1 (25)	0	4 (13.3)
Spitzoid melanoma	2 (1.4)	0	0	1 (50)	4 (13.3)
Lentigo maligna melanoma	17 (12.2)	0	0	0	0
Acral melanoma	4 (2.9)	0	0	0	1 (3.3)
Spitzoid morphology, No. (%) [§]	2 (1.4)	2 (50)	3 (75)	1 (50)	23 (76.7)
Breslow thickness (mm), No. (%)					
in situ	29 (20.9)	0	0	0	5 (16.7)
0.1-0.7	59 (42.4)	3 (75)	2 (50)	2 (100)	10 (33.3)
0.8-2.0	29 (20.9)	1 (25)	2 (50)	0	6 (20)
2.1-4.0	14 (10.1)	0	0	0	8 (26.7)
>4.0	8 (5.8)	0	0	0	1 (3.3)
Ulceration	13 (19.1)	0	1 (25)	0	5 (16.7)
No. (%) of invasive melanomas	110 (100)	4 (100)	4 (100)	2 (100)	25 (100)
Mitotic rate (#/mm ²) for invasive melanomas					
0	73 (66.4)	4 (100)	3 (75)	2 (100)	11 (44)
1-5	27 (24.5)	0	1 (25)	0	11 (44)
>5	10 (9.1)	0	0	0	3 (12)

NCI, National Cancer Institute.

*Pathogenic variants in melanoma susceptibility genes (*CDKN2A*, *CDK4*, *BAP1*, *POT1*, *TERT*, *ACD*, *TERF2IP*) were excluded by whole exome sequencing analysis.

[†]All European noncarrier cases were from Italy ($n = 81$). European variant carrier cases were from Spain ($n = 2$ from *TERF2IP* variant carriers; $n = 1$ from a *TERT* variant carrier; $n = 12$ from *POT1* variant carriers) and the United Kingdom ($n = 9$ from *POT1* variant carriers).

[‡]All melanomas from individuals with germline variants in telomere maintenance genes (*POT1*, *TERF2IP*, *ACD*, and *TERT*) were independently reviewed by 4 dermatopathologists (MRS, EC, DM, and DEE), and melanoma subtype was based on the majority opinion of the reviewers. Histologic subtype for noncarrier cases was assessed by a single dermatopathologist (MRS).

[§]Spitzoid morphology was defined as large epithelioid or spindled melanocytes with a cytoplasm to nuclear ratio of greater than or equal to 1. Melanomas from individuals with germline variants in telomere maintenance genes (*POT1*, *TERF2IP*, *ACD*, and *TERT*) were classified as spitzoid if at least 3 of the 4 dermatopathologists (MRS, EC, DM, and DEE) reported spitzoid morphology in 25% or more of the tumor cells. Dermatopathologist agreement on spitzoid classification was slight to moderate (vs NCI dermatopathologist [MRS]: non-NCI reviewer 1, kappa = 0.42; non-NCI reviewer 2, kappa = 0.16; and non-NCI reviewer 3, kappa = 0.38), consistent with a previous study (PubMed ID: 34757982) evaluating interobserver agreement for spitzoid classification. Spitzoid morphology in noncarrier cases was assessed by a single dermatopathologist (MRS).

^{||}Information was ascertained from pathology reports and confirmed by histopathology review (MRS).

[¶]All noncarriers were white. All variant carriers were white except for one individual with a germline *TERF2IP* variant with unknown race.

unknown in 10 cases from *POT1* carriers and excluding these cases did not impact associations (Table II).

Spitzoid morphology was also more prevalent in melanomas from individuals with *TERF2IP*, *ACD*, and *TERT* variants compared to noncarriers

Table II. Odds of spitzoid morphology and subtype among individuals with germline variants in telomere maintenance genes

Group	Spitzoid morphology*			Spitzoid subtype†		
	No. (%)	OR (95% CI)‡	P-value	No. (%)	OR (95% CI)‡	P-value
Noncarriers (N = 139)§	2 (1.4)	Reference		2 (1.4)	Reference	
Any <i>POT1</i> variant (N = 30)	23 (76.7)	225.1 (51.7-980.5)	<.001	4 (13.3)	10.5 (2.1-51.7)	.004
Likely pathogenic or pathogenic <i>POT1</i> variant (N = 20)¶	15 (75.0)	205.5 (8.7-1089.9)	<.001	3 (15.0)	12.1 (2.2-65.7)	.004
Any <i>POT1</i> variant from individual with negative testing for germline <i>BAP1</i> variants (N = 20)	17 (85.0)	388.2 (69.2- 2177.5)	<.001	2 (10)	7.6 (1.0-55.6)	.046
Any <i>ACD</i> , <i>TERF2IP</i> , or <i>TERT</i> variant (N = 10)¶	6 (60.0)	82.4 (21.3-494.6)	<.001	1 (10.0)	8.7 (2.1-51.7)	.004

ACMG, American College of Medical Genetics and Genomics; AMP, Association for Molecular Pathology.

*Spitzoid morphology was defined as large epithelioid or spindled melanocytes with a cytoplasm to nuclear ratio of greater than or equal to 1. Melanomas from individuals with germline variants in telomere maintenance genes (*POT1*, *TERF2IP*, *ACD*, and *TERT*) were classified as spitzoid if at least 3 of the 4 dermatopathologists (MRS, EC, DM, and DEE) reported spitzoid morphology in 25% or more of the tumor cells. Spitzoid morphology for noncarrier cases was assessed by a single dermatopathologist (MRS).

†All melanomas from individuals with germline variants in telomere maintenance genes (*POT1*, *TERF2IP*, *ACD*, and *TERT*) were independently reviewed by 4 dermatopathologists (MRS, EC, DM, and DEE), and melanoma subtype was based on the majority opinion of the reviewers. Histologic subtype for noncarrier cases was assessed by a single dermatopathologist (MRS).

‡Confidence intervals accommodated within person and within family correlations by using a generalized estimating equation approach with a working independence covariance matrix.

§The noncarrier group includes the United States and Italian familial melanoma cases from individuals with no identifiable germline pathogenic variants in melanoma predisposition genes (*CDKN2A*, *CDK4*, *BAP1*, *POT1*, *TERT*, *ACD*, and *TERF2IP*) based on whole exome sequencing analysis.

¶Variant classification based on ACMG and AMP standards and guidelines for the interpretation of sequence variants (PubMed ID: 25741868).

¶No variants were predicted to be likely pathogenic or pathogenic based on ACMG/AMP standards and guidelines for the interpretation of sequence variants (PubMed ID: 25741868). ACMG/AMP predictions for this group included variants of uncertain significance (n = 5 cases) and benign variants (n = 5 cases). All variant carriers had at least one first, second, or third degree relative with melanoma.

(OR = 82.4, 95% CI: 21.3-494.6; *P* < .001) (Table II). Representative photomicrographs of spitzoid morphology in cutaneous melanomas from individuals with germline *POT1* variants are shown in Figs 1 to 3. Additional photomicrographs of spitzoid morphology in cutaneous melanomas from individuals with germline variants in other TMG are shown in Fig 4.

The American College of Medical Genetics and Genomics and Association for Molecular Pathology predictions for variant pathogenicity are provided in Table III.

DISCUSSION

Identification of individuals with GPV in TMG is critically important for counseling and cancer screening. Our analysis identified spitzoid morphology ($\geq 25\%$ of tumor) in 77% of cutaneous melanomas from *POT1* carriers, consistent with a recent case series from our group where 60% of cutaneous melanomas from *POT1* carriers exhibited this histology.¹² In the current study, all melanomas exhibiting spitzoid morphology from *POT1* carriers occurred in adulthood (range: 35-77 years), although

spitzoid subtype melanomas with complete spitzoid morphology (100% of tumor) have been reported in *POT1* carriers as young as age 16 years.¹² We also identified spitzoid morphology in melanomas from individuals with variants in TMG other than *POT1* (*TERF2IP*, *ACD*, and *TERT*). All cases classified as having spitzoid morphology were independently observed to have this histologic finding by at least 3 of 4 dermatopathologists suggesting that spitzoid cells can be differentiated from other morphologies.

BAP1 variants have also been associated with spitzoid pathology.¹⁰ Inactivation of *BAP1* increases *TERT* expression and telomere elongation, suggesting that it may induce spitzoid differentiation via dysfunctional telomere maintenance.¹³ In our study, *BAP1* variants were not detected in any individuals with *TERF2IP*, *ACD*, or *TERT* variants. Furthermore, exclusion of cases from *POT1* carriers with unknown *BAP1* germline status did not significantly impact associations.

Our study findings support the inclusion of TMG (*POT1*, *TERF2IP*, *ACD*, and *TERT*) on genetic testing panels when a familial melanoma case exhibits spitzoid morphology in at least 25% of




Fig 1. Cutaneous melanomas exhibiting spitzoid morphology from individuals with germline *POT1* variants. **A**, Low magnification (4.9 \times) of spitzoid subtype melanoma (case 14-B) from an individual with germline *POT1* p.Asp598Serfs*22 variant. **B**, Higher magnification (25.3 \times) of case 14-B shows atypical melanocytes with abundant eosinophilic cytoplasm consistent with spitzoid morphology. **C**, Low magnification (2.9 \times) of spitzoid subtype melanoma (case 3) from an individual with germline *POT1* c.1164-1G>A splice acceptor variant. **D**, Higher magnification (21.1 \times) of case 3 shows epidermal and dermal nests of epithelioid and spindled melanocytes with abundant cytoplasm; some junctional nests show separation from the surrounding epidermis (arrows), which is a feature of Spitz tumors. **E**, Low magnification (3.7 \times) of a melanoma (case 9) from an individual with germline *POT1* p.Lys427Arg variant; the tumor exhibits areas of spitzoid (outlined in red) and nonspitzoid differentiation. **F**, Higher magnification (40 \times) photomicrograph of spitzoid area in case 9 showing enlarged epithelioid melanocytes with abundant eosinophilic cytoplasm. **G**, Higher magnification (40 \times) photomicrograph of nonspitzoid area in case 9 showing epithelioid and spindled melanocytes that lack abundant cytoplasm.




Fig 2. Cutaneous melanoma exhibiting spitzoid morphology with acantholytic features from individual with germline *POT1* variant. Case 6 is from an individual with a germline *POT1* variant (p.Arg117His) and pathology shows a sheet-like proliferation of spitzoid cells with loss of cohesion, consistent with an acantholytic growth pattern.

the tumor cells. Since genetic testing is most commonly ordered by dermatologists, oncologists, or geneticists who evaluate melanoma-prone families, dermatopathologists should report the

presence ($\geq 25\%$ of tumor) or absence of spitzoid morphology in the synoptic or microscopic description sections of all melanoma pathology reports regardless of the melanoma subtype (superficial spreading, nodular, lentigo maligna, spitzoid, acral, and desmoplastic) or family medical history, which could be incomplete or unavailable at the time of biopsy. Routine reporting of this information could assist with identifying individuals with GPVs in TMG, and detecting these variants will inform cancer surveillance for the patient, and possibly family members, based on the associated cancer risks.

LIMITATIONS

Although this case series is the largest histopathologic assessment of cutaneous melanomas from individuals with germline variants in TMG, our study ascertained cases from GenoMEL melanoma-prone families, and therefore, the study findings may not be




Fig 3. Cutaneous melanoma exhibiting spitzoid morphology in the dermis from individual with germline *POT1* variant. **A**, Case 1 is from an individual with a germline variant in *POT1* (p.Arg137His) and pathology shows a superficial spreading growth pattern (nesting, pagetoid scatter, and increased pigmentation) in the epidermis and an invasive component with spitzoid morphology (*asterisk*). **B**, Higher magnification (40 \times) of spitzoid area in case 1.

applicable to nonfamilial melanoma cases. Additionally, while cases were independently reviewed by multiple expert dermatopathologists, confirmation bias cannot be excluded. Therefore, case-control studies are needed to confirm the observed findings in this study and assess interobserver agreement for spitzoid classification. Noncarrier cases were reviewed by a single dermatopathologist and were not matched by country, which could have also impacted association estimates. However, previous GenoMEL studies of familial melanoma, involving histopathology review by multiple expert pathologists, did not identify spitzoid morphology in *CDKN2A* carrier and noncarrier familial melanomas.^{14,15} Lastly, tissue was not available for cases with spitzoid morphology to evaluate for somatic alterations (ie, *HRAS* variants and gene fusions involving *ALK*, *ROS1*, *NTRK1*, *NTRK3*, *MET*, *RET*, *BRAF*, *MAP3K8*, and *CRTC1::TRIM11*) that have been associated with pediatric spitzoid tumors.^{16,17} Therefore, tumor studies are needed to understand the biological mechanisms linking dysfunctional telomere maintenance to spitzoid differentiation in familial melanoma cases.




Fig 4. Cutaneous melanomas exhibiting spitzoid morphology from individuals with *TERT*, *TERF2IP*, and *ACD* variants. **A**, Case 24 is from an individual with a germline variant in *TERT* (p.Ala202Thr) and pathology shows nests and sheets of large epithelioid melanocytes with abundant eosinophilic cytoplasm consistent with spitzoid morphology (40 \times magnification). **B**, Case 18 is from an individual with a germline variant in *TERF2IP* (p.Glu304del) and pathology shows nests and single cells of melanocytes with abundant eosinophilic cytoplasm in the papillary dermis consistent with spitzoid morphology (40 \times magnification). **C**, Case 20-A is from an individual with a germline variant in *TERF2IP* (p.Pro285Ser) and pathology shows a compound melanocytic lesion with epidermal and dermal nests of epithelioid melanocytes with abundant eosinophilic cytoplasm consistent with spitzoid morphology (40 \times magnification). **D**, Case 21-C is from an individual with a germline variant in *ACD* (p.Val432Ala) and pathology shows epidermal and dermal nests of melanocytes with abundant eosinophilic cytoplasm, consistent with spitzoid morphology, and a dense lymphocytic infiltrate (40 \times magnification).

Table III. Classification of variants using ACMG/AMP criteria

Cases	Gene	Genomic location (human genome reference GRCh38)	Transcripts	Nucleotide change	Amino acid change	ClinVar classification	ClinVar ID	dbSNP	ACMG/AMP variant classification	ACMG/AMP criteria
1*	<i>POT1</i>	chr7: 124863486	NM_015450.3 ENST00000357628.8	c.410G>A	p.Arg137His	VUS	139523	rs587777475	LP	PM2, PS3, PS4
2*	<i>POT1</i>	chr7: 124851918	NM_015450.3 ENST00000357628.8	c.903G>T	p.Gln301His	LB; B	475112	rs116916706	LB	PM2, BS2, BP6
3*,5*	<i>POT1</i>	chr7: 124841179	NM_015450.3 ENST00000357628.8	c.1164-1G>A	splice acceptor	LP; P	475026	rs866612394	P	PVS1, PM2, PP5
4*	<i>POT1</i>	chr7: 124842843	NM_015450.3 ENST00000357628.8	c.1127A>G	p.Gln376Arg	Conflicting (VUS; LB)	475020	rs143635917	VUS	PM2, BP6
6*	<i>POT1</i>	chr7: 124863546	NM_015450.3 ENST00000357628.8	c.350G>A	p.Arg117His	VUS	654409	rs1385542313	VUS	PM2, PM5
7*, 12 [†] , 13 [‡]	<i>POT1</i>	chr7: 124858989	NM_015450.3 ENST00000357628.8	c.670G>A	p.Asp224Asn	VUS	139527	rs202187871	LP	PM2, PP3, PS3, PS4
8*	<i>POT1</i>	chr7: 124841131	NM_015450.3 ENST00000357628.8	c.1211G>T	p.Gly404Val	B	475029	rs35536751	B	BA1, BS1, BS2, BP4, BP6
9*	<i>POT1</i>	chr7: 124841062	NM_015450.3 ENST00000357628.8	c.1280A>G	p.Lys427Arg	Not in ClinVar			VUS	PM2, BP4
10 [‡]	<i>POT1</i>	chr7: 124870933	NM_015450.3 ENST00000357628.8	c.233T>C	p.Ile78Thr	VUS	475073	rs947005337	LP	PM2, PS4
11-A-F [‡]	<i>POT1</i>	chr7: 124825358	NM_015450.3 ENST00000357628.8	c.1687-1G>A	splice acceptor	Conflicting (VUS; risk factor)	139521	rs587777473	P	PVS1, PM2, PS4
14-A-G [‡]	<i>POT1</i>	chr7: 124825252	NM_015450.3 ENST00000357628.8	c.1792G>A	p.Asp598Serfs*22	Not in ClinVar		rs1487204320	LP	PM2, PS3
15-A-C [§]	<i>POT1</i>	chr7: 124840979	NM_015450.3 ENST00000357628.8	c.1363A>G	p.Ile455Val	VUS	475039	rs776965979	VUS	PM2
16,17 [‡]	<i>POT1</i>	chr7: 124851884	NM_015450.3 ENST00000357628.8	c.937G>A	p.Asp313Asn	VUS	1359398	rs770779418	VUS	PM2, BP4
18 [‡]	<i>TERF2IP</i>	chr16: 75656305	NM_018975.4 ENST00000300086.5	c.910_912 delGAA	p.Glu304del	Not in ClinVar		rs747026590	VUS	PM2, BP3
19 [‡]	<i>TERF2IP</i>	chr16: 75656381	NM_018975.4 ENST00000300086.5	c.970A>G	p.Lys324Glu	Not in ClinVar		rs4888444	B	BS1, BS2, BP4
20-A-B [§]	<i>TERF2IP</i> ^e	chr16: 75656264	NM_018975.4 ENST00000300086.5	c.853C>T	p.Pro285Ser	Not in ClinVar		rs756850727	VUS	PM2, BP4
21-A-C [‡] , 22 [‡]	<i>ACD</i>	chr16: 67657765	NM_001082486.2 ENST00000620761.6	c.1295T>C	p.Val432Ala	B	1167321	rs6979	B	BA1, BS2, BP4, BP6
23 [‡]	<i>TERT</i>	chr5: 1254479	NM_198253.3 ENST00000310581.10	c.3184G>A	p.Ala1062Thr	Conflicting (VUS; LB; B)	39121	rs35719940	VUS	PM2, BP3
24 [‡]	<i>TERT</i>	chr5: 1294282	NM_198253.3 ENST00000310581.10	c.604G>A	p.Ala202Thr	Conflicting (VUS; LB; B)	12729	rs121918661	VUS	PP2, BS2, BP6

ACMG, American College of Medical Genetics and Genomics; AMP, Association for Molecular Pathology; B, benign; LB, likely benign; LP, likely pathogenic; P, pathogenic; VUS, variant of uncertain significance.

*Germline variant identified by panel testing of the following genes: *CDKN2A*, *CDK4*, and *POT1*.

[†]Germline *POT1* variant identified by Sanger sequencing of the gene. Germline whole genome sequencing of a sibling of case 12 with the family *POT1* variant that did not identify variants in other melanoma susceptibility genes including *BAP1*.

[‡]Germline variant identified by whole exome/genome sequencing analysis.

[§]Germline variant identified by panel testing of the following genes: *CDKN2A*, *CDK4*, *BAP1*, *MITF*, *POT1*, *ACD*, *TERF2IP*, *TERT* promoter, *BRCA1*, *BRCA2*, *MSH2*, and *MSH6*.

^{||}ClinVar and ACMG and AMP classifications.

CONCLUSIONS

In conclusion, we identified that a high proportion of cutaneous melanomas from individuals with germline variants in TMG exhibit spitzoid morphology. Therefore, panel testing for *POT1*, *TERF2IP*, *ACD*, and *TERT* should be considered when familial melanoma cases exhibit spitzoid differentiation.

We would like to thank Marc Tischkowitz (Department of Medical Genetics, University of Cambridge, Cambridge, United Kingdom) for reviewing variant classifications. Queensland acknowledges Hayley Hamilton's contribution in coordinating confirmation of participating family members' melanoma status along with the multiple pathology centers and Australian cancer registries involved. We would also like to thank the families who have agreed to participate in familial melanoma research studies as well as the nurses and research assistants who have supported this research.

Conflicts of interest

None disclosed.

REFERENCES

- Potrony M, Badenas C, Aguilera P, et al. Update in genetic susceptibility in melanoma. *Ann Transl Med*. 2015;3(15):210. <https://doi.org/10.3978/j.issn.2305-5839.2015.08.11>
- Goldstein AM, Chan M, Harland M, et al. Features associated with germline CDKN2A mutations: a GenoMEL study of melanoma-prone families from three continents. *J Med Genet*. 2007;44(2):99-106. <https://doi.org/10.1136/jmg.2006.043802>
- Shi J, Yang XR, Ballew B, et al. Rare missense variants in POT1 predispose to familial cutaneous malignant melanoma. *Nat Genet*. 2014;46(5):482-486. <https://doi.org/10.1038/ng.2941>
- Robles-Espinoza CD, Harland M, Ramsay AJ, et al. POT1 loss-of-function variants predispose to familial melanoma. *Nat Genet*. 2014;46(5):478-481. <https://doi.org/10.1038/ng.2947>
- Aoude LG, Pritchard AL, Robles-Espinoza CD, et al. Nonsense mutations in the shelterin complex genes ACD and TERF2IP in familial melanoma. *J Natl Cancer Inst*. 2015;107(2):dju408. <https://doi.org/10.1093/jnci/dju408>
- Horn S, Figl A, Rachakonda PS, et al. TERT promoter mutations in familial and sporadic melanoma. *Science*. 2013;339(6122):959-961. <https://doi.org/10.1126/science.1230062>
- Potrony M, Puig-Butille JA, Ribera-Sola M, et al. POT1 germline mutations but not TERT promoter mutations are implicated in melanoma susceptibility in a large cohort of Spanish melanoma families. *Br J Dermatol*. 2019;181(1):105-113. <https://doi.org/10.1111/bjd.17443>
- Calvete O, Garcia-Pavia P, Dominguez F, et al. The wide spectrum of POT1 gene variants correlates with multiple cancer types. *Eur J Hum Genet*. 2017;25(11):1278-1281. <https://doi.org/10.1038/ejhg.2017.134>
- Lim TL, Lieberman DB, Davis AR, et al. Germline POT1 variants can predispose to myeloid and lymphoid neoplasms. *Leukemia*. 2022;36(1):283-287. <https://doi.org/10.1038/s41375-021-01335-w>
- Haugh AM, Njauw CN, Bublely JA, et al. Genotypic and phenotypic features of BAP1 cancer syndrome: a report of 8 new families and review of cases in the literature. *JAMA Dermatol*. 2017;153(10):999-1006. <https://doi.org/10.1001/jamadermatol.2017.2330>
- Wiesner T, Obenauf AC, Murali R, et al. Germline mutations in BAP1 predispose to melanocytic tumors. *Nat Genet*. 2011;43(10):1018-1021. <https://doi.org/10.1038/ng.910>
- Sargen MR, Calista D, Elder DE, et al. Histologic features of melanoma associated with germline mutations of CDKN2A, CDK4, and POT1 in melanoma-prone families from the United States, Italy, and Spain. *J Am Acad Dermatol*. 2020;83(3):860-869. <https://doi.org/10.1016/j.jaad.2020.03.100>
- Linne H, Yasaei H, Marriott A, et al. Functional role of SETD2, BAP1, PARP-3 and PBRM1 candidate genes on the regulation of hTERT gene expression. *Oncotarget*. 2017;8(37):61890-61900. <https://doi.org/10.18632/oncotarget.18712>
- Sargen MR, Kanetsky PA, Newton-Bishop J, et al. Histologic features of melanoma associated with CDKN2A genotype. *J Am Acad Dermatol*. 2015;72(3):496-507.e7. <https://doi.org/10.1016/j.jaad.2014.11.014>
- Taylor NJ, Handorf EA, Mitra N, et al. Phenotypic and histopathological tumor characteristics according to CDKN2A mutation status among affected members of melanoma families. *J Invest Dermatol*. 2016;136(5):1066-1069. <https://doi.org/10.1016/j.jid.2016.01.009>
- Elder DE, Bastian BC, Cree IA, Massi D, Scolyer RA. The 2018 World Health Organization classification of cutaneous, mucosal, and uveal melanoma: detailed analysis of 9 distinct subtypes defined by their evolutionary pathway. *Arch Pathol Lab Med*. 2020;144(4):500-522. <https://doi.org/10.5858/arpa.2019-0561-RA>
- Vest BE, Harview CL, Liu V, et al. Cutaneous melanocytic tumor with CRTC1::TRIM11 fusion and prominent epidermal involvement: a case report. *J Cutan Pathol*. 2022;49(12):1025-1030. <https://doi.org/10.1111/cup.14287>