

UC Merced

UC Merced Previously Published Works

Title

Permanent draft genome of *Thiobacillus thioparus* DSM 505T, an obligately chemolithoautotrophic member of the Betaproteobacteria

Permalink

<https://escholarship.org/uc/item/02q3p4vv>

Journal

Environmental Microbiome, 12(1)

ISSN

2524-6372

Authors

Hutt, Lee P

Huntemann, Marcel

Clum, Alicia

et al.

Publication Date

2017

DOI

10.1186/s40793-017-0229-3

Peer reviewed

SHORT GENOME REPORT

Open Access



Permanent draft genome of *Thiobacillus thioparus* DSM 505^T, an obligately chemolithoautotrophic member of the *Betaproteobacteria*

Lee P. Hutt^{1,2}, Marcel Huntemann³, Alicia Clum³, Manoj Pillay³, Krishnaveni Palaniappan³, Neha Varghese³, Natalia Mikhailova³, Dimitrios Stamatis³, Tatiparthi Reddy³, Chris Daum³, Nicole Shapiro³, Natalia Ivanova³, Nikos Kyrpides³, Tanja Woyke³ and Rich Boden^{1,2*} 

Abstract

Thiobacillus thioparus DSM 505^T is one of first two isolated strains of inorganic sulfur-oxidising *Bacteria*. The original strain of *T. thioparus* was lost almost 100 years ago and the working type strain is Culture C^T (=DSM 505^T = ATCC 8158^T) isolated by Starkey in 1934 from agricultural soil at Rutgers University, New Jersey, USA. It is an obligate chemolithoautotroph that conserves energy from the oxidation of reduced inorganic sulfur compounds using the Kelly-Trudinger pathway and uses it to fix carbon dioxide. It is not capable of heterotrophic or mixotrophic growth. The strain has a genome size of 3,201,518 bp. Here we report the genome sequence, annotation and characteristics. The genome contains 3,135 protein coding and 62 RNA coding genes. Genes encoding the transaldolase variant of the Calvin-Benson-Bassham cycle were also identified and an operon encoding carboxysomes, along with Smith's biosynthetic horseshoe *in lieu* of Krebs' cycle *sensu stricto*. Terminal oxidases were identified, *viz.* cytochrome *c* oxidase (*ccb3*, EC 1.9.3.1) and ubiquinol oxidase (*bd*, EC 1.10.3.10). There is a partial *sox* operon of the Kelly-Friedrich pathway of inorganic sulfur-oxidation that contains *soxXYZAB* genes but lacking *soxCDEF*, there is also a lack of the DUF302 gene previously noted in the *sox* operon of other members of the '*Proteobacteria*' that can use trithionate as an energy source. In spite of apparently not growing anaerobically with denitrification, the *nar*, *nir*, *nor* and *nos* operons encoding enzymes of denitrification are found in the *T. thioparus* genome, in the same arrangements as in the true denitrifier *T. denitrificans*.

Keywords: *Thiobacillus thioparus*, *Betaproteobacteria*, Sulfur oxidation, Chemolithoautotroph, Carboxysome, Denitrification

Introduction

In 1902, *Thiobacillus thioparus* was one of the first two obligately chemolithoautotrophic sulfur-oxidizing *Bacteria* to be isolated, (along with what is now *Halothiobacillus neapolitanus*), and was named in 1904 [1, 2]. The original isolates were lost, but *Thiobacillus thioparus* is now the type species of the genus (a member of the

Betaproteobacteria) and the type strain is an isolate from Starkey (1934) (= Culture C^T = DSM 505^T = ATCC 8158^T = CIP 104484^T = JCM 3859^T = NBRC 103402^T) [3, 4]. Originally, the characteristic of utilizing inorganic sulfur compounds as an energy source was thought to be a taxonomic trait unique to *Thiobacillus* and at its height the genus contained in at least 32 different 'species' [4]. With phylogenetic methods however, many of these strains have since been reassigned to different, often new genera with *T. thioparus* DSM 505^T being one of four species with validly published names left in the genus. Compared to other inorganic sulfur-oxidisers, surprisingly little research has been conducted on *T.*

* Correspondence: rich.boden@plymouth.ac.uk

¹School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK

²Sustainable Earth Institute, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK

Full list of author information is available at the end of the article

thioparus DSM 505^T in terms of physiology and biochemistry or genetics, possibly due to the low growth yields of this species when compared to *T. denitrificans* and *T. aquaesulis* [5, 6] making it more challenging to study. It may, however, give extended and contrasted insights into autotrophic sulfur-oxidation in the *Bacteria*. It was selected for genome sequencing as part of the Department of Energy DOE-CSP 2012 initiative – as type species of a genus.

Organism information

Classification and features

Thiobacillus thioparus DSM 505^T was isolated from sandy loam soil from the New Jersey Agricultural Experimental Station planted with unspecified vegetable crops using 20mM thiosulfate as sole energy source in basal medium at pH 8.5 by Starkey (1934) [3] and is currently one of four species with validly published names within the genus. It forms small white colonies of 1–3 mm diameter after 2–3 days that turn pink or brown with age and which become coated with yellow-ish elementary sulfur. When grown in liquid media, finely divided (white) elementary sulfur is formed during early stages of growth, particularly when thiosulfate is used as the energy source. This disappears when growth approaches stationary phase. Thiosulfate is oxidized stoichiometrically to tetrathionate after 24 h accompanied by a rise in pH, characteristic of the Kelly-Trudinger pathway. Tetrathionate is subsequently oxidized to sulfate with culture pH falling to pH 4.8 by stationary phase. During continuous culture, no intermediates are detected in the medium once steady-state has been established. If the dilution rate of a thiosulfate limited chemostat is increased, a large production of elementary sulfur is observed, which disappears as a new steady-state is established at the faster dilution rate. General features of *T. thioparus* DSM 505^T are summarized in Table 1. A phylogenetic tree based on the 16S rRNA gene sequence showing this organisms position within the *Betaproteobacteria* and rooted with *Thermithiobacillus tepidarius* is given in Fig. 1.

Cells are 1.5–2.0 by 0.6 to 0.8 μm and stain Gram negative. They are motile by means of a single polar flagellum between 6 and 10 μm in length, as shown in Fig. 2. The dominant respiratory quinone is ubiquinone-8 [7–10] and fix carbon dioxide using the Calvin-Benson-Bassham cycle at the expense of inorganic sulfur oxidation. Cells accumulate polyphosphate ('volutin') granules in batch culture and to a lesser extent in an energy-limited chemostat [8]. Carboxysomes are present in cells regardless of growth conditions employed (Fig. 1), but are apparently not found in *T. denitrificans* cells, at least not under growth conditions employed in previous studies [9]. *T. thioparus* can only use oxygen as a terminal

electron acceptor – nitrate, nitrite, thiosulfate, sulfate, elementary sulfur and ferric iron do not support growth as terminal electron acceptors. The genomic DNA G + C content has been estimated using the thermal denaturation [11] at 61–62 mol% [7, 10]. *T. thioparus* DSM 505^T does not grow on any organic carbon compound tested, including sugars (glucose, ribose, fructose, sucrose), intermediates of Krebs' cycle (citrate, succinate, fumarate, malate, oxaloacetate), carboxylates (glycolate, formate, acetate, propionate, pyruvate), one-carbon compounds (monomethylamine, dimethylamine, trimethylamine, methanol, methane), structural amino acids (all 20), substituted thiophenes (thiophene-2-carboxylate, thiophene-3-carboxylate) or complex media (yeast extract, nutrient broth, brain-heart infusion, Columbia sheep or horse blood agar, chocolate agar). Energy sources that support autotrophic growth of DSM 505^T include thiosulfate, trithionate, tetrathionate, pentathionate, hexathionate, thiocyanate and dithionate. Some *bone fide* strains of *T. thioparus* (Tk-m and E6) grow autotrophically on carbon disulfide, dimethylsulfide, dimethyldisulfide and Admidate (*O,O*-dimethylphosphoramidothioate) [4], but it is not known if the type strain DSM 505^T is capable of this. Autotrophic growth is not supported by Fe(II), Mn(II), Cu(I), U(IV), sulfite, dimethylsulfoxide, dimethylsulfone, pyrite or formate. During batch growth on thiosulfate the intermediate production of tetrathionate is observed during early stages of growth, indicative of the Kelly-Trudinger pathway [12]. Compared to the other two members of the genus [5, 6], *T. thioparus* DSM 505^T has relatively low growth yields on thiosulfate (Hutt and Boden, *manuscript in preparation*) which may give insight into the physiological variances of Kelly-Trudinger pathway organisms even within one genus.

Genome sequencing information

Genome project history

This organism was selected for sequencing on the basis of its role in sulfur cycling, physiological, biochemical, evolutionary and biogeochemical importance, and is part of the GEBA-KMG project at the U.S. Department of Energy JGI. The genome project is deposited in the Genomes OnLine Database [13] and a high-quality permanent draft genome sequence in IMG (the annotated genome is publically available in IMG under Genome ID 2515154076) [14]. Sequencing, finishing and annotation were performed by the JGI using state of the art sequencing technology [15]. A summary of the project information is given in Table 2.

Growth conditions and genomic DNA preparation

T. thioparus DSM 505^T DNA was obtained from Dr Hans-Peter Klenk at the DSMZ, having been grown on

Table 1 Classification and general features of *Thiobacillus thioparus* DSM 505^T according to MIGS recommendations [43]

MIGS ID	Property	Term	Evidence code ^a
	Classification	Domain <i>Bacteria</i>	TAS [44]
		Phylum ' <i>Proteobacteria</i> '	TAS [45]
		Class <i>Betaproteobacteria</i>	TAS [46]
		Order <i>Hydrogenophilales</i>	TAS [47]
		Family <i>Hydrogenophilaceae</i>	TAS [48]
		Genus <i>Thiobacillus</i>	TAS [2, 3]
		Species <i>Thiobacillus thioparus</i>	TAS [2, 3]
		(Type) strain: DSM 505 ^T	TAS [49]
		Gram stain	<i>Negative</i>
	Cell shape	<i>Rod</i>	TAS [2, 3]
	Motility	<i>Motile</i>	TAS [2, 3]
	Sporulation	<i>None</i>	TAS [3]
	Temperature range	<i>N.D.</i>	NAS
	Optimum temperature	28 °C	TAS [3, 50]
	pH range; Optimum	<i>N.D.; 7.0</i>	NAS
	Carbon source	<i>Carbon dioxide</i>	TAS [3, 50]
MIGS-6	Habitat	<i>Agricultural soil (sandy loam)</i>	TAS [3]
MIGS-6.3	Salinity	<i>N.D.</i>	NAS
MIGS-22	Oxygen requirement	<i>Aerobic</i>	TAS [3, 50]
MIGS-15	Biotic relationship	<i>Free-living</i>	TAS [3, 50]
MIGS-14	Pathogenicity	<i>Non-pathogen</i>	NAS
MIGS-4	Geographic location	<i>New Jersey, United States of America</i>	TAS [3]
MIGS-5	Sample collection	<i>1934</i>	TAS [3]
MIGS-4.1	Latitude	<i>40° 28' 57" N</i>	TAS [3]
MIGS-4.2	Longitude	<i>74° 26' 14" E</i>	TAS [3]
MIGS-4.4	Altitude	<i>28 m</i>	TAS [3]

^a Evidence codes - *IDA* Inferred from Direct Assay, *TAS* Traceable Author Statement (i.e., a direct report exists in the literature), *NAS* Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [51, 52]

basal salts medium pH 6.6, supplemented with 40 mM thiosulfate as the sole energy source (DSM Medium 36), under air at 26 °C for 72 h. DNA was extracted using the JETFLEX Genomic DNA Purification Kit from Genomed (Löhrne, Germany) into TE Buffer. Quality was checked by agarose gel electrophoresis.

Genome sequencing and assembly

The draft genome of *Thiobacillus thioparus* DSM 505^T was generated at the DOE JGI using the Illumina technology [16]. An Illumina standard shotgun library was constructed and sequenced using the Illumina HiSeq 2000 platform which generated 11,161,382 reads totalling 1,674.2 Mbp. All general aspects of library construction and sequencing performed at the JGI can be found at <http://www.jgi.doe.gov>. All raw Illumina sequence data was passed through DUK, a filtering program developed at JGI, which removes known Illumina sequencing and library preparation artifacts (Mingkun L, Copeland A, Han J, Unpublished). Following steps were then performed for assembly: (1) filtered Illumina reads were assembled using Velvet (version 1.1.04) (Mingkun, L., Unpublished), (2) 1–3 Kbp simulated paired end reads were created from Velvet contigs using wgsim [17], (3) Illumina reads were assembled with simulated read pairs using Allpaths-LG (version r41043) [18]. Parameters for assembly steps were: 1) Velvet (velveth: 63 –shortPaired and velvetg: –very clean yes –export-Filtered yes –min contig lgth 500 –scaffolding no –cov cutoff 10) 2) wgsim (–e 0–1 100–2 100 –r 0 –R 0 –X 0) 3) Allpaths-LG (PrepareAllpathsInputs: PHRED 64 = 1 PLOIDY = 1 FRAG COVERAGE = 125 JUMP COVERAGE = 25 LONG JUMP COV = 50, RunAllpathsLG: THREADS = 8 RUN = std shredpairs TARGETS = standard VAPI WARN ONLY = True OVERWRITE = OVERWRITE = True). The final draft assembly contained 20 contigs in 20 scaffolds. The total size of the genome is 3.2 Mbp and the final assembly is based on 392.6 Mbp of Illumina data, which provides an average 122.7× coverage of the genome.

Genome annotation

Genes were identified using Prodigal [19], followed by a round of manual curation using GenePRIMP [20] for finished genomes and Draft genomes in fewer than 10 scaffolds. The predicted CDSs were translated and used to search the NCBI nonredundant database, UniProt, TIGR-Fam, Pfam, KEGG, COG, and InterPro databases. The tRNAScanSE tool [21] was used to find tRNA genes, whereas ribosomal RNA genes were found by searches against models of the ribosomal RNA genes built from SILVA [22]. Other non-coding RNAs such as the RNA components of the protein secretion complex and the RNase P were identified by searching the genome for the corresponding Rfam profiles using INFERNAL [23]. Additional gene prediction analysis and manual functional annotation was performed within the IMG platform [24] developed by the JGI, Walnut Creek, CA, USA [25, 26].

Genome properties

The genome of *T. thioparus* DSM 505^T is 3,201,518 bp-long with a 62.3 mol% G + C content (Table 3). Of the 3,197 predicted genes, 3,135 were protein-coding genes




Fig. 1 Maximum-likelihood phylogenetic tree based on MUSCLE alignment of 16S rRNA gene sequences of the genus *Thiobacillus* and the closely related members of the *Betaproteobacteria*. Type strains of each species are used and only species with validly published names are shown. Sequences pertaining to organisms for which a publically available genome sequence exists are underlined. Accession numbers for the GenBank database are in parentheses. Alignment and tree were constructed in MEGA 6 [53]. Tree was drawn using the Tamura-Nei model for maximum-likelihood trees [54]. Values at nodes are based on 5,000 bootstrap replicates, with values <70% omitted. Scale-bar indicates 2 substitutions per 100. *Thermithiobacillus tepidarius* DSM 3134^T from the *Acidithiobacillia* is used as the outgroup

and 62 were RNA genes. A total of 2,597 genes (81.23%) have predicted function. A total of 538 (16.83%) were identified as pseudogenes – the remainder annotated as hypothetical proteins. The properties and the statistics of the genome are given in Table 3, the distribution of genes into COG functional categories is given in Table 4. The genome is the second largest genome of obligate chemolithoautotrophs sequenced to date and is 89% (bp/bp) or 90% (protein coding genes/protein coding genes) of the size of that of *T. denitrificans* DSM 12475^T [12].

Insights from the genome sequence

As an obligate autotroph, it would be expected for a complete Calvin-Benson-Bassham cycle and, *in lieu* of

Krebs' cycle, Smith's biosynthetic horseshoe [12, 27–29]. Smith's horseshoe representing a very near-complete Krebs' cycle was identified, in which citrate synthase (EC 2.3.3.16), aconitase (EC 4.2.1.3), isocitrate dehydrogenase (NADP⁺, EC 1.1.1.42), succinyl coenzyme A synthase (ADP-forming, EC 2.6.1.5), succinate dehydrogenase (EC 1.3.5.1) and malate dehydrogenase (oxaloacetate decarboxylating, NADP⁺, EC 1.1.1.40) genes are present, but fumarase (EC 4.2.1.2) and the E3 subunit of α -ketoglutarate dehydrogenase (NAD⁺, EC 1.2.4.2) genes are absent – E1 and E2 are present. This is a comparatively large version of Smith's horseshoe [29], which varies between Classes of the '*Proteobacteria*', for example, genes for fumarase, succinate dehydrogenase and the E1 subunit of




Fig. 2 Transmission electron micrographs of *T. thiooparus* DSM 505^T cells obtained from a thiosulfate-limited chemostat (20 mM, $D=0.07\text{ h}^{-1}$) visualized in a JEOL JEM-1400Plus transmission electron microscope, operating at 120 kV. **a** Negatively stained cells. Cells were applied to Formvar[®] and carbon coated copper grid before washing with saline and staining in 50 mM uranyl acetate for 5 mins and washing again. **b** Sectioned cells showing the presence of an electron dense polyphosphate (volutin) granule and numerous polyhedral carboxysomes that are paler in comparison

Table 2 Project information

MIGS ID	Property	Term
MIGS 31	Finishing quality	Improved High-Quality Draft
MIGS-28	Libraries used	Illumina Standard PE
MIGS 29	Sequencing platforms	Illumina
MIGS 31.2	Fold coverage	122.7
MIGS 30	Assemblers	Allpaths/Velvet
MIGS 32	Gene calling method	NCBI Prokaryotic Genome Annotation Pipeline
	Locus Tag	B058
	Genbank ID	ARDU00000000
	GenBank Date of Release	April 16th, 2013
	GOLD ID	Ga0025551
	BIOPROJECT	PRJNA169730
MIGS 13	Source Material Identifier	DSM 505 ^T
	Project relevance	GEBA-KMG

α -ketoglutarate dehydrogenase were recently found to be missing from *Thermithiobacillus tepidarius* DSM 3134^T [12, 30, 31] of the *Acidithiobacillia*. Activities of Krebs' cycle enzymes assayed by Smith et al. [29] in cell-free extracts of *T. thioparus* found activities of all enzymes except for α -ketoglutarate dehydrogenase, indicating the E1 and E2 subunits alone were not sufficient for activity. Interestingly, Smith detected fumarase activity; however, this strain of *T. thioparus* was isolated by the authors themselves and was not DSM 505^T, so there may be further variation of Smith's horseshoe at strain level. With many species having been erroneously classified as strains of *T. thioparus* in the past that have subsequently been proven to belong to other genera [4] it is

Table 3 Genome statistics

Attribute	Value	% of Total
Genome size (bp)	3,201,518	100.00
DNA coding (bp)	2,937,381	91.75
DNA G + C (bp)	1,994,510	62.30
DNA scaffolds	18	100.00
Total genes	3,197	100.00
Protein coding genes	3,135	98.06
RNA genes	62	1.94
Pseudo genes	538	16.83
Genes in internal clusters	267	8.35
Genes with function prediction	2,597	81.23
Genes assigned to COGs	2,258	70.63
Genes with Pfam domains	2,700	84.45
Genes with signal peptides	376	11.76
Genes with transmembrane helices	767	23.99
CRISPR repeats	2	0.04

Table 4 Number of genes associated with general COG functional categories

Code	Value	%age	Description
J	205	8.2	Translation, ribosomal structure and biogenesis
A	1	0.0	RNA processing and modification
K	120	4.8	Transcription
L	83	3.3	Replication, recombination and repair
B	1	0.0	Chromatin structure and dynamics
D	38	1.5	Cell cycle control, Cell division, chromosome partitioning
V	61	2.4	Defense mechanisms
T	156	6.2	Signal transduction mechanisms
M	225	9.0	Cell wall/membrane biogenesis
N	101	4.0	Cell motility
U	56	2.2	Intracellular trafficking and secretion
O	144	5.8	Posttranslational modification, protein turnover, chaperones
C	207	8.3	Energy production and conversion
G	86	3.4	Carbohydrate transport and metabolism
E	153	6.1	Amino acid transport and metabolism
F	63	2.5	Nucleotide transport and metabolism
H	144	5.8	Coenzyme transport and metabolism
I	76	3.0	Lipid transport and metabolism
P	206	8.2	Inorganic ion transport and metabolism
Q	37	1.5	Secondary metabolites biosynthesis, transport and catabolism
R	165	6.6	General function prediction only
S	143	5.7	Function unknown
-	939	29.4	Not in COGs

The total is based on the total number of protein coding genes in the genome

also plausible that Smith's strain was from another specie, genus or even Class. The E1, E2 and E3 subunit genes for α -ketoglutarate dehydrogenase were identified in the genome of *T. denitrificans* ATCC 25259^T [32], though enzyme activity was also absent in this strain [29, 33]. It is important to stress that the full suite of enzymes of Krebs' cycle or Smith's horseshoe have never been assayed in *T. thioparus* DSM 505^T - clearly this is needed in order to identify if genomic data reflect true activities in vivo [27].

A complete Calvin-Benson-Bassham cycle is present, with a single copy of the large (*cbbL*) and small (*cbbS*) form I RuBisCO (EC 4.1.1.39) subunits and, owing to the presence of a transaldolase (EC 2.2.1.2) and absence of a sedoheptulose-1,7-bisphosphatase (EC 3.1.3.37) gene, we can conclude that it uses the transaldolase-

variant Calvin-Benson-Bassham cycle [34]. Adjacent to these genes are *cbbO* and *cbbQ*, consistent with form IA_q RuBisCO [35], which is canonically cytoplasmic, rather than carboxysomal. A cluster of 11 genes that encode carboxysome shell proteins and a carboxysome carbonic anhydrase were also identified. This cluster is located on the forward strand while on the reverse strand the RuBisCO cluster is located *c.*9 kb upstream, between which is a divergently transcribed transcriptional regulator (*cbbR*) [32]. Further evidence of carboxysome expression can be seen in transmission electron micrographs (Fig. 1). The carboxysome gene cluster (experimental data would be required to demonstrate if it is an operon or not) in *T. thioparus* DSM 505^T does not start with *cbbLS*, (as would be found in *Halothiobacillus*, *Acidithiobacillus* and *Thermithiobacillus* spp. [36]) - the lack of these genes has also been observed in *T. denitrificans* ATCC 25259^T [32] and may be a diagnostic property of this genus.

Whilst *T. thioparus* DSM 505^T cannot grow anaerobically with denitrification but *T. denitrificans* does, putative operons encoding the nitrate reductase (*nar*), nitrite reductase (*nir*), nitric oxide reductase (*nor*) and nitrous oxide reductase (*nos*) proteins of canonical denitrification are found in both *T. denitrificans* and *T. thioparus* - this may indicate a potential for the latter to grow with denitrification albeit not under conditions previously employed - for example, it may occur only under micro-oxic conditions rather than fully anoxic conditions, alternatively, some unknown factor may prevent detection of nitrate and/or the expression of these genes. No evidence for any alternative denitrification pathways [37] was found

With regard to respiration, five cytochromes *c*₅₅₃, one cytochrome *c*₅₅₆ and three cytochrome *b* were present. The two high-affinity terminal oxidases were both found - with multiple gene copies of cytochrome *c* oxidase (*cbb*₃, EC 1.9.3.1) and a single copy of cytochrome *bd*-type quinol oxidase (EC 1.10.3.10). It is worth noting that *T. denitrificans* also possess the *aa*₃ variant of cytochrome *c* oxidase (EC 1.9.3.1), which may permit it greater metabolic diversity, for example, growth under more variable oxygen partial pressures - this could potentially explain why *T. thioparus* cannot denitrify under anoxic conditions, even though it possesses the operons encoding enzymes of denitrification. It is known that microaerophilic organisms usually employ *cbb*₃ or *bd*-type high-affinity terminal oxidases [37] and this may further evidence that *T. thioparus* and *T. denitrificans* may grow under such conditions, potentially with denitrification, though we cannot find any studies that demonstrate this in vivo thus far.

Extended insights

Enzymes of the Kelly-Trudinger pathway remain poorly understood and many of the genes are yet to be

identified [12]. The oxidation of thiosulfate to tetrathionate takes place *via* a cytochrome *c*-linked thiosulfate dehydrogenase (EC 1.8.2.2), one gene (*tsdA*) for which was identified in *Allochrochromatium vinosum*, a member of the *Gammaproteobacteria* [38]. However, *tsdA* is not present in *T. thioparus* DSM 505^T supporting the hypothesis that more than one thiosulfate dehydrogenase is present in the 'Proteobacteria' or could vary at Class level. Numerous Kelly-Friedrich pathway genes (*soxXYZAB*) were present but the remaining of the conserved *soxTRS-VW-XYZABCDEFGH* genes being absent [39–41]. In *Paracoccus* spp. (*Alphaproteobacteria*) *soxYZ* encodes a protein complex that binds thiosulfate *via* a cysteine residue in the initial stage of the Kelly-Friedrich pathway while *soxXA* encode cytochromes *c*₅₅₁ and *c*_{552.5} which capture two electrons from thiosulfate oxidation. Finally *soxB* encodes a hydrolase that removes the terminal sulfone group as sulfate. Missing the SoxCD or sulfur dehydrogenase protein from the multi-enzyme system would leave a sulfur atom attached to the SoxYZ residue, preventing the action of SoxB to liberate this residue to act further with another thiosulfate molecule. An unidentified protein may participate in the release of these sulfur atoms and potentially may explain the deposits of sulfur seen during initial stages of growth. This does not explain the production of tetrathionate which still would require the activity of thiosulfate dehydrogenase. If *soxXYZAB* genes are being expressed then they may be required as a functional part of the Kelly-Trudinger pathway. The action of Sox proteins (if any) in *T. thioparus* DSM 505^T in conjunction and potentially collaboration with additional Kelly-Trudinger pathway proteins would undoubtedly be essential in resolving its chemolithoautotrophic metabolism, which remains poorly understood. An unidentified gene encoding a putative DUF302-family protein is present between *soxA* and *soxB* genes of *Thermithiobacillus tepidarius* DSM 3134^T, *Acidithiobacillus thiooxidans* ATCC 19377^T and *Acidithiobacillus caldus* ATCC 51756^T, and *Thiohalorhabdus denitrificans* DSM 15699^T, the function of which may be important in the Kelly-Trudinger pathway [12]; however, DUF302 is not present on the *sox* operon of *T. thioparus* DSM 505^T, although six unidentified genes annotated as DUF302 family proteins are present elsewhere in the genome. The *soxEF* genes encode a flavo-cytochrome *c* sulfide dehydrogenase (EC 1.8.2.3) and are both found in the *T. denitrificans* genome, separate from the main *sox* gene cluster, but they are not found in *T. thioparus*, whereas dissimilatory sulfite reductase (*dsr*) genes are found in both genomes, as are adenyl sulfite reductase (*aprAB*, EC 1.8.99.2) genes, the presence of which has also been confirmed in *T. aquaesulis* [42].

It was previously noted that 5.9% of the genome (178 genes) of *Thermithiobacillus tepidarius* DSM 3134^T were potential horizontally transferred genes from

Thiobacillus thioparus, *Thiobacillus denitrificans* and *Sulfuricella denitrificans* of the *Betaproteobacteria* [12]; 96 of these genes came from the two *Thiobacillus* spp. However, very little gene transfer has taken place from members of *Acidithiobacillia* to *T. thioparus* DSM 505^T with only 6 genes from *Ttb. tepidarius* DSM 3134^T, 4 from *Acidithiobacillus* spp. This is perhaps not surprising due to the thermophilic and acidophilic nature of these three *Acidithiobacillia* compared to the mesophilic requirements of *T. thioparus* and the unlikelihood of these species co-inhabiting the same environments. A far larger portion (143 genes; 4.82%) of genes were attributed to transfer from members of *Gammaproteobacteria*, many of which grow at more neutral pH and mesophilic temperatures. There was no distinct pattern of any particular metabolism pathways or resistances etc. being encoded by these potentially transferred genes.

Conclusions

The genome of *Thiobacillus thioparus* DSM 505^T gives insights into many aspects of its physiology, biochemistry and evolution. This organism uses the transaldolase variant of the Calvin-Benson-Bassham cycle and produces carboxysomes for carbon dioxide fixation, evident from both the genome and transmission electron microscopy. The expression of both carboxysomes and Ru-BisCO may be regulated by the same divergently transcribed transcriptional regulator. Smith's biosynthetic horseshoe is present *in lieu* of Krebs' cycle *sensu stricto*, but this is unusually large as only 2 genes are missing, though *T. denitrificans* [32] is only missing 1 – this may in part explain the heterotrophic growth of *T. aquaesulis* since just one additional gene would convert Smith's horseshoe into a functional version of Krebs' cycle [6]. Many inorganic sulfur-oxidation genes of the *sox* cluster were found but *soxC*, *soxD*, *soxE* and *soxF* are absent. The *tsdA* gene for a thiosulfate dehydrogenase identified in *Allochromatium vinosum* is absent and confirmation of the presence of a different thiosulfate dehydrogenase enzyme and gene will require further study. The genome sequence will enable evolutionary studies into the nature of *Thiobacillus* and chemolithoautotrophs in general, in particular reigniting the obligate *versus* facultative and the autotrophic *versus* mixotrophic debates that have been largely absent from the literature in recent years, but genome sequences becoming available will now answer many questions proposed over 10 years ago [27, 29].

Abbreviations

ATCC: American type culture collection; BLAST: Basic linear alignment search tool; CIP: Collection de L'institut Pasteur; COG: Clusters of orthologous groups; DSMZ: Leibniz-Institut DSMZ - Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH; GEBA: Genomic encyclopedia of *Bacteria* and *Archaea*; IMG: Integrated microbial genomes database; JCM: Japan collection of microorganisms; JGI: Joint Genome Institute;

KMG: 1,000 microbial genomes project; NBRC: NITE biological resource center; NCBI: National center for biotechnology information; RuBisCO: Ribulose-1,5-bisphosphate carboxylase/oxygenase

Acknowledgements

We acknowledge Dr Hans-Peter Klenk at the DSMZ for the provision of genomic DNA.

Funding

The sequencing and annotation was performed under the auspices of the United States Department of Energy JGI, a DOE Office of Science User Facility and is supported by the Office of Science of the United States Department of Energy under Contract Number DE-AC02-05CH11231. The authors wish to acknowledge the School of Biological and Marine Sciences, University of Plymouth for studentship funding to LH that supported the analysis of the genome and to the Royal Society Research Grant RG120444 awarded to RB to support the analysis of this genome.

Authors' contributions

LPH and RB analysed and mined the genome data in public databases for genes of interest and performed BLASTn/BLASTp searches to verify and validate the annotation etc. and made comparisons of the sulfur oxidation, denitrification etc. operons with those in other organisms. RB constructed the phylogenetic tree. LPH grew the organism and performed electron microscopy at the Electron Microscopy Unit, University of Plymouth. All other authors contributed to the sequencing, assembly and annotation of the genome sequence. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Author details

¹School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK. ²Sustainable Earth Institute, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK. ³DOE Joint Genome Institute, 94598 Walnut Creek, CA, USA.

Received: 12 October 2016 Accepted: 8 January 2017

Published online: 19 January 2017

References

- Nathanson A. Über eine neue Gruppe von Schwefelbakterien und ihren Stoffwechsel. *Mitt zool Stn Neapel*. 1902;15:655–80.
- Beijerinck MW. Ueber die Bakterien, welche sich im Dunkeln mit Kohlensäure als Kohlenstoffquelle ernähren können. *Centralbl Bakteriol Parasitenkd Infektionskr Hyg Abt II*. 1904;11:592–599.
- Starkey RL. Isolation of some bacteria which oxidise thiosulfate. *Soil Sci*. 1934;39:197–219.
- Boden R, Cleland D, Green PN, Katayama Y, Uchino Y, Murrell JC, Kelly DP. Phylogenetic assessment of culture collection strains of *Thiobacillus thioparus*, and definitive 16S rRNA gene sequences for *T. thioparus*, *T. denitrificans*, and *Halothiobacillus neapolitanus*. *Arch Microbiol*. 2011;194:187–95.
- Justin P, Kelly DP. Growth kinetics of *Thiobacillus denitrificans* in anaerobic and aerobic chemostat culture. *J Gen Microbiol*. 1978;107:123–30.
- Wood AP, Kelly DP. Isolation and physiological characterisation of *Thiobacillus aquaesulis* sp. nov., a novel facultatively autotrophic moderate thermophile. *Arch Microbiol*. 1988;149:339–43.
- Katayama-Fujimura Y, Tsuzaki N, Kuraishi H. Ubiquinone, fatty acid and DNA base composition determination as a guild to the taxonomy of the genus *Thiobacillus*. *J Gen Microbiol*. 1982;128:1599–611.
- Katayama-Fujimura Y, Enokizono Y, Kaneko T, Kuraishi H. Deoxyribonucleic acid homologies among species of the genus *Thiobacillus*. *J Gen Appl Microbiol*. 1983;29:287–95.
- Katayama-Fujimura Y, Tsuzaki N, Hirata A, Kuraishi H. Polyhedral inclusion bodies (carboxysomes) in *Thiobacillus* species with reference to the taxonomy of the Genus *Thiobacillus*. *J Gen Appl Microbiol*. 1984;30:211–22.
- Kelly DP, Wood AP. Confirmation of *Thiobacillus denitrificans* as a species of the genus *Thiobacillus*, in the β -subclass of the Proteobacteria, with strain NCIMB 9548 as the type strain. *Int J Syst Evol Microbiol*. 2000;50:547–50.

11. Marmur J, Doty P. Determination of the base composition of deoxyribonucleic acid from its thermal denaturation temperature. *J Mol Biol*. 1962;5:109–18.
12. Boden R, Hutt LP, Huntemann M, Clum A, Pillay M, Palaniappan K, Varghese N, Mikhailova N, Stamatis D, Reddy T, Ngan CY, Daum C, Shapiro N, Markowitz V, Ivanova N, Woyke T, Kyrpides N. Permanent draft genome of *Thermithiobacillus tepidarius* DSM 3134^T, a moderately thermophilic, obligately chemolithoautotrophic member of the *Acidithiobacillia*. *Standards in Genomic Sciences*. 2016; 11. DOI: 10.1186/s40793-016-0188-0
13. Pagani I, Liolios K, Jansson J, Chen IM, Smirnova T, Nosrat B, et al. The Genomes OnLine Database (GOLD) v. 4: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res*. 2012;40:D571–9.
14. Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M, et al. IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic Acids Res*. 2014;42:D560–7.
15. Mavromatis K, Land ML, Brettin TS, Quest DJ, Copeland A, Clum A, et al. The fast changing landscape of sequencing technologies and their impact on microbial genome assemblies and annotation. *PLoS One*. 2012;7:e48837.
16. Bennett S. Solexa Ltd. *Pharmacogenomics*. 2004;5(4):433–8.
17. Zerbino D, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res*. 2008;18:821–9.
18. Reads simulator wgsim. [<https://github.com/lh3/wgsim>].
19. Gnerre S, MacCallum I. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *PNAS*. 2011;108(4):1513–8.
20. Hyatt D, Chen GL, Lacascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*. 2010;11:119.
21. Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods*. 2010;7:455–7.
22. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res*. 1997;25:955–64.
23. Pruesse E, Quast C, Fuchs BM, Knittel K, Ludwig W, Peplies J, Glickner FO. SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nuc Acids Res*. 2007;35:2188–7196.
24. INFERNAL. Inference of RNA alignments. [<http://infernal.janelia.org>].
25. The Integrated Microbial Genomes (IMG) platform. [<http://img.jgi.doe.gov>].
26. Markowitz VM, Mavromatis K, Ivanova NN, Chen IMA, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics*. 2009;25:2271–8.
27. Wood AP, Aurikko JP, Kelly DP. A challenge for 21st century molecular biology and biochemistry: what are the causes of obligate autotrophy and methanotrophy. *FEMS Microbiol Rev*. 2004;28:335–52.
28. Peeters TL, Liu MS, Aleem MH. The tricarboxylic acid cycle in *Thiobacillus denitrificans* and *Thiobacillus-A2*. *J Gen Microbiol*. 1970;64:29–35.
29. Smith AJ, London J, Stanier RY. Biochemical basis of obligate autotrophy in blue-green algae and thiobacilli. *J Bacteriol*. 1967;94:972–83.
30. Kelly DP, Wood AP. Reclassification of some species of *Thiobacillus* to the newly designated genera *Acidithiobacillus* gen. nov., *Haloithiobacillus* gen. nov. and *Thermithiobacillus* gen. nov. *Int J Syst Evol Microbiol*. 2000;50:511–6.
31. Hudson C, Williams K, Kelly DP. Definitive assignment by multigenome analysis of the Gammaproteobacterial Genus *Thermithiobacillus* to the Class *Acidithiobacillia*. *Pol J Microbiol*. 2014;63:245–7.
32. Beller HR, Chain PSG, Letain TE, Chakicherla A, Larimer FW, Richardson PM, Coleman MA, Wood AP, Kelly DP. The genome sequence of the obligately chemolithoautotrophic, facultatively anaerobic bacterium *Thiobacillus denitrificans*. *J Bacteriol*. 2006;188:1473–88.
33. Taylor BF, Hoare DS. *Thiobacillus denitrificans* as an obligate chemolithoautotroph. Cell suspension and enzymic studies. *Arch Mikrobiol*. 1971;80:262–76.
34. Anthony C. *The biochemistry of methyloproteobacteria*. 1st ed. London: Academic Press; 1982.
35. Badger MR, Bek EJ. Multiple RuBisCO forms in proteobacteria: their functional significance in relation to CO₂ acquisition by the CBB cycle. *J Exp Bot*. 2008;59:1525–41.
36. Sutter M, Roberts EW, Gonzalez RC, Bates C, Dawoud S, Landry K, Cannon GC, Heinhorst S, Kerfeld CA. Structural characterization of a newly identified component of α -carboxysomes: the AAA+ domain protein CsoCBBQ. *Sci Rep*. 2015;5:16243.
37. Chen J, Strous M. Denitrification and aerobic respiration, hybrid electron transport chains and co-evolution. *Biochim Biophys Acta Bioenergetics*. 2013;1827:136–44.
38. Denkmann K, Grein F, Zigann R, Siemen A, Bergmann J, van Helmont S, Nicolai A, Pereira IAC, Dahl C. Thiosulfate dehydrogenase: a widespread unusual acidophilic c-type cytochrome. *Environ Microbiol*. 2012;14:2673–88.
39. Wodara C, Kostka S, Egert M, Kelly DP, Friedrich CG. (1994) Identification and sequence analysis of the *soxB* gene essential for sulfur oxidation of *Paracoccus denitrificans* GB17. *J Bacteriol*. 1994;176:6188–91.
40. Wodara C, Bardischewsky F, Friedrich CG. Cloning and characterization of sulfite dehydrogenase, two c-type cytochromes, and a flavoprotein of *Paracoccus denitrificans* GB17: essential role of sulfite dehydrogenase in lithotrophic sulfur oxidation. *J Bacteriol*. 1997;179:5014–23.
41. Friedrich CG, Quentmeier A, Bardischewsky F, Rother D, Kraft R, Kostka S, Prinz H. Novel genes coding for lithotrophic sulfur oxidation of *Paracoccus pantotrophus* GB17. *J Bacteriol*. 2000;182:4677–87.
42. Mayer B, Kuever J. Molecular analysis of the distribution and phylogeny of dissimilatory adenosine 5'-phosphosulfate reductase-encoding genes (*aprBA*) among sulfur-oxidizing prokaryotes. *Microbiol (UK)*. 2007;153:3478–98.
43. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol*. 2008;26:541–7.
44. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci U S A*. 1990;87:4576 [PubMed].
45. Garrity GM, Bell JA, Lilburn T, Phylum XIV. *Proteobacteria* phyl. nov. In: Garrity GM, Brenner DJ, Krieg NR, Staley JT, editors. *Bergey's manual of systematic bacteriology*, vol. 2. Second ed. New York: Springer; 2005. p. 1. Part B.
46. Garrity GM, Bell JA, Lilburn T. Class II. *Betaproteobacteria* class. nov. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, editors. *Bergey's Manual of Systematic Bacteriology*, second edition, vol. 2 (The Proteobacteria) part C (The Alpha-, Beta-, Delta-, and Epsilonproteobacteria). New York: Springer; 2005. p. 575.
47. Garrity GM, Bell JA, Lilburn T. Order II. *Hydrogenophilales* ord. nov. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, editors. *Bergey's Manual of Systematic Bacteriology*, second edition, vol. 2 (The Proteobacteria), part C (The Alpha-, Beta-, Delta-, and Epsilonproteobacteria). New York: Springer; 2005. p. 763.
48. Garrity GM, Bell JA, Lilburn T. "Family I. *Hydrogenophilaceae* fam. nov.". In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, editors. *Bergey's Manual of Systematic Bacteriology*, second edition, vol. 2 (The Proteobacteria), part C (The Alpha-, Beta-, Delta-, and Epsilonproteobacteria). New York: Springer; 2005. p. 763.
49. Skerman VBD, McGowan V, Sneath PHA. Approved list of bacterial names. *Int J Syst Evol Microbiol*. 1980;30:225–420.
50. Kelly DP, Wood AP, Stackebrandt E. Genus II. *Thiobacillus* Beijerinck 1904. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM, editors. *The Proteobacteria*, Part C, The Alpha-, Beta-, Delta-, and Epsilonproteobacteria, 2. 2nd ed. New York: Springer; 2005. p. 764–9.
51. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet*. 2000;25:25–9.
52. Guide to GO evidence codes. [<http://www.geneontology.org/GO.evidence.shtml>].
53. Tamura K, Stecher G, Peterson D, Filipitski A, Sumar S. MEGA6: Molecular Evolutionary Genomics Analysis version 6.0. *Mol Biol Evol*. 2013;30:2725–19.
54. Tamura K, Nei M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol*. 1993;10:512–26.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

