

UC Santa Cruz

UC Santa Cruz Previously Published Works

Title

From genomics to functions: preclinical mouse models for understanding oncogenic pathways in prostate cancer.

Permalink

<https://escholarship.org/uc/item/049047qt>

Journal

American Journal of Cancer Research, 9(10)

ISSN

2156-6976

Authors

Yu, Chuan
Hu, Kevin
Nguyen, Daniel
[et al.](#)

Publication Date

2019

Peer reviewed

Review Article

From genomics to functions: preclinical mouse models for understanding oncogenic pathways in prostate cancer

Chuan Yu, Kevin Hu, Daniel Nguyen, Zhu A Wang

Department of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, CA 95064, USA

Received August 13, 2019; Accepted September 10, 2019; Epub October 1, 2019; Published October 15, 2019

Abstract: Next-generation sequencing has revealed numerous genomic alterations that induce aberrant signaling activities in prostate cancer (PCa). Among them are pathways affecting multiple cancer types, including the PI3K/AKT/mTOR, p53, Rb, Ras/Raf/MAPK, Myc, FGF, and Wnt signaling pathways, as well as ones that are prominent in PCa, including alterations in genes of AR signaling, the ETS family, *NKX3.1*, and *SPOP*. Cross talk among the oncogenic pathways can confer PCa resistance to therapy, particularly in advanced tumors, which are castration-resistant or show neuroendocrine features. Various experimental models, such as cancer cell lines, animal models, and patient-derived xenografts and organoids have been utilized to dissect PCa progression mechanisms. Here, we review the current preclinical mouse models for studying the most commonly altered pathways in PCa, with an emphasis on their interplays. We highlight the power of genetically engineered mouse models (GEMMs) in translating genomic discoveries into understanding of the functions of these oncogenic events in vivo. Developing and analyzing PCa mouse models will undoubtedly continue to offer new insights into tumor biology and guide novel rationalized therapy.

Keywords: Prostate, cancer progression, mouse models, castration-resistance, neuroendocrine, PI3K, AR, p53, Wnt, ETS

Introduction

Prostate cancer (PCa) is one of the most common malignancies in men of developed countries and is continuously increasing in developing countries [1]. Prostatic intraepithelial neoplasia (PIN), the precursor lesion to adenocarcinoma, can be present in men of young age. Most patients with early stage PCa undergo active surveillance, or are treated with radiation or radical prostatectomy. Androgen-deprivation therapy was first adopted by Huggins and Hodges in 1941 [2], and remains the first line treatment for advanced PCa. However, the inevitable recurrence of castration-resistant prostate cancer (CRPC) after androgen-deprivation therapy remains a major challenge, as treatment options are limited (**Figure 1**). Tumors that become androgen receptor (AR) independent may develop into neuroendocrine prostate cancer (NEPC), which is one of the most lethal subtypes of PCa [3]. Understanding

the molecular mechanism of PCa progression is crucial for the development of effective therapies.

The prostate gland forms through budding from the urogenital sinus during development [4]. The mature prostate contains ductal structures consisting of a stromal compartment and an epithelium, from which adenocarcinomas arise. The normal prostate epithelium is primarily comprised of stratified basal cells and luminal cells identifiable by specific markers, as well as interspersed rare neuroendocrine cells [5]. Multiple cancer genomics studies have revealed the recurrent mutational events involved in PCa progression, such as *MYC* and *AR* amplification/overexpression, ETS family gene fusions, alterations in the TP53, PTEN, MAPK, and WNT pathways, and various mutations in *FOXA1*, *SPOP*, and genes of DNA repair and chromatin remodeling (**Figure 1**). Among them, TP53, WNT and AR pathway alterations are par-




Figure 1. Overview of PCa stages, treatment options, and molecular alterations. Cancer progression in the prostate undergoes different stages from hyperplasia to metastatic tumor. Timing and the temporary effect on the tumor volume are shown for different treatment options at various stages. Genomics studies have revealed major molecular alterations in PCa, shown in their likely order during cancer progression. Color code indicates whether the alteration is loss-of-function or gain-of-function.

ticularly enriched in metastatic CRPC compared to primary PCa, and are believed to be late events [6-12].

In order to understand how various mutations and dysregulated signaling pathways affect PCa progression, different experimental models have been utilized. PCa cell lines are traditionally the most widely used models in PCa research and have been discussed in detail elsewhere [13-16]. Recent development of the organoid technology has enabled better in vitro modeling of prostate tumors from patient biopsies which recapitulate the molecular diversity of prostate cancer subtypes [17, 18]. In addition, xenograft models, in particular patient-derived xenografts (PDX) provide valuable resources to study PCa treatment responses and resistance [19, 20]. Finally, over the past two decades, genetically engineered mouse models (GEMMs) have been instrumental in revealing the functions of genetic alterations in promoting PCa progression in vivo. The NIH established the Mouse Models of Human Cancer Consortium (MMCC) in 1999. Since then, GE-

MMs of PCa have evolved from simple knockout and transgenic mice to complex conditional manipulation on multiple genes of interest in prostate-specific or cell-type-specific fashions. Similarities between mouse and human PCa progression are abundant, although differences such as the propensity of bone metastasis also exist. Below, we summarize GEMMs for the most frequently altered pathways found in human PCa (listed in **Table 1**) following the likely order of events during cancer progression, and discuss how they have contributed to our understanding of pathway interactions and therapeutic resistance mechanisms.

PI3K signaling pathway

PCa mouse models investigating the role of the PI3K/Akt/mTOR axis

The PI3K/Akt/mTOR axis is an important intracellular signaling pathway that drives cellular growth and survival, and its hyper-activation is common across many types of cancers. Nearly 50% of primary and metastatic PCas have aberrant PI3K signaling activities, commonly through copy-number loss of the *PTEN* tumor-suppressor gene [7, 10, 11]. Although PI3K activation is not the earliest event in PCa initiation, we discuss this pathway first since its activation is the single most extensively modeled event in PCa mouse studies, often serving as a baseline model for studying oncogenic cooperation.

Pten loss results in constitutive activation of downstream targets including the Akt kinase family [21]. Initially, knockout mice were generated to investigate the function of *Pten* in cancer. While *Pten*^{-/-} mice are embryonic lethal, *Pten*^{+/-} mice developed multiple types of tumors, and hyperplasia/dysplasia in the prostate [22, 23]. Later, the prostate-specific *Pb-Cre4* line [24] was used for *Pten* conditional knockout, and the *Pb-Cre; Pten*^{f/f} model reli-

Prostate cancer mouse model review

Table 1. List of GEMMs analyzing the functions and interactions of major altered signaling pathways in PCa

Pathways	References	Mouse model	Histology and interpretation	
PI3K/Akt/mTOR	[22, 23]	<i>Pten</i> ^{-/+}	Hyperplasia/dysplasia, decreasing Pten level associated with worse phenotypes	
	[30]	<i>Pten</i> ^{fl/+} <i>Pten</i> ^{fl/-}		
	[25]	<i>Pb-Cre; Pten</i> ^{fl/fl}	Adenocarcinoma	
	[28, 29]	<i>ARR2PB-CreER</i> ^{T2} ; <i>Pten</i> ^{fl/fl} <i>PSA-CreER</i> ^{T2} ; <i>Pten</i> ^{fl/fl}	PIN/adenocarcinoma	
	[62, 63]	<i>CK5 (or CK14, CK8, Nkx3.1)-CreER</i> ^{T2} ; <i>Pten</i> ^{fl/fl}	PIN/adenocarcinoma, with more aggressive phenotypes using luminal drivers	
	[34]	<i>Pb-myrAkt1 (MPAKT)</i>	PIN	
	[32]	<i>Pb-Cre; p110α</i> ^{fl/fl} ; <i>Pten</i> ^{fl/fl} <i>Pb-Cre; p110β</i> ^{fl/fl} ; <i>Pten</i> ^{fl/fl}	PIN Impairment of PIN formation	
	[33]	<i>ARR2PB-MF-p110β</i>	PIN	
	[37]	<i>Pten</i> ^{-/+} ; <i>Rictor</i> ^{-/+} <i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Rictor</i> ^{fl/fl}	Impairment of hyperplasia or PIN formation	
	[38]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>elF4E</i> ^{S209A}	Impairment of hyperplasia or PIN formation	
	[40]	<i>Pten</i> ^{-/+} ; <i>Pml</i> ^{-/+}	Accelerated progression to PIN or adenocarcinoma	
	[41]	<i>Pten</i> ^{-/+} ; <i>Tsc2</i> ^{-/+}	Accelerated progression to PIN or adenocarcinoma	
	[39]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>4EBP1</i> ^{-/-} ; <i>4EBP2</i> ^{-/-}	Accelerated progression to PIN or adenocarcinoma	
	[42]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Pik3ca</i> ^{H1047R/+}	Accelerated progression to PIN or adenocarcinoma	
	PI3K & p27	[45]	<i>Pten</i> ^{-/+} ; <i>p27</i> ^{Kip1} ^{-/+}	Accelerated PIN
[46]		<i>MPAKT; p27</i> ^{Kip1} ^{-/-}	Adenocarcinoma, overcome of senescence	
PI3K & TGFβ	[52]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Smad4</i> ^{fl/fl}	Metastatic tumor	
	[53]	<i>CK8-CreER</i> ^{T2} ; <i>Pten</i> ^{fl/fl} ; <i>Tgfb2</i> ^{fl/fl}	Metastatic tumor	
Other PI3K interactions	[47]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Her2</i> ^{Ki}	Accelerated adenocarcinoma, overcome of senescence	
	[48]	<i>Pb-Cre; Pten</i> ^{fl/+} ; <i>Z/Sox9</i>	Accelerated PIN	
	[49-51]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Zbtb7a</i> ^{fl/fl}	Overcome of senescence, resistance to castration	
	[56]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Whsc1</i> ^{fl/fl} <i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Whsc1</i> ^{OE/+}	Impairment of PIN formation Metastatic tumor	
Myc	[70]	<i>Lo-Myc, Hi-Myc</i>	PIN/Adenocarcinoma	
	[72]	<i>Pb-Cre; Z-Myc</i>	PIN	
PI3K & Myc		<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Z-Myc</i>	Accelerated PIN/Adenocarcinoma	
	[75]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>Hi-Myc</i>	Accelerated PIN/Adenocarcinoma	
PI3K & N-Myc	[79]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>R26</i> ^{LSL-MYCN}	Neuroendocrine prostate cancer	
Nkx3.1	[84-90]	<i>Nkx3.1</i> ^{-/+} or <i>Nkx3.1</i> ^{-/-} <i>PSA-Cre; Nkx3.1</i> ^{fl/fl}	PIN	
PI3K & Nkx3.1	[89-92]	<i>Nkx3.1</i> ^{-/-} ; <i>Pten</i> ^{+/-}	Adenocarcinoma in aged mice	
ETS	[98, 99]	<i>ARR2PB-ETV1</i>	PIN	
	[100, 101]	<i>ARR2Pb-ERG</i>	PIN of various degrees	
	[102, 103]	<i>ARR2Pb-ERG</i>	Mostly normal morphology	
	[104]	<i>Pb-Cre; R26</i> ^{ERG/ERG}	Mostly normal morphology	
	[105-107]	<i>Tg (TMPRSS2-ERG)</i>	Mostly normal morphology	
	PI3K & ETS	[103]	<i>ARR2Pb-ERG; Pten</i> ^{+/-}	Accelerated progression to PIN/Adenocarcinoma
		[104]	<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>R26</i> ^{ERG/ERG}	Accelerated progression to PIN/Adenocarcinoma
[105]		<i>KI (TMPRSS2-ERG); Pten</i> ^{-/+} <i>KI (TMPRSS2-ERG); Pb-Cre; Pten</i> ^{fl/fl}	Accelerated progression to PIN/Adenocarcinoma	
[106]		<i>ARR2Pb-TMPRSS2-ERG; Pten</i> ^{+/-}	Accelerated progression to PIN/Adenocarcinoma	
[111]		<i>Pb-Cre; Ets2</i> ^{fl/fl} ; <i>Pten</i> ^{fl/fl}	Accelerated progression to PIN/Adenocarcinoma	
SPOP	[115]	<i>Pb-Cre; Spop</i> ^{fl/fl}	PIN, upregulation of AR and Myc	
	[116, 119]	<i>Pb-Cre; R26</i> ^{SPOP-F133V}	Normal	
PI3K & SPOP		<i>Pb-Cre; Pten</i> ^{fl/fl} ; <i>R26</i> ^{SPOP-F133V}	Accelerated PIN	
Wnt	[123-128]	<i>MMTV-Cre, Nkx3.1-Cre, Pb-Cre or p63</i> ^{CreERT2/+} ; <i>Ctnnb1</i> ^{L(ex3)/L(ex3)}	Squamous metaplasia, PIN of varied degrees	
	[130]	<i>Pb-Cre; Apc</i> ^{fl/fl}	Adenocarcinoma	
	[131]	<i>Ubi-Cat</i>	Adenocarcinoma	

Prostate cancer mouse model review

PI3K & Wnt	[129]	<i>Pb-Cre; Ctnnb^{fl/fl}; Pten^{fl/fl}</i>	No impairment of progression
		<i>Pb-Cre; Ctnnb1^{L(ex3)/L(ex3)}; Pten^{fl/fl}</i>	Accelerated adenocarcinoma
Wnt & p53/Rb	[126]	<i>Pb-Cre; Ctnnb1^{L(ex3)/L(ex3)}; LPB-Tag</i>	Adenocarcinoma with neuroendocrine feature
MAPK	[136]	<i>Pb-H-Ras^{G12V}</i>	Low grade PIN
Wnt & MAPK	[127]	<i>Pb-Cre; K-Ras^{LSL-V12/+}</i>	Adenocarcinoma
		<i>Pb-Cre; ctnnb1^{+lox(ex3)}; K-Ras^{LSL-V12/+}</i>	
MAPK & p16	[137]	<i>iBRAF[*]</i>	Adenocarcinoma
PI3K & MAPK	[138, 141, 142]	<i>Pb-Cre; Pten^{fl/fl}; K-Ras^{LSL-G12D/+}</i>	Metastatic tumor
	[143]	<i>Nkx3.1^{CreERT2/+}; Pten^{fl/fl}; K-Ras^{LSL-G12D/+}</i>	Metastatic tumor
FGF	[152, 153]	<i>Pb-Fgf7, Pb-Fgf8, Pb-Fgfr2iib</i>	PIN, Fgfr1 and Fgfr2iib synergize to promote high grade PIN
	[154-157]	<i>Pb-Fgfr1</i>	PIN, Fgfr1 and Fgfr2iib synergize to promote high grade PIN
PI3K & FGF	[158]	<i>ARR2PB-Fgf8b; Pb-Cre; Pten^{fl/+}</i>	Metastatic tumor
Wnt & FGF	[159]	<i>Ubi-Cat; JOCK1</i>	Accelerated adenocarcinoma with reactive stroma
Tp53/Rb	[43, 76, 169]	<i>Pb-Cre; p53^{fl/fl}</i>	Normal
	[170]	<i>Pb-p53^{R273H}</i>	PIN
	[171]	<i>Nkx3.1-Cre; p53^{LSL-R270H/R270H}</i>	PIN
	[176]	<i>Pb-Cre; Rb^{fl/fl}</i>	Hyperplasia
	[177-179]	<i>TRAMP, LADY</i>	Metastatic tumor with neuroendocrine feature
PI3K & p53/Rb	[169, 185]	<i>Pb-Cre; p53^{fl/fl}; Rb^{fl/fl}</i>	Neuroendocrine tumor
	[43, 50]	<i>Pb-Cre; Pten^{fl/fl}; p53^{fl/fl}</i>	Accelerated adenocarcinoma, bypass of senescence and castration-induced apoptosis
	[186, 187]	<i>Pb-Cre; Pten^{fl/fl}; Rb1^{fl/fl}</i> <i>Pb-Cre; Pten^{fl/fl}; Rb1^{fl/fl}; p53^{fl/fl}</i>	Metastatic tumor with neuroendocrine feature
AR	[190]	<i>Pb-mAR</i>	Low grade PIN at old age
	[191]	<i>Osr1-Cre; R26-LSL-AR</i>	PIN/adenocarcinoma
	[192]	<i>Pb-AR-E231G</i>	PIN
	[194]	<i>Pb-AR^{Δ567es}</i>	Adenocarcinoma at old age
	[195]	<i>ARR2PB-AR-V7</i>	PIN
Wnt & AR	[128, 215]	<i>Pb-Cre or p63^{CreERT2/+}; Ctnnb1^{L(ex3)/+}; R26hAR^{+/+}</i>	Accelerated adenocarcinoma
PI3K & AR	[212, 213]	<i>Pb-Cre; Pten^{fl/fl}; AR^{fl/Y}</i>	AR dispensable for Pten tumor progression. PI3K & AR form reciprocal negative feedback loop.
	[204, 214]	<i>Osr1-Cre; Pten^{fl/fl}; AR^{fl/Y}</i> <i>CK5-CreERT2; Pten^{fl/fl}; AR^{fl/Y}</i> <i>Nkx3.1^{CreERT2}; Pten^{fl/fl}; AR^{fl/Y}</i>	AR dispensable for Pten tumor progression

ably developed invasive adenocarcinoma [25]. Studying murine cell lines derived from this model suggested that Pten loss bestows tumors androgen-independent proliferation [26]. In addition, this model also revealed that epithelial *Pten* deletion may trigger the secretion of inflammatory cytokines to promote the expansion of a subset of Gr-1⁺CD11b⁺ myeloid-derived suppressor cells (MDSCs), leading to immune suppression and tumor progression [27]. It is worth noting that in this model *Pten* deletion occurs during prostate postnatal development due to the early timing of *Pb-Cre* activation. Indeed, when the inducible *ARR2PB-CreERT²* or *PSA-CreERT²* lines were used, *Pten* loss at the adult stage yielded less aggressive prostate tumors [28, 29]. Besides timing, *Pten* expression level can heavily influence PCa progression. Using hypomorphic *Pten* mutant mice *Pten^{hy}*, it was shown that decreasing Pten

levels were correlated with progressively worse phenotypes (*Pten^{hy/+}* > *Pten^{+/-}* > *Pten^{hy/-}* > *Pten^{prostate-knockout}*) [30]. One possible mechanism of regulating Pten protein level is through the ubiquitin ligase NEDD4-1, which is strongly expressed in neoplastic areas of *Pten^{hy/-}* prostates [31].

Other components of the PI3K/Akt/mTOR axis have been modeled (**Figure 2**). PI3K class IA catalytic subunit p110 α (PIK3CA) or p110 β (PIK3CB) is frequently overexpressed in PCa [11]. Interestingly, conditional knockout of p110 β , but not p110 α , impeded Akt phosphorylation and Pten-loss-induced tumorigenesis [32], while overexpressing activated p110 β in transgenic mice induced PIN phenotypes [33]. A transgenic mouse model overexpressing activated Akt1 under the probasin promoter (MPAKT) was developed, and mice only




Figure 2. Mouse models used to study the PI3K/AKT/mTOR axis in PCa. Major components of the PI3K/AKT/mTOR pathway and their molecular signaling relationships are shown. Pioneer papers that described the GEM models of the individual components or their interactions are listed.

developed PIN in the ventral lobes [34]. Akt-induced cancer initiation is dependent on downstream mTOR activation since treatment with the mTOR inhibitor RAD001 reversed the PIN phenotype in MPAKT mice [35]. mTOR is a protein kinase that exists in two distinct complexes, the mTOR complex 1 (mTORC1) and mTORC2, and serves as a central regulator of cell metabolism and growth. The two complexes share certain protein subunits, with Rictor being one of the components in mTORC2 [36]. Double mutant mice *Pten*^{-/+}; *Rictor*^{-/+} and *Pb-Cre*; *Pten*^{fl/fl}; *Rictor*^{fl/fl} showed dramatically reduced tumor phenotypes compared to their respective models of single *Pten* mutant, suggesting an essential role of mTORC2 in transducing the PI3K signal in *Pten*-loss tumors [37]. On the other hand, mTORC1 can phosphorylate 4E-BP1 and 4E-BP2, preventing them from binding and inhibiting eIF4E, a factor pro-

moting protein synthesis. To study the role of eIF4E, a knock-in model *eIF4E*^{S209A} was built to mutate its only phosphorylation site, and this was sufficient to inhibit the *Pten*-loss-induced PCa progression [38]. Interestingly, in *Pb-Cre*; *Pten*^{fl/fl}; *4EBP1*^{-/-}; *4EBP2*^{-/-} mice, 4E-BP1/2 knockout significantly accelerated PCa progression from *Pten* loss, suggesting 4E-BP1/2 are tumor-suppressing even under constitutive mTORC1 activation, possibly due to their dephosphorylation under hypoxia conditions [39]. The accelerated PCa progression compared to *Pten* loss alone was also seen in *Pten*^{-/+}; *Pml*^{-/+} mice, which had reduced Akt antagonist *Pml* [40], *Pten*^{-/+}; *Tsc2*^{-/+} mice, which had reduced mTOR antagonist *Tsc2* [41], and *Pten* conditional knockout mice that overexpressed a mutant p110α (*Pik3ca*^{H1047R}) [42]. Overall, these findings suggest that PCa progression is very sensitive to the activity lev-

els of Akt and mTOR, and alterations of other components in the PI3K pathway are not functionally redundant to Pten loss.

Models of other regulators that cooperate with the PI3K pathway

Although the PI3K/Akt/mTOR axis plays a pivotal role in driving prostate adenocarcinoma formation, Pten loss can induce p53-dependent cell senescence response to prevent further progression [43]. We will discuss this and other major pathway interactions in later individual sections. Here, we briefly list a few other regulators that help keep the *Pten*-null tumors in check.

In many tissues, p27^{Kip1} functions as a negative regulator of cyclin-CDK activity to inhibit cell cycle progression. Disruption of p27^{Kip1} led to prostatic hyperplasia in p27^{Kip1}^{-/-} mice [44], and *Pten*^{-/+}; p27^{Kip1}^{-/+} mice displayed accelerated neoplastic transformation [45]. Notably, p27^{Kip1} and senescence marker levels increased in MPAKT mice, and MPAKT; p27^{Kip1}^{-/-} mice developed invasive PCa, suggesting that a p27^{Kip1}-driven checkpoint limits progression from PIN to PCa [46]. Another event that can overcome Pten-loss-induced cell senescence is the activation of the receptor tyrosine kinase Her2, as conditional expression of activated Her2 in *Pb-Cre*; *Pten*^{fl/fl}; *Her2*^{KI} mice activated the MAPK pathway and led to faster progression [47].

Transcription factor Sox9 is expressed in certain human PCa specimens and is correlated with decreased survival. Prostate-specific Sox9 overexpression was shown to cooperate with one copy loss of *Pten* in transformation in *Pb-Cre*; *Pten*^{fl/+}; *Z/Sox9* mice [48]. One factor that can inhibit Sox9-dependent oncogenic pathways is Zbtb7a, as its homozygous deletion led to bypass of *Pten*-loss-induced cellular senescence [49] and resistance to castration in the *Pten*-null tumors [50]. Recently, loss of Zbtb7a was also shown to trigger the infiltration of Gr1⁺CD11b⁺ immune cells through CXCL5 up-regulation in *Pb-Cre*; *Pten*^{fl/fl}; *Zbtb7a*^{fl/fl} mice to promote tumor progression [51].

Regarding PCa metastasis, TGFβ/Smad4 signaling has been shown to serve as a barrier for *Pten*-null tumors. *Pb-Cre*; *Pten*^{fl/fl}; *Smad4*^{fl/fl} mice developed metastasis to the lymph nod-

es and lung with high penetrance [52], and inactivating TGFβ receptor in luminal cells in *CK8-CreER^{T2}*; *Pten*^{fl/fl}; *Tgfb2*^{fl/fl} mice enhanced metastasis and luminal cell dedifferentiation [53]. Another gene that was shown to drive *Pten*-null tumors towards metastasis is *Whsc1*, a histone methyltransferase overexpressed in a number of metastatic tumors [54, 55]. Prostate-specific ablation of *Whsc1* prevented tumor progression in *Pb-Cre*; *Pten*^{fl/fl}; *Whsc1*^{fl/fl} mice, while its overexpression promoted tumor metastasis in *Pb-Cre*; *Pten*^{fl/fl}; *Whsc1*^{OE/+} mice [56]. Mechanistically, Pten-loss-induced Akt activation stabilizes *Whsc1* from degradation via phosphorylation, and allows *Whsc1* to transcriptionally upregulate Rictor, forming a positive-feedback loop through mTORC2 to further enhance Akt activity [56].

PI3K-based mouse models for studying PCa cell of origin

It has been appreciated in multiple cancers that the cell of origin, defined as a normal cell that can give rise to a tumor upon transformation, plays a major role in determining tumor subtype and outcome [57]. Although PCa shows a predominantly luminal phenotype, tumors originating from basal and luminal cells may be distinguishable by their aggressiveness and such information may be useful for prognosis [58]. Transplantation-based studies, in which basal and luminal cells were renal-grafted in immunodeficient mice, mostly reported basal cells to be more aggressive in cancer initiation [59-61]. However, the renal-graft assay intrinsically disfavors luminal cell growth, and lacks the intact tumor microenvironment in GEMMs. To test the cell of origin in situ, cell-type-specific inducible Cre lines have been used to knockout *Pten* in either basal or luminal cells of the mouse prostate. In one study, basal-lineage-traced *K14-CreER^{T2}*; *Pten*^{fl/fl}; *mTmG* mice and luminal-lineage-traced *K8-CreER^{T2}*; *Pten*^{fl/fl}; *mTmG* mice were analyzed. Luminal cells were found to be more responsive to Pten-null induced mitogenic signaling, while basal cells underwent luminal differentiation before transformation [62]. Similar results were also observed by comparing *CK5-CreER^{T2}*; *Pten*^{fl/fl}; *R26R-YFP/+* and *Nkx3.1^{CreERT2/+}*; *Pten*^{fl/fl}; *R26R-YFP/+* mice, where luminal-derived tumors progressed faster and showed a gene signature predictive of poor outcomes in patients [63].

Notably, the relative susceptibility of luminal cells to oncogenic transformation is not restricted to the Pten-loss models, as a comparative lineage-tracing study showed that luminal cells are favored to be the cell of origin for PCa in multiple genetic and chemical-induced models [64].

Early oncogenic events in PCa initiation

Recent research suggests that prostate carcinogenesis favors the dysregulation of cancer genes in defined orders [65]. In this section, we discuss how modeling some of the early oncogenic events has led to a deeper understanding of the mechanisms underlying the transitions from normal prostate to PIN and subsequently to adenocarcinoma.

Myc up-regulation

The Myc transcription factor responds to diverse mitogenic and developmental signals to coordinate expression of diverse genes important for the orderly proliferation of somatic cells [66]. Myc expression is highly regulated, and changes of its expression level can influence somatic cell proliferation and oncogenesis [67, 68]. Up-regulation of c-Myc appears to be an early event in human PIN [69], and the gene is within the most commonly amplified loci on 8q24.21 in advanced PCa [10]. As a functional confirmation, transgenic mice of Myc overexpression under the probasin promoters (*Lo-Myc* and *Hi-Myc*) displayed PIN and progressed to invasive adenocarcinoma [70]. Two other models, the transgenic *C(3)1-c-Myc* [71] and the inducible *Pb-Cre; Z-Myc* mice [72] showed much milder PIN phenotypes, possibly due to the different expression levels or mouse backgrounds. Nevertheless, up-regulation of Myc needs to cooperate with other oncogenic pathways for PCa to progress. Myc overexpression synergized with PIM1 kinase expression [73] and myrAkt expression [59, 74] in lentiviral-transfected renal grafts. In GEMMs, *Pb-Cre; Pten^{fl/fl}; Hi-Myc* bigenic mice showed faster tumor progression than either of the single lesion models [75]. Z-Myc also cooperates with Pten loss [72]. Interestingly, in *Pb-Cre; Z-Myc; Pten^{fl/+}; p53^{fl/+}* triple mutant mice, loss of the wild-type Pten allele occurred prior to p53 [76]. The order of these oncogenic events is consistent with findings in humans, and suggests a

greater selective advantage to the tumor cell by Pten loss than p53 loss.

Unlike c-Myc, N-Myc was proposed to be involved in the development of NEPC [77]. Indeed, introduction of N-Myc and activated AKT1 into human prostate cells was sufficient to transform them into NEPC in renal grafts [78]. In *Pb-Cre; Pten^{fl/fl}; R26^{LSL-MYC}* mice, overexpression of N-Myc when Pten is lost led to the development of poorly differentiated, invasive NEPC through the Ezh2-mediated transcriptional program [79]. This study, and others which are discussed later in the section of the p53 and Rb pathways, provide excellent examples of using GEMM to understand the mechanism behind transition from CRPC to NEPC as a treatment resistance.

Nkx3.1 down-regulation

The homeobox gene *Nkx3.1* is a key regulator in prostate development and tumorigenesis and is located within a region on chromosome 8p21.2 that frequently undergoes loss-of-heterozygosity in PIN lesions [80-82]. Although *Nkx3.1* is rarely mutated in primary PCa [9], its down-regulation is correlated with PCa initiation and progression [83]. *NKX3.1^{+/-}, Nkx3.1^{-/-}, and PSA-Cre; Nkx3.1^{fl/fl}* mice all developed PIN lesions [84-90], supporting an initiating role of *Nkx3.1* loss-of-function in prostate tumorigenesis. *Nkx3.1^{-/-}; Pten^{+/-}* mice were prone to develop adenocarcinoma in aged mice [89-92], emphasizing the importance of the PI3K pathway in driving the progression of *Nkx3.1*-deficient tumors. In fact, loss of Pten alone leads to *Nkx3.1* down-regulation at the transcriptional level [25, 93, 94]. Decreased *Nkx3.1* expression was also observed in prostate tumors but not PINs of *Hi-Myc* mice, raising the possibility that Myc gain and *Nkx3.1* loss may be cooperating events in the PIN to adenocarcinoma transition [70]. Mechanistically, in cancer cell line and xenograft models, *Nkx3.1* has been shown to negatively modulate AR transcription and stabilize p53 to impede cancer progression [94].

Genomic rearrangement of ETS family genes

Fusions of the *TMPRSS2* gene to the ETS family transcription factor genes *ERG*, *ETV1*, *ETV4* and *ETV5* are frequent chromosome rearran-

gement events in human PCa, accounting for 50-60% of all cases [95]. The most common gene fusion is *TMPRSS2-ERG*, which allows androgens to drive expression of the N-terminally truncated ERG protein in the prostate [95-97]. The presence of this fusion in benign prostatic hyperplasia and PIN cases indicates that it can be a relatively early event in prostate tumorigenesis.

Transgenic mouse models were established to functionally test the role of ETS genes. The *ARR2Pb* promoter was used to drive full-length and truncated ETV1, and these mice developed PIN with high penetrance [98, 99]. Mixed results have been reported for mice overexpressing truncated ERG. Some studies showed that *ARR2Pb-ERG* mice displayed phenotypes ranging from PIN to adenocarcinoma at old age, with higher levels of ERG expression correlating with more severe phenotypes [100, 101]. Mechanistically, ERG was suggested to activate the YAP1/Hippo transcriptional program, and a mouse model with prostate-specific activation of YAP1 developed phenotypes that were similar to *ARR2Pb-ERG* mice [101]. However, others reported largely normal prostate morphology for the *ARR2Pb-ERG* model [102, 103], the *Pb-Cre; R26^{ERG/ERG}* conditional expression model [104], and models in which the *TMPRSS2-ERG* fusion product is expressed [105-107]. Mouse strains, timing of analyses, the precise portion of the expressed protein, and transgene integration sites are all possible factors contributing to these varied outcomes. Rigorous quantitation of ERG levels and better control for mouse genetic background may help clarify, but *TMPRSS2-ETS* fusions certainly need to cooperate with other pathways for PCa to progress. Analysis of genetic alterations in human PCa showed that ERG rearrangement was significantly associated with PTEN loss [103, 106, 108]. Compound mouse models with ERG overexpression and Pten loss, such as *ARR2Pb-ERG; Pten^{+/-}* [103], *ARR2Pb-TMPRSS2-ERG; Pten^{+/-}* [106], *TMPRSS2-ERG knock-in; Pten^{-/+}* and *TMPRSS2-ERG knock-in; Pb-Cre; Pten^{fl/fl}* [105], and *Pb-Cre; Pten^{fl/fl}; R26^{ERG/ERG}* [104] mice, all showed accelerated PIN and adenocarcinomas progression compared to *Pten*-mutant mice, suggesting synergy between ERG overexpression and PI3K pathway activation. Mechanistically, ERG may restore AR transcriptional output in the *Pten*-loss background, and upregulate genes involv-

ed in cell migration and angiogenesis [104], but differences in ERG- and ETV1-regulated transcriptional programs have also been reported [105].

Interestingly, *TMPRSS2-ERG* fusion can be accompanied by the deletion of a highly conserved interstitial region between these two genes [109, 110], raising the possibility that this event may have other oncogenic functions besides ERG overexpression. Indeed, in a recent study comparing two *TMPRSS2-ERG* knock-in models, the model containing the interstitial deletion cooperated more strongly with *Pten*-loss [111]. One of the deleted genes is *Ets2*, and *Pb-Cre; Ets2^{fl/fl}; Pten^{fl/fl}* mice displayed more aggressive phenotypes than *Pb-Cre; Pten^{fl/fl}* mice, suggesting the delete genes may have tumor suppressor functions [111].

SPOP mutations

Recurrent mutations in the gene *SPOP*, which encodes a Cullin-based E3 ubiquitin ligase subunit, are found in ~10% of primary and advanced PCa cases [8, 112]. Interestingly, *SPOP* mutations are mutually exclusive with ETS family gene rearrangements, defining a distinct molecular subtype of PCa [8, 9, 113]. *SPOP* mutations are significantly enriched in primary tumors [6], indicating that they are early events during PCa development [114].

In a loss-of-function model, *Pb-Cre; Spop^{fl/fl}* mice developed PIN through upregulation of AR and c-Myc [115]. In another model, *SPOP-F133V*, a common missense mutation found in human PCa, was conditionally overexpressed in *Pb-Cre; R26^{SPOP-F133V}* mice [116]. Although *SPOP-F133V* alone was insufficient to drive tumorigenesis, it synergized with *Pten* loss, through increasing PI3K/mTOR signaling activity and maintaining AR activity against PI3K-mediated feedback inhibition [116], indicating *SPOP-F133V* has dominant negative function. Notably, there has been some controversy regarding the tumor-suppressing mechanism of *SPOP*. Some studies reported that *SPOP* facilitates the ubiquitination and degradation of ERG, but not *TMPRSS2-ERG* [117, 118]. However, the vast majority of human *SPOP*-mutant cancers do not express ERG, and *Pb-Cre; Pten^{fl/fl} or +/-; R26^{SPOP-F133V}* mice showed no evidence of ERG expression [119], questioning such regulation. Analysis of human PCa datasets and zebrafish models of *SPOP-F133V* sug-

gested that SPOP mutations drive prostate tumorigenesis at least partly through impaired DNA double strand break repair and increased genomic instability [114].

Wnt signaling pathway

The canonical Wnt signaling pathway via nuclear β -catenin regulates numerous processes in animal development, tissue homeostasis, and diseases [120, 121]. A recent integrated analysis of primary and metastatic PCas found that 10% of the cases harbored genomic alterations in the Wnt/ β -catenin pathway [6], and Wnt-related alterations are significantly enriched in CRPCs [7, 11].

To investigate the potential role of the canonical Wnt pathway in prostate tumorigenesis, many groups used the *Ctnnb1*^{L(ex3)/+} mouse strain, in which Cre-mediated deletion of the third exon of the β -catenin gene results in a dominant stable protein [122]. Driven by the *MMTV-Cre*, *Nkx3.1-Cre*, *Pb-Cre* or *p63*^{CreERT2/+}, such mouse models developed squamous metaplasia and varied degrees of PIN [123-128]. These phenotypes progressed to adenocarcinoma when β -catenin stabilization is combined with SV40-large T-antigen overexpression [126] or *Pten* deletion [129], suggesting synergy with the p53/Rb and PI3K pathways, respectively. Notably, although β -catenin level seems to increase in *Pb-Cre; Pten*^{fl/fl} tumors, there is no phenotypic difference between *Pb-Cre; Pten*^{fl/fl} and *Pb-Cre; Ctnnb1*^{fl/fl}; *Pten*^{fl/fl} tumors, suggesting that β -catenin is dispensable for Pten loss-driven PCa [129]. Besides *Ctnnb1*^{L(ex3)/+}-based models, adenocarcinomas were also observed in *Pb-Cre; Apc*^{fl/fl} mice where the Wnt pathway negative regulator Apc is inactivated [130], and in *Ubi-Cat* mice where β -catenin is activated through a genetically modified inducible Wnt co-receptor LRP-5 [131]. Taken together, these data imply Wnt/ β -catenin pathway as a driving force of prostate tumorigenesis. Further evidence comes from renal-graft studies showing that manipulation of the stromal compartment by overexpressing *Hmga2* or knocking out *Tgfb2* enhances paracrine Wnt signaling to promote PCa initiation [132, 133]. Despite these progresses, the role of Wnt activation in promoting PCa cell metastasis remains largely elusive. Developing GEMMs that activate Wnt in advanced tumors should shed light onto this important question.

Ras/Raf/MAPK signaling pathway

The mitogen-activated protein kinase (MAPK) cascade relays extracellular cues to activate Ras and Raf for subsequent phosphorylation and activation of MEK1/2 and ERK1/2 (MAPKs), which phosphorylate multiple targets regulating cell survival, proliferation, and differentiation [134]. Although RAS and RAF mutations are not frequent in PCa, the activity of the MAPK signaling pathway is often upregulated in both primary and metastatic PCas [9, 10, 135]. Ras^{G12V} and BRAF^{V600E} activating mutations are common in human cancers and are also present in PCa patients [10]. To model these oncogenic events in vivo, *Pb-H-Ras*^{G12V} transgenic mice [136] and *Pb-Cre; K-Ras*^{LSL-V12/+} conditional activating mice [127] were developed, and both models exhibited low grade PIN. In the iBRAF^{*} model where BRAF^{V600E} expression is doxycycline-inducible under the *Ink4a/Arf*^{-/-} background, mice developed prostate adenocarcinomas [137]. Another hotspot mutation, K-Ras^{G12D}, is not found in human PCa. Nevertheless, it has been modeled in *Pb-Cre; K-Ras*^{LSL-G12D/+} mice, which showed no sign of cancer development [138].

Aberrant Wnt signaling has been shown to synergize with Ras/Raf/MAPK activation, as *Pb-Cre; ctnnb1*^{+/lox(ex3); K-Ras^{LSL-V12/+} mice developed invasive carcinoma [127]. But it is the cooperation of the PI3K and MAPK pathway activations in PCa progression that has been best characterized [139, 140]. For example, PCa in *Pb-Cre; Pten*^{fl/fl}; *K-Ras*^{LSL-G12D/+} mice displayed epithelial to mesenchymal transition (EMT) and metastatic phenotypes [138, 141], which could be suppressed by inhibition of the chromatin remodeling protein HMGA2 [142]. Metastasis also occurred in *Nkx3.1*^{CreERT2/+}; *Pten*^{fl/fl}; *K-Ras*^{LSL-G12D/+} mice, and the ETS transcription factor family member ETV4 was activated in response to activation of MAPK and PI3K pathways to promote this process [143]. Importantly, inhibition of mTOR by the drug RAD001 activates the MAPK pathway in both human samples and the *Pb-Cre; Pten*^{fl/fl} model, suggesting a feedback loop between PI3K and MAPK pathways [144]. Recently, loss of Pml in *Pten*-null mouse tumors was shown to relieve this feedback inhibition and activate the MAPK pathway, resulting in an SREBP-dependent lipogenic program to promote metastasis [145]. Collectively, these studies suggest the combined therapy using both mTOR and}

MAPK inhibitors as a potential strategy for treating patients with advanced PCa.

FGF/FGFR signaling pathway

The fibroblast growth factor (FGF) and its receptor FGFR are one of the key stromal paracrine signals that regulate prostate epithelial development and homeostasis. The pathway consists of 18 FGF ligands, 4 receptor tyrosine kinases (FGFRs), and a pool of heparan sulfate proteoglycans (HSPGs) [146]. Although genomic mutations in the FGF pathway are rare in human PCa, research using xenograft and cell line models has suggested the existence of abnormal autocrine and paracrine FGF loops in PCa [147]. For instance, enhancing paracrine FGF10 in urogenital sinus mesenchyme led to the formation of PIN and adenocarcinoma [60, 148-151].

In earlier GEM models, FGF7 or FGF8 expression driven by the probasin promoter induced PIN formation [152, 153], while enforced expression of a dominant negative FGFR2iib induced hyperplasia with neuroendocrine feature [152]. PIN was also observed when FGFR1 or its mutant activated form was overexpressed using the probasin promoter [154-157], and combined expression of FGFR1 and FGFR2iib synergistically promoted high-grade PIN [155]. These models clearly demonstrated that aberrant FGF signaling environment is conducive to PCa initiation. Possible mechanisms include the activation of downstream MAPK/Erk signaling [156], increase in EMT-associated Sox9 and changes in the Wnt signaling pathway [157].

Synergy of FGF signaling with the PI3K pathway was shown in *ARR2PB-Fgf8b; Pb-Cre; Pten^{fl/+}* mice, which developed metastatic PCa [158]. Induced FGFR1 expression also synergized with induced β -catenin expression to accelerate PCa progression, potentially through elevating stromal TGF- β signaling [159]. Recently, a molecular study in human PCa cells and xenografts discovered that FGF signaling was able to sustain a portion of AR-null CRPC [160]. It will be interesting to develop new preclinical GEMMs to test the possibility that inhibition of FGF and MAPK pathways together may be efficacious against PCas that are resistant to AR-directed therapies.

Tp53 and Rb pathways

The p53 and Rb pathways are almost universally involved in cancer [161]. The p53 pathway responds to stresses to initiate programs of cell cycle arrest, cellular senescence or apoptosis [162]. Pathological analyses of clinical samples indicated that *TP53* alterations are closely associated with PCa progression and occur mostly at later stages [163-167]. Comprehensive molecular analyses also showed that *TP53* mutations are highly enriched in metastatic PCa compared to primary PCa [9, 11].

It is important to point out that p53 mutations can provoke activities involved in cell invasion, metastasis, proliferation and survival that are different to those resulting from simply loss of wild-type p53 [168]. Therefore, different p53 mouse models may yield different phenotypes. Simple prostate-specific knockout of p53 in *Pb-Cre; p53^{fl/fl}* mice largely had no tumorigenic effect [43, 76, 169], while overexpressing a human hotspot mutant R273H (mouse R270H) in *Pb-p53^{R273H}* mice [170] or *Nkx3.1-Cre; p53^{LSL-R270H/R270H}* mice [171] induced PIN. More mouse models have focused on how p53 alterations drive cancer progression. It has been proposed that p53 presents a PCa progression barrier and its alterations lead to genomic instability and evasion of senescence or apoptosis in clones that will become aggressive [65]. In support of this idea, p53-dependent cellular senescence and castration-induced apoptosis were found to be bypassed in *Pb-Cre; Pten^{fl/fl}; p53^{fl/fl}* mice [43, 50], and analyzing *Pb-Cre; Z-Myc; Pten^{fl/fl}; p53^{fl/fl}* mice revealed that c-Myc expression shifted the p53 response from senescence to apoptosis by repressing the p53 target gene *p21^{Cip1}* [72, 76]. Other potential oncogenic mechanisms induced by p53 loss include upregulation of hexokinase 2-mediated Warburg effect [172] as well as increased immune cell infiltration through CXCL17 upregulation [51]. Therefore, p53 serves as an essential checkpoint preventing Pten-loss-induced PCa progression.

The tumor suppressor gene Rb plays a pivotal role in cell-cycle regulation. It often undergoes copy number loss in both localized and advanced PCas [10, 11, 173, 174], but is more commonly mutated in neuroendocrine prosta-

te cancers [175]. Prostate-specific deletion of Rb alone in *Pb-Cre; Rb^{fl/fl}* mice only showed hyperplasia [176]. However, it is the NEPC phenotypes induced by the combined loss of Rb and p53 that should be brought to attention. The pioneer models of PCa, including the TRAMP model [177, 178] and the LADY model [179] used the *Pb* promoter to drive the expression of SV40 large T-antigen, which binds and inactivates p53 and Rb [180-182]. These mice showed rapid PCa progression with metastasis, and tumors often had “small cell” and synaptophysin positive phenotypes, suggesting they were neuroendocrine [183, 184]. Carcinomas from the conditional double knockout model *Pb-Cre; p53^{fl/fl}; Rb^{fl/fl}* also displayed neuroendocrine differentiation [169, 185]. Recently, two important studies characterizing the *Pb-Cre; Pten^{fl/fl}; Rb1^{fl/fl}* and *Pb-Cre; Pten^{fl/fl}; Rb1^{fl/fl}; p53^{fl/fl}* mice suggested that Rb and p53 loss depresses epigenetic reprogramming factors Ezh2 and Sox2, thereby facilitating lineage plasticity and metastasis of prostate adenocarcinoma initiated by *Pten* loss [186, 187]. These findings have important therapeutic implications, since tumors cells can acquire lineage plasticity as a means to escape luminal-targeted androgen deprivation therapy, and epigenetic modulations such as Ezh2 inhibition may provide new approaches to treat NEPC.

AR signaling pathway

The androgen receptor (AR), a member of the steroid hormone receptor superfamily, is essential for normal prostate development, and supports growth of PCa. In the past decade, aberrant AR signaling has emerged as the central driver of CRPC [3]. Alterations of the AR signaling through AR gene mutations, gene/enhancer copy-number amplification, AR splicing variants, and mutations of AR binding partners are particularly frequent in CRPC [7, 9-11, 188, 189].

GEMMs of prostate AR loss- or gain-of-function experiments are generally in agreement with human sequencing data, showing that manipulation of AR signaling alone has subdued effect on early cancer initiation. In the *Pb-mAR* transgenic model where the AR transgene is driven by the rat probasin promoter, PIN lesions only appeared in old mice (> 1 year) [190]. PIN was also observed in an *Osr1-Cre* induced AR-

overexpressing mouse model [191]. In another study, overexpressing either wild-type AR or AR-T857A (human T877A), an AR mutant present in LNCaP cells that is responsive to non-classical ligands, had no obvious phenotypes, while overexpressing AR-E231G, a mutant with increased ligand-independent activity, induced PCa formation [192]. However, it should be noted that AR-E231G was discovered in the TRAMP model [193], and has not been observed in human PCa. Recently, an AR^{v567es} splice variant frequently found in human CRPC was shown to induce invasive adenocarcinoma in *Pb-AR^{v567es}* transgenic mice by one year of age [194], and overexpression of AR-V7, a constitutively active and androgen-independent AR splice variant, by the *ARR2PB* promoter induced PIN in mice [195]. These models offer opportunities for studying the mechanisms of AR-variant-driven PCa progression. Notably, in these studies AR overexpression started at the developmental stage, and the probasin promoter itself is regulated by androgen. Modeling AR overexpression in later cancer stages may prove to be beneficial.

On the other hand, AR loss-of-function modeling has also been performed by AR-floxed conditional knockout using various Cre or CreER lines targeting the whole prostate or specific epithelial or stromal cell types [196-205]. The phenotypes were mixed and sometimes controversial, possibly due to the complex timing and spatial specificities of different Cre expression. Overall, although decreased androgen levels in aging men has been shown to correlate with the increase of PCa incidence [206], the AR gene is not lost in human PCa. Therefore, while these loss-of-function models have shed light onto prostate development and normal homeostasis, their cancer-relevance is less clear.

The interplay of AR signaling with other pathways in PCa progression has been under extensive investigation. In renal-graft experiments, overexpression of AR synergized with Akt, K-ras, FGF10, ERG, Src family kinases, and Hmga2 to promote PCa initiation and progression [60, 61, 132, 149, 207-211]. In GEMMs, the PI3K pathway and AR signaling pathway have been shown to form a reciprocal negative feedback loop to drive PCa survival. In *Pb-Cre; Pten^{fl/fl}* mice and human PCa cells, drug inhibi-

Prostate cancer mouse model review




Figure 3. Using GEMMs to study pathway interactions in PCa progression. Diagram summarizes pathway interactions shown to promote PCa progression by studying GEMMs. The PI3K pathway activation has been the most extensively modeled event in PCa GEMMs and usually serves as a baseline model. Cooperation of PI3K pathway with other pathways is required for PCa to overcome various barriers and checkpoints to progress. Each line connecting two pathways indicates that at least one GEMM has been used to establish a functional correlation of the two pathways, through analyzing either phenotypes of double mutant mice or pathway activities in single mutant mice. Pathway interactions revealed by cancer cell line and xenograft studies are not drawn.

tion of the PI3K pathway activates the AR signaling output through increased HER3 kinase expression [212]. *Pb-Cre; Pten^{fl/fl}* tumors were also shown to suppress androgen-responsive genes by upregulating AR signaling co-regulators EGR1, c-JUN, and EZH2, thereby rendering cancer cells less dependent on androgen [213]. On the other side, while AR is dispensable for tumor formation in *Pb-Cre; Pten^{fl/fl}; AR^{fl/y}* mice, deletion of AR or AR blockade upregulated Akt Activity in *Pten*-null cells by decreasing FKBP5- and PHLPP-mediated Akt dephosphorylation [212, 213]. These findings were also corroborated by models using the *Osr1-Cre* and basal- and luminal-specific CreER lines [204, 214]. Together, these studies provide strong rationale for combined therapy inhibiting both the PI3K and AR signaling pathways in PCa patients. Recently, AR overexpression was shown to accelerate PCa progression in *Pb-Cre* or *p63^{CreERT2/+}* driven *Ctnnb1^{L(ex3)/+;} R26hAR^{L/+}* mice, where β -catenin was stabilized to activate the Wnt signaling pathway [128, 215]. In these Wnt and AR compound mice, tumor- and metastasis-promoting genes such as *Spp1*, *Egr1*, c-Myc, and *Sp6* were upregulated. Future GEMMs that turn on AR signaling at later stages in combination with multiple other path-

way perturbations should continue to reveal the key to castration resistance in CRPC.

Conclusion

Development of the transgenic, knock-out/in, and Cre-lox conditional expression techniques in mice paved the way for building GEMMs for answering key biological questions in PCa progression. The intact tumor microenvironment and fully functional immune system in GEMMs offer unique advantages to other experimental models. Although mice rarely develop spontaneous PCa, modeling the recurrent genetic alterations of human patients has provided a much clearer picture of PCa progression. It is increasingly appreciated that many of the molecular events, such as p53

or Rb loss of function and *TMPRSS2-ERG* fusion, are insufficient to initiate PCa if they occur early individually. However, these events may prime prostate cells for future oncogenic transformation by altering their downstream transcriptional programs. PI3K pathway activation in the form of *Pten* loss, on the other hand, has proved to be an essential step in many models for the transition from PIN to adenocarcinoma. Further progression to CRPC appears to require PI3K signaling cooperation with other pathways (Figure 3). For example, the discovery of the reciprocal negative feedback loop between PI3K and AR signaling argues for the combined treatment targeting both pathways. The finding that loss of p53 and Rb at later stage further drives the tumor cells to undergo lineage switch provides new potential therapeutic avenues to inhibit disease progression towards neuroendocrine cancer. In comparing various GEMM studies, it is important to bear in mind that mouse strain background could be a complicating factor contributing to some of the controversies. For example, mice of the commonly used C57BL/6 genetic background are relatively tumor resistant [216]. PCa metastasis remains an underaddressed topic in GEMMs, as very few models show bone me-

tastasis, which is common in patients. While this may very well reflect intrinsic species differences, developing new models, such as late-stage activation of Wnt and AR variants models, may lead to new discoveries on the mechanisms of PCa metastasis. With the incorporation of new technologies such as CRISPR and single cell techniques, we envision that the next-generation mouse models will help dissect the signaling cross talks in PCa with more sophisticated modeling of molecular events in more refined tissue space.

Acknowledgements

Z.A.W. is supported by NIH grant R01GM116872 and D.N. is supported by the Santa Cruz Cancer Benefit Group.

Disclosure of conflict of interest

None.

Address correspondence to: Zhu A Wang, Department of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, CA 95064, USA. E-mail: zwang36@ucsc.edu

References

[1] Global Burden of Disease Cancer Collaboration, Fitzmaurice C, Dicker D, Pain A, Hamavid H, Moradi-Lakeh M, MacIntyre MF, Allen C, Hansen G, Woodbrook R, Wolfe C, Hamadeh RR, Moore A, Werdecker A, Gessner BD, Te Ao B, McMahon B, Karimkhani C, Yu C, Cooke GS, Schwebel DC, Carpenter DO, Pereira DM, Nash D, Kazi DS, De Leo D, Plass D, Ukwaja KN, Thurston GD, Yun Jin K, Simard EP, Mills E, Park EK, Catalá-López F, deVeber G, Gotay C, Khan G, Hosgood HD 3rd, Santos IS, Leasher JL, Singh J, Leigh J, Jonas JB, Sanabria J, Beardsley J, Jacobsen KH, Takahashi K, Franklin RC, Ronfani L, Montico M, Naldi L, Tonelli M, Geleijnse J, Petzold M, Shrimme MG, Younis M, Yonemoto N, Breitborde N, Yip P, Pourmalek F, Lotufo PA, Esteghamati A, Hankey GJ, Ali R, Lunevicius R, Malekzadeh R, Dellavalle R, Weintraub R, Lucas R, Hay R, Rojas-Rueda D, Westerman R, Sepanlou SG, Nolte S, Patten S, Weichenthal S, Abera SF, Fereshtehnejad SM, Shiue I, Driscoll T, Vasankari T, Alsharif U, Rahimi-Movaghar V, Vlassov VV, Marcenes WS, Mekonnen W, Melaku YA, Yano Y, Artaman A, Campos I, MacLachlan J, Mueller U, Kim D, Trilini M, Eshrati B, Williams HC, Shibuya K, Dandona R, Murthy K, Cowie B, Amare AT, Antonio CA, Castañeda-Orjuela C, van Gool CH, Violan-

te F, Oh IH, Deribe K, Soreide K, Knibbs L, Kereselidze M, Green M, Cardenas R, Roy N, Tillmann T, Li Y, Krueger H, Monasta L, Dey S, Sheikhbahaei S, Hafezi-Nejad N, Kumar GA, Sreeramareddy CT, Dandona L, Wang H, Vollset SE, Mokdad A, Salomon JA, Lozano R, Vos T, Forouzanfar M, Lopez A, Murray C and Naghavi M. The global burden of cancer 2013. *JAMA Oncol* 2015; 1: 505-27.

[2] Huggins C. Studies on prostatic cancer. II. The effect of castration on clinical patients with carcinoma of the prostate. *Arch Surg* 1941; 43: 209.

[3] Watson PA, Arora VK and Sawyers CL. Emerging mechanisms of resistance to androgen receptor inhibitors in prostate cancer. *Nat Rev Cancer* 2015; 15: 701-11.

[4] Toivanen R and Shen MM. Prostate organogenesis: tissue induction, hormonal regulation and cell type specification. *Development* 2017; 144: 1382-98.

[5] Shen MM and Abate-Shen C. Molecular genetics of prostate cancer: new prospects for old challenges. *Genes Dev* 2010; 24: 1967-2000.

[6] Armenia J, Wankowicz SAM, Liu D, Gao J, Kundra R, Reznik E, Chatila WK, Chakravarty D, Han GC, Coleman I, Montgomery B, Pritchard C, Morrissey C, Barbieri CE, Beltran H, Sboner A, Zafeiriou Z, Miranda S, Bielski CM, Penson AV, Tolonen C, Huang FW, Robinson D, Wu YM, Lonigro R, Garraway LA, Demichelis F, Kantoff PW, Taplin ME, Abida W, Taylor BS, Scher HI, Nelson PS, de Bono JS, Rubin MA, Sawyers CL, Chinnaiyan AM; PCF/SU2C International Prostate Cancer Dream Team, Schultz N, Van Allen EM. The long tail of oncogenic drivers in prostate cancer. *Nat Genet* 2018; 50: 645-51.

[7] Grasso CS, Wu YM, Robinson DR, Cao X, Dhannasekaran SM, Khan AP, Quist MJ, Jing X, Lonigro RJ, Brenner JC, Asangani IA, Ateeq B, Chun SY, Siddiqui J, Sam L, Anstett M, Mehra R, Prensner JR, Palanisamy N, Ryslik GA, Vandin F, Raphael BJ, Kunju LP, Rhodes DR, Pienta KJ, Chinnaiyan AM and Tomlins SA. The mutational landscape of lethal castration-resistant prostate cancer. *Nature* 2012; 487: 239-43.

[8] Barbieri CE, Baca SC, Lawrence MS, Demichelis F, Blattner M, Theurillat JP, White TA, Stojanov P, Van Allen E, Stransky N, Nickerson E, Chae SS, Boysen G, Auclair D, Onofrio RC, Park K, Kitabayashi N, MacDonald TY, Sheikh K, Vuong T, Guiducci C, Cibulskis K, Sivachenko A, Carter SL, Saksena G, Voet D, Hussain WM, Ramos AH, Winckler W, Redman MC, Ardlie K, Tewari AK, Mosquera JM, Rupp N, Wild PJ, Moch H, Morrissey C, Nelson PS, Kantoff PW, Gabriel SB, Golub TR, Meyerson M, Lander ES, Getz G, Rubin MA and Garraway LA. Exome sequencing identifies recurrent SPOP, FOXA1 and

Prostate cancer mouse model review

- MED12 mutations in prostate cancer. *Nat Genet* 2012; 44: 685-9.
- [9] Cancer Genome Atlas Research Network. The molecular taxonomy of primary prostate cancer. *Cell* 2015; 163: 1011-25.
- [10] Taylor BS, Schultz N, Hieronymus H, Gopalan A, Xiao Y, Carver BS, Arora VK, Kaushik P, Cerami E, Reva B, Antipin Y, Mitsiades N, Landers T, Dolgalev I, Major JE, Wilson M, Socci ND, Lash AE, Heguy A, Eastham JA, Scher HI, Reuter VE, Scardino PT, Sander C, Sawyers CL and Gerald WL. Integrative genomic profiling of human prostate cancer. *Cancer Cell* 2010; 18: 11-22.
- [11] Robinson D, Van Allen EM, Wu YM, Schultz N, Lonigro RJ, Mosquera JM, Montgomery B, Taplin ME, Pritchard CC, Attard G, Beltran H, Abida W, Bradley RK, Vinson J, Cao X, Vats P, Kunju LP, Hussain M, Feng FY, Tomlins SA, Cooney KA, Smith DC, Brennan C, Siddiqui J, Mehra R, Chen Y, Rathkopf DE, Morris MJ, Solomon SB, Durack JC, Reuter VE, Gopalan A, Gao J, Loda M, Lis RT, Bowden M, Balk SP, Gaviola G, Sougnez C, Gupta M, Yu EY, Mostaghel EA, Cheng HH, Mulcahy H, True LD, Plymate SR, Dvinge H, Ferraldeschi R, Flohr P, Miranda S, Zafeiriou Z, Tunariu N, Mateo J, Perez-Lopez R, Demichelis F, Robinson BD, Sboner A, Schiffman M, Nanus DM, Tagawa ST, Sigaras A, Eng KW, Elemento O, Sboner A, Heath EI, Scher HI, Pienta KJ, Kantoff P, de Bono JS, Rubin MA, Nelson PS, Garraway LA, Sawyers CL and Chinnaiyan AM. Integrative clinical genomics of advanced prostate cancer. *Cell* 2015; 161: 1215-28.
- [12] Quigley DA, Dang HX, Zhao SG, Lloyd P, Aggarwal R, Alumkal JJ, Foye A, Kothari V, Perry MD, Bailey AM, Playdle D, Barnard TJ, Zhang L, Zhang J, Youngren JF, Cieslik MP, Parolia A, Beer TM, Thomas G, Chi KN, Gleave M, Lack NA, Zoubeidi A, Reiter RE, Rettig MB, Witte O, Ryan CJ, Fong L, Kim W, Friedlander T, Chou J, Li H, Das R, Li H, Moussavi-Baygi R, Goodarzi H, Gilbert LA, Lara PN Jr, Evans CP, Goldstein TC, Stuart JM, Tomlins SA, Spratt DE, Cheetham RK, Cheng DT, Farh K, Gehring JS, Hakenberg J, Liao A, Febbo PG, Shon J, Sickler B, Batzoglou S, Knudsen KE, He HH, Huang J, Wyatt AW, Dehm SM, Ashworth A, Chinnaiyan AM, Maher CA, Small EJ and Feng FY. Genomic hallmarks and structural variation in metastatic prostate cancer. *Cell* 2018; 174: 758-69, e9.
- [13] van Bokhoven A, Varella-Garcia M, Korch C, Johannes WU, Smith EE, Miller HL, Nordeen SK, Miller GJ and Lucia MS. Molecular characterization of human prostate carcinoma cell lines. *Prostate* 2003; 57: 205-25.
- [14] Sobel RE and Sadar MD. Cell lines used in prostate cancer research: a compendium of old and new lines—part 2. *J Urol* 2005; 173: 360-72.
- [15] Sobel RE and Sadar MD. Cell lines used in prostate cancer research: a compendium of old and new lines—part 1. *J Urol* 2005; 173: 342-59.
- [16] Cunningham D and You Z. In vitro and in vivo model systems used in prostate cancer research. *J Biol Methods* 2015; 2.
- [17] Gao D, Vela I, Sboner A, laquinta PJ, Karthaus WR, Gopalan A, Dowling C, Wanjala JN, Undvall EA, Arora VK, Wongvipat J, Kossai M, Ramazanoglu S, Barboza LP, Di W, Cao Z, Zhang QF, Sirota I, Ran L, MacDonald TY, Beltran H, Mosquera JM, Touijer KA, Scardino PT, Laudone VP, Curtis KR, Rathkopf DE, Morris MJ, Danila DC, Slovin SF, Solomon SB, Eastham JA, Chi P, Carver B, Rubin MA, Scher HI, Clevers H, Sawyers CL and Chen Y. Organoid cultures derived from patients with advanced prostate cancer. *Cell* 2014; 159: 176-87.
- [18] Puca L, Bareja R, Prandi D, Shaw R, Benelli M, Karthaus WR, Hess J, Sigouros M, Donoghue A, Kossai M, Gao D, Cyrta J, Sailer V, Vosoughi A, Pauli C, Churakova Y, Cheung C, Deonaraine LD, McNary TJ, Rosati R, Tagawa ST, Nanus DM, Mosquera JM, Sawyers CL, Chen Y, Inghirami G, Rao RA, Grandori C, Elemento O, Sboner A, Demichelis F, Rubin MA and Beltran H. Patient derived organoids to model rare prostate cancer phenotypes. *Nat Commun* 2018; 9: 2404.
- [19] Lin D, Wyatt AW, Xue H, Wang Y, Dong X, Haggert A, Wu R, Brahmhatt S, Mo F, Jong L, Bell RH, Anderson S, Hurtado-Coll A, Fazli L, Sharma M, Beltran H, Rubin M, Cox M, Gout PW, Morris J, Goldenberg L, Volik SV, Gleave ME, Collins CC and Wang Y. High fidelity patient-derived xenografts for accelerating prostate cancer discovery and drug development. *Cancer Res* 2014; 74: 1272-83.
- [20] Tentler JJ, Tan AC, Weekes CD, Jimeno A, Leong S, Pitts TM, Arcaroli JJ, Messersmith WA and Eckhardt SG. Patient-derived tumour xenografts as models for oncology drug development. *Nat Rev Clin Oncol* 2012; 9: 338-50.
- [21] Vazquez F and Sellers WR. The PTEN tumor suppressor protein: an antagonist of phosphoinositide 3-kinase signaling. *Biochim Biophys Acta* 2000; 1470: M21-35.
- [22] Podsypanina K, Ellenson LH, Nemes A, Gu J, Tamura M, Yamada KM, Cordon-Cardo C, Cattoretti G, Fisher PE and Parsons R. Mutation of Pten/Mmac1 in mice causes neoplasia in multiple organ systems. *Proc Natl Acad Sci U S A* 1999; 96: 1563-8.
- [23] Di Cristofano A, Pesce B, Cordon-Cardo C and Pandolfi PP. Pten is essential for embryonic development and tumour suppression. *Nat Genet* 1998; 19: 348-55.
- [24] Wu X, Wu J, Huang J, Powell WC, Zhang J, Matusik RJ, Sangiorgi FO, Maxson RE, Sucov HM

Prostate cancer mouse model review

- and Roy-Burman P. Generation of a prostate epithelial cell-specific Cre transgenic mouse model for tissue-specific gene ablation. *Mech Dev* 2001; 101: 61-9.
- [25] Wang S, Gao J, Lei Q, Rozengurt N, Pritchard C, Jiao J, Thomas GV, Li G, Roy-Burman P, Nelson PS, Liu X and Wu H. Prostate-specific deletion of the murine Pten tumor suppressor gene leads to metastatic prostate cancer. *Cancer Cell* 2003; 4: 209-21.
- [26] Jiao J, Wang S, Qiao R, Vivanco I, Watson PA, Sawyers CL and Wu H. Murine cell lines derived from Pten null prostate cancer show the critical role of PTEN in hormone refractory prostate cancer development. *Cancer Res* 2007; 67: 6083-91.
- [27] Garcia AJ, Ruscetti M, Arenzana TL, Tran LM, Bianci-Frias D, Sybert E, Priceman SJ, Wu L, Nelson PS, Smale ST and Wu H. Pten null prostate epithelium promotes localized myeloid-derived suppressor cell expansion and immune suppression during tumor initiation and progression. *Mol Cell Biol* 2014; 34: 2017-28.
- [28] Luchman HA, Benediktsson H, Villemaire ML, Peterson AC and Jirik FR. The pace of prostatic intraepithelial neoplasia development is determined by the timing of Pten tumor suppressor gene excision. *PLoS One* 2008; 3: e3940.
- [29] Ratnacaram CK, Teletin M, Jiang M, Meng X, Chambon P and Metzger D. Temporally controlled ablation of PTEN in adult mouse prostate epithelium generates a model of invasive prostatic adenocarcinoma. *Proc Natl Acad Sci U S A* 2008; 105: 2521-6.
- [30] Trotman LC, Niki M, Dotan ZA, Koutcher JA, Di Cristofano A, Xiao A, Khoo AS, Roy-Burman P, Greenberg NM, Van Dyke T, Cordon-Cardo C and Pandolfi PP. Pten dose dictates cancer progression in the prostate. *PLoS Biol* 2003; 1: E59.
- [31] Wang X, Trotman LC, Koppie T, Alimonti A, Chen Z, Gao Z, Wang J, Erdjument-Bromage H, Tempst P, Cordon-Cardo C, Pandolfi PP and Jiang X. NEDD4-1 is a proto-oncogenic ubiquitin ligase for PTEN. *Cell* 2007; 128: 129-39.
- [32] Jia S, Liu Z, Zhang S, Liu P, Zhang L, Lee SH, Zhang J, Signoretti S, Loda M, Roberts TM and Zhao JJ. Essential roles of PI(3)K-p110beta in cell growth, metabolism and tumorigenesis. *Nature* 2008; 454: 776-9.
- [33] Lee SH, Poulgiannis G, Pyne S, Jia S, Zou L, Signoretti S, Loda M, Cantley LC and Roberts TM. A constitutively activated form of the p110beta isoform of PI3-kinase induces prostatic intraepithelial neoplasia in mice. *Proc Natl Acad Sci U S A* 2010; 107: 11002-7.
- [34] Majumder PK, Yeh JJ, George DJ, Febbo PG, Kum J, Xue Q, Bikoff R, Ma H, Kantoff PW, Golub TR, Loda M and Sellers WR. Prostate intraepithelial neoplasia induced by prostate restricted Akt activation: the MPAKT model. *Proc Natl Acad Sci U S A* 2003; 100: 7841-6.
- [35] Majumder PK, Febbo PG, Bikoff R, Berger R, Xue Q, McMahon LM, Manola J, Brugarolas J, McDonnell TJ, Golub TR, Loda M, Lane HA and Sellers WR. mTOR inhibition reverses Akt-dependent prostate intraepithelial neoplasia through regulation of apoptotic and HIF-1-dependent pathways. *Nat Med* 2004; 10: 594-601.
- [36] Laplante M and Sabatini DM. mTOR signaling in growth control and disease. *Cell* 2012; 149: 274-93.
- [37] Guertin DA, Stevens DM, Saitoh M, Kinkel S, Crosby K, Sheen JH, Mullholland DJ, Magnusson MA, Wu H and Sabatini DM. mTOR complex 2 is required for the development of prostate cancer induced by Pten loss in mice. *Cancer Cell* 2009; 15: 148-59.
- [38] Furic L, Rong L, Larsson O, Koumakpayi IH, Yoshida K, Brueschke A, Petroulakis E, Robichaud N, Pollak M, Gaboury LA, Pandolfi PP, Saad F and Sonenberg N. eIF4E phosphorylation promotes tumorigenesis and is associated with prostate cancer progression. *Proc Natl Acad Sci U S A* 2010; 107: 14134-9.
- [39] Ding M, Van der Kwast TH, Vellanki RN, Foltz WD, McKee TD, Sonenberg N, Pandolfi PP, Koritzinsky M and Wouters BG. The mTOR targets 4E-BP1/2 restrain tumor growth and promote hypoxia tolerance in PTEN-driven prostate cancer. *Mol Cancer Res* 2018; 16: 682-95.
- [40] Trotman LC, Alimonti A, Scaglioni PP, Koutcher JA, Cordon-Cardo C and Pandolfi PP. Identification of a tumour suppressor network opposing nuclear Akt function. *Nature* 2006; 441: 523-7.
- [41] Ma L, Teruya-Feldstein J, Behrendt N, Chen Z, Noda T, Hino O, Cordon-Cardo C and Pandolfi PP. Genetic analysis of Pten and Tsc2 functional interactions in the mouse reveals asymmetrical haploinsufficiency in tumor suppression. *Genes Dev* 2005; 19: 1779-86.
- [42] Pearson HB, Li J, Meniel VS, Fennell CM, Wang P, Montgomery KG, Rebello RJ, Macpherson AA, Koushyar S, Furic L, Cullinane C, Clarkson RW, Smalley MJ, Simpson KJ, Phesse TJ, Shepherd PR, Humbert PO, Sansom OJ and Phillips WA. Identification of Pik3ca mutation as a genetic driver of prostate cancer that cooperates with pten loss to accelerate progression and castration-resistant growth. *Cancer Discov* 2018; 8: 764-779.
- [43] Chen Z, Trotman LC, Shaffer D, Lin HK, Dotan ZA, Niki M, Koutcher JA, Scher HI, Ludwig T, Gerald W, Cordon-Cardo C and Pandolfi PP. Crucial role of p53-dependent cellular senescence in suppression of Pten-deficient tumorigenesis. *Nature* 2005; 436: 725-30.

Prostate cancer mouse model review

- [44] Cordon-Cardo C, Koff A, Drobnjak M, Capodiceci P, Osman I, Millard SS, Gaudin PB, Fazzari M, Zhang ZF, Massague J and Scher HI. Distinct altered patterns of p27KIP1 gene expression in benign prostatic hyperplasia and prostatic carcinoma. *J Natl Cancer Inst* 1998; 90: 1284-91.
- [45] Di Cristofano A, De Acetis M, Koff A, Cordon-Cardo C and Pandolfi PP. Pten and p27KIP1 cooperate in prostate cancer tumor suppression in the mouse. *Nat Genet* 2001; 27: 222-4.
- [46] Majumder PK, Grisanzio C, O'Connell F, Barry M, Brito JM, Xu Q, Guney I, Berger R, Herman P, Bikoff R, Fedele G, Baek WK, Wang S, Ellwood-Yen K, Wu H, Sawyers CL, Signoretti S, Hahn WC, Loda M and Sellers WR. A prostatic intraepithelial neoplasia-dependent p27 Kip1 checkpoint induces senescence and inhibits cell proliferation and cancer progression. *Cancer Cell* 2008; 14: 146-55.
- [47] Ahmad I, Patel R, Singh LB, Nixon C, Seywright M, Barnetson RJ, Brunton VG, Muller WJ, Edwards J, Sansom OJ and Leung HY. HER2 overcomes PTEN (loss)-induced senescence to cause aggressive prostate cancer. *Proc Natl Acad Sci U S A* 2011; 108: 16392-7.
- [48] Thomsen MK, Ambroisine L, Wynn S, Cheah KS, Foster CS, Fisher G, Berney DM, Møller H, Reuter VE, Scardino P, Cuzick J, Ragavan N, Singh PB, Martin FL, Butler CM, Cooper CS, Swain A; Transatlantic Prostate Group. SOX9 elevation in the prostate promotes proliferation and cooperates with PTEN loss to drive tumor formation. *Cancer Res* 2010; 70: 979-87.
- [49] Wang G, Lunardi A, Zhang J, Chen Z, Ala U, Webster KA, Tay Y, Gonzalez-Billalabeitia E, Egia A, Shaffer DR, Carver B, Liu XS, Taulli R, Kuo WP, Nardella C, Signoretti S, Cordon-Cardo C, Gerald WL and Pandolfi PP. Zbtb7a suppresses prostate cancer through repression of a Sox9-dependent pathway for cellular senescence bypass and tumor invasion. *Nat Genet* 2013; 45: 739-46.
- [50] Lunardi A, Ala U, Epping MT, Salmena L, Clohessy JG, Webster KA, Wang G, Mazzucchelli R, Bianconi M, Stack EC, Lis R, Patnaik A, Cantley LC, Bublely G, Cordon-Cardo C, Gerald WL, Montironi R, Signoretti S, Loda M, Nardella C and Pandolfi PP. A co-clinical approach identifies mechanisms and potential therapies for androgen deprivation resistance in prostate cancer. *Nat Genet* 2013; 45: 747-55.
- [51] Bezzi M, Seitzer N, Ishikawa T, Reschke M, Chen M, Wang G, Mitchell C, Ng C, Katon J, Lunardi A, Signoretti S, Clohessy JG, Zhang J and Pandolfi PP. Diverse genetic-driven immune landscapes dictate tumor progression through distinct mechanisms. *Nat Med* 2018; 24: 165-75.
- [52] Ding Z, Wu CJ, Chu GC, Xiao Y, Ho D, Zhang J, Perry SR, Labrot ES, Wu X, Lis R, Hoshida Y, Hiller D, Hu B, Jiang S, Zheng H, Stegh AH, Scott KL, Signoretti S, Bardeesy N, Wang YA, Hill DE, Golub TR, Stampfer MJ, Wong WH, Loda M, Mucci L, Chin L and DePinho RA. SMAD4-dependent barrier constrains prostate cancer growth and metastatic progression. *Nature* 2011; 470: 269-73.
- [53] Hao Y, Bjerke GA, Pietrzak K, Melhuish TA, Han Y, Turner SD, Frierson HF Jr and Wotton D. TGF-beta signaling limits lineage plasticity in prostate cancer. *PLoS Genet* 2018; 14: e1007409.
- [54] Hudlebusch HR, Santoni-Rugiu E, Simon R, Ralfkiaer E, Rossing HH, Johansen JV, Jørgensen M, Sauter G and Helin K. The histone methyltransferase and putative oncoprotein MMSET is overexpressed in a large variety of human tumors. *Clin Cancer Res* 2011; 17: 2919-33.
- [55] Ezponda T, Popovic R, Shah MY, Martinez-Garcia E, Zheng Y, Min DJ, Will C, Neri A, Kelleher NL, Yu J and Licht JD. The histone methyltransferase MMSET/WHSC1 activates TWIST1 to promote an epithelial-mesenchymal transition and invasive properties of prostate cancer. *Oncogene* 2013; 32: 2882-90.
- [56] Li N, Xue W, Yuan H, Dong B, Ding Y, Liu Y, Jiang M, Kan S, Sun T, Ren J, Pan Q, Li X, Zhang P, Hu G, Wang Y, Wang X, Li Q and Qin J. AKT-mediated stabilization of histone methyltransferase WHSC1 promotes prostate cancer metastasis. *J Clin Invest* 2017; 127: 1284-302.
- [57] Visvader JE. Cells of origin in cancer. *Nature* 2011; 469: 314-22.
- [58] Lee SH and Shen MM. Cell types of origin for prostate cancer. *Curr Opin Cell Biol* 2015; 37: 35-41.
- [59] Park JW, Lee JK, Phillips JW, Huang P, Cheng D, Huang J and Witte ON. Prostate epithelial cell of origin determines cancer differentiation state in an organoid transformation assay. *Proc Natl Acad Sci U S A* 2016; 113: 4482-7.
- [60] Lawson DA, Zong Y, Memarzadeh S, Xin L, Huang J and Witte ON. Basal epithelial stem cells are efficient targets for prostate cancer initiation. *Proc Natl Acad Sci U S A* 2010; 107: 2610-5.
- [61] Goldstein AS, Huang J, Guo C, Garraway IP and Witte ON. Identification of a cell of origin for human prostate cancer. *Science* 2010; 329: 568-71.
- [62] Choi N, Zhang B, Zhang L, Ittmann M and Xin L. Adult murine prostate basal and luminal cells are self-sustained lineages that can both serve as targets for prostate cancer initiation. *Cancer Cell* 2012; 21: 253-65.
- [63] Wang ZA, Mitrofanova A, Bergren SK, Abate-Shen C, Cardiff RD, Califano A and Shen MM. Lineage analysis of basal epithelial cells re-

Prostate cancer mouse model review

- veals their unexpected plasticity and supports a cell-of-origin model for prostate cancer heterogeneity. *Nat Cell Biol* 2013; 15: 274-83.
- [64] Wang ZA, Toivanen R, Bergren SK, Chambon P and Shen MM. Luminal cells are favored as the cell of origin for prostate cancer. *Cell Rep* 2014; 8: 1339-46.
- [65] Baca SC, Prandi D, Lawrence MS, Mosquera JM, Romanel A, Drier Y, Park K, Kitabayashi N, MacDonald TY, Ghandi M, Van Allen E, Kryukov GV, Sboner A, Theurillat JP, Soong TD, Nickerson E, Auclair D, Tewari A, Beltran H, Onofrio RC, Boysen G, Guiducci C, Barbieri CE, Cibulskis K, Sivachenko A, Carter SL, Saksena G, Voet D, Ramos AH, Winckler W, Cipicchio M, Ardlie K, Kantoff PW, Berger MF, Gabriel SB, Golub TR, Meyerson M, Lander ES, Elemento O, Getz G, Demichelis F, Rubin MA and Garraway LA. Punctuated evolution of prostate cancer genomes. *Cell* 2013; 153: 666-77.
- [66] Dang CV. MYC on the path to cancer. *Cell* 2012; 149: 22-35.
- [67] Hofmann JW, Zhao X, De Cecco M, Peterson AL, Pagliaroli L, Manivannan J, Hubbard GB, Ikeno Y, Zhang Y, Feng B, Li X, Serre T, Qi W, Van Remmen H, Miller RA, Bath KG, de Cabo R, Xu H, Neretti N and Sedivy JM. Reduced expression of MYC increases longevity and enhances healthspan. *Cell* 2015; 160: 477-88.
- [68] Murphy DJ, Junttila MR, Pouyet L, Karnezis A, Shchors K, Bui DA, Brown-Swigart L, Johnson L and Evan GI. Distinct thresholds govern Myc's biological output in vivo. *Cancer Cell* 2008; 14: 447-57.
- [69] Gurel B, Iwata T, Koh CM, Jenkins RB, Lan F, Van Dang C, Hicks JL, Morgan J, Cornish TC, Sutcliffe S, Isaacs WB, Luo J and De Marzo AM. Nuclear MYC protein overexpression is an early alteration in human prostate carcinogenesis. *Mod Pathol* 2008; 21: 1156-67.
- [70] Ellwood-Yen K, Graeber TG, Wongvipat J, Iruela-Arispe ML, Zhang J, Matusik R, Thomas GV and Sawyers CL. Myc-driven murine prostate cancer shares molecular features with human prostate tumors. *Cancer Cell* 2003; 4: 223-38.
- [71] Zhang X, Lee C, Ng PY, Rubin M, Shabsigh A and Buttyan R. Prostatic neoplasia in transgenic mice with prostate-directed overexpression of the c-myc oncoprotein. *Prostate* 2000; 43: 278-85.
- [72] Kim J, Eltoun IE, Roh M, Wang J and Abdulkadir SA. Interactions between cells with distinct mutations in c-MYC and Pten in prostate cancer. *PLoS Genet* 2009; 5: e1000542.
- [73] Wang J, Kim J, Roh M, Franco OE, Hayward SW, Wills ML and Abdulkadir SA. Pim1 kinase synergizes with c-MYC to induce advanced prostate carcinoma. *Oncogene* 2010; 29: 2477-87.
- [74] Stoyanova T, Cooper AR, Drake JM, Liu X, Armstrong AJ, Pienta KJ, Zhang H, Kohn DB, Huang J, Witte ON and Goldstein AS. Prostate cancer originating in basal cells progresses to adenocarcinoma propagated by luminal-like cells. *Proc Natl Acad Sci U S A* 2013; 110: 20111-6.
- [75] Clegg NJ, Couto SS, Wongvipat J, Hieronymus H, Carver BS, Taylor BS, Ellwood-Yen K, Gerald WL, Sander C and Sawyers CL. MYC cooperates with AKT in prostate tumorigenesis and alters sensitivity to mTOR inhibitors. *PLoS One* 2011; 6: e17449.
- [76] Kim J, Roh M, Doubinskaia I, Algarroba GN, Eltoun IE and Abdulkadir SA. A mouse model of heterogeneous, c-MYC-initiated prostate cancer with loss of Pten and p53. *Oncogene* 2012; 31: 322-32.
- [77] Beltran H, Rickman DS, Park K, Chae SS, Sboner A, MacDonald TY, Wang Y, Sheikh KL, Terry S, Tagawa ST, Dhir R, Nelson JB, de la Taille A, Allory Y, Gerstein MB, Perner S, Pienta KJ, Chinnaiyan AM, Wang Y, Collins CC, Gleave ME, Demichelis F, Nanus DM and Rubin MA. Molecular characterization of neuroendocrine prostate cancer and identification of new drug targets. *Cancer Discov* 2011; 1: 487-95.
- [78] Lee JK, Phillips JW, Smith BA, Park JW, Stoyanova T, McCaffrey EF, Baertsch R, Sokolov A, Meyerowitz JG, Mathis C, Cheng D, Stuart JM, Shokat KM, Gustafson WC, Huang J and Witte ON. N-myc drives neuroendocrine prostate cancer initiated from human prostate epithelial cells. *Cancer Cell* 2016; 29: 536-47.
- [79] Dardenne E, Beltran H, Benelli M, Gayvert K, Berger A, Puca L, Cyrta J, Sboner A, Noorzad Z, MacDonald T, Cheung C, Yuen KS, Gao D, Chen Y, Eilers M, Mosquera JM, Robinson BD, Elemento O, Rubin MA, Demichelis F and Rickman DS. N-Myc induces an EZH2-mediated transcriptional program driving neuroendocrine prostate cancer. *Cancer Cell* 2016; 30: 563-77.
- [80] He WW, Scivolino PJ, Wing J, Augustus M, Hudson P, Meissner PS, Curtis RT, Shell BK, Bostwick DG, Tindall DJ, Gelmann EP, Abate-Shen C and Carter KC. A novel human prostate-specific, androgen-regulated homeobox gene (NKX3. 1) that maps to 8p21, a region frequently deleted in prostate cancer. *Genomics* 1997; 43: 69-77.
- [81] Emmert-Buck MR, Vocke CD, Pozzatti RO, Duray PH, Jennings SB, Florence CD, Zhuang Z, Bostwick DG, Liotta LA and Linehan WM. Allelic loss on chromosome 8p12-21 in microdissected prostatic intraepithelial neoplasia. *Cancer Res* 1995; 55: 2959-62.
- [82] Asatiani E, Huang WX, Wang A, Rodriguez Ortner E, Cavalli LR, Haddad BR and Gelmann EP. Deletion, methylation, and expression of the NKX3. 1 suppressor gene in primary human prostate cancer. *Cancer Res* 2005; 65: 1164-73.

Prostate cancer mouse model review

- [83] Debelec-Butuner B, Bostancı A, Özcan F, Singin O, Karamil S, Aslan M, Roggenbuck D and Korkmaz KS. Loss of NKX3. 1 expression in human prostate cancers correlates with tumor progression. *Cancer Invest* 2019; 37: 113-126.
- [84] Bhatia-Gaur R, Donjacour AA, Scivolino PJ, Kim M, Desai N, Young P, Norton CR, Gridley T, Cardiff RD, Cunha GR, Abate-Shen C and Shen MM. Roles for Nkx3.1 in prostate development and cancer. *Genes Dev* 1999; 13: 966-77.
- [85] Kim MJ, Bhatia-Gaur R, Banach-Petrosky WA, Desai N, Wang Y, Hayward SW, Cunha GR, Cardiff RD, Shen MM and Abate-Shen C. Nkx3.1 mutant mice recapitulate early stages of prostate carcinogenesis. *Cancer Res* 2002; 62: 2999-3004.
- [86] Abdulkadir SA, Magee JA, Peters TJ, Kaleem Z, Naughton CK, Humphrey PA and Milbrandt J. Conditional loss of Nkx3.1 in adult mice induces prostatic intraepithelial neoplasia. *Mol Cell Biol* 2002; 22: 1495-503.
- [87] Song H, Zhang B, Watson MA, Humphrey PA, Lim H and Milbrandt J. Loss of Nkx3.1 leads to the activation of discrete downstream target genes during prostate tumorigenesis. *Oncogene* 2009; 28: 3307-19.
- [88] Magee JA, Abdulkadir SA and Milbrandt J. Haploinsufficiency at the Nkx3.1 locus. A paradigm for stochastic, dosage-sensitive gene regulation during tumor initiation. *Cancer Cell* 2003; 3: 273-83.
- [89] Kim MJ, Cardiff RD, Desai N, Banach-Petrosky WA, Parsons R, Shen MM and Abate-Shen C. Cooperativity of Nkx3.1 and Pten loss of function in a mouse model of prostate carcinogenesis. *Proc Natl Acad Sci U S A* 2002; 99: 2884-9.
- [90] Ouyang X, DeWeese TL, Nelson WG and Abate-Shen C. Loss-of-function of Nkx3.1 promotes increased oxidative damage in prostate carcinogenesis. *Cancer Res* 2005; 65: 6773-9.
- [91] Abate-Shen C, Banach-Petrosky WA, Sun X, Economides KD, Desai N, Gregg JP, Borowsky AD, Cardiff RD and Shen MM. Nkx3.1; Pten mutant mice develop invasive prostate adenocarcinoma and lymph node metastases. *Cancer Res* 2003; 63: 3886-90.
- [92] Gao H, Ouyang X, Banach-Petrosky WA, Shen MM and Abate-Shen C. Emergence of androgen independence at early stages of prostate cancer progression in Nkx3.1; Pten mice. *Cancer Res* 2006; 66: 7929-33.
- [93] Xie Q and Wang ZA. Transcriptional regulation of the Nkx3.1 gene in prostate luminal stem cell specification and cancer initiation via its 3' genomic region. *J Biol Chem* 2017; 292: 13521-13530.
- [94] Lei Q, Jiao J, Xin L, Chang CJ, Wang S, Gao J, Gleave ME, Witte ON, Liu X and Wu H. NKX3.1 stabilizes p53, inhibits AKT activation, and blocks prostate cancer initiation caused by PTEN loss. *Cancer Cell* 2006; 9: 367-78.
- [95] Kumar-Sinha C, Tomlins SA and Chinnaiyan AM. Recurrent gene fusions in prostate cancer. *Nat Rev Cancer* 2008; 8: 497-511.
- [96] Demichelis F, Fall K, Perner S, Andr n O, Schmidt F, Setlur SR, Hoshida Y, Mosquera JM, Pawitan Y, Lee C and Adami HO. TMPRSS2: ERG gene fusion associated with lethal prostate cancer in a watchful waiting cohort. *Oncogene* 2007; 26: 4596.
- [97] Tomlins SA, Rhodes DR, Perner S, Dhanasekaran SM, Mehra R, Sun XW, Varambally S, Cao X, Tchinda J, Kuefer R, Lee C, Montie JE, Shah RB, Pienta KJ, Rubin MA and Chinnaiyan AM. Recurrent fusion of TMPRSS2 and ETS transcription factor genes in prostate cancer. *Science* 2005; 310: 644-8.
- [98] Tomlins SA, Laxman B, Dhanasekaran SM, Helgeson BE, Cao X, Morris DS, Menon A, Jing X, Cao Q, Han B, Yu J, Wang L, Montie JE, Rubin MA, Pienta KJ, Roulston D, Shah RB, Varambally S, Mehra R and Chinnaiyan AM. Distinct classes of chromosomal rearrangements create oncogenic ETS gene fusions in prostate cancer. *Nature* 2007; 448: 595-9.
- [99] Shin S, Kim TD, Jin F, van Deursen JM, Dehm SM, Tindall DJ, Grande JP, Munz JM, Vasmatzis G and Janknecht R. Induction of prostatic intraepithelial neoplasia and modulation of androgen receptor by ETS variant 1/ETS-related protein 81. *Cancer Res* 2009; 69: 8102-10.
- [100] Klezovitch O, Risk M, Coleman I, Lucas JM, Null M, True LD, Nelson PS and Vasioukhin V. A causal role for ERG in neoplastic transformation of prostate epithelium. *Proc Natl Acad Sci U S A* 2008; 105: 2105-10.
- [101] Nguyen LT, Tretiakova MS, Silvis MR, Lucas J, Klezovitch O, Coleman I, Bolouri H, Kutuyavin VI, Morrissey C, True LD, Nelson PS and Vasioukhin V. ERG activates the YAP1 transcriptional program and induces the development of age-related prostate tumors. *Cancer Cell* 2015; 27: 797-808.
- [102] Carver BS, Tran J, Chen Z, Carracedo-Perez A, Alimonti A, Nardella C, Gopalan A, Scardino PT, Cordon-Cardo C, Gerald W and Pandolfi PP. ETS rearrangements and prostate cancer initiation. *Nature* 2009; 457: E1; discussion E2-3.
- [103] Carver BS, Tran J, Gopalan A, Chen Z, Shaikh S, Carracedo A, Alimonti A, Nardella C, Varmeh S, Scardino PT, Cordon-Cardo C, Gerald W and Pandolfi PP. Aberrant ERG expression cooperates with loss of PTEN to promote cancer progression in the prostate. *Nat Genet* 2009; 41: 619-24.
- [104] Chen Y, Chi P, Rockowitz S, laquinta PJ, Shamu T, Shukla S, Gao D, Sirota I, Carver BS, Wongvipat J, Scher HI, Zheng D and Sawyers CL. ETS

Prostate cancer mouse model review

- factors reprogram the androgen receptor cisrome and prime prostate tumorigenesis in response to PTEN loss. *Nat Med* 2013; 19: 1023-9.
- [105] Baena E, Shao Z, Linn DE, Glass K, Hamblen MJ, Fujiwara Y, Kim J, Nguyen M, Zhang X, Godinho FJ, Bronson RT, Mucci LA, Loda M, Yuan GC, Orkin SH and Li Z. ETV1 directs androgen metabolism and confers aggressive prostate cancer in targeted mice and patients. *Genes Dev* 2013; 27: 683-98.
- [106] King JC, Xu J, Wongvipat J, Hieronymus H, Carver BS, Leung DH, Taylor BS, Sander C, Cardiff RD, Couto SS, Gerald WL and Sawyers CL. Cooperativity of TMPRSS2-ERG with PI3-kinase pathway activation in prostate oncogenesis. *Nature genetics*. *Nat Genet* 2009; 41: 524-6.
- [107] Casey OM, Fang L, Hynes PG, Abou-Kheir WG, Martin PL, Tillman HS, Petrovics G, Awwad HO, Ward Y, Lake R, Zhang L and Kelly K. TMPRSS2-driven ERG expression in vivo increases self-renewal and maintains expression in a castration resistant subpopulation. *PLoS One* 2012; 7: e41668.
- [108] Yoshimoto M, Joshua AM, Cunha IW, Coudry RA, Fonseca FP, Ludkovski O, Zielenska M, Soares FA and Squire JA. Absence of TMPRSS2: ERG fusions and PTEN losses in prostate cancer is associated with a favorable outcome. *Mod Pathol* 2008; 21: 1451-60.
- [109] Hermans KG, van Marion R, van Dekken H, Jenster G, van Weerden WM and Trapman J. TMPRSS2: ERG fusion by translocation or interstitial deletion is highly relevant in androgen-dependent prostate cancer, but is bypassed in late-stage androgen receptor-negative prostate cancer. *Cancer Res* 2006; 66: 10658-63.
- [110] Perner S, Demichelis F, Beroukhir R, Schmidt FH, Mosquera JM, Setlur S, Tchinda J, Tomlins SA, Hofer MD, Pienta KG, Kuefer R, Vessella R, Sun XW, Meyerson M, Lee C, Sellers WR, Chinnaiyan AM and Rubin MA. TMPRSS2:ERG fusion-associated deletions provide insight into the heterogeneity of prostate cancer. *Cancer Res* 2006; 66: 8337-41.
- [111] Linn DE, Penney KL, Bronson RT, Mucci LA and Li Z. Deletion of interstitial genes between TMPRSS2 and ERG promotes prostate cancer progression. *Cancer Res* 2016; 76: 1869-81.
- [112] Kan Z, Jaiswal BS, Stinson J, Janakiraman V, Bhatt D, Stern HM, Yue P, Haverty PM, Bourgon R, Zheng J, Moorhead M, Chaudhuri S, Tomsho LP, Peters BA, Pujara K, Cordes S, Davis DP, Carlton VE, Yuan W, Li L, Wang W, Eigenbrot C, Kaminker JS, Eberhard DA, Waring P, Schuster SC, Modrusan Z, Zhang Z, Stokoe D, de Sauvage FJ, Faham M and Seshagiri S. Diverse somatic mutation patterns and pathway alterations in human cancers. *Nature* 2010; 466: 869-73.
- [113] Blattner M, Lee DJ, O'Reilly C, Park K, MacDonald TY, Khani F, Turner KR, Chiu YL, Wild PJ, Dolgalev I, Heguy A, Sboner A, Ramazangolu S, Hieronymus H, Sawyers C, Tewari AK, Moch H, Yoon GS, Known YC, Andr n O, Fall K, Demichelis F, Mosquera JM, Robinson BD, Barbieri CE and Rubin MA. SPOP mutations in prostate cancer across demographically diverse patient cohorts. *Neoplasia* 2014; 16: 14-20.
- [114] Boysen G, Barbieri CE, Prandi D, Blattner M, Chae SS, Dahija A, Nataraj S, Huang D, Marotz C, Xu L, Huang J, Lecca P, Chhangawala S, Liu D, Zhou P, Sboner A, de Bono JS, Demichelis F, Houvras Y and Rubin MA. SPOP mutation leads to genomic instability in prostate cancer. *Elife* 2015; 4.
- [115] Geng C, Kaochar S, Li M, Rajapakshe K, Fiskus W, Dong J, Foley C, Dong B, Zhang L, Kwon OJ, Shah SS, Bolaki M, Xin L, Ittmann M, O'Malley BW, Coarfa C and Mitsiades N. SPOP regulates prostate epithelial cell proliferation and promotes ubiquitination and turnover of c-MYC oncoprotein. *Oncogene* 2017; 36: 4767-77.
- [116] Blattner M, Liu D, Robinson BD, Huang D, Poliakov A, Gao D, Nataraj S, Deonaraine LD, Augello MA, Sailer V, Ponnala L, Ittmann M, Chinnaiyan AM, Sboner A, Chen Y, Rubin MA and Barbieri CE. SPOP mutation drives prostate tumorigenesis in vivo through coordinate regulation of PI3K/mTOR and AR signaling. *Cancer Cell* 2017; 31: 436-51.
- [117] Gan W, Dai X, Lunardi A, Li Z, Inuzuka H, Liu P, Varmeh S, Zhang J, Cheng L, Sun Y, Asara JM, Beck AH, Huang J, Pandolfi PP and Wei W. SPOP promotes ubiquitination and degradation of the ERG oncoprotein to suppress prostate cancer progression. *Mol Cell* 2015; 59: 917-30.
- [118] An J, Ren S, Murphy SJ, Dalangood S, Chang C, Pang X, Cui Y, Wang L, Pan Y, Zhang X, Zhu Y, Wang C, Halling GC, Cheng L, Sukov WR, Karnes RJ, Vasmatzis G, Zhang Q, Zhang J, Cheville JC, Yan J, Sun Y and Huang H. Truncated ERG oncoproteins from TMPRSS2-ERG fusions are resistant to SPOP-mediated proteasome degradation. *Mol Cell* 2015; 59: 904-16.
- [119] Shoag J, Liu D, Blattner M, Sboner A, Park K, Deonaraine L, Robinson BD, Mosquera JM, Chen Y, Rubin MA and Barbieri CE. SPOP mutation drives prostate neoplasia without stabilizing oncogenic transcription factor ERG. *J Clin Invest* 2018; 128: 381-6.
- [120] Nelson WJ and Nusse R. Convergence of Wnt, beta-catenin, and cadherin pathways. *Science* 2004; 303: 1483-7.
- [121] Anastas JN and Moon RT. WNT signalling pathways as therapeutic targets in cancer. *Nat Rev Cancer* 2013; 13: 11-26.

Prostate cancer mouse model review

- [122] Harada N, Tamai Y, Ishikawa T, Sauer B, Takaku K, Oshima M and Taketo MM. Intestinal polypoidism in mice with a dominant stable mutation of the beta-catenin gene. *EMBO J* 1999; 18: 5931-42.
- [123] Gounari F, Signoretti S, Bronson R, Klein L, Sellers WR, Kum J, Siermann A, Taketo MM, von Boehmer H and Khazaie K. Stabilization of beta-catenin induces lesions reminiscent of prostatic intraepithelial neoplasia, but terminal squamous transdifferentiation of other secretory epithelia. *Oncogene* 2002; 21: 4099-107.
- [124] Bierie B, Nozawa M, Renou JP, Shillingford JM, Morgan F, Oka T, Taketo MM, Cardiff RD, Miyoshi K, Wagner KU, Robinson GW and Hennighausen L. Activation of beta-catenin in prostate epithelium induces hyperplasias and squamous transdifferentiation. *Oncogene* 2003; 22: 3875-87.
- [125] Yu X, Wang Y, Jiang M, Bierie B, Roy-Burman P, Shen MM, Taketo MM, Wills M and Matusik RJ. Activation of beta-Catenin in mouse prostate causes HGPIN and continuous prostate growth after castration. *Prostate* 2009; 69: 249-62.
- [126] Yu X, Wang Y, DeGraff DJ, Wills ML and Matusik RJ. Wnt/beta-catenin activation promotes prostate tumor progression in a mouse model. *Oncogene* 2011; 30: 1868-79.
- [127] Pearson HB, Phesse TJ and Clarke AR. K-ras and Wnt signaling synergize to accelerate prostate tumorigenesis in the mouse. *Cancer Res* 2009; 69: 94-101.
- [128] He Y, Hooker E, Yu EJ, Cunha GR, Liao L, Xu J, Earl A, Wu H, Gonzalgo ML and Sun Z. Androgen signaling is essential for development of prostate cancer initiated from prostatic basal cells. *Oncogene* 2019; 38: 2337-50.
- [129] Francis JC, Thomsen MK, Taketo MM and Swain A. Beta-catenin is required for prostate development and cooperates with Pten loss to drive invasive carcinoma. *PLoS Genet* 2013; 9: e1003180.
- [130] Bruxvoort KJ, Charbonneau HM, Giambardi TA, Goolsby JC, Qian CN, Zylstra CR, Robinson DR, Roy-Burman P, Shaw AK, Buckner-Berghuis BD, Sigler RE, Resau JH, Sullivan R, Bushman W and Williams BO. Inactivation of Apc in the mouse prostate causes prostate carcinoma. *Cancer Res* 2007; 67: 2490-6.
- [131] Shahi P, Park D, Pond AC, Seethamagari M, Chiou SH, Cho K, Carstens JL, Decker WK, McCrea PD, Ittmann MM, Rosen JM and Spencer DM. Activation of Wnt signaling by chemically induced dimerization of LRP5 disrupts cellular homeostasis. *PLoS One* 2012; 7: e30814.
- [132] Zong Y, Huang J, Sankarasharma D, Morikawa T, Fukayama M, Epstein JI, Chada KK and Witte ON. Stromal epigenetic dysregulation is sufficient to initiate mouse prostate cancer via paracrine Wnt signaling. *Proc Natl Acad Sci U S A* 2012; 109: E3395-404.
- [133] Li X, Placencio V, Iturregui JM, Uwamariya C, Sharif-Afshar AR, Koyama T, Hayward SW and Bhowmick NA. Prostate tumor progression is mediated by a paracrine TGF-beta/Wnt3a signaling axis. *Oncogene* 2008; 27: 7118-30.
- [134] McKay MM and Morrison DK. Integrating signals from RTKs to ERK/MAPK. *Oncogene* 2007; 26: 3113-21.
- [135] Agell L, Hernández S, Salido M, de Muga S, Juanpere N, Arumí-Uria M, Menendez S, Lorenzo M, Lorente JA, Serrano S and Lloreta J. PI3K signaling pathway is activated by PIK3CA mRNA overexpression and copy gain in prostate tumors, but PIK3CA, BRAF, KRAS and AKT1 mutations are infrequent events. *Mod Pathol* 2011; 24: 443-52.
- [136] Scherl A, Li JF, Cardiff RD and Schreiber-Agus N. Prostatic intraepithelial neoplasia and intestinal metaplasia in prostates of probasin-RAS transgenic mice. *Prostate* 2004; 59: 448-59.
- [137] Jeong JH, Wang Z, Guimaraes AS, Ouyang X, Figueiredo JL, Ding Z, Jiang S, Guney I, Kang GH, Shin E, Hahn WC, Loda MF, Abate-Shen C, Weissleder R and Chin L. BRAF activation initiates but does not maintain invasive prostate adenocarcinoma. *PLoS One* 2008; 3: e3949.
- [138] Mulholland DJ, Kobayashi N, Ruscetti M, Zhi A, Tran LM, Huang J, Gleave M and Wu H. Pten loss and RAS/MAPK activation cooperate to promote EMT and metastasis initiated from prostate cancer stem/progenitor cells. *Cancer Res* 2012; 72: 1878-89.
- [139] Castellano E and Downward J. RAS Interaction with PI3K: more than just another effector pathway. *Genes Cancer* 2011; 2: 261-74.
- [140] Pylayeva-Gupta Y, Grabocka E and Bar-Sagi D. RAS oncogenes: weaving a tumorigenic web. *Nat Rev Cancer* 2011; 11: 761-74.
- [141] Ruscetti M, Quach B, Dadashian EL, Mulholland DJ and Wu H. Tracking and functional characterization of epithelial-mesenchymal transition and mesenchymal tumor cells during prostate cancer metastasis. *Cancer Res* 2015; 75: 2749-59.
- [142] Ruscetti M, Dadashian EL, Guo W, Quach B, Mulholland DJ, Park JW, Tran LM, Kobayashi N, Bianchi-Frias D, Xing Y, Nelson PS and Wu H. HDAC inhibition impedes epithelial-mesenchymal plasticity and suppresses metastatic, castration-resistant prostate cancer. *Oncogene* 2016; 35: 3781-95.
- [143] Aytes A, Mitrofanova A, Kinkade CW, Lefebvre C, Lei M, Phelan V, LeKaye HC, Koutcher JA, Cardiff RD, Califano A, Shen MM and Abate-Shen C. ETV4 promotes metastasis in response to activation of PI3-kinase and Ras

Prostate cancer mouse model review

- signaling in a mouse model of advanced prostate cancer. *Proc Natl Acad Sci U S A* 2013; 110: E3506-15.
- [144] Carracedo A, Ma L, Teruya-Feldstein J, Rojo F, Salmena L, Alimonti A, Egia A, Sasaki AT, Thomas G, Kozma SC, Papa A, Nardella C, Cantley LC, Baselga J and Pandolfi PP. Inhibition of mTORC1 leads to MAPK pathway activation through a PI3K-dependent feedback loop in human cancer. *J Clin Invest* 2008; 118: 3065-74.
- [145] Chen M, Zhang J, Sampieri K, Clohessy JG, Mendez L, Gonzalez-Billalabeitia E, Liu XS, Lee YR, Fung J, Katon JM, Menon AV, Webster KA, Ng C, Palumbieri MD, Diolombi MS, Breitkopf SB, Teruya-Feldstein J, Signoretti S, Bronson RT, Asara JM, Castillo-Martin M, Cordon-Cardo C and Pandolfi PP. An aberrant SREBP-dependent lipogenic program promotes metastatic prostate cancer. *Nat Genet* 2018; 50: 206-218.
- [146] Turner N and Grose R. Fibroblast growth factor signalling: from development to cancer. *Nat Rev Cancer* 2010; 10: 116-29.
- [147] Babina IS and Turner NC. Advances and challenges in targeting FGFR signalling in cancer. *Nat Rev Cancer* 2017; 17: 318-32.
- [148] Memarzadeh S, Xin L, Mulholland DJ, Mansukhani A, Wu H, Teitell MA and Witte ON. Enhanced paracrine FGF10 expression promotes formation of multifocal prostate adenocarcinoma and an increase in epithelial androgen receptor. *Cancer Cell* 2007; 12: 572-85.
- [149] Memarzadeh S, Cai H, Janzen DM, Xin L, Lukacs R, Riedinger M, Zong Y, DeGendt K, Verhoeven G, Huang J and Witte ON. Role of autonomous androgen receptor signaling in prostate cancer initiation is dichotomous and depends on the oncogenic signal. *Proc Natl Acad Sci U S A* 2011; 108: 7962-7.
- [150] Li Q, Ingram L, Kim S, Beharry Z, Cooper JA and Cai H. Paracrine fibroblast growth factor initiates oncogenic synergy with epithelial FGFR/ Src transformation in prostate tumor progression. *Neoplasia* 2018; 20: 233-43.
- [151] Lukacs RU, Memarzadeh S, Wu H and Witte ON. Bmi-1 is a crucial regulator of prostate stem cell self-renewal and malignant transformation. *Cell Stem Cell* 2010; 7: 682-93.
- [152] Foster BA, Evangelou A, Gingrich JR, Kaplan PJ, DeMayo F and Greenberg NM. Enforced expression of FGF-7 promotes epithelial hyperplasia whereas a dominant negative FGFR2iib promotes the emergence of neuroendocrine phenotype in prostate glands of transgenic mice. *Differentiation* 2002; 70: 624-32.
- [153] Song Z, Wu X, Powell WC, Cardiff RD, Cohen MB, Tin RT, Matusik RJ, Miller GJ and Roy-Burman P. Fibroblast growth factor 8 isoform B overexpression in prostate epithelium: a new mouse model for prostatic intraepithelial neoplasia. *Cancer Res* 2002; 62: 5096-105.
- [154] Wang F, McKeenan K, Yu C, Ittmann M and McKeenan WL. Chronic activity of ectopic type 1 fibroblast growth factor receptor tyrosine kinase in prostate epithelium results in hyperplasia accompanied by intraepithelial neoplasia. *Prostate* 2004; 58: 1-12.
- [155] Jin C, McKeenan K, Guo W, Jauma S, Ittmann MM, Foster B, Greenberg NM, McKeenan WL and Wang F. Cooperation between ectopic FGFR1 and depression of FGFR2 in induction of prostatic intraepithelial neoplasia in the mouse prostate. *Cancer Res* 2003; 63: 8784-90.
- [156] Freeman KW, Welm BE, Gangula RD, Rosen JM, Ittmann M, Greenberg NM and Spencer DM. Inducible prostate intraepithelial neoplasia with reversible hyperplasia in conditional FGFR1-expressing mice. *Cancer Res* 2003; 63: 8256-63.
- [157] Acevedo VD, Gangula RD, Freeman KW, Li R, Zhang Y, Wang F, Ayala GE, Peterson LE, Ittmann M and Spencer DM. Inducible FGFR-1 activation leads to irreversible prostate adenocarcinoma and an epithelial-to-mesenchymal transition. *Cancer Cell* 2007; 12: 559-71.
- [158] Zhong C, Saribekyan G, Liao CP, Cohen MB and Roy-Burman P. Cooperation between FGF8b overexpression and PTEN deficiency in prostate tumorigenesis. *Cancer Res* 2006; 66: 2188-94.
- [159] Carstens JL, Shahi P, Van Tsang S, Smith B, Creighton CJ, Zhang Y, Seamans A, Seethamagari M, Vedula I, Levitt JM, Ittmann MM, Rowley DR and Spencer DM. FGFR1-WNT-TGF-beta signaling in prostate cancer mouse models recapitulates human reactive stroma. *Cancer Res* 2014; 74: 609-20.
- [160] Bluemn EG, Coleman IM, Lucas JM, Coleman RT, Hernandez-Lopez S, Tharakan R, Bianchi-Frias D, Dumpit RF, Kaipainen A, Corella AN, Yang YC, Nyquist MD, Mostaghel E, Hsieh AC, Zhang X, Corey E, Brown LG, Nguyen HM, Pienta K, Ittmann M, Schweizer M, True LD, Wise D, Rennie PS, Vessella RL, Morrissey C and Nelson PS. Androgen receptor pathway-independent prostate cancer is sustained through FGF signaling. *Cancer Cell* 2017; 32: 474-89, e6.
- [161] Sherr CJ and McCormick F. The RB and p53 pathways in cancer. *Cancer Cell* 2002; 2: 103-12.
- [162] Harris SL and Levine AJ. The p53 pathway: positive and negative feedback loops. *Oncogene* 2005; 24: 2899-908.
- [163] Bookstein R, MacGrogan D, Hilsenbeck SG, Sharkey F and Allred DC. p53 is mutated in a subset of advanced-stage prostate cancers. *Cancer Res* 1993; 53: 3369-73.

Prostate cancer mouse model review

- [164] Schlomm T, Iwers L, Kirstein P, Jessen B, Köllermann J, Minner S, Passow-Drolet A, Mirlacher M, Milde-Langosch K, Graefen M, Haese A, Steuber T, Simon R, Huland H, Sauter G and Erbersdobler A. Clinical significance of p53 alterations in surgically treated prostate cancers. *Mod Pathol* 2008; 21: 1371-8.
- [165] Qian J, Hirasawa K, Bostwick DG, Bergstralh EJ, Slezak JM, Anderl KL, Borell TJ, Lieber MM and Jenkins RB. Loss of p53 and c-myc overrepresentation in stage T(2-3)N(1-3)M(0) prostate cancer are potential markers for cancer progression. *Mod Pathol* 2002; 15: 35-44.
- [166] Navone NM, Labate ME, Troncoso P, Pisters LL, Conti CJ, von Eschenbach AC and Logothetis CJ. p53 mutations in prostate cancer bone metastases suggest that selected p53 mutants in the primary site define foci with metastatic potential. *J Urol* 1999; 161: 304-8.
- [167] Mirchandani D, Zheng J, Miller GJ, Ghosh AK, Shibata DK, Cote RJ and Roy-Burman P. Heterogeneity in intratumor distribution of p53 mutations in human prostate cancer. *Am J Pathol* 1995; 147: 92-101.
- [168] Muller PA and Vousden KH. p53 mutations in cancer. *Nat Cell Biol* 2013; 15: 2-8.
- [169] Zhou Z, Flesken-Nikitin A, Corney DC, Wang W, Goodrich DW, Roy-Burman P and Nikitin AY. Synergy of p53 and Rb deficiency in a conditional mouse model for metastatic prostate cancer. *Cancer Res* 2006; 66: 7889-98.
- [170] Elgavish A, Wood PA, Pinkert CA, Eltoun IE, Cartee T, Wilbanks J, Mentor-Marcel R, Tian L and Scroggins SE. Transgenic mouse with human mutant p53 expression in the prostate epithelium. *Prostate* 2004; 61: 26-34.
- [171] Vinall RL, Chen JQ, Hubbard NE, Sulaimon SS, Shen MM, Devere White RW and Borowsky AD. Initiation of prostate cancer in mice by Tp53R270H: evidence for an alternative molecular progression. *Dis Model Mech* 2012; 5: 914-20.
- [172] Wang L, Xiong H, Wu F, Zhang Y, Wang J, Zhao L, Guo X, Chang LJ, Zhang Y, You MJ, Koochekpour S, Saleem M, Huang H, Lu J and Deng Y. Hexokinase 2-mediated Warburg effect is required for PTEN- and p53-deficiency-driven prostate cancer growth. *Cell Rep* 2014; 8: 1461-74.
- [173] Phillips SM, Barton CM, Lee SJ, Morton DG, Wallace DM, Lemoine NR and Neoptolemos JP. Loss of the retinoblastoma susceptibility gene (RB1) is a frequent and early event in prostatic tumorigenesis. *Br J Cancer* 1994; 70: 1252-7.
- [174] Ittmann MM and Wiczorek R. Alterations of the retinoblastoma gene in clinically localized, stage B prostate adenocarcinomas. *Hum Pathol* 1996; 27: 28-34.
- [175] Beltran H, Prandi D, Mosquera JM, Benelli M, Puca L, Cyrta J, Marotz C, Giannopoulou E, Chakravarthi BV, Varambally S, Tomlins SA, Nanus DM, Tagawa ST, Van Allen EM, Elemento O, Sboner A, Garraway LA, Rubin MA and Demichelis F. Divergent clonal evolution of castration-resistant neuroendocrine prostate cancer. *Nat Med* 2016; 22: 298-305.
- [176] Maddison LA, Sutherland BW, Barrios RJ and Greenberg NM. Conditional deletion of Rb causes early stage prostate cancer. *Cancer Res* 2004; 64: 6018-25.
- [177] Greenberg NM, DeMayo F, Finegold MJ, Medina D, Tilley WD, Aspinall JO, Cunha GR, Donjacour AA, Matusik RJ and Rosen JM. Prostate cancer in a transgenic mouse. *Proc Natl Acad Sci U S A* 1995; 92: 3439-43.
- [178] Gingrich JR, Barrios RJ, Morton RA, Boyce BF, DeMayo FJ, Finegold MJ, Angelopoulou R, Rosen JM and Greenberg NM. Metastatic prostate cancer in a transgenic mouse. *Cancer Res* 1996; 56: 4096-102.
- [179] Kasper S, Sheppard PC, Yan Y, Pettigrew N, Borowsky AD, Prins GS, Dodd JG, Duckworth ML and Matusik RJ. Development, progression, and androgen-dependence of prostate tumors in probasin-large T antigen transgenic mice: a model for prostate cancer. *Lab Invest* 1998; 78: 319-33.
- [180] Lane DP and Crawford LV. T antigen is bound to a host protein in SV40-transformed cells. *Nature* 1979; 278: 261-3.
- [181] Linzer DI and Levine AJ. Characterization of a 54K dalton cellular SV40 tumor antigen present in SV40-transformed cells and uninfected embryonal carcinoma cells. *Cell* 1979; 17: 43-52.
- [182] DeCaprio JA, Ludlow JW, Figge J, Shew JY, Huang CM, Lee WH, Marsilio E, Paucha E and Livingston DM. SV40 large tumor antigen forms a specific complex with the product of the retinoblastoma susceptibility gene. *Cell* 1988; 54: 275-83.
- [183] Chiaverotti T, Couto SS, Donjacour A, Mao JH, Nagase H, Cardiff RD, Cunha GR and Balmain A. Dissociation of epithelial and neuroendocrine carcinoma lineages in the transgenic adenocarcinoma of mouse prostate model of prostate cancer. *Am J Pathol* 2008; 172: 236-46.
- [184] Kaplan-Lefko PJ, Chen TM, Ittmann MM, Barrios RJ, Ayala GE, Huss WJ, Maddison LA, Foster BA and Greenberg NM. Pathobiology of autochthonous prostate cancer in a pre-clinical transgenic mouse model. *Prostate* 2003; 55: 219-37.
- [185] Zhou Z, Flesken-Nikitin A and Nikitin AY. Prostate cancer associated with p53 and Rb deficiency arises from the stem/progenitor cell-enriched proximal region of prostatic ducts. *Cancer Res* 2007; 67: 5683-90.

Prostate cancer mouse model review

- [186] Ku SY, Rosario S, Wang Y, Mu P, Seshadri M, Goodrich ZW, Goodrich MM, Labbé DP, Gomez EC, Wang J, Long HW, Xu B, Brown M, Loda M, Sawyers CL, Ellis L and Goodrich DW. Rb1 and Trp53 cooperate to suppress prostate cancer lineage plasticity, metastasis, and antiandrogen resistance. *Science* 2017; 355: 78-83.
- [187] Mu P, Zhang Z, Benelli M, Karthaus WR, Hoover E, Chen CC, Wongvipat J, Ku SY, Gao D, Cao Z, Shah N, Adams EJ, Abida W, Watson PA, Prandi D, Huang CH, de Stanchina E, Lowe SW, Ellis L, Beltran H, Rubin MA, Goodrich DW, Demichelis F and Sawyers CL. SOX2 promotes lineage plasticity and antiandrogen resistance in TP53- and RB1-deficient prostate cancer. *Science* 2017; 355: 84-8.
- [188] Xu J and Qiu Y. Role of androgen receptor splice variants in prostate cancer metastasis. *Asian J Urol* 2016; 3: 177-184.
- [189] Takeda DY, Spisák S, Seo JH, Bell C, O'Connor E, Korthauer K, Ribli D, Csabai I, Solymosi N, Szállási Z, Stillman DR, Cejas P, Qiu X, Long HW, Tisza V, Nuzzo PV, Rohanizadegan M, Pomerantz MM, Hahn WC and Freedman ML. A somatically acquired enhancer of the androgen receptor is a noncoding driver in advanced prostate cancer. *Cell* 2018; 174: 422-32, e13.
- [190] Stanbrough M, Leav I, Kwan PW, Bubley GJ and Balk SP. Prostatic intraepithelial neoplasia in mice expressing an androgen receptor transgene in prostate epithelium. *Proc Natl Acad Sci U S A* 2001; 98: 10823-8.
- [191] Zhu C, Luong R, Zhuo M, Johnson DT, McKenney JK, Cunha GR and Sun Z. Conditional expression of the androgen receptor induces oncogenic transformation of the mouse prostate. *J Biol Chem* 2011; 286: 33478-88.
- [192] Han G, Buchanan G, Ittmann M, Harris JM, Yu X, Demayo FJ, Tilley W and Greenberg NM. Mutation of the androgen receptor causes oncogenic transformation of the prostate. *Proc Natl Acad Sci U S A* 2005; 102: 1151-6.
- [193] Han G, Foster BA, Mistry S, Buchanan G, Harris JM, Tilley WD and Greenberg NM. Hormone status selects for spontaneous somatic androgen receptor variants that demonstrate specific ligand and cofactor dependent activities in autochthonous prostate cancer. *J Biol Chem* 2001; 276: 11204-13.
- [194] Liu G, Sprenger C, Sun S, Epilepsia KS, Haugk K, Zhang X, Coleman I, Nelson PS and Plymate S. AR variant ARv567es induces carcinogenesis in a novel transgenic mouse model of prostate cancer. *Neoplasia* 2013; 15: 1009-17.
- [195] Sun F, Chen HG, Li W, Yang X, Wang X, Jiang R, Guo Z, Chen H, Huang J, Borowsky AD and Qiu Y. Androgen receptor splice variant AR3 promotes prostate cancer via modulating expression of autocrine/paracrine factors. *J Biol Chem* 2014; 289: 1529-39.
- [196] Niu Y, Altuwajiri S, Lai KP, Wu CT, Ricke WA, Messing EM, Yao J, Yeh S and Chang C. Androgen receptor is a tumor suppressor and proliferator in prostate cancer. *Proc Natl Acad Sci U S A* 2008; 105: 12182-7.
- [197] Wu CT, Altuwajiri S, Ricke WA, Huang SP, Yeh S, Zhang C, Niu Y, Tsai MY and Chang C. Increased prostate cell proliferation and loss of cell differentiation in mice lacking prostate epithelial androgen receptor. *Proc Natl Acad Sci U S A* 2007; 104: 12679-84.
- [198] Chua CW, Epsi NJ, Leung EY, Xuan S, Lei M, Li Bl, Bergren SK, Hibshoosh H, Mitrofanova A and Shen MM. Differential requirements of androgen receptor in luminal progenitors during prostate regeneration and tumor initiation. *Elife* 2018; 7.
- [199] Zhang B, Kwon OJ, Henry G, Malewska A, Wei X, Zhang L, Brinkley W, Zhang Y, Castro PD, Titus M, Chen R, Sayeeduddin M, Raj GV, Mauck R, Roehrborn C, Creighton CJ, Strand DW, Ittmann MM and Xin L. Non-cell-autonomous regulation of prostate epithelial homeostasis by androgen receptor. *Mol Cell* 2016; 63: 976-89.
- [200] Lee SO, Tian J, Huang CK, Ma Z, Lai KP, Hsiao H, Jiang M, Yeh S and Chang C. Suppressor role of androgen receptor in proliferation of prostate basal epithelial and progenitor cells. *J Endocrinol* 2012; 213: 173-82.
- [201] Yu S, Zhang C, Lin CC, Niu Y, Lai KP, Chang HC, Yeh SD, Chang C and Yeh S. Altered prostate epithelial development and IGF-1 signal in mice lacking the androgen receptor in stromal smooth muscle cells. *Prostate* 2011; 71: 517-24.
- [202] Welsh M, Moffat L, McNeilly A, Brownstein D, Saunders PT, Sharpe RM and Smith LB. Smooth muscle cell-specific knockout of androgen receptor: a new model for prostatic disease. *Endocrinology* 2011; 152: 3541-51.
- [203] Niu Y, Wang J, Shang Z, Huang SP, Shyr CR, Yeh S and Chang C. Increased CK5/CK8-positive intermediate cells with stromal smooth muscle cell atrophy in the mice lacking prostate epithelial androgen receptor. *PLoS One* 2011; 6: e20202.
- [204] Xie Q, Liu Y, Cai T, Horton C, Stefanson J and Wang ZA. Dissecting cell-type-specific roles of androgen receptor in prostate homeostasis and regeneration through lineage tracing. *Nat Commun* 2017; 8: 14284.
- [205] Lai KP, Yamashita S, Vitkus S, Shyr CR, Yeh S and Chang C. Suppressed prostate epithelial development with impaired branching morphogenesis in mice lacking stromal fibromuscular androgen receptor. *Mol Endocrinol* 2012; 26: 52-66.
- [206] Wilson JD. The pathogenesis of benign prostatic hyperplasia. *Am J Med* 1980; 68: 745-56.

Prostate cancer mouse model review

- [207] Xin L, Teitell MA, Lawson DA, Kwon A, Mellinghoff IK and Witte ON. Progression of prostate cancer by synergy of AKT with genotropic and nongenotropic actions of the androgen receptor. *Proc Natl Acad Sci U S A* 2006; 103: 7789-94.
- [208] Cai H, Memarzadeh S, Stoyanova T, Beharry Z, Kraft AS and Witte ON. Collaboration of Kras and androgen receptor signaling stimulates EZH2 expression and tumor-propagating cells in prostate cancer. *Cancer Res* 2012; 72: 4672-81.
- [209] Zong Y, Xin L, Goldstein AS, Lawson DA, Teitell MA and Witte ON. ETS family transcription factors collaborate with alternative signaling pathways to induce carcinoma from adult murine prostate cells. *Proc Natl Acad Sci U S A* 2009; 106: 12465-70.
- [210] Cai H, Babic I, Wei X, Huang J and Witte ON. Invasive prostate carcinoma driven by c-Src and androgen receptor synergy. *Cancer Res* 2011; 71: 862-72.
- [211] Kim S, Alsaidan OA, Goodwin O, Li Q, Sulejmani E, Han Z, Bai A, Albers T, Beharry Z, Zheng YG, Norris JS, Szulc ZM, Bielawska A, Lebedyeva I, Pegan SD and Cai H. Blocking myristoylation of Src inhibits its kinase activity and suppresses prostate cancer progression. *Cancer Res* 2017; 77: 6950-62.
- [212] Carver BS, Chapinski C, Wongvipat J, Hieronymus H, Chen Y, Chandralapaty S, Arora VK, Le C, Koutcher J, Scher H, Scardino PT, Rosen N and Sawyers CL. Reciprocal feedback regulation of PI3K and androgen receptor signaling in PTEN-deficient prostate cancer. *Cancer Cell* 2011; 19: 575-86.
- [213] Mulholland DJ, Tran LM, Li Y, Cai H, Morim A, Wang S, Plaisier S, Garraway IP, Huang J, Graeber TG and Wu H. Cell autonomous role of PTEN in regulating castration-resistant prostate cancer growth. *Cancer Cell* 2011; 19: 792-804.
- [214] Lee SH, Johnson D, Luong R and Sun Z. Cross-talking between androgen and PI3K/AKT signaling pathways in prostate cancer cells. *J Biol Chem* 2015; 290: 2759-68.
- [215] Lee SH, Luong R, Johnson DT, Cunha GR, Rivina L, Gonzalgo ML and Sun Z. Androgen signaling is a confounding factor for beta-catenin-mediated prostate tumorigenesis. *Oncogene* 2016; 35: 702-14.
- [216] Svensson RU, Haverkamp JM, Thedens DR, Cohen MB, Ratliff TL and Henry MD. Slow disease progression in a C57BL/6 pten-deficient mouse model of prostate cancer. *Am J Pathol* 2011; 179: 502-12.