

UC Irvine

UC Irvine Previously Published Works

Title

One-Year Mortality after Contemporary Laparoscopic Bariatric Surgery: An Analysis of the Bariatric Outcomes Longitudinal Database

Permalink

<https://escholarship.org/uc/item/06d0m7mk>

Journal

Journal of the American College of Surgeons, 226(6)

ISSN

1072-7515

Authors

Inaba, Colette S
Koh, Christina Y
Sujatha-Bhaskar, Sarath
[et al.](#)

Publication Date

2018-06-01

DOI

10.1016/j.jamcollsurg.2018.02.013

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



HHS Public Access

Author manuscript

J Am Coll Surg. Author manuscript; available in PMC 2019 June 01.

Published in final edited form as:

J Am Coll Surg. 2018 June ; 226(6): 1166–1174. doi:10.1016/j.jamcollsurg.2018.02.013.

Low One-Year Mortality After Contemporary Laparoscopic Bariatric Surgery: An Analysis of the Bariatric Outcomes Longitudinal Database

Colette S. Inaba, MD¹, Christina Y. Koh, MD¹, Sarath Sujatha-Bhaskar, MD¹, Jack P. Silva, BS¹, Yanjun Chen, MS², Danh V. Nguyen, PhD^{2,3}, and Ninh T. Nguyen, MD, FACS¹

¹Department of Surgery, University of California Irvine Medical Center

²Institute for Clinical and Translational Science, University of California Irvine

³Department of Medicine, University of California Irvine

BRIEF TITLE

One-Year Mortality After Bariatric Surgery

PRECIS

The 1-year mortality rate after contemporary laparoscopic Roux-en-Y gastric bypass and laparoscopic sleeve gastrectomy is much lower than previously reported. Older age, higher body mass index, and the presence of 30-day leak and pulmonary embolus increase the odds of 1-year mortality.

INTRODUCTION

The risk of death after bariatric surgery has been decreasing over the past two decades since the introduction of laparoscopic bariatric surgery. Early (30-day) mortality after bariatric surgery is now a relatively rare event with contemporary rates typically <0.2%. (1–4) In a study examining the mortality rate in patients who underwent bariatric surgery over an 8-year period, Nguyen and colleagues (5) reported that the in-hospital mortality rate for bariatric operations decreased from 0.4% in 2002 to 0.06% in 2009. Factors leading to reduction of mortality over time include transitioning from open to laparoscopic bariatric surgery (5, 6), initiation of laparoscopic bariatric surgery training by national societies (7), implementation of a national bariatric surgery accreditation system (8, 9), and improvement in surgeons' skills after overcoming the learning curve for bariatric surgery (10).

Corresponding Author: Ninh T. Nguyen, MD, FACS, Department of Surgery, University of California, Irvine Medical Center, 333 City Blvd West, Suite 1600, Orange, CA 92868, ninhn@uci.edu, Tel: (714) 456-8598, Fax: (714) 456-6027.

DISCLOSURES

The American Society for Metabolic and Bariatric Surgery, the surgeons and the hospitals participating in the Centers of Excellence program from 2007 to 2012 are the source of the data used herein; they have not verified and are not responsible for the statistical validity of the data analysis or the conclusions derived by the authors.

MEETING PRESENTATION

This research is scheduled to be presented as a "Top Paper" at the Annual Meeting of the Southern California Chapter of the American College of Surgeons in Santa Barbara, CA, January 2018. (Note—this meeting has been postponed)

Most studies examining the risk of mortality after bariatric surgery have been limited to either in-hospital or 30-day data, which may underrepresent the true risk of bariatric surgery. (11–14) In a study examining early mortality among Medicare beneficiaries who underwent open bariatric surgery between 1997–2002, Flum and colleagues (11) found that the 30-day and 1-year mortality were vastly different at 2.0% and 4.6%, respectively. In a study examining mortality after laparoscopic and open bariatric operations performed in Veterans Affairs Medical Centers, Arterburn and colleagues (12) similarly reported a high 30-day and 1-year mortality of 1.3% and 3.4%, respectively. These studies suggest that about 60% of deaths within the first year following bariatric surgery occur after 30 days, meaning the true risk of mortality after bariatric surgery may be beyond the 30-day mark and better evaluated out to one year after surgery. Additionally, the high 1-year mortality rates reported from these studies may represent outcomes from the era of open bariatric surgery. However, few studies have examined 1-year mortality after contemporary laparoscopic bariatric surgery within the past decade. (15, 16)

The Bariatric Outcomes Longitudinal Database (BOLD) was initiated in 2007 by the American Society of Metabolic and Bariatric Surgery (ASMBS) with a unique aspect of reporting of 1-year outcomes, providing the opportunity to examine 1-year mortality after contemporary laparoscopic bariatric surgery. The aim of this study was to use the BOLD to examine 30-day and 1-year mortality rates, as well as risk factors for 1-year mortality after contemporary laparoscopic bariatric surgery.

METHODS

Data Source

The BOLD is managed by the ASMBS and contains HIPAA-compliant, patient-level data prospectively collected from over 700 centers participating in the ASMBS Bariatric Surgery Centers of Excellence program. The ASMBS and the American College of Surgeons Bariatric Surgery Center Network merged their respective bariatric accreditation programs in 2012 to form the Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program (MBSAQIP). At that time, the BOLD was discontinued and a more refined MBSAQIP database was started. However, the BOLD database was selected for use in this study because it contains data from accredited bariatric centers with 1-year outcomes after index bariatric operations, whereas MBSAQIP data are limited to 30-day outcomes.

This study was exempt from Institutional Review Board approval, as the BOLD contains only de-identified patient data and any specific patient intervention or interaction was not possible.

Data Analysis

All patients who underwent laparoscopic Roux-en-Y gastric bypass (LRYGB) or laparoscopic sleeve gastrectomy (LSG) between 2008–2012 were examined. To control for the effect of the type of procedure on outcomes, data were analyzed separately according to whether LRYGB or LSG was performed. Open cases were excluded. The primary outcome of interest was the mortality rate within one year after a patient's index bariatric operation.

Statistical Analysis

Statistical analysis was performed using SAS version 9.4 (SAS Institute, Cary, NC). Multivariate logistic regression was used to determine the adjusted odds ratio (AOR) for the association between 1-year mortality and patients' demographics; comorbidities; and the presence of major complications including 30-day leak, 30-day pulmonary embolism (PE), and 30-day hemorrhage. The statistical analysis was limited to these three complications as they were the only variables available for specific complications. Other complications were grouped as a composite endpoint that included mortality, and hence could not be used in our model to determine risk factors for mortality. A parsimonious model was then obtained using backward selection of variables with $P < 0.1$ from the multivariate logistic regression model. Results were reported as AOR with 95% confidence intervals (CI). Patients with missing preoperative condition variables were excluded from the regression analysis. Two-sided tests with $P < 0.05$ were considered statistically significant.

RESULTS

Data on 158,606 laparoscopic bariatric operations were examined, including 128,349 (80.9%) LRYGB and 30,257 (19.1%) LSG procedures. Patient demographics and comorbidities are summarized in Table 1. For patients undergoing LRYGB, mean age was 45.5 (standard deviation, SD, 11.7) years old; race was predominately Caucasian (71.8%); mean body mass index (BMI) was 47.5 (SD 8.1) kg/m²; and the most common preoperative comorbidities included hypertension (54.8%), obstructive sleep apnea (43.4%), depression (36.4%), and diabetes (32.9%). For patients undergoing LSG, mean age was 44.6 (SD 11.3) years old; race was predominately Caucasian (68.3%); mean BMI was 46.4 (SD 8.6) kg/m²; and the most common preoperative comorbidities included hypertension (50.4%), obstructive sleep apnea (39.7%), depression (31.7%), and diabetes (27.1%).

Outcomes after LRYGB and LSG are summarized in Table 2. For LRYGB, the 30-day mortality rate was 0.13%, and the 1-year mortality rate was 0.23%, representing a 76.9% relative increase. Among the 294 cases of 1-year mortality, 124 (42.2%) occurred between 30 days and 1 year after surgery. Postoperative rates of 30-day leak, PE, and hemorrhage were 0.45%, 0.16%, and 1.54%, respectively. For LSG, the 30-day mortality rate was 0.06%, and the 1-year mortality rate was 0.11%, representing an 83.3% relative increase. Among the 32 cases of 1-year mortality, 17 (53.1%) occurred between 30 days and 1 year after surgery. Postoperative rates of 30-day leak, PE, and hemorrhage were 0.26%, 0.11%, and 0.59%, respectively.

Table 3 lists patients' demographics, comorbidities, and the presence of 30-day complications, along with their associated odds of 1-year mortality after laparoscopic bariatric surgery after adjustment for all covariates. A parsimonious model was then obtained using backward selection of variables with $P < 0.1$ from the multivariate logistic regression model, with results as listed in Table 4. For LRYGB, significant risk factors independently associated with higher odds of 1-year mortality included older age (AOR 1.05 per year; $P < 0.001$); male gender (AOR 1.88; $P < 0.001$); higher BMI (AOR 1.04 per unit; $P < 0.001$); history of venous thromboembolism (VTE; AOR 2.04; $P = 0.006$); any functional impairment (i.e., bedridden or needing an assistance device to mobilize; AOR 1.76; $P =$

0.003); and the presence of 30-day leak (AOR 25.4; $P < 0.001$), 30-day PE (AOR 34.5; $P < 0.001$), and 30-day hemorrhage (AOR 2.34; $P = 0.001$). For LSG, significant risk factors independently associated with higher odds of 1-year mortality included older age (AOR 1.08 per year; $P < 0.001$); higher BMI (AOR 1.05 per unit; $P = 0.009$); and the presence of 30-day leak (AOR 35.8; $P < 0.001$) or 30-day PE (AOR 252.0; $P < 0.001$). For both LRYGB and LSG, the effect of substance abuse on death could not be determined, as none of the patients with a history of substance abuse died in this series.

DISCUSSION

In this study examining the outcomes of contemporary laparoscopic bariatric surgery performed at Bariatric Surgery Centers of Excellence, we found an overall low risk of 30-day mortality (0.13% for LRYGB and 0.06% for LSG) that increased by 1.8-fold by one year (0.23% for LRYGB and 0.11% for LSG). Factors associated with higher 1-year mortality included older age; higher BMI; and the presence of 30-day leak and PE for both LRYGB and LSG. For LRYGB specifically, other factors associated with higher 1-year mortality included male gender, any functional impairment, history of VTE, and the presence of 30-day hemorrhage.

The most important finding from the current study is the low contemporary 1-year mortality rate after laparoscopic bariatric surgery, which is much lower than previously reported. Prior studies examining open and laparoscopic bariatric surgery have reported 1-year mortality rates ranging between 0.38%–4.6%. (11, 12, 15, 16) In a study of bariatric surgeries recorded in Swedish national registries between 1980–2010, Tao and colleagues (15) reported a low 1-year mortality of 0.38%. In a study examining outcome of open and laparoscopic gastric bypass in the state of California between 1995 and 2009, Weiss and colleagues (16) reported a 1-year mortality rate of 2.2%. Flum and colleagues (11) reported a 1-year mortality rate as high as 4.6% in their study on over 16,000 patients who underwent bariatric surgery from 1997–2002. The low 1-year mortality of 0.11–0.23% in the current study is likely multifactorial. One contributing factor may be the transition from open to laparoscopic bariatric surgery, which has been shown to be associated with lower mortality. (5, 6) Additionally, the implementation of training programs by national societies to minimize the learning curve of laparoscopic bariatric surgery may also account for the decreased mortality over time. (7, 10) Furthermore, the development of the bariatric surgery accreditation system may have also contributed to the reduction in mortality rates, since accredited centers in bariatric surgery have been shown to have improved outcomes compared to nonaccredited centers. (8, 9)

Our study also confirmed that limiting analysis to 30-day mortality significantly underestimates the mortality rate after laparoscopic bariatric surgery. For LRYGB, 42.2% of the mortality events within the first year of surgery occurred after 30 days. Similarly, for LSG, 53.1% of the mortality events within the first year occurred after 30 days. This is consistent with the finding that limiting analysis to 30 days after bariatric surgery excludes about 60% of mortality events that occur within the first year after surgery. (11–13) Furthermore, in a study examining mortality after bariatric surgery using a Pennsylvania statewide database, Omalu and colleagues (13) found that deaths due to traumatic causes

(i.e., motor vehicle accidents, homicides, falls, suicides, drug overdoses) comprised only 10% of the mortality events recorded over a nine-year period. This suggests that the majority of deaths occurring after 30 days are due to complications of surgery or underlying comorbidities, and that evaluating 1-year mortality may provide a more accurate picture of the true risk of bariatric surgery. The higher 1-year mortality rates also suggest that patients would benefit from comprehensive, long-term follow-up, such as is required by the MBSAQIP for accreditation of bariatric centers. This calls into question the 2013 decision by Medicare to reverse the National Coverage Determination that required bariatric centers to be accredited to serve Medicare beneficiaries. Without accreditation, there is no formal, mandatory review process to ensure that Medicare patients receive long-term follow-up after bariatric surgery, which may place them at undue risk of long-term complications.

Factors associated with increased odds of 1-year mortality after LRYGB in this study include older age, male gender, and higher BMI. These findings are consistent with results from previously published studies on mortality after laparoscopic bariatric surgery (Table 5). Additionally, our study identified that a history of VTE and impaired functional status are both independently associated with higher 1-year mortality. We also found that the presence of 30-day leak, PE, and hemorrhage events were associated with increased odds of 1-year mortality after LRYGB. In a study examining the effect of short-term complications on early mortality in 580 LRYGB cases, Fernandez and colleagues (17) similarly found that the presence of postoperative leak and PE were associated with higher early mortality. The effect of postoperative hemorrhage on the risk of mortality was not assessed in that study.

Few studies have examined risk factors for mortality after LSG. In an American College of Surgeons National Surgical Quality Improvement Program (NSQIP) study, 5,871 LSG cases were used to develop a risk calculator for serious adverse events that was then validated in a series of 3,130 cases from the same database. (1) The authors found that 30-day serious adverse events were correlated with male gender (OR 1.68; 95% CI 1.03–2.72) and increased BMI (OR 1.03; 95% CI 1.01–1.05), along with various other preoperative comorbidities. In our study, which has a similar population of patients undergoing LSG, we likewise found an association between higher BMI and increased odds of 1-year mortality (AOR 1.05 per unit; $P=0.013$). We also found that the presence of 30-day complications including leak and PE were associated with an increase in odds of 1-year mortality.

A final point to note is the discrepancy between the large number of LRYGB cases compared to LSG cases performed in the current study. One possible reason that over four times as many LRYGB cases were performed is that the learning curve for LRYGB was reaching a plateau during the study period, whereas the curve for LSG was still developing. In a study using the Nationwide Inpatient Sample from 2008–2012, a time period similar to the one used in the current study, Khan et al (18) found that the percentage of LRYGB cases increased from 58.2% of total bariatric cases in 2008 to a peak of 63.2% in 2009, and then decreased to 49.4% by 2012. In contrast, the authors showed that the percentage of LSG cases increased from 8.2% in 2011 to 39.6% in 2012. Similarly, the percentage of LRYGB cases in the current study decreased from 95.3% in 2008 to 62.6% in 2012, while the percentage of LSG cases increased from 4.7% in 2008 to 37.4% in 2012. This suggests a

transition from LRYGB toward LSG over the study period, a trend that has been demonstrated in a number of other recent studies. (19–21)

There are several limitations to our study. First, this was a retrospective study based on a national database and was thus subject to possible selection bias, inaccurate data input, coding errors, and missing data. Secondly, we are unable to determine the actual cause of death and whether it was related to the index bariatric operation. Despite these limitations, this study included a large sample size from a national cohort of patients. In addition, analysis in this study was stratified by type of procedure, either LRYBG or LSG, allowing us to control for both the effect of operative approach (i.e., no open cases were analyzed that could influence our results) and the effect of procedure type. In addition, our study is one of few that have analyzed risk factors for mortality out to one year after laparoscopic bariatric surgery, allowing us to include a significant percentage of patients who are not evaluated in analyses limited to short-term outcomes.

CONCLUSION

The risk of 1-year mortality after contemporary laparoscopic bariatric surgery is much lower than previously reported, likely due to overcoming the learning curve of laparoscopic surgery, the development of specialized bariatric training programs, and the establishment of an accreditation program for bariatric surgery. It is important to continually refine surgical techniques and perioperative management in order to minimize leaks and pulmonary emboli after laparoscopic bariatric surgery, as these complications increase the risk for mortality. In addition, limiting the analysis of mortality to a 30-day endpoint may underrepresent the true risk of mortality after laparoscopic bariatric surgery. Future studies should consider using 1-year mortality as a more meaningful and accurate endpoint for analysis of outcomes after laparoscopic bariatric surgery.

FUNDING

This statistical analysis for this study was partially supported by grant UL1 TR001414 from the National Center for Advancing Translational Sciences, National Institutes of Health (NIH), through the Biostatistics, Epidemiology and Research Design Unit at the University of California Irvine. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. The NIH had no role in the study design; the collection, analysis, or interpretation of data; the writing of the report; or the decision to submit the report for publication.

ABBREVIATIONS

AOR	adjusted odds ratio
ASMBS	American Society of Metabolic and Bariatric Surgery
BMI	body mass index
BOLD	Bariatric Outcomes Longitudinal Database
LRYGB	laparoscopic Roux-en-Y gastric bypass
LSG	laparoscopic sleeve gastrectomy

MBSAQIP	Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program
NSQIP	National Surgical Quality Improvement Program
PE	pulmonary embolism
VTE	venous thromboembolism

REFERENCES

1. Aminian A, Brethauer SA, Sharafkhan M, Schauer PR. Development of a sleeve gastrectomy risk calculator. *Surg Obes Relat Dis* 2015;11:758–64. [PubMed: 26117166]
2. Young MT, Gebhart A, Phelan MJ, Nguyen NT. Use and Outcomes of Laparoscopic Sleeve Gastrectomy vs Laparoscopic Gastric Bypass: Analysis of the American College of Surgeons NSQIP. *J Am Coll Surg* 2015;220:880–5. [PubMed: 25907869]
3. Benotti P, Wood GC, Winegar DA, et al. Risk factors associated with mortality after Roux-en-Y gastric bypass surgery. *Ann Surg* 2014;259:123–30. [PubMed: 23470583]
4. Lazzati A, Audureau E, Hemery F, et al. Reduction in early mortality outcomes after bariatric surgery in France between 2007 and 2012: A nationwide study of 133,000 obese patients. *Surgery* 2016;159:467–74. [PubMed: 26376758]
5. Nguyen NT, Nguyen B, Smith B, et al. Proposal for a bariatric mortality risk classification system for patients undergoing bariatric surgery. *Surg Obes Relat Dis* 2013;9:239–46. [PubMed: 22336492]
6. Rausa E, Bonavina L, Asti E, et al. Rate of Death and Complications in Laparoscopic and Open Roux-en-Y Gastric Bypass. A Meta-analysis and Meta-regression Analysis on 69,494 Patients. *Obes Surg* 2016;26:1956–63. [PubMed: 27189352]
7. Ali MR, Tichansky DS, Kothari SN, et al. Validation that a 1-year fellowship in minimally invasive and bariatric surgery can eliminate the learning curve for laparoscopic gastric bypass. *Surg Endosc* 2010;24:138–44. [PubMed: 19517173]
8. Nguyen NT, Nguyen B, Nguyen VQ, et al. Outcomes of bariatric surgery performed at accredited vs nonaccredited centers. *J Am Coll Surg* 2012;215:467–74. [PubMed: 22727608]
9. Young MT, Jafari MD, Gebhart A, et al. A decade analysis of trends and outcomes of bariatric surgery in Medicare beneficiaries. *J Am Coll Surg* 2014;219:480–8. [PubMed: 25067804]
10. Doumouras AG, Saleh F, Anvari S, et al. Mastery in Bariatric Surgery: The Long-term Surgeon Learning Curve of Roux-en-Y Gastric Bypass. *Ann Surg* 2018; 267(3):489–94. [PubMed: 28230663]
11. Flum DR, Salem L, Elrod JA, et al. Early mortality among Medicare beneficiaries undergoing bariatric surgical procedures. *Jama* 2005;294:1903–8. [PubMed: 16234496]
12. Arterburn D, Livingston EH, Schiffner T, et al. Predictors of long-term mortality after bariatric surgery performed in Veterans Affairs medical centers. *Arch Surg* 2009;144:914–20. [PubMed: 19841358]
13. Omalu BI, Ives DG, Buhari AM, et al. Death rates and causes of death after bariatric surgery for Pennsylvania residents, 1995 to 2004. *Arch Surg* 2007;142:923–8. [PubMed: 17938303]
14. Debs T, Petruccianni N, Iannelli A, et al. Postoperative Mortality After Bariatric Surgery: Do the Numbers Reflect the Reality? *Obes Surg* 2016;26:1944–5. [PubMed: 27324134]
15. Tao W, Plecka-Ostlund M, Lu Y, et al. Causes and risk factors for mortality within 1 year after obesity surgery in a population-based cohort study. *Surg Obes Relat Dis* 2015;11:399–405. [PubMed: 25604834]
16. Weiss AC, Parina R, Horgan S, et al. Quality and safety in obesity surgery-15 years of Roux-en-Y gastric bypass outcomes from a longitudinal database. *Surg Obes Relat Dis* 2016;12:33–40. [PubMed: 26164113]
17. Fernandez AZ, Jr., Demaria EJ, Tichansky DS, et al. Multivariate analysis of risk factors for death following gastric bypass for treatment of morbid obesity. *Ann Surg* 2004;239:698–702. [PubMed: 15082974]

18. Khan S, Rock K, Baskara A, et al. Trends in bariatric surgery from 2008 to 2012. *Am J Surg* 2016;211:1041–6. [PubMed: 26762828]
19. Koh CY, Inaba CS, Sujatha-Bhaskar S, et al. Laparoscopic Adjustable Gastric Band Explantation and Implantation at Academic Centers. *J Am Coll Surg* 2017;225:532–7. [PubMed: 28754410]
20. English WJ, Demaria EJ, Brethauer SA, et al. American Society for Metabolic and Bariatric Surgery estimation of metabolic and bariatric procedures performed in the United States in 2016. *Surg Obes Relat Dis* 2017. doi: 10.1016/j.soard.2017.12.013. [Epub ahead of print].
21. Kizy S, Jahansouz C, Downey MC, et al. National Trends in Bariatric Surgery 2012–2015: Demographics, Procedure Selection, Readmissions, and Cost. *Obes Surg* 2017;27:2933–9. [PubMed: 28534189]
22. Morton JM, Winegar D, Blackstone R, Wolfe B. Is ambulatory laparoscopic Roux-en-Y gastric bypass associated with higher adverse events? *Ann Surg* 2014;259:286–92. [PubMed: 24169190]
23. Morino M, Toppino M, Forestieri P, et al. Mortality after bariatric surgery: analysis of 13,871 morbidly obese patients from a national registry. *Ann Surg* 2007;246:1002–7. [PubMed: 18043102]
24. Nguyen NT, Masoomi H, Laugenour K, et al. Predictive factors of mortality in bariatric surgery: data from the Nationwide Inpatient Sample. *Surgery* 2011;150:347–51. [PubMed: 21801970]
25. Dorman RB, Abraham AA, Al-Refaie WB, et al. Bariatric Surgery Outcomes in the Elderly: An ACS NSQIP Study. *Journal of Gastrointestinal Surgery* 2012;16:35–44. [PubMed: 22038414]
26. Ramanan B, Gupta PK, Gupta H, et al. Development and validation of a bariatric surgery mortality risk calculator. *J Am Coll Surg* 2012;214:892–900. [PubMed: 22521443]
27. Khan MA, Grinberg R, Johnson S, et al. Perioperative risk factors for 30-day mortality after bariatric surgery: is functional status important? *Surg Endosc* 2013;27:1772–7. [PubMed: 23299129]
28. Nguyen NT, Nguyen B, Smith B, et al. Proposal for a bariatric mortality risk classification system for patients undergoing bariatric surgery. *Surgery for Obesity and Related Diseases* 2013;9:239–46. [PubMed: 22336492]
29. Telem DA, Talamini M, Shroyer AL, et al. Long-term mortality rates (>8-year) improve as compared to the general and obese population following bariatric surgery. *Surg Endosc* 2015;29:529–36. [PubMed: 25037725]

Table 1.

Patient Demographics and Characteristics According to Type of Procedure

Characteristic	LRYGB (n = 128,349)	LSG (n = 30,257)
Age, y, mean (SD)	45.5 (11.7)	44.6(11.3)
Age, y, median (IQR)	45.0 (37–54)	44.0 (36–53)
Female sex, n (%)	57,944 (77.5)	13,757 (75.2)
Race, n (%)		
Caucasian	92,111 (71.8)	20,675 (68.3)
Hispanic	10,879 (8.48)	2,959 (9.78)
African American	14,841 (11.6)	3,748 (12.4)
Other	10,518 (8.19)	2,875 (9.50)
BMI, m/kg ² , mean (SD)	47.5 (8.10)	46.4 (8.60)
BMI, m/kg ² , median (IQR)	46.0 (41.7–51.7)	44.7 (40.4–50.7)
ASA Class, n (%)		
I	3,512 (2.74)	996 (3.29)
II	27,989 (21.8)	8,828 (29.2)
III	91,150 (71.0)	19,548 (64.6)
IV	5,698 (4.44)	885 (2.92)
Comorbidities		
Congestive heart disease	2,560 (2.01)	440 (1.47)
Depression	46,452 (36.4)	9,516 (31.7)
Diabetes	41,958 (32.9)	8,134 (27.1)
Hypertension	69,831 (54.8)	15,122 (50.4)
Ischemic heart disease	3,511 (2.75)	622 (2.07)
Liver disease	9,020 (7.07)	1,654 (5.51)
Sleep apnea	55,378 (43.4)	11,916 (39.7)
Peripheral vascular disease	1,432 (1.12)	278 (0.93)
Venous thromboembolism	2,346 (1.83)	435 (1.44)
Any functional impairment	4,351 (3.41)	672 (2.24)
Alcohol use	17,597 (13.8)	5,469 (18.2)
Tobacco use	6,196 (4.86)	1,494 (4.98)
Substance abuse	383 (0.30)	75 (0.25)

Values are presented as frequencies (%) unless otherwise indicated.

ASA, American Society of Anesthesiologists; IQR, interquartile range; LRYGB, laparoscopic Roux-en-Y gastric bypass; LSG, laparoscopic sleeve gastrectomy.

Table 2.

Operative Outcomes According to Type of Procedure

Outcome	LRYGB (n = 128,349)		LSG (n = 30,257)	
	n	%	n	%
30-d Leak	575	(0.45)	80	(0.26)
30-d Pulmonary embolus	208	(0.16)	33	(0.11)
30-d Hemorrhage	1,979	(1.54)	179	(0.59)
30-d Mortality	170	(0.13)	17	(0.06)
1-y Mortality	294	(0.23)	32	(0.11)

Values are presented as frequencies (%) unless otherwise indicated.

LRYGB, laparoscopic Roux-en-Y gastric bypass; LSG, laparoscopic sleeve gastrectomy

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

Factors Predictive of 1-Year Mortality after Laparoscopic Bariatric Surgery According to Type of Procedure, Multivariate Logistic Regression Model

Variable	LRYGB			LSG		
	AOR	95% CI	p value	AOR	95% CI	p value
Age, per 1-year increase	1.05	(1.04, 1.07)	<0.001	1.08	(1.04, 1.13)	<0.001
Male gender (ref: Female)	1.93	(1.50, 2.48)	<0.001	1.39	(0.63, 3.06)	0.416
Race (ref: Caucasian)						
Hispanic	1.16	(0.71, 1.88)	0.554	0.67	(0.09, 5.21)	0.703
African American	1.28	(0.89, 1.86)	0.184	2.08	(0.74, 5.82)	0.163
Other	1.22	(0.79, 1.86)	0.368	0.89	(0.23, 3.46)	0.865
BMI, per 1-unit increase	1.04	(1.03, 1.05)	<0.001	1.05	(1.01, 1.08)	0.013
ASA Class (ref: Class I)						
II	0.69	(0.29, 1.66)	0.409	0.24	(0.04, 1.36)	0.106
III	0.81	(0.35, 1.84)	0.610	0.21	(0.04, 0.99)	0.048
IV	1.67	(0.70, 4.02)	0.249	2.34	(0.46, 11.7)	0.304
Comorbidities (ref: absence of comorbidity)						
Congestive heart disease	1.25	(0.72, 2.16)	0.431	0.67	(0.08, 5.57)	0.713
Depression	1.09	(0.85, 1.39)	0.523	1.07	(0.47, 2.44)	0.868
Diabetes	0.91	(0.70, 1.18)	0.478	0.68	(0.29, 1.59)	0.379
Hypertension	1.29	(0.98, 1.69)	0.067	1.36	(0.58, 3.18)	0.477
Ischemic heart disease	0.95	(0.56, 1.60)	0.846	2.74	(0.77, 9.75)	0.120
Liver disease	0.82	(0.50, 1.35)	0.438	2.88	(0.95, 8.77)	0.062
Sleep apnea	1.09	(0.85, 1.39)	0.500	1.00	(0.45, 2.18)	0.989
Peripheral vascular disease	1.28	(0.61, 2.67)	0.512	0.88	(0.09, 8.27)	0.913
Venous thromboembolism	2.00	(1.19, 3.34)	0.008	0.88	(0.09, 7.42)	0.844
Any functional impairment	1.71	(1.17, 2.50)	0.005	1.07	(0.28, 4.05)	0.922
Alcohol use	0.81	(0.56, 1.18)	0.276	1.10	(0.40, 3.04)	0.852
Tobacco use	1.47	(0.90, 2.41)	0.126	1.95	(0.44, 8.57)	0.375
Complications, 30d (ref: no complication)						
Leak	24.67	(16.7, 36.6)	<0.001	37.25	(8.50, 163.2)	<0.001
Pulmonary embolus	33.73	(20.5, 55.7)	<0.001	233.5	(60.4, 901.9)	<0.001
Hemorrhage	2.30	(1.37, 3.86)	0.002	6.47	(1.05, 39.7)	0.044

AOR, adjusted odds ratio; ASA, American Society of Anesthesiologists; BMI, body mass index; CI, confidence interval; LRYGB, laparoscopic Roux-en-Y gastric bypass; LSG, laparoscopic sleeve gastrectomy; ref, reference

Table 4.

Factors Predictive of 1-Year Mortality after Laparoscopic Bariatric Surgery According to Type of Procedure, Parsimonious Model*

Variable	LRYGB			LSG		
	AOR	95% CI	p-value	AOR	95% CI	p-value
Age, per 1-year increase	1.05	(1.04, 1.07)	<0.001	1.08	(1.04, 1.13)	<0.001
Male gender (ref: Female)	1.88	(1.48, 2.40)	<0.001	--	--	--
BMI, per 1-unit increase	1.04	(1.03, 1.05)	<0.001	1.05	(1.01, 1.08)	0.009
ASA Class III						
Comorbidities (ref: absence of comorbidity)						
Hypertension	1.29	(1.00, 1.66)	0.050	--	--	--
Liver disease	--	--	--	2.88	(0.98, 8.50)	0.055
Venous thromboembolism	2.04	(1.23, 3.38)	0.006	--	--	--
Functional status--any impairment	1.76	(1.21, 2.56)	0.003	--	--	--
Complications, 30d (ref: no complication)						
Leak	25.4	(17.2, 37.5)	<0.001	35.8	(8.61, 157.3)	<0.001
Pulmonary embolism	34.5	(20.9, 56.8)	<0.001	252	(71.0, 895.2)	<0.001
Hemorrhage	2.34	(1.39, 3.92)	0.001	5.24	(0.88, 31.4)	0.070

* Parsimonious model was obtained using backward selection of variables with $P < 0.1$ from the multivariate logistic regression model

AOR, adjusted odds ratio; ASA, American Society of Anesthesiologists; BMI, body mass index; CI, confidence interval; LRYGB, laparoscopic Roux-en-Y gastric bypass; LSG, laparoscopic sleeve gastrectomy; PE, pulmonary embolism; ref, reference

Table 5.

Summary of Selected Studies Published over the Last Decade Examining Risk Factors for Mortality After Laparoscopic Roux-en-Y Gastric Bypass and/or Laparoscopic Sleeve Gastrectomy

Procedure, article	Database	Groups Studied	Group size n	Mortality rate, % (follow-up time)	Risk factors for mortality
LRYGB or LSG analyzed as separate groups					
Current Study	BOLD 2008–2012	LRYGB LSG	N = 128,349 N = 30,257	0.23% (1 y) 0.11% (1 y)	Both: older age, higher BMI, 30-d leak, 30-day PE, 30-day hemorrhage LRYGB only: male gender, history of VTE, decreased functional status
Morton 2014 (22)	BOLD 2007–2010	LRYGB	N = 51,788	0.1% (30 d)	Age > 50; male gender; BMI 50–70; comorbidities > 5; LOS 0, 0/1, or 4
Aminian 2015 (1)	NSQIP 2012	LSG	N = 5871	0.05% (30 d)	Male sex, higher BMI, CHF, chronic steroid use, DM, preoperative serum total bilirubin level, preoperative hematocrit level
LRYGB or LSG analyzed in combination with other bariatric procedures					
Morino 2007 (23)	SICOB Registry 1996–2006	LRYGB+ ORYGB	N = 1106	0.54% (60 d)	Laparoscopic-to-open conversion, HTN
Arterburn 2009 (12)	VAMC NSQIP 2000–2006	LRYGB + ORYGB + LAGB+ OAGB + LSG + VBG + OSG + other	N = 856	1.3% (30 d) 2.1% (90 d) 3.4% (1 y)	BMI > 50, Diagnostic Cost Group score > 2, open approach
Nguyen 2011 (24)	NIS 2006–2008	LRYGB + ORYGB + LAGB + VBG	N = 304,515	0.12% (inpatient)	Age > 50, male gender, CHF, CRF, PVD, open vs. laparoscopic approach, gastric bypass vs. other bariatric procedures
Dorman 2012 (25)	NSQIP 2005–2009	LRYGB + ORYGB + LAGB + VBG + BPD-DS	Laparoscopic: N = 43,361 Open: N = 5,017	0.15% (30 d)	Laparoscopic bariatric procedures only: BMI > 60, male gender, DM, cardiac comorbidities Open bariatric procedures only: BMI > 60
Ramanan 2012 (26)	NSQIP 2006–2008	LRYGB + ORYGB + LAGB + VBG + BPD-DS + other	N = 21,891	0.14% (30 d)	Older age, higher BMI, PVD, dyspnea, previous PCI, chronic steroid use, type of bariatric procedure
Khan 2013 (27)	NSQIP 2007–2009	LRYGB + ORYGB + LAGB + VBG + BPD-DS	N = 44,408	0.14% (30 d)	Age > 45, BMI > 50, male gender, DM, totally dependent functional status, PCI, dyspnea, >10% weight loss in 6 mo, bleeding disorder, open vs. laparoscopic approach
Nguyen 2013 (28)	UHC database 2002–2009	LRYGB + ORYGB + LAGB + VBG	N = 105,287	0.17% (inpatient)	Age > 60, male gender, Medicare payer, DM, open approach, gastric bypass vs. nongastric bypass bariatric procedure
Benotti 2014 (3)	BOLD 2007–2011	LRYGB+ ORYGB	N = 81,751	0.15% (30 d)	Older age, male gender, higher BMI, pulmonary HTN, CHF, liver disease

Procedure, article	Database	Groups Studied	Group size n	Mortality rate, % (follow-up time)	Risk factors for mortality
Tao 2015 (15)	Swedish Patient Registry 1980–2010	LRYGB + ORYGB + LAGB + VBG + BPD-DS + other	N = 22,487	0.38% (1 y)	Male gender, CHF, DM, open approach
Telem 2015 (29)	SPARCS 1999–2005	LRYGB + LAGB + VBG + sleeve gastrectomy	N = 7862	2.5% (8–14 y)	Older age, male gender, Medicare/Medicaid payment source, CHF, HTN, DM, chronic pulmonary disorder, rheumatoid arthritis
Lazzati 2016 (4)	French National Health Service Database 2007–2012	LRYGB + ORYGB + LAGB + OAGB + LSG + OSG	N = 133,804	0.12% (90 d)	Age >50, BMI > 50, male gender, DM, HTN, open approach, procedure type, lower procedure volume
Weiss 2016 (16)	California OSHPD 1995–2009	LRYGB + ORYGB	N = 129,432	2.2% (1 y); 4.4% (5 y); 8.1% (10 y)	Older age, male gender, open approach, DM, smoker, alcohol abuse, higher Charlson Index, insurance status, LOS < 4 or 7

ASA, American Society of Anesthesiologists; BMI, body mass index; BOLD, Bariatric Outcomes Longitudinal Database; CAD, coronary artery disease; CHF, congestive heart failure; CRF, chronic renal failure; DM, diabetes mellitus; HTN, hypertension; LAGB, laparoscopic adjustable gastric banding; LOS, length of stay; LRYGB, laparoscopic Roux-en-Y gastric bypass; LSG, laparoscopic sleeve gastrectomy; NIS, National Inpatient Sample; NSQIP, National Surgical Quality Improvement Program; OAGB, open adjustable gastric banding; ORYGB, open Roux-en-Y gastric bypass; OS-MRS, Obesity Surgery Mortality Risk Score; OSG, open sleeve gastrectomy; OSHPD, Office of Statewide Health Planning and Development; PCI, percutaneous coronary intervention; PE, pulmonary embolus; PVD, peripheral vascular disease; SICOB, Italian Society of Obesity Surgery; SPARCS, New York Statewide Planning and Research Cooperative System; UHC, University HealthSystem Consortium; VAMC, Veterans Affairs Medical Centers; VBG, vertical banded gastroplasty; VTE, venous thromboembolism