

UCSF

UC San Francisco Previously Published Works

Title

Pseudomonas aeruginosa Stimulates Inflammation and Enhances Kaposi's Sarcoma Herpesvirus-Induced Cell Proliferation and Cellular Transformation through both Lipopolysaccharide and Flagellin

Permalink

<https://escholarship.org/uc/item/06d8n7f9>

Journal

mBio, 11(6)

ISSN

2161-2129

Authors

Markazi, Ashley
Bracci, Paige M
McGrath, Michael
et al.

Publication Date

2020-12-22

DOI

10.1128/mbio.02843-20

Peer reviewed



Pseudomonas aeruginosa Stimulates Inflammation and Enhances Kaposi's Sarcoma Herpesvirus-Induced Cell Proliferation and Cellular Transformation through both Lipopolysaccharide and Flagellin

Ashley Markazi,^{a,b} Paige M. Bracci,^c Michael McGrath,^d Shou-Jiang Gao^{a,b}

^aCancer Virology Program, UPMC Hillman Cancer Center, Pittsburgh, Pennsylvania, USA

^bDepartment of Microbiology and Molecular Genetics, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA

^cDepartment of Epidemiology and Biostatistics, University of California at San Francisco, San Francisco, California, USA

^dThe AIDS and Cancer Specimen Resource and Department of Laboratory Medicine, Pathology, and Medicine, University of California at San Francisco, San Francisco, California, USA

ABSTRACT Inflammation triggered by innate immunity promotes carcinogenesis in cancer. Kaposi's sarcoma (KS), a hyperproliferative and inflammatory tumor caused by Kaposi's sarcoma-associated herpesvirus (KSHV) infection, is the most common cancer in AIDS patients. KSHV infection sensitizes cells to pathogen-associated molecular patterns (PAMPs). We examined the role of *Pseudomonas aeruginosa*, an opportunistic bacterium that can affect AIDS patients, in inflammation and cell proliferation of KSHV-transformed cells. *P. aeruginosa* stimulation increased cell proliferation and efficiency of colony formation in soft agar of KSHV-transformed rat primary mesenchymal precursor (KMM) cells but had no significant effect on the untransformed (MM) cells. *P. aeruginosa* stimulation also increased cell proliferation of KSHV-infected human B cells, BJAB, but not the uninfected cells. Mechanistically, *P. aeruginosa* stimulation resulted in increased inflammatory cytokines and activation of p38, ERK1/2, and JNK mitogen-activated protein kinase (MAPK) pathways in KMM cells while having no obvious effect on MM cells. *P. aeruginosa* induction of inflammation and MAPKs was observed with and without inhibition of the Toll-like receptor 4 (TLR4) pathway, while a flagellin-deleted mutant of *P. aeruginosa* required a functional TLR4 pathway to induce inflammation and MAPKs. Furthermore, treatment with either lipopolysaccharide (LPS) or flagellin alone was sufficient to induce inflammatory cytokines, activate MAPKs, and increase cell proliferation and efficiency of colony formation in soft agar of KMM cells. These results demonstrate that both LPS and flagellin are PAMPs that contribute to *P. aeruginosa* induction of inflammation in KSHV-transformed cells. Because AIDS-KS patients are susceptible to *P. aeruginosa* infection, our work highlights the preventive and therapeutic potential of targeting *P. aeruginosa* infection in these patients.

IMPORTANCE Kaposi's sarcoma (KS), caused by infection with Kaposi's sarcoma-associated herpesvirus (KSHV), is one of the most common cancers in AIDS patients. KS is a highly inflammatory tumor, but how KSHV infection induces inflammation remains unclear. We have previously shown that KSHV infection upregulates Toll-like receptor 4 (TLR4), sensitizing cells to lipopolysaccharide (LPS) and *Escherichia coli*. In the current study, we examined the role of *Pseudomonas aeruginosa*, an opportunistic bacterium that can affect AIDS patients, in inflammation and cell proliferation of KSHV-transformed cells. *P. aeruginosa* stimulation increased cell proliferation, inflammatory cytokines, and activation of growth and survival pathways in KSHV-transformed cells through two pathogen-associated molecular patterns, LPS and

Citation Markazi A, Bracci PM, McGrath M, Gao S-J. 2020. *Pseudomonas aeruginosa* stimulates inflammation and enhances Kaposi's sarcoma herpesvirus-induced cell proliferation and cellular transformation through both lipopolysaccharide and flagellin. mBio 11:e02843-20. <https://doi.org/10.1128/mBio.02843-20>.

Editor Blossom Damania, University of North Carolina, Chapel Hill

Copyright © 2020 Markazi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Shou-Jiang Gao, gaos8@upmc.edu.

This article is a direct contribution from Shou-Jiang Gao, a Fellow of the American Academy of Microbiology, who arranged for and secured reviews by Kenneth Kaye, Harvard Medical School /Brigham and Women's Hospital, and Richard Longnecker, Northwestern University Feinberg School of Medicine.

Received 6 October 2020

Accepted 12 October 2020

Published 10 November 2020

flagellin. Because AIDS-KS patients are susceptible to *P. aeruginosa* infection, our work highlights the preventive and therapeutic potential of targeting *P. aeruginosa* infection in these patients.

KEYWORDS Kaposi's sarcoma-associated herpesvirus, KSHV, Kaposi's sarcoma, *Pseudomonas aeruginosa*, lipopolysaccharide, LPS, flagellin, inflammation, MAPK, cellular transformation

Inflammation is the immune system's response to tissue damage or microbial infection. When unregulated, inflammation can cause or exacerbate carcinogenesis (1). For example, the continuous induction of inflammatory cytokines and cytotoxic mediators, such as reactive oxygen or nitrogen species, can, over time, cause damage to the cellular genome, resulting in cellular mutations that lead to dysregulated cell proliferation and cancer development (2). Inflammatory cytokines can further enhance cancer progression by activating pro-oncogenic and -survival cellular pathways, including NF- κ B and STAT3 pathways (3). By further elucidating the mechanisms by which inflammation promotes cancer cell growth, proliferation, and survival, we can gain important insights into the mechanism of oncogenesis and identify novel therapeutic targets and biomarkers.

Interestingly, patients with HIV/AIDS are at a much higher risk for developing AIDS-associated cancer (AAC) than those with healthy immune systems (4). HIV patients are 500 times more likely to develop Kaposi's sarcoma (KS), 12 times more likely to be diagnosed with non-Hodgkin's lymphoma, and, among women, 3 times more likely to be diagnosed with cervical cancer (4). Moreover, HIV patients are at a higher risk for non-AIDS-defining cancer (NADC) as well, including anal cancer, liver cancer, lung cancer, and Hodgkin's lymphoma (5). HIV enhances cellular transformation through its protein, Tat, which represses tumor suppressor gene p53, promotes cell cycle progression, and induces inflammation (6, 7). For cancers caused by infections of oncogenic viruses, HIV regulates both the replication of these viruses and the progression of their associated cancers (8). For example, HIV-encoded products Tat, Nef, and Vpr regulate KSHV replication and the functions of KSHV genes, resulting in enhanced cell migration, invasion, and angiogenesis (9–15). Moreover, long-term use of antiretroviral drugs in HIV patients is associated with an elevated risk of several cancers (16).

A number of other mechanisms also contribute to the chronic inflammation and increased cancer risk in HIV-infected patients. One of these factors is immunodeficiency associated with HIV infection, such as declined CD4⁺ T cell count (17). Because CD4⁺ T cells stimulate B cells and CD8⁺ T cells, patients with a low CD4⁺ T cell count are less capable of eliminating infections, including viruses, bacteria, and fungi, thereby increasing pathogen-associated molecular patterns (PAMPs) (18), which continuously stimulate immune and cancer cells to secrete inflammatory cytokines, resulting in chronic inflammation. Indeed, many reports show that a lower CD4⁺ cell count is correlated with an increase in inflammatory cytokines (19). Moreover, chronic stress, such as metabolic stress of the immune and cancer cells induced by HIV infection, as well as the long-term use of antiretroviral drugs, triggers damage-associated molecular patterns (DAMPs), which further promote chronic inflammation in the same fashion as PAMPs (20).

KS, the most common cancer in HIV-infected patients, is a hyperproliferative and inflammatory cancer caused by infection with Kaposi's sarcoma-associated herpesvirus (KSHV) (21). KSHV infection provides an excellent model for examining the complex interactions of HIV, a cancer-causing virus (KSHV), innate immunity, inflammation, and cancer. KSHV encodes numerous genes that directly contribute to cellular transformation (21), and KSHV infection alone is sufficient to induce inflammatory cytokines, which can stimulate cell proliferation and survival and regulate KSHV replication (22–28). Furthermore, we have shown that KSHV infection sensitizes the infected cells to PAMPs, leading to the activation of Toll-like receptor 4 (TLR4) and alternative complement pathways, which induce inflammatory cytokines and promote cell survival, prolifera-

tion, and cellular transformation (22, 29). The *Escherichia coli*- and lipopolysaccharide (LPS)-activated TLR4 pathway stimulates cell proliferation, cellular transformation, and tumorigenesis by increasing interleukin-6 (IL-6) expression to activate the STAT3 pathway (22). Indeed, in a recent study, we showed the impoverishment of oral microbial diversity and enrichment of specific microbiota in oral KS in HIV-infected patients (30). However, the precise mechanism of how specific microbiota promote KS remains to be elucidated.

Pseudomonas aeruginosa is normally considered a commensal bacterium. However, *P. aeruginosa* can cause severe infection in individuals with immunosuppression (31). HIV/AIDS patients with CD4⁺ T cell counts below 200 cell/mm³ are at a significantly higher risk for *P. aeruginosa* infection (32). *P. aeruginosa* consists of PAMPs, such as LPS and flagellin, which activate TLR4 and TLR5, respectively (33). Hence, *P. aeruginosa* infection might induce inflammatory cytokines of KSHV-infected cells and promote cell proliferation and cellular transformation.

In the current study, we analyzed the effects of *P. aeruginosa* on cell proliferation and cellular transformation in a KS-like model of KSHV-induced cellular transformation of rat primary embryonic metanephric mesenchymal precursor cells (MM) (34). We observed that *P. aeruginosa* stimulation increased both cell proliferation and cellular transformation in KSHV-transformed MM cells (KMM) yet had no significant effect on MM cells. Moreover, we observed similar results of increased cell proliferation in a KSHV-infected human B cell line, KSHV-BJAB, compared to the BJAB uninfected control. In KMM cells, *P. aeruginosa* stimulation resulted in increased expression of inflammatory cytokines and activation of p38, ERK1/2, and JNK pathways. Interestingly, we observed the induction of inflammatory cytokines and activation of the p38 and ERK1/2 pathways, even after the inhibition of the TLR4 pathway in KMM cells stimulated by *P. aeruginosa*, and that this effect disappeared when KMM cells were stimulated with a flagellin-deleted mutant of *P. aeruginosa*, indicating that *P. aeruginosa* mediated inflammation and cellular transformation of KSHV-transformed cells through both LPS and flagellin.

RESULTS

***P. aeruginosa* stimulation enhances cell proliferation and cellular transformation of KMM cells but has no significant effect on MM cells.** To examine the effect of *P. aeruginosa* on the proliferation of KSHV-transformed cells, we treated the cells with 1×10^7 CFU/ml *P. aeruginosa* (ATCC 15442) or 1 μ g/ml LPS. *P. aeruginosa* increased the proliferation of KMM cells but did not have any significant effect on MM cells (Fig. 1A). Similar results were observed with LPS, as expected (22). Both *P. aeruginosa* and LPS also increased the sizes and efficiency of colony formation in soft agar of KMM cells (Fig. 1B and C). As previously reported, MM cells did not form any significant colonies (34). These results indicated that, similar to LPS, *P. aeruginosa* stimulated the proliferation and cellular transformation of KMM cells (22). To assess the effects of *P. aeruginosa* and LPS stimulation on KSHV-infected human B cells, we treated BJAB and KSHV-BJAB cells with 1×10^7 CFU/ml *P. aeruginosa* (ATCC 15442) or 1 μ g/ml LPS. Although less pronounced than that in KMM cells, *P. aeruginosa* stimulation also increased proliferation of KSHV-BJAB cells while having no significant effects in BJAB cells (Fig. 1D). Because KMM cells can form colonies in soft agar, permitting the evaluation of the transforming potential of the cells, we chose to further examine the effect of *P. aeruginosa* on KMM cells and the control MM cells in subsequent experiments (34).

***P. aeruginosa* stimulation increases the expression levels of inflammatory cytokines in KMM cells while having minimal effect on MM cells.** We previously showed that purified *E. coli* LPS induced the inflammatory cytokines interleukin-6 (IL-6), IL-1 β , and IL-18 in KMM cells but had only a weak effect on MM cells. *P. aeruginosa* (ATCC 15442) stimulation resulted in higher mRNA levels of IL-6 and IL-1 β but had no significant effect on IL-18 in KMM cells (Fig. 2A). Additionally, we analyzed the cytokines tumor necrosis factor alpha (TNF- α) and CXCL-1, as *P. aeruginosa* increased levels of these inflammatory cytokines in mice (35, 36). *P. aeruginosa* stimulation also resulted in




FIG 1 *P. aeruginosa* (PA) stimulation enhances cell proliferation and cellular transformation of KSHV-infected cells but has no significant effect on the uninfected cells. (A) Cell proliferation of MM and KMM cells treated with PBS, 1 μ g/ml LPS, or 1×10^7 CFU/ml *P. aeruginosa* (ATCC 15442), analyzed by cell counting. (B and C) Formation of colonies of KMM cells in soft agar treated with PBS, 1 μ g/ml LPS, or 1×10^6 to 1×10^8 CFU/ml *P. aeruginosa* (ATCC 15442), shown by representative pictures (B) and results of statistical analysis from 3 wells, each with 5 representative fields (C). (D) Cell proliferation of BJAB and KSHV-BJAB cells treated with PBS, 1 μ g/ml LPS, or 1×10^7 CFU/ml *P. aeruginosa* (ATCC 15442), analyzed by cell counting. *, $P \leq 0.05$. NS, not significant. PA, *P. aeruginosa*.

higher mRNA levels of TNF- α and CXCL-1 in KMM cells (Fig. 2A). In contrast, cytokines IL-6, IL-1 β , IL-18, TNF- α , and CXCL-1 were not significantly upregulated in MM cells by *P. aeruginosa* (Fig. 2A).

We further examined the induction kinetics of inflammatory cytokines in KMM cells by *P. aeruginosa* by stimulating the cells with 1×10^7 CFU/ml *P. aeruginosa* (ATCC 15442) and analyzing them at 0, 0.5, 1, 2, 4, 6, and 8 h poststimulation. IL-6,




FIG 2 *P. aeruginosa* (PA) stimulation increases the expression of inflammatory cytokines in KMM cells while having minimal effect on MM cells. (A) MM and KMM cells were treated with PBS, 1 μ g/ml LPS, or 1 \times 10⁶ to 1 \times 10⁸ CFU/ml *P. aeruginosa* (ATCC 15442) for 1 h and analyzed for cytokines IL-6, TNF- α , IL-1 β , IL-18, and CXCL-1 by qRT-PCR. (B) MM and KMM cells were treated with PBS or 1 \times 10⁷ CFU/ml *P. aeruginosa* (ATCC 15442) at the specified time points and analyzed for cytokines IL-6, TNF- α , IL-1 β , IL-18, and CXCL-1 by qRT-PCR. *, $P \leq 0.05$.

IL-1 β , TNF- α , and CXCL-1 had the highest mRNA levels at 1 h after *P. aeruginosa* stimulation in KMM cells (Fig. 2B). No significant induction of inflammatory cytokines at different time points in MM cells was observed following *P. aeruginosa* stimulation (Fig. 2B).




FIG 3 *P. aeruginosa* (PA) stimulation activates the p38, JNK, and ERK1/2 MAPK pathways in KMM cells but has no obvious effect on MM cells. MM and KMM cells were stimulated with 1×10^7 CFU/ml *P. aeruginosa* (ATCC 15442) for the specific times and analyzed by Western blotting.

***P. aeruginosa* stimulation activates the MAPK pathways in KMM cells but has no obvious effect in MM cells.** The p38, ERK1/2, and JNK mitogen-activated protein kinase (MAPK) pathways have been implicated in the induction of inflammatory cytokines and are commonly activated in cancer cells (37). We examined the activation of these pathways following stimulation with 1×10^7 CFU/ml *P. aeruginosa* (ATCC 15442) in MM and KMM cells. We detected the activation of p38, ERK1/2, and JNK pathways, which peaked at 15 min after *P. aeruginosa* stimulation in KMM cells (Fig. 3). In contrast, no increased activation of p38, ERK1/2, or JNK was observed in MM cells (Fig. 3).

***P. aeruginosa* stimulation increases the expression of inflammatory cytokines by both TLR4-dependent and -independent mechanisms in KSHV-transformed cells.** We have previously shown that LPS can induce inflammatory cytokines in KMM cells. Since *P. aeruginosa* contains other PAMPs in addition to LPS, to determine if other *P. aeruginosa* PAMPs also contributed to the *P. aeruginosa*-induced inflammation in KMM cells, we stimulated KMM cells with 1×10^7 CFU/ml *P. aeruginosa* (ATCC 15442) in the presence of 10 μ g/ml TLR4 inhibitor CLI095. The levels of induced proinflammatory cytokines IL-6, IL-1 β , TNF- α , and CXCL-1 by *P. aeruginosa* were significantly decreased in KMM cells by CLI095; however, they remained significantly higher than those of the unstimulated KMM cells (Fig. 4). As expected, induction of inflammatory cytokines by LPS was completely blocked by CLI095 in KMM cells, with levels similar to those of the unstimulated cells or cells treated with CLI095 alone (Fig. 4). No obvious change of inflammatory cytokines in MM cells was observed under these treatments. These results indicated that, besides LPS, another *P. aeruginosa* PAMP(s) also contributes to *P. aeruginosa*-induced inflammation in KMM cells.

***P. aeruginosa* flagellin contributes to *P. aeruginosa* induction of inflammatory cytokines in KMM cells but has no significant effect on MM cells.** As flagellin is *P. aeruginosa*'s second most immunogenic PAMP, we stimulated KMM cells with a *P. aeruginosa* strain with the *fliC* gene deleted from its genome, the PAO1 Δ *fliC* strain, which rendered it defective in flagellin expression, and its parallel wild type, *P. aeruginosa* PAO1 (38). While PAO1 and PAO1 Δ *fliC* strains at 1×10^7 CFU/ml induced




FIG 4 *P. aeruginosa* (PA) stimulation increases the expression of inflammatory cytokines by both TLR4-dependent and -independent mechanisms in KSHV-transformed cells. MM and KMM cells were treated with 10 μg/ml TLR4 inhibitor (CLI095) for 1 h and then treated with PBS, 1 μg/ml LPS, or 1 × 10⁷ CFU/ml *P. aeruginosa* (ATCC 15442) for 1 h and analyzed for the expression of cytokines by RT-qPCR. *, P ≤ 0.05.

inflammatory cytokines in KMM cells, the levels of induction were significantly lower in KMM cells stimulated with the PAO1Δ*flhC* than PAO1 strain (Fig. 5A). As expected, 1 × 10⁷ CFU/ml PAO1 induced inflammatory cytokines at levels similar to 1 × 10⁷ CFU/ml *P. aeruginosa* (ATCC 15442) (Fig. 5A). We then treated MM and KMM cells with CLI095 for 1 h before stimulating with the PAO1 or PAO1Δ*flhC* strain or LPS. Similar to *P. aeruginosa* (ATCC 15442), the PAO1 induction of inflammatory cytokines was reduced in KMM cells by a TLR4 inhibitor, CLI095, but remained at significantly higher levels than those in the unstimulated cells (Fig. 5B). In contrast, PAO1Δ*flhC* strain induction of inflammatory cytokines was completely abolished by CLI095 (Fig. 5B). No significant effect was observed on MM cells with either the PAO1Δ*flhC* or PAO1 strain (Fig. 5B).

To further confirm the role of flagellin in *P. aeruginosa* induction of inflammation in KMM cells, we stimulated the cells with 0.3 μg/ml purified *P. aeruginosa* flagellin. Treatment with purified *P. aeruginosa* flagellin alone was sufficient to induce inflammatory cytokines in KMM cells at levels similar to those of LPS alone (Fig. 5C). In contrast, purified *P. aeruginosa* flagellin stimulation had no significant effect on MM cells (Fig. 5C).

Taken together, these results indicated that flagellin contributed to *P. aeruginosa* induction of inflammatory cytokines in KMM cells. Hence, at least LPS and flagellin contributed to *P. aeruginosa* induction of inflammation in KMM cells.

***P. aeruginosa* flagellin activates p38 and ERK1/2 pathways in KMM cells but has no obvious effect on MM cells.** The results described above indicated that the patterns of inflammatory cytokines induced by LPS and flagellin were not entirely the same, which could be due to distinct activation of the signaling pathways by the two PAMPs. While all three MAPK pathways were activated in KMM cells by PAO1, the activation levels were reduced in cells stimulated with the PAO1Δ*flhC* strain (Fig. 6A). Treatment with TLR4 inhibitor CLI095 reduced PAO1 activation of p38 and ERK1/2 pathways but completely abolished that of JNK pathway and PAO1Δ*flhC* strain activation of all three MAPK pathways (Fig. 6A). These results indicated that LPS mediated *P.*




FIG 5 *P. aeruginosa* (PA) flagellin contributes to *P. aeruginosa* induction of inflammatory cytokines in KMM cells but has no significant effect on MM cells. (A) MM and KMM cells were treated with PBS, 1 μ g/ml LPS, 1 $\times 10^7$ CFU/ml *P. aeruginosa* (ATCC 15442), 1 $\times 10^7$ CFU/ml PAO1, or 1 $\times 10^7$ CFU/ml PAO1 Δ flhC strain for 1 h and analyzed for the expression of cytokines by qRT-PCR. (B) MM and KMM cells were treated with 10 μ g/ml TLR4 inhibitor (CLI095) for 1 h and then treated with PBS, 1 $\times 10^7$ CFU/ml wild-type PAO1, or 1 $\times 10^7$ CFU/ml PAO1 Δ flhC strain for 1 h and analyzed for the expression of cytokines by qRT-PCR. (C) MM and KMM cells were treated with PBS, 1 μ g/ml LPS, 1 $\times 10^7$ CFU/ml *P. aeruginosa* (ATCC 15442), or 0.3 μ g/ml *P. aeruginosa* flagellin for 1 h and analyzed for the expression of cytokines by qRT-PCR. *, $P \leq 0.05$. NS, not significant.

aeruginosa activation of all three pathways while flagellin mediated the activation of p38 and ERK1/2 pathways. We then investigated whether flagellin alone was sufficient to activate the p38 and ERK1/2 pathways in KMM cells by treating the cells with purified *P. aeruginosa* flagellin. As expected, flagellin alone indeed activated the p38 and ERK1/2 pathways, and treatment with CLI095 had no effect on flagellin activation of these two pathways, although the reduction of ERK1/2 activation by CLI095 in *P. aeruginosa*-stimulated KMM cells was less pronounced in this experiment, indicating possible variation of *P. aeruginosa* LPS effect on ERK1/2 activation (Fig. 6B). Together, these results indicated that flagellin contributed to *P. aeruginosa* activation of p38 and ERK1/2 pathways, and LPS contributed to *P. aeruginosa* activation of all three MAPK pathways in KMM cells.

***P. aeruginosa* flagellin stimulation increases cell proliferation and cellular transformation of KMM cells but has no significant effect on MM cells.** Because flagellin participated in *P. aeruginosa*-induced inflammation and activation of MAPK




FIG 6 *P. aeruginosa*-flagellin induces p38 and ERK1/2 activation in KMM cells but has no obvious effect on MM cells. (A) MM and KMM cells were treated for 1 h with 10 μg/ml TLR4 inhibitor CLI095 and then treated with PBS, 1 × 10⁷ CFU/ml wild-type PAO1, or 1 × 10⁷ CFU/ml PAO1ΔflgC strain for 15 min and analyzed for the activation of p38, ERK1/2, and JNK pathways by Western blotting. (B) MM and KMM cells were treated for 1 h with 10 μg/ml TLR4 inhibitor CLI095 and then treated for 15 min with PBS, 1 × 10⁷ CFU/ml *P. aeruginosa* (PA), or 0.3 μg/ml *P. aeruginosa* flagellin and then analyzed for the activation of p38 and ERK1/2 pathways by Western blotting.

pathways, we examined whether flagellin could promote KSHV-induced cell proliferation and cellular transformation. KMM cells were treated with purified *P. aeruginosa* flagellin or phosphate-buffered saline (PBS) as a control and analyzed for cell proliferation. Treatment with *P. aeruginosa* flagellin alone was sufficient to induce faster cell proliferation in KMM cells than in the untreated cells (Fig. 7A). In contrast, *P. aeruginosa* flagellin had no effect on the proliferation of MM cells (Fig. 7A).

We further examined the effect of flagellin on cellular transformation of KMM cells. Treatment with purified *P. aeruginosa* flagellin increased the numbers and sizes of colonies of KMM cells compared to those of the untreated cells (Fig. 7B and C). MM cells did not form any observable colonies in soft agar in both flagellin-treated and untreated cells (results not shown).

Simultaneous inhibition of p38 and ERK1/2 pathways decreases flagellin-induced inflammation and cell proliferation of KMM cells but has no obvious effect on MM cells. To determine whether p38 and ERK1/2 pathways mediated flagellin-induced inflammation in KMM cells, we stimulated the cells with *P. aeruginosa* flagellin in the presence of specific inhibitors of these pathways. p38 inhibitor (SB203580) or ERK1/2 inhibitor (U0126) alone had no effect on flagellin induction of inflammatory cytokines (Fig. 8A). However, treatment with both inhibitors significantly decreased the flagellin induction of all inflammatory cytokines (Fig. 8A), confirming the important roles of p38 and ERK1/2 pathways in flagellin induction of inflammatory cytokines. Hence, both p38 and ERK1/2 pathways mediated flagellin induction of inflammatory cytokines.

We further assessed the roles of p38 and ERK1/2 MAPK pathways on flagellin-induced cell proliferation. While *P. aeruginosa* flagellin stimulated the proliferation of KMM cells by 20%, SB203580 or U0126 alone or in combination completely abolished this effect (Fig. 8B). SB203580 or U0126 alone or in combination had no significant effect on the proliferation of MM cells (Fig. 8B). These results indicated that both p38 and ERK1/2 MAPK pathways were essential for flagellin-stimulated proliferation of KMM cells.




FIG 7 *P. aeruginosa* flagellin stimulation increases cell proliferation and cellular transformation of KMM cells but has no significant effect on MM cells. (A) Cell proliferation of MM and KMM cells treated with PBS or 1 $\mu\text{g/ml}$ *P. aeruginosa* flagellin. (B and C) Formation of colonies of KMM cells in soft agar treated with PBS or 1 $\mu\text{g/ml}$ *P. aeruginosa* flagellin, shown by representative pictures (B) and results of statistical analysis from 3 wells, each with 5 representative fields (C). *, $P \leq 0.05$. NS, not significant.

DISCUSSION

It has been reported that the diversity of overall bacterial species is decreased while the number of pathogenic species is increased in the microbiomes of immunosuppressed patients compared to that in healthy individuals (39). The effects of pathogenic bacteria on cancer pathology have been studied extensively in numerous cancers (40). Pathogenic bacteria such as *Fusobacterium nucleatum*, *Helicobacter pylori*, and *Salmonella* spp. can exacerbate cancer pathogenesis by producing bacterial toxin and effector proteins that induce host cell damage and interfere with cell signaling pathways involved in cell proliferation (40). Additionally, bacterial metabolic products, including short-chain fatty acids, can reactivate oncogenic viruses, such as Epstein-Barr virus and KSHV, resulting in dissemination of cancer cells (41). Our study specifically focuses on bacterial PAMPs, which activate TLRs, resulting in pro-survival pathway activation and increased cell proliferation (22).

In the current study, we investigated the role of *P. aeruginosa* in KSHV induction of chronic inflammation and KSHV-induced cell proliferation and cellular transformation. *P. aeruginosa* infection can occur in HIV-infected patients and can cause serious complications in immunocompetent hosts despite being a commensal bacterium (32). The results showed that *P. aeruginosa* stimulation enhanced cell proliferation, and the sizes and efficiency of colony formation in soft agar of KSHV-transformed cells, but had no effect on the uninfected primary cells. Mechanistically, *P. aeruginosa* stimulation increased the expression of inflammatory cytokines and activated multiple MAPK pathways. Importantly, besides LPS, we found that *P. aeruginosa* flagellin also contributed to the induction of inflammatory cytokines and cell proliferation of KSHV-transformed cells, and that LPS and flagellin differentially activated MAPK pathways, but both induced similar profiles of inflammatory cytokines (Fig. 8C). Specifically, LPS activated p38, JNK, and ERK1/2 pathways while flagellin activated the p38 and ERK1/2 pathways, despite both PAMPs inducing similar levels of inflammatory cytokines IL-6, IL-1 β , TNF- α , and CXCL-1. Our results indicate that KSHV-transformed cells are more susceptible to *P. aeruginosa*-induced inflammation through the PAMPs LPS and flagellin




FIG 8 Simultaneous inhibition of p38 and ERK1/2 pathways decreases flagellin-induced inflammation and cell proliferation of KMM cells but has no obvious effect on MM cells. (A) MM and KMM cells were treated for 1 h with PBS, 10 μ M p38 inhibitor (SB 203580), 10 μ M ERK1/2 inhibitor (U0126), or a combination of 10 μ M p38 inhibitor and 10 μ M ERK1/2 inhibitor. Cells were then treated with PBS or 0.3 μ g/ml flagellin for 1 h and analyzed for cytokines by qRT-PCR. (B) MM and KMM cells were treated for 1 h with PBS, 10 μ M p38 inhibitor (SB 203580), 10 μ M ERK1/2 inhibitor (U0126), or a combination of 10 μ M p38 inhibitor and 10 μ M ERK1/2 inhibitor. Cells were then treated with 1 μ g/ml flagellin and examined for cell number at day 3 posttreatment. (C) Schematic illustration summarizing *P. aeruginosa* ligands, and their receptors and activated downstream pathways, that might enhance inflammation and cell proliferation of KSHV-transformed cells during *P. aeruginosa* infection. *, $P \leq 0.05$. NS, not significant.

than the uninfected primary cells. It would be interesting to further confirm the roles of *P. aeruginosa* in KS development in AIDS-KS patients.

It has been shown that TLR activation leads to activation of downstream prosurvival pathways, resulting in increased cancer pathogenesis (42, 43). We previously showed

that KSHV infection upregulated numerous TLRs, including TLR4 and TLR5 (22). LPS and flagellin are highly immunogenic ligands for TLR4 and TLR5, respectively (33). Indeed, TLR4 was upregulated close to 50-fold in KSHV-transformed cells, sensitizing the cells to TLR4 ligands (22). Activation of the TLR4 pathway by *E. coli* and LPS stimulated cell proliferation, cellular transformation, and tumorigenesis by increasing IL-6 expression to activate the STAT3 pathway (22). In this study, we demonstrate that both *P. aeruginosa* LPS and flagellin activate the MAPK pathways and that inhibition of MAPK pathways decreases expression of inflammatory cytokines, cell proliferation, and cellular transformation (Fig. 8C).

We previously demonstrated that LPS stimulation increased cytokines IL-1 β , IL-6, and IL-18 in KMM cells (22). In our current study, we showed that *P. aeruginosa* stimulation increased cytokines, including IL-1 β , IL-6, TNF- α , and CXCL-1, in KMM cells. Inflammatory cytokines such as TNF- α act synergistically with IL-6 to activate the STAT3 pathway (22, 44). Surprisingly, we did not see an increase in IL-18, despite previous studies reporting IL-18 upregulation correlating with *P. aeruginosa* infection (45, 46). It is possible that we had missed the peak time point for the induction of IL-18, as we focused on the effect of acute stimulation. As previously reported, we confirmed the induction of TNF- α and CXCL-1 by *P. aeruginosa* (35, 36, 47). Our results showed that both *P. aeruginosa* LPS and flagellin contributed to the induction of IL-1 β , IL-6, TNF- α , and CXCL-1. It would be interesting to further assess the specific effects of LPS and flagellin in stimulating specific cytokines and their downstream inflammatory pathways in future studies.

The relationship between TLR activation and cancer cell proliferation has been extensively studied (42). Studies have demonstrated that LPS- and flagellin-induced inflammation increases cell proliferation in several cancer cell lines (13, 48, 49). Moreover, some studies suggest that LPS and flagellin blood levels are correlated with a higher cancer risk (50, 51). However, contradictory studies show decreased cell proliferation in LPS- or flagellin-stimulated cells (52–54). In fact, bacteria and the PAMPS have been studied as antitumor agents (55, 56). It is worth mentioning that multiple factors might affect how cells respond to TLR activation. The differences in cell response to TLR activation may depend on the cellular location of the TLRs, as increased TLR expression in the cytoplasm may result in more chronic inflammation and cell proliferation while TLR expression on the cell membrane may result in antitumor effects (57). The proportions of the types of TLRs expressed (TLR2 versus TLR4, for example) may also affect the cellular response (52, 58). There is no doubt that the tumor microenvironment, including the composition of immune cells, can substantially affect the response and proliferation of tumor cells (59). Activation of TLRs in immune cells can result in a Th1 response, causing apoptosis (59). Further elucidation of TLRs and their downstream pathways is essential for understanding the complex interactions between cell proliferation and inflammation.

Lastly, we focused on the effects of *P. aeruginosa* on MAPK activation, showing that *P. aeruginosa* stimulation of KMM cells resulted in the activation of p38, ERK1/2, and JNK pathways. *P. aeruginosa* flagellin contributed to cell proliferation by activating the p38 and ERK1/2 pathways, and inhibition of the p38 and ERK1/2 pathway abrogated the enhanced cell proliferation in flagellin-stimulated KMM cells (Fig. 8C). The MAPK pathways play an important role in KSHV biology and likely in KS pathology. Primary infection of KSHV results in activation of p38, ERK1/2, and JNK activation, and inhibition of the MAPK pathways can reduce KSHV infectivity and induction of IL-6 (24, 60). The MAPK pathways also mediate KSHV reactivation (61). Our results identify the important role of *P. aeruginosa*-induced inflammation in cell proliferation and cellular transformation while providing further evidence on the therapeutic value of inhibiting the MAPK pathways in KS patients.

Overall, our study utilizes a clinically relevant, opportunistic bacterium to study the effect of inflammation on KSHV-transformed cells. We report that *P. aeruginosa* stimulation increases cell proliferation and cellular transformation in KSHV-transformed cells while having no significant effect on MM cells. These results emphasize that KSHV

TABLE 1 Sequences of primers used for qRT-PCR

Primer	Sequence (5' to 3')	
	Forward	Reverse
Rat β -actin	GGAAATCGTGCCTGACATTA	AGGAAGGAAGGCTGGAAGAG
Rat IL-6	TCCTACCCCAACTTCCAATGC	TTGGATGGTCTTGGTCTTAG
Rat IL-1 β	CATTGTGGCTGTGGAGAAG	ATCATCCCACGAGTCACAGA
Rat IL-18	GGCTCTTGCTCAACTTCAAA	TTATCAGTCTGGTCTGGGATT
Rat TNF- α	TGAACTTCGGGGTGATCG	GGGCTTGTCACTCGAGTTTT
Rat CXCL-1	CATTAATATTTAACGATGTGGATGCGTTTCA	GCCTACCATCTTTAAACTGCACAAT

cellular transformation results in enhanced sensitivity to external stimuli, which may further increase cell proliferation and cellular transformation. As KS and HIV/AIDS patients are at an increased risk for opportunistic infection, it is critical to understand the effects of bacteria on KS pathogenesis. Our results indicate that elimination of certain bacterial infections identified to promote inflammation have a preventive value for KSHV-infected AIDS patients who are at a high risk for developing KS as well as a therapeutic value for AIDS-KS patients. Moreover, our study further dissects the specific *P. aeruginosa* PAMPs that contribute to cell proliferation, demonstrating that both LPS and flagellin can induce inflammation in KSHV-transformed cells. Besides LPS and flagellin, *P. aeruginosa* consists of many additional PAMPs that can induce inflammation, such as peptidoglycans and lipoproteins (62, 63). Analyzing the effects of these additional PAMPs on cell proliferation, by exploring the specific pathways and mechanisms of inflammatory induction, may clarify the correlative, additive, or synergistic effects as well as independent effects of these other PAMPs in inflammation processes in general as well as for inflammation and KS pathogenesis specifically.

MATERIALS AND METHODS

Cell culture. Early passages (<20) of MM, KMM, BJAB, and KSHV-BJAB cells were grown as previously described (34, 64). All cell lines were routinely tested for mycoplasma contamination using a LookOut Mycoplasma quantitative PCR (qPCR) detection kit (MP0035-1KT; Sigma).

Reagents. Purified flagellin from *P. aeruginosa* (tlrl-pafla; InvivoGen) and ultrapure LPS from *E. coli* K-12 (tlrl-pek1ps; InvivoGen) were resuspended in water. CLI095 (tlrl-cli095; Thermo Fisher Scientific), SB203850 (NC9041893; Fisher), and U0126 (19-147; Sigma) were resuspended in dimethyl sulfoxide.

Bacterial preparation. Three *P. aeruginosa* strains were used for this study: *P. aeruginosa* (ATCC 15442), PAO1 Δ flhC strain (JH325), and PAO1. PAO1 Δ flhC and PAO1 strains were kindly provided by Jennifer Bomberger (University of Pittsburgh), Joe Harrison (University of Calgary), and Matthew Parsek (University of Washington). *P. aeruginosa* strains were grown in LB broth (Sigma-Aldrich) overnight until an optical density (OD) value of 1.1. The culture was then washed three times in PBS by centrifuging at 3,200 \times g for 10 min and then diluted with PBS to the specified concentrations (CFU per milliliter) for experiments. To confirm the accuracy of the bacterial concentration, the culture was serially diluted, grown on agar plates overnight, and analyzed for the number of colonies.

Cell proliferation assay. MM/KMM cells and BJAB/KSHV-BJAB cells were plated at a density of 20,000 and 10,000 cells/well (respectively) in 24-well plates for 16 h, treated with the indicated reagents, and counted using a Malassez chamber.

Soft agar assay. Colony formation in soft agar was carried out as previously described (34).

RNA extraction and qRT-PCR. Total RNAs were extracted with TRI Reagent (T9424; Sigma). Reverse transcription (RT) was performed with 1 μ g of total RNA using SsoAdvanced universal SYBR green supermix (172-5272; Bio-Rad). cDNAs diluted 2 times were examined by qPCR with the KAPA SYBR fast qPCR kit (K4602; Kapa Biosystems) using specific primers for β -actin, IL-6, IL-1 β , IL-18, TNF- α , and CXCL-1. The β -actin gene was used for loading calibration. All of the sequences of primers used for quantitative RT-PCR (qRT-PCR) are listed in Table 1.

Western blot analysis. Western blot analysis was performed as previously described (65). Primary antibodies included mouse monoclonal antibodies to β -tubulin and rabbit polyclonal antibodies to glyceraldehyde-3-phosphate dehydrogenase (5174S; Cell Signaling Technology), JNK (9252S; Cell Signaling Technology), phospho-JNK (4668S; Cell Signaling Technology), p38 (8690S; Cell Signaling Technology), phospho-p38 (4511S; Cell Signaling Technology), ERK1/2 (4695S; Cell Signaling Technology), and phospho-ERK1/2 (4370S; Cell Signaling Technology).

Statistical analysis. Results are expressed as means \pm standard errors from at least three independent experiments. Statistical analysis was performed using two-tailed *t* test, and a *P* value of ≤ 0.05 was considered significant.

ACKNOWLEDGMENTS

We thank Jennifer Bomberger of the University of Pittsburgh, Joe Harrison of the University of Calgary, and Matthew Parsek of the University of Washington for providing *P. aeruginosa* strains.

This work was supported by grants from the National Institutes of Health (CA096512, CA124332, CA132637, CA213275, DE025465 and CA197153 to S.-J. Gao). This work was supported in part by award P30CA047904.

We thank members of S.-J. Gao's laboratory for technical assistance and helpful discussions.

REFERENCES

- Shaked H, Hofseth LJ, Chumanevich A, Chumanevich AA, Wang J, Wang Y, Taniguchi K, Guma M, Shenouda S, Clevers H, Harris CC, Karin M. 2012. Chronic epithelial NF- κ B activation accelerates APC loss and intestinal tumor initiation through iNOS up-regulation. *Proc Natl Acad Sci U S A* 109:14007–14012. <https://doi.org/10.1073/pnas.1211509109>.
- Coussens LM, Werb Z. 2002. Inflammation and cancer. *Nature* 420:860–867. <https://doi.org/10.1038/nature01322>.
- Suarez-Carmona M, Lesage J, Cataldo D, Gilles C. 2017. EMT and inflammation: inseparable actors of cancer progression. *Mol Oncol* 11:805–823. <https://doi.org/10.1002/1878-0261.12095>.
- Hernández-Ramírez RU, Shiels MS, Dubrow R, Engels EA. 2017. Cancer risk in HIV-infected people in the USA from 1996 to 2012: a population-based, registry-linkage study. *Lancet HIV* 4:e495–e504. [https://doi.org/10.1016/S2352-3018\(17\)30125-X](https://doi.org/10.1016/S2352-3018(17)30125-X).
- Franzetti M, Ricci E, Bonfanti P. 2019. The pattern of non-AIDS-defining cancers in the HIV population: epidemiology, risk factors and prognosis. A review. *Curr HIV Res* 17:1–12. <https://doi.org/10.2174/1570162X17666190327153038>.
- Ariumi Y, Kaida A, Hatanaka M, Shimotohno K. 2001. Functional cross-talk of HIV-1 Tat with p53 through its C-terminal domain. *Biochem Biophys Res Commun* 287:556–561. <https://doi.org/10.1006/bbrc.2001.5626>.
- Nyagol J, Leucci E, Onnis A, De Falco G, Tigli C, Sanseverino F, Torricelli M, Palumbo N, Pacenti L, Santopietro R, Spina D, Gichangi P, Muchiri L, Lazzi S, Petraglia F, Leoncini L, Giordano A. 2006. The effects of HIV-1 Tat protein on cell cycle during cervical carcinogenesis. *Cancer Biol Ther* 5:684–690. <https://doi.org/10.4161/cbt.5.6.2907>.
- Vangipuram R, Tyring SK. 2019. AIDS-associated malignancies. *Cancer Treat Res* 177:1–21. https://doi.org/10.1007/978-3-030-03502-0_1.
- Yao S, Hu M, Hao T, Li W, Xue X, Xue M, Zhu X, Zhou F, Qin D, Yan Q, Zhu J, Gao SJ, Lu C. 2019. miRNA-891a-5p mediates HIV-1 Tat and KSHV Orf-K1 synergistic induction of angiogenesis by activating NF- κ B signaling. *Nucleic Acids Res* 47:2700. <https://doi.org/10.1093/nar/gkz088>.
- Yan Q, Shen C, Qin J, Li W, Hu M, Lu H, Qin D, Zhu J, Gao SJ, Lu C. 2016. HIV-1 Vpr inhibits Kaposi's sarcoma-associated herpesvirus lytic replication by inducing microRNA miR-942-5p and activating NF- κ B signaling. *J Virol* 90:8739–8753. <https://doi.org/10.1128/JVI.00797-16>.
- Yan Q, Zhao R, Shen C, Wang F, Li W, Gao SJ, Lu C. 2018. Upregulation of microRNA 711 mediates HIV-1 Vpr promotion of Kaposi's sarcoma-associated herpesvirus latency and induction of pro-proliferation and pro-survival cytokines by targeting the Notch/NF- κ B-signaling axis. *J Virol* 92:e00580-18.
- Xue M, Yao S, Hu M, Li W, Hao T, Zhou F, Zhu X, Lu H, Qin D, Yan Q, Zhu J, Gao SJ, Lu C. 2014. HIV-1 Nef and KSHV oncogene K1 synergistically promote angiogenesis by inducing cellular miR-718 to regulate the PTEN/AKT/mTOR signaling pathway. *Nucleic Acids Res* 42:9862–9879. <https://doi.org/10.1093/nar/gku583>.
- Yang H, Wang B, Wang T, Xu L, He C, Wen H, Yan J, Su H, Zhu X. 2014. Toll-like receptor 4 prompts human breast cancer cells invasiveness via lipopolysaccharide stimulation and is overexpressed in patients with lymph node metastasis. *PLoS One* 9:e109980. <https://doi.org/10.1371/journal.pone.0109980>.
- Zhou F, Xue M, Qin D, Zhu X, Wang C, Zhu J, Hao T, Cheng L, Chen X, Bai Z, Feng N, Gao SJ, Lu C. 2013. HIV-1 Tat promotes Kaposi's sarcoma-associated herpesvirus (KSHV) vIL-6-induced angiogenesis and tumorigenesis by regulating PI3K/PTEN/AKT/GSK-3 β signaling pathway. *PLoS One* 8:e53145. <https://doi.org/10.1371/journal.pone.0053145>.
- Zeng Y, Zhang X, Huang Z, Cheng L, Yao S, Qin D, Chen X, Tang Q, Lv Z, Zhang L, Lu C. 2007. Intracellular Tat of human immunodeficiency virus type 1 activates lytic cycle replication of Kaposi's sarcoma-associated herpesvirus: role of JAK/STAT signaling. *J Virol* 81:2401–2417. <https://doi.org/10.1128/JVI.02024-06>.
- Chao C, Leyden WA, Xu L, Horberg MA, Klein D, Towner WJ, Quesenberry CP, Jr, Abrams DI, Silverberg MJ. 2012. Exposure to antiretroviral therapy and risk of cancer in HIV-infected persons. *AIDS* 26:2223–2231. <https://doi.org/10.1097/QAD.0b013e32835935b3>.
- Okoye AA, Picker LJ. 2013. CD4(+) T-cell depletion in HIV infection: mechanisms of immunological failure. *Immunol Rev* 254:54–64. <https://doi.org/10.1111/immr.12066>.
- Nowroozalazadeh S, Månsson F, da Silva Z, Repits J, Dabo B, Pereira C, Biague A, Albert J, Nielsen J, Aaby P, Fenyö EM, Norrgren H, Holmgren B, Jansson M. 2010. Microbial translocation correlates with the severity of both HIV-1 and HIV-2 infections. *J Infect Dis* 201:1150–1154. <https://doi.org/10.1086/651430>.
- Shebl FM, Yu K, Landgren O, Goedert JJ, Rabkin CS. 2012. Increased levels of circulating cytokines with HIV-related immunosuppression. *AIDS Res Hum Retroviruses* 28:809–815. <https://doi.org/10.1089/aid.2011.0144>.
- Drozd DR, Graham SM, Crane HM, Delaney JA, Kitahata MM, Liles WC. 2016. Short communication: effect of antiretroviral therapy on circulating damage-associated molecular pattern molecules and CD4 immune reconstitution in HIV-infected individuals. *AIDS Res Hum Retroviruses* 32:876–878. <https://doi.org/10.1089/AID.2016.0059>.
- He M, Cheng F, da Silva SR, Tan B, Sorel O, Gruffaz M, Li T, Gao SJ. 2019. Molecular biology of KSHV in relation to HIV/AIDS-associated oncogenesis. *Cancer Treat Res* 177:23–62. https://doi.org/10.1007/978-3-030-03502-0_2.
- Gruffaz M, Vasan K, Tan B, Ramos da Silva S, Gao SJ. 2017. TLR4-mediated inflammation promotes KSHV-induced cellular transformation and tumorigenesis by activating the STAT3 pathway. *Cancer Res* 77:7094–7108. <https://doi.org/10.1158/0008-5472.CAN-17-2321>.
- Park MK, Cho H, Roh SW, Kim SJ, Myoung J. 2019. Cell type-specific interferon- γ -mediated antagonism of KSHV lytic replication. *Sci Rep* 9:2372. <https://doi.org/10.1038/s41598-019-38870-7>.
- Xie J, Pan H, Yoo S, Gao SJ. 2005. Kaposi's sarcoma-associated herpesvirus induction of AP-1 and interleukin 6 during primary infection mediated by multiple mitogen-activated protein kinase pathways. *J Virol* 79:15027–15037. <https://doi.org/10.1128/JVI.79.24.15027-15037.2005>.
- Qian LW, Xie J, Ye F, Gao SJ. 2007. Kaposi's sarcoma-associated herpesvirus infection promotes invasion of primary human umbilical vein endothelial cells by inducing matrix metalloproteinases. *J Virol* 81:7001–7010. <https://doi.org/10.1128/JVI.00016-07>.
- Ye FC, Blackburn DJ, Mengel M, Xie JP, Qian LW, Greene W, Yeh IT, Graham D, Gao SJ. 2007. Kaposi's sarcoma-associated herpesvirus promotes angiogenesis by inducing angiopoietin-2 expression via AP-1 and Ets1. *J Virol* 81:3980–3991. <https://doi.org/10.1128/JVI.02089-06>.
- Xu Y, Ganem D. 2007. Induction of chemokine production by latent Kaposi's sarcoma-associated herpesvirus infection of endothelial cells. *J Gen Virol* 88:46–50. <https://doi.org/10.1099/vir.0.82375-0>.
- Ye F, Zhou F, Bedolla RG, Jones T, Lei X, Kang T, Guadalupe M, Gao SJ. 2011. Reactive oxygen species hydrogen peroxide mediates Kaposi's sarcoma-associated herpesvirus reactivation from latency. *PLoS Pathog* 7:e1002054. <https://doi.org/10.1371/journal.ppat.1002054>.
- Lee MS, Jones T, Song DY, Jang JH, Jung JU, Gao SJ. 2014. Exploitation of the complement system by oncogenic Kaposi's sarcoma-associated

- herpesvirus for cell survival and persistent infection. *PLoS Pathog* 10: e1004412. <https://doi.org/10.1371/journal.ppat.1004412>.
30. Gruffaz M, Zhang T, Marshall V, Gonçalves P, Ramaswami R, Labo N, Whitby D, Uldrick TS, Yarchoan R, Huang Y, Gao SJ. 2020. Signatures of oral microbiome in HIV-infected individuals with oral Kaposi's sarcoma and cell-associated KSHV DNA. *PLoS Pathog* 16:e1008114. <https://doi.org/10.1371/journal.ppat.1008114>.
 31. Riccardi N, Rotulo GA, Favilli F, Loy A, Moratto D, Giliani S, Mesini A, Romanini MV, Volpi S, Moscatelli A, Castagnola E. 2019. *Pseudomonas aeruginosa* severe skin infection in a toddler with X-linked agammaglobulinemia due to a novel BTK mutation. *Infez Med* 27:73–76.
 32. Huang L, Crothers K. 2009. HIV-associated opportunistic pneumonias. *Respirology* 14:474–485. <https://doi.org/10.1111/j.1440-1843.2009.01534.x>.
 33. Sun Y, Karmakar M, Roy S, Ramadan RT, Williams SR, Howell S, Shive CL, Han Y, Stopford CM, Rietsch A, Pearlman E. 2010. TLR4 and TLR5 on corneal macrophages regulate *Pseudomonas aeruginosa* keratitis by signaling through MyD88-dependent and -independent pathways. *J Immunol* 185:4272–4283. <https://doi.org/10.4049/jimmunol.1000874>.
 34. Jones T, Ye F, Bedolla R, Huang Y, Meng J, Qian L, Pan H, Zhou F, Moody R, Wagner B, Arar M, Gao SJ. 2012. Direct and efficient cellular transformation of primary rat mesenchymal precursor cells by KSHV. *J Clin Investig* 122:1076–1081. <https://doi.org/10.1172/JCI58530>.
 35. Yang YW, Jiang YZ, Hsu CM, Chen LW. 2017. *Pseudomonas aeruginosa* ventilator-associated pneumonia induces lung injury through TNF- α /c-Jun NH2-terminal kinase pathways. *PLoS One* 12:e0169267. <https://doi.org/10.1371/journal.pone.0169267>.
 36. Liu Y, Di ME, Chu HW, Liu X, Wang L, Wenzel S, Di YP. 2013. Increased susceptibility to pulmonary *Pseudomonas* infection in Splunc1 knockout mice. *J Immunol* 191:4259–4268. <https://doi.org/10.4049/jimmunol.1202340>.
 37. Kim EK, Choi EJ. 2010. Pathological roles of MAPK signaling pathways in human diseases. *Biochim Biophys Acta* 1802:396–405. <https://doi.org/10.1016/j.bbadis.2009.12.009>.
 38. Harrison JJ, Almlad H, Irie Y, Wolter DJ, Eggleston HC, Randall TE, Kitzman JO, Stackhouse B, Emerson JC, McNamara S, Larsen TJ, Shendure J, Hoffman LR, Wozniak DJ, Parsek MR. 2020. Elevated exopolysaccharide levels in *Pseudomonas aeruginosa* flagellar mutants have implications for biofilm growth and chronic infections. *PLoS Genet* 16:e1008848. <https://doi.org/10.1371/journal.pgen.1008848>.
 39. Bose M, Mukherjee P. 2019. Role of microbiome in modulating immune responses in cancer. *Mediators Inflamm* 2019:4107917. <https://doi.org/10.1155/2019/4107917>.
 40. van Elsland D, Neeffjes J. 2018. Bacterial infections and cancer. *EMBO Rep* 19:e46632.
 41. Gorres KL, Daigle D, Mohanram S, Miller G. 2014. Activation and repression of Epstein-Barr Virus and Kaposi's sarcoma-associated herpesvirus lytic cycles by short- and medium-chain fatty acids. *J Virol* 88: 8028–8044. <https://doi.org/10.1128/JVI.00722-14>.
 42. Bagratuni T, Sklirou AD, Kastiris E, Liacos CI, Spilioti C, Eleutherakis-Papaikovou E, Kanellias N, Gavriatopoulou M, Terpos E, Trougakos IP, Dimopoulos MA. 2019. Toll-Like receptor 4 activation promotes multiple myeloma cell growth and survival via suppression of the endoplasmic reticulum stress factor chop. *Sci Rep* 9:3245. <https://doi.org/10.1038/s41598-019-39672-7>.
 43. Ran J, Zhou J. 2019. Targeted inhibition of histone deacetylase 6 in inflammatory diseases. *Thorac Cancer* 10:405–412. <https://doi.org/10.1111/1759-7714.12974>.
 44. De Simone V, Franzè E, Ronchetti G, Colantoni A, Fantini MC, Di Fusco D, Sica GS, Sileri P, MacDonald TT, Pallone F, Monteleone G, Stolfi C. 2015. Th17-type cytokines, IL-6 and TNF- α synergistically activate STAT3 and NF- κ B to promote colorectal cancer cell growth. *Oncogene* 34: 3493–3503. <https://doi.org/10.1038/onc.2014.286>.
 45. Cendra MDM, Christodoulides M, Hossain P. 2017. Signaling mediated by Toll-like receptor 5 sensing of *Pseudomonas aeruginosa* flagellin influences IL-1 β and IL-18 production by primary fibroblasts derived from the human cornea. *Front Cell Infect Microbiol* 7:130. <https://doi.org/10.3389/fcimb.2017.00130>.
 46. Sahoo M, Ceballos-Olvera I, del Barrio L, Re F. 2011. Role of the inflammasome, IL-1 β , and IL-18 in bacterial infections. *ScientificWorldJournal* 11:2037–2050. <https://doi.org/10.1100/2011/212680>.
 47. Metruccio MME, Tam C, Evans DJ, Xie AL, Stern ME, Fleiszig SMJ. 2017. Contributions of MyD88-dependent receptors and CD11c-positive cells to corneal epithelial barrier function against *Pseudomonas aeruginosa*. *Sci Rep* 7:13829. <https://doi.org/10.1038/s41598-017-14243-w>.
 48. Li N, Xu H, Ou Y, Feng Z, Zhang Q, Zhu Q, Cai Z. 2019. LPS-induced CXCR7 expression promotes gastric cancer proliferation and migration via the TLR4/MD-2 pathway. *Diagn Pathol* 14:3. <https://doi.org/10.1186/s13000-019-0780-x>.
 49. Song EJ, Kang MJ, Kim YS, Kim SM, Lee SE, Kim CH, Kim DJ, Park JH. 2011. Flagellin promotes the proliferation of gastric cancer cells via the Toll-like receptor 5. *Int J Mol Med* 28:115–119. <https://doi.org/10.3892/ijmm.2011.656>.
 50. Fedirko V, Tran HQ, Gewirtz AT, Stepien M, Trichopoulos A, Aleksandrova K, Olsen A, Tjønneland A, Overvad K, Carbonnel F, Boutron-Ruault MC, Severi G, Kühn T, Kaaks R, Boeing H, Bamia C, Lagiou P, Grioni S, Panico S, Palli D, Tumino R, Naccarati A, Peeters PH, Bueno-de-Mesquita HB, Weiderpass E, Castaño JM, Barricarte A, Sánchez MJ, Dorronsoro M, Quirós JR, Agudo A, Sjöberg K, Ohlsson S, Hemmingsson O, Werner M, Bradbury KE, Khaw KT, Wareham N, Tsilidis KK, Aune D, Scalbert A, Romieu I, Riboli E, Jenab M. 2017. Exposure to bacterial products lipopolysaccharide and flagellin and hepatocellular carcinoma: a nested case-control study. *BMC Med* 15:72. <https://doi.org/10.1186/s12916-017-0830-8>.
 51. Kong SY, Tran HQ, Gewirtz AT, McKeown-Eyssen G, Fedirko V, Romieu I, Tjønneland A, Olsen A, Overvad K, Boutron-Ruault M-C, Bastide N, Affret A, Ku Hn T, Kaaks R, Boeing H, Aleksandrova K, Trichopoulos A, Kritikou M, Vasilopoulou E, Palli D, Krogh V, Mattiello A, Tumino R, Naccarati A, Bueno-de-Mesquita HB, Peeters PH, Weiderpass E, Quiros JR, Salá N, Sanchez M-J, Castano JMH, Barricarte A, Dorronsoro M, Werner M, Wareham NJ, Khaw K-T, Bradbury KE, Freisling H, Stavropoulou F, Ferrari P, Gunter MJ, Cross AJ, Riboli E, Bruce WR, Jenab M. 2016. Serum endotoxins and flagellin and risk of colorectal cancer in the european prospective investigation into cancer and nutrition (EPIC) cohort. *Cancer Epidemiol Biomarkers Prev* 25:291–301. <https://doi.org/10.1158/1055-9965.EPI-15-0798>.
 52. Shi J, Zhao Y, Wang Y, Gao W, Ding J, Li P, Hu L, Shao F. 2014. Inflammatory caspases are innate immune receptors for intracellular LPS. *Nature* 514:187–192. <https://doi.org/10.1038/nature13683>.
 53. Fried S, Tosun S, Troost G, Keil S, Zaenker KS, Dittmar T. 2016. Lipopolysaccharide (LPS) promotes apoptosis in human breast epithelial \times breast cancer hybrids, but not in parental cells. *PLoS One* 11:e0148438. <https://doi.org/10.1371/journal.pone.0148438>.
 54. Cai Z, Sanchez A, Shi Z, Zhang T, Liu M, Zhang D. 2011. Activation of Toll-like receptor 5 on breast cancer cells by flagellin suppresses cell proliferation and tumor growth. *Cancer Res* 71:2466–2475. <https://doi.org/10.1158/0008-5472.CAN-10-1993>.
 55. Shetab Boushehri MA, Lamprecht A. 2019. Challenges of using lipopolysaccharides for cancer immunotherapy and potential delivery-based solutions thereto. *Ther Deliv* 10:165–187. <https://doi.org/10.4155/tde-2018-0076>.
 56. Duong MT, Qin Y, You SH, Min JJ. 2019. Bacteria-cancer interactions: bacteria-based cancer therapy. *Exp Mol Med* 51:1–15. <https://doi.org/10.1038/s12276-019-0297-0>.
 57. Jouhi L, Renkonen S, Atula T, Mäkitie A, Haglund C, Hagström J. 2014. Different Toll-like receptor expression patterns in progression toward cancer. *Front Immunol* 5:638. <https://doi.org/10.3389/fimmu.2014.00638>.
 58. Khan AA, Khan Z, Warnakulasuriya S. 2016. Cancer-associated toll-like receptor modulation and insinuation in infection susceptibility: association or coincidence? *Ann Oncol* 27:984–997. <https://doi.org/10.1093/annonc/mdw053>.
 59. Li J, Yang F, Wei F, Ren X. 2017. The role of toll-like receptor 4 in tumor microenvironment. *Oncotarget* 8:66656–66667. <https://doi.org/10.18632/oncotarget.19105>.
 60. Pan H, Xie J, Ye F, Gao SJ. 2006. Modulation of Kaposi's sarcoma-associated herpesvirus infection and replication by MEK/ERK, JNK, and p38 multiple mitogen-activated protein kinase pathways during primary infection. *J Virol* 80:5371–5382. <https://doi.org/10.1128/JVI.02299-05>.
 61. Xie J, Ajibade AO, Ye F, Kuhne K, Gao SJ. 2008. Reactivation of Kaposi's sarcoma-associated herpesvirus from latency requires MEK/ERK, JNK and p38 multiple mitogen-activated protein kinase pathways. *Virology* 371: 139–154. <https://doi.org/10.1016/j.virol.2007.09.040>.
 62. Torrens G, Escobar-Salom M, Pol-Pol E, Camps-Munar C, Cabot G, López-Causapé C, Rojo-Moliner E, Oliver A, Juan C. 2019. Comparative analysis of peptidoglycans from *Pseudomonas aeruginosa* isolates recovered

- from chronic and acute infections. *Front Microbiol* 10:1868. <https://doi.org/10.3389/fmicb.2019.01868>.
63. Firoved AM, Ornatowski W, Deretic V. 2004. Microarray analysis reveals induction of lipoprotein genes in mucoid *Pseudomonas aeruginosa*: implications for inflammation in cystic fibrosis. *Infect Immun* 72: 5012–5018. <https://doi.org/10.1128/IAI.72.9.5012-5018.2004>.
64. Chen L, Lagunoff M. 2005. Establishment and maintenance of Kaposi's sarcoma-associated herpesvirus latency in B cells. *J Virol* 79: 14383–14391. <https://doi.org/10.1128/JVI.79.22.14383-14391.2005>.
65. Gao SJ, Kingsley L, Hoover DR, Spira TJ, Rinaldo CR, Saah A, Phair J, Detels R, Parry P, Chang Y, Moore PS. 1996. Seroconversion to antibodies against Kaposi's sarcoma-associated herpesvirus-related latent nuclear antigens before the development of Kaposi's sarcoma. *N Engl J Med* 335:233–241. <https://doi.org/10.1056/NEJM199607253350403>.