

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

Sites of Azaserine Inhibition During Photosynthesis by Scenedesmus

### Permalink

<https://escholarship.org/uc/item/06h8z5q4>

### Authors

Barker, S Alan

Bassham, James A

Calvin, M

et al.

### Publication Date

1956-05-01

UNIVERSITY OF  
CALIFORNIA

*Radiation  
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory  
Berkeley, California

Contract No. W-7405-eng-48

SITES OF AZASERINE INHIBITION  
DURING PHOTOSYNTHESIS BY SCENEDESMUS

S. Alan Barker, James A. Bassham, M. Calvin,  
and Ursula C. Quarck

May 8, 1956

**SITES OF AZASERINE INHIBITION  
DURING PHOTOSYNTHESIS BY SCENEDESMUS**

**S. Alan Barker, James A. Bassham, M. Calvin  
and Ursula C. Quarck**

**Radiation Laboratory  
University of California  
Berkeley, California**

**May 8, 1956**

**ABSTRACT**

**L-azaserine has been found to have a profound effect on the reservoir sizes of many of the metabolic intermediates produced during photosynthesis by Scenedesmus. Marked increases in levels of glutamine and the acids of the Krebs cycle were accompanied by a corresponding depletion of the amino acid reservoirs, indicating that one of the major sites of azaserine action is in reactions involving transamination. In contrast, the photosynthetic carbon cycle is virtually unaffected and the rate of formation of sucrose is increased.**

SITES OF AZASERINE INHIBITION  
DURING PHOTOSYNTHESIS BY SCENEDESMUS

S. Alan Barker, \* James A. Bassham, M. Calvin,  
And Ursula C. Quarck

Radiation Laboratory  
University of California  
Berkeley, California

May 8, 1956

INTRODUCTION

The success attending the use of azaserine<sup>1, 2</sup> as a specific inhibitor of one stage in the metabolic pathway leading to the synthesis of inosinic acid in pigeon liver prompted us to use this antibiotic in a similar attack on purine synthesis in Scenedesmus. However, investigation of the products produced during photosynthesis by suspensions of these algae in the presence of azaserine showed that a more widespread interference with metabolism had occurred. The purpose of this communication is to describe the nature of these effects and to attempt to assess their importance in a general picture of the metabolic effects of azaserine.

EXPERIMENTAL

Experimental Procedure

Two suspensions, each containing washed Scenedesmus cells (packed volume, 0.2 cc) and  $\text{KH}_2\text{PO}_4$  solution (0.4 cc;  $3.2 \times 10^{-6} \text{ M}$ ) in 21 cc and one with added L-azaserine (4 mg), were left for 1 hour in thin glass containers illuminated on each side by a 150-watt light (reflector flood) to achieve steady states with 4%  $\text{CO}_2$  in air. Each suspension was then allowed to photosynthesize for 5 minutes with  $\text{NaHC}^{14}\text{O}_3$  solution (0.9 cc; 360  $\mu\text{c}$ ) and then flushed with air for 1 minute. The cells were killed by pouring them into boiling ethanol (88 cc), and the resulting 80% ethanol extract was separated from insoluble material which was then reextracted with 20% ethanol (100 cc). The total fixation of radioactivity was determined in each case by uniformly distributing and drying 50- $\mu\text{l}$  aliquots of the 80% ethanol suspensions plus

---

\* Rockefeller Research Fellow, 1955-56.

<sup>1</sup> Hartman, Levenberg, and Buchanan, J. Am. Chem. Soc. 77, 501 (1955).

<sup>2</sup> B. Levenberg and J. M. Buchanan, J. Am. Chem. Soc. 78, 504 (1956).

50  $\mu$ l 6 N acetic acid on aluminum discs, and counting the radioactivity with a Scott large-window Geiger-Mueller tube. The radioactivities extracted from each suspension with 80% ethanol and with 20% ethanol were determined in a similar manner. After the combined extracts of each suspension were concentrated to 3 to 4 cc, aliquots (calculated to contain  $1 \times 10^6$  counts/min. each) were applied to several washed Whatman No. 4 papers and separated first in phenol-water and then in butanol-propionic acid in the manner described by Wilson and Calvin.<sup>3</sup> After radioautographs of the chromatograms had been made, the various components detected on the papers were counted (Table I). Since the correction for self-absorption of radiation would be the same for each compound, no correction was applied. The results of a duplicate experiment, in which 1 mg of azaserine was used, are also presented (Table I).

Other experiments carried out in a manner identical with those described above were (1) a repetition of the 1-mg-azaserine experiment in which an intense photospot light was substituted for one of the reflector floods for 50 minutes of the 1-hour steady-state period (Table II), and (2) an experiment in which one suspension contained 4 mg azaserine and the other contained both azaserine (4 mg) and glutamine (5 mg) (Table III).

Chromatograms from each of the above experiments were sprayed with ninhydrin to obtain a qualitative estimate of the concentrations of amino acids present. Glutamine obtained from azaserine chromatograms, as well as the glutamic acid produced from it by hydrolysis with 1 N HCl at 100° for 2 hours, were characterized by cochromatography on two-way paper chromatograms and by ionophoresis on paper for 3 hours at 600 v in 0.1 M propionate, pH 5.6. Sucrose, amino acids, and carboxylic acids mentioned in Table I were likewise characterized by cochromatography on two-way paper chromatograms and most of them also by ionophoretic separations. The lipids, phospholipids, and pigments all were moved at a similar distance in phenol-water but were separated in butanol-propionic acid into two mobile areas designated lipid (containing some pigment), phospholipid (containing no visible pigment), and a stationary area designated X (containing some pigment). No radioactivity could be extracted from these areas by water elution, but most of the activity was extractable with a mixture of petroleum ether and ethanol. The compounds in Area X ran faster than diphosphopyridine nucleotide in phenol-water, but showed the same lack of mobility in butanol-propionic acid.

<sup>3</sup> A. T. Wilson and M. Calvin, J. Am. Chem. Soc. 77, 5948 (1955).

Table I

The effect of azaserine on Scenedesmus

|   | Control I              | Azaserine I (4 mg)   | Control II             | Azaserine II (1 mg)  |
|---|------------------------|----------------------|------------------------|----------------------|
| <u>Determination on aluminum disc (counts/min)</u>  |                        |                      |                        |                      |
| Total fixation  | 22 x 10 <sup>6</sup>   | 27 x 10 <sup>6</sup> | 29.2 x 10 <sup>6</sup> | 33 x 10 <sup>6</sup> |
| 80% ethanol extract   | 45.5%                  | 47%                  | 47.3%                  | 47.3%                |
| 20% ethanol extract   | 5.9%                   | 7.5%                 | 7.5%                   | 8.5%                 |
| <u>Determination on paper (counts/min equal total radioactivity placed on each paper)</u> |                        |                      |                        |                      |
| Glutamine   | 332                    | 1192                 | 390                    | 1312                 |
| Glutamic acid   | 2066                   | 1227                 | 2632                   | 1139                 |
| Aspartic acid   | 4436                   | 989                  | 7020                   | 762                  |
| Serine  | 1839                   | 849                  | 2411                   | 990                  |
| Alanine   | 1166                   | 272                  | 2000                   | 314                  |
| Glycine }<br>Threonine }  | 400                    | 574                  | 1045                   | 908                  |
| Tyrosine  | 261                    | 171                  | 444                    | 295                  |
| Valine  | 1051                   | 787                  | 1468                   | 1053                 |
| Sucrose   | 923 x 9.5 <sup>a</sup> | 2380 x 9.5           | 937                    | 2072                 |
| Malic Acid  | 3922 x 9.5             | 5814 x 9.5           | 2691 x 9.5             | 4263 x 9.5           |
| Citric acid   | 466 x 9.5              | 1118 x 9.5           | 145 x 9.5              | 400 x 9.5            |
| $\alpha$ -Ketoglutaric acid   | 40                     | 200                  | 270                    | 1054                 |
| Fumaric acid  | 269                    | 108                  | 381                    | 344                  |
| Succinic acid   | 1117                   | 1689                 | 822                    | 922                  |
| Glyceric acid   | 576                    | 966                  | 1349                   | 845                  |
| Glycolic acid   | 200                    | 3182                 | 240                    | 880                  |
| Lipids  | 1650 x 9.5             | 1259 x 9.5           | 2540 x 9.5             | 1923 x 9.5           |
| Phospholipids   | 1229 x 9.5             | 1071 x 9.5           | 1350 x 9.5             | 853 x 9.5            |
| Area X  | 1001 x 9.5             | 751 x 9.5            | 621 x 9.5              | 538 x 9.5            |
| Sugar phosphates,<br>PGA and origin   | 14,040 x 9.5           | 16,379 x 9.5         | 15,071 x 9.5           | 14,424 x 9.5         |

<sup>a</sup> The factor 9.5 was used whenever the radioactivity was counted through aluminum foil.



Table II

---



---

 The effect of azaserine and strong light on Scenedesmus


---

|  | Control III         | Azaserine III (1 mg) |
|--|---------------------|----------------------|
| <u>Determination on aluminum disc (counts/min)</u> |                     |                      |
| Total fixation                                     | $20.9 \times 10^6$  | $14 \times 10^6$     |
| 80% ethanol extract                                | 47.8%               | 57.8%                |
| 20% ethanol extract                                | 2.5%                | 4.4%                 |
| <u>Determination on paper (counts/min)</u>         |                     |                      |
| Glutamine  | 529                 | 4964                 |
| Glutamic acid                                      | 4951                | 2945                 |
| Aspartic acid                                      | 9089                | 2262                 |
| Serine   | 2834                | 1591                 |
| Alanine  | 2345                | 951                  |
| Glycine }<br>Threonine }                           | 949                 | 929                  |
| Sucrose  | $1357 \times 9.5$   | $1907 \times 9.5$    |
| Malic acid   | $3743 \times 9.5$   | $6479 \times 9.5$    |
| Citric acid  | $254 \times 9.5$    | $1919 \times 9.5$    |
| $\alpha$ -Ketoglutaric acid                        | 381                 | 1851                 |
| Fumaric acid                                       | 344                 | 540                  |
| Succinic acid                                      | 1621                | 3180                 |
| Glyceric acid                                      | 1553                | 1312                 |
| Glycolic acid                                      | 285                 | 5 06                 |
| Lipids   | $3582 \times 9.5$   | $2491 \times 9.5$    |
| Phospholipids                                      | $2259 \times 9.5$   | $1665 \times 9.5$    |
| Area X   | $725 \times 9.5$    | $506 \times 9.5$     |
| Sugar Phosphates,<br>PGA and origin                | $11,800 \times 9.5$ | $9312 \times 9.5$    |

---



---

Table III

---

 Attempted reversal of azaserine inhibition with glutamine
 

---

|  | Azaserine (4 mg)<br>alone | Azaserine (4 mg)+<br>glutamine (5 mg) |
|--|---------------------------|---------------------------------------|
| <u>Determination on aluminum disc (counts/min)</u> |                           |                                       |
| Total fixation                                     | $33.6 \times 10^6$        | $35.8 \times 10^6$                    |
| 80% ethanol extract                                | 41.2%                     | 52.9%                                 |
| 20% ethanol extract                                | 1.15%                     | 2.22%                                 |
| <u>Determination on paper (counts/min)</u>         |                           |                                       |
| Glutamine  | 2182                      | 6572                                  |
| Glutamic acid                                      | 1353                      | 2704                                  |
| Aspartic acid                                      | 666                       | 939                                   |
| Serine   | 735                       | 690                                   |
| Alanine  | 393                       | 473                                   |
| Glycine }<br>Threonine }                           | 503                       | 527                                   |
| Sucrose  | $1496 \times 9.5$         | $1111 \times 9.5$                     |
| Malic acid   | $3697 \times 9.5$         | $4631 \times 9.5$                     |
| Citric acid  | $586 \times 9.5$          | $446 \times 9.5$                      |
| $\alpha$ -Ketoglutaric acid                        | 696                       | 1052                                  |
| Fumaric acid                                       | 242                       | 249                                   |
| Succinic acid                                      | 870                       | 1308                                  |
| Glycolic acid                                      | 3424                      | 3138                                  |
| Lipids   | $1639 \times 9.5$         | $2192 \times 9.5$                     |
| Phospholipids                                      | $541 \times 9.5$          | $1310 \times 9.5$                     |
| Area X   | $797 \times 9.5$          | $1370 \times 9.5$                     |
| Sugar Phosphates<br>PGA and origin                 | $11,970 \times 9.5$       | $12,383 \times 9.5$                   |

---

The triose phosphate, pentose monophosphate, glucose cyclic 1, 2-phosphate, hexose monophosphate, phosphoglyceric acid, and phosphoenolpyruvate were eluted from control and azaserine chromatograms, and treated with purified "Polidase" phosphatase, and the products were analyzed by paper chromatography and radioautography.<sup>3</sup>

The effect of azaserine (4 mg) on *Chlorella* was also determined in two experiments (Table IV) identical with that described for the *Scenedesmus* model experiment except that in one case the time of flushing with air was 30 sec and in the other, 60 sec.

## DISCUSSION

It is evident from Table I that with *Scenedesmus* the azaserine causes a build-up of the acids (e. g., citric, malic,  $\alpha$ -ketoglutaric, succinic) of the Krebs tricarboxylic acid cycle and a decrease in radioactivity and amount (revealed by ninhydrin) of the amino acids (glutamic acid, aspartic acid) derived therefrom by amination or transamination. Alanine, serine, valine and tyrosine, which are also known to be formed by transamination of their corresponding keto-acids, were likewise depleted. Because of the low activity of glycine and threonine and the presence of other compounds in this area the effect of azaserine on these amino acids could not be assessed. Glutamine,<sup>4</sup> which is a donor of amino groups during transamination, increases markedly in activity and amount (approximately 1 mg/g cells, as estimated by ninhydrin) in the presence of azaserine. It is therefore suggested that azaserine interferes with transamination reactions, either directly by inhibiting the actual process of transamination or indirectly by interfering with the synthesis of pyridoxal phosphate or pyridoxalamine phosphate, which are necessary as coenzymes of such reactions. The second possibility seems less likely because of the short time available for that inhibition to appear for which these compounds are coenzymes. Figure 1 indicates some of the suggested sites of azaserine inhibition. Kaplan and Stock<sup>5</sup> have also concluded that azaserine is an inhibitor of amino acid synthesis, on the basis of their observation that inhibition of growth in *E. coli* by azaserine can be overcome by addition of certain amino acids. Two azaserine-inhibited reactions, which have recently been reported, would also fit into this scheme. It has been shown that azaserine inhibits the conversion of  $\alpha$ -N-formyl glycinamide ribotide to  $\alpha$ -N-formylglycinamide ribotide<sup>2</sup> and the amination of xanthosine-5'-phosphate to guanosine-5'-phosphate.<sup>6</sup> L-glutamine is the donor of amino groups in both these reactions.

<sup>4</sup> A. Meister and S. V. Tice, *J. Biol. Chem.* **187**, 173 (1950).

<sup>5</sup> L. Kaplan and C. C. Stock, *Federation Proc.* **13**, 239 (1954).

<sup>6</sup> M. Bentley and R. Adams, *Federation Proc.* **15**, 218 (1956).

Table IV

The Effect of Azaserine on Chlorella

|  | Control IV             | Azaserine IV<br>(4 mg)-air-<br>flushed 30 sec | Control V              | Azaserine V<br>(4 mg)-air-<br>flushed 60 sec |
|--|------------------------|---|------------------------|--|
| <b>Determination on aluminum disc (counts/min)</b> |                        |   |                        |  |
| Total fixation                                     | 25.4 x 10 <sup>6</sup> | 30.8 x 10 <sup>6</sup>                        | 23.4 x 10 <sup>6</sup> | 34.8 x 10 <sup>6</sup>                       |
| 80% ethanol extract                                | 60.7%                  | 66.4%   | 62%                    | 62.8%  |
| 20% ethanol extract                                | 8.3%                   | 8.25%   | 10.6%                  | 8.1%   |
| <b>Determination on paper (counts/min)</b>         |                        |   |                        |  |
| Glutamine  | 250                    | 300   | 623                    | 530  |
| Glutamic acid                                      | 2203                   | 2290  | 2370                   | 2000   |
| Aspartic acid                                      | 2541                   | 2700  | 1900                   | 2180   |
| Serine   | 2866                   | 3361  | 3890                   | 4770   |
| Alanine  | 9635                   | 8232  | 10,300                 | 9670   |
| Glycine }<br>Threonine }                           | 800                    | 960   | 780                    | 830  |
| Sucrose  | 4458 x 9.5             | 3933 x 9.5                                    | 4350 x 9.5             | 4330 x 9.5                                   |
| Malic acid   | 2243                   | 2409  | 1980                   | 1970   |
| Citric acid  | 245                    | 250   | 200                    | 220  |
| Fumaric acid                                       | 571                    | 428   | 380                    | 300  |
| Lipids }<br>Phospholipids }                        | 2101 x 9.5             | 1895 x 9.5                                    | 2800 x 9.5             | 2750 x 9.5                                   |
| Area X   | 880 x 9.5              | 858 x 9.5                                     | 630 x 9.5              | 640 x 9.5                                    |
| Sugar phosphates,<br>PGA and origin                | 11,583 x 9.5           | 12,085 x 9.5                                  | 18,690 x 9.5           | 19,140 x 9.5                                 |




Fig. 1. Possible sites of azaserine inhibition.

An interesting feature is the build-up of glutamine despite the fact that the formation of glutamic acid (its precursor) is partially inhibited by azaserine. It is to be noted, however, that the decrease in glutamic acid is by no means so marked as the decreased formation of aspartic acid in the presence of azaserine. One explanation of this may be that whereas it is known<sup>7</sup> that glutamic acid can be synthesized from  $\text{NH}_4^+$  and  $\alpha$ -ketoglutaric acid, the synthesis of aspartic acid by a similar reaction of  $\text{NH}_4^+$  and oxalacetic acid has not been proved. This alternative pathway of glutamic acid synthesis, not involving transamination, probably operates in Scenedesmus.

Since the formation of serine is blocked partially by azaserine, smaller amounts of ethanalamine and choline (which are derived from it) are available for incorporation into phospholipids, and this may account for the decreased radioactivity found in this region of the chromatograms in the presence of azaserine. The lipid and area X also show decreased activity. This might be accounted for, at least in part, by blockage of a transamination reaction leading to the glycine that is necessary for the synthesis of the heterocyclic rings in the porphyrin pigments.<sup>8</sup>

Glycolic acid, which generally accumulates<sup>3</sup> at low  $\text{CO}_2$  pressures, was considerably increased at higher levels of azaserine. This could not be caused by a lower rate of  $\text{CO}_2$  fixation, since this is increased by azaserine during photosynthesis. The increased levels of acids in the Krebs cycle may cause an increased formation of glyoxylic acid,<sup>9</sup> which with an inhibited glycine formation would be available for glycolic acid production. The increased rate of  $\text{CO}_2$  fixation in the presence of azaserine is probably responsible for the increased level of sucrose drained from the photosynthetic cycle.

Examination of the components of the photosynthetic cycle, by analysis of the products produced by phosphatase action, revealed the presence of glyceric acid and all the usual sugars. Several additional unidentified components were detected in the phosphatased triose phosphate and glucose cyclic 1, 2 phosphate areas eluted from azaserine chromatograms.

The total fixations in the experiments, in which an intense photospot light was substituted for one of the reflector floods for 50 minutes of the 1-hour steady-state period (Table II), indicate that a larger proportion of the algae were killed by the bright light in the presence of azaserine than in

<sup>7</sup> H. A. Krebs and P. D. Cohen, *Biochem. J.* 33, 1895 (1939).

<sup>8</sup> Della Rosa, Altman, and Salomon, *J. Biol. Chem.* 202, 771-9 (1953).

<sup>9</sup> Smith, Stamer, and Gunsalus, *Biochem. Biophys. Acta* 19, 567 (1956).

its absence. However, control and azaserine chromatograms, each carrying a total activity of  $1 \times 10^6$  counts/min, showed spots (Table II) with the same order of relative activities as those given in Table I.

An attempt to reverse the effects of azaserine inhibition with glutamine was only partially successful (Table III). Some increase in radioactivity in the lipid, phospholipid, area X, and glutamic acid was observed. The apparent continued build-up of glutamine can be explained by suggesting that while the rate of conversion of  $C^{14}$ -labeled precursors to glutamine remains almost the same, the introduction of inactive glutamine causes a dilution of radiocarbon in the glutamine pool, with a consequent lowering in the specific activity of the glutamine being converted to further products and hence an increase in the residual  $C^{14}$  glutamine observed. It is interesting to observe that the combined effect of azaserine and added inactive glutamine has caused almost a twentyfold increase of  $C^{14}$  glutamine.

Although azaserine caused a marked increase in  $CO_2$  fixation by Chlorella (Table IV), analysis of aliquots containing equal amounts of radioactivity showed that the radioactivity of most of the metabolites relative to one another remained the same.

This work was done under the auspices of the U. S. Atomic Energy Commission.