

UC Davis

UC Davis Previously Published Works

Title

Noninvasive Stereotactic Radiation for Refractory Ventricular Tachycardia After Failure of Cardiac Sympathetic Denervation.

Permalink

<https://escholarship.org/uc/item/07k2t3g5>

Journal

JACC. Case reports, 4(18)

ISSN

2666-0849

Authors

Bernstein, Hannah M
Leon, William
Daly, Megan E
et al.

Publication Date

2022-09-01

DOI

10.1016/j.jaccas.2022.06.025

Peer reviewed

CASE REPORT

ADVANCED

CLINICAL CASE

Noninvasive Stereotactic Radiation for Refractory Ventricular Tachycardia After Failure of Cardiac Sympathetic Denervation



Hannah M. Bernstein, MD, MPH,^a William Leon, MD,^a Megan E. Daly, MD,^b Peter Park, PhD,^b Ahmadreza Ghasemiesfe, MD,^c Neil Beri, MD,^a Maia Eng, MD,^a Uma N. Srivatsa, MBBS, MAS^a

ABSTRACT

Stereotactic body radiation therapy is a novel treatment option for refractory ventricular tachycardia. We present a case of ventricular tachycardia, with epicardial origin located in large inferior infarct scar, that recurred despite treatment with multiple antiarrhythmic drugs, catheter ablation, and cardiac sympathetic denervation. Stereotactic body radiation therapy safely and effectively terminated the arrhythmia. (**Level of Difficulty: Advanced.**) (J Am Coll Cardiol Case Rep 2022;4:1189-1194) © 2022 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

HISTORY OF PRESENTATION

A 75-year-old man presented with palpitations and shortness of breath. He was afebrile, with a pulse of 112 beats/min, blood pressure of 102/56 mm Hg, respiratory rate of 28 breaths/min, and oxygen saturation of 95% on room air. He was alert and responding to questions appropriately. Cardiac examination revealed tachycardia with regular rhythm, and jugular

venous distension was present; pulmonary examination revealed bilateral crackles. Electrocardiogram demonstrated wide-complex regular tachycardia (**Figure 1**). He was admitted to the intensive care unit. Home metoprolol and amiodarone were continued; mexiletine was discontinued and lidocaine infusion was started.

PAST MEDICAL HISTORY

Medical history includes diabetes, hypertension, coronary artery disease with prior inferior infarct status post-5-vessel coronary artery bypass graft and multiple percutaneous coronary interventions to his right coronary artery, and heart failure with reduced ejection fraction.

Three months before this presentation, the patient experienced multiple episodes of ventricular

LEARNING OBJECTIVES

- To understand different treatment options for refractory ventricular tachycardia.
- To review the role of stereotactic body radiation therapy as a new salvage therapy for refractory ventricular arrhythmia.

From the ^aDivision of Cardiovascular Medicine, University of California Davis Medical Center, Sacramento, California, USA; ^bDivision of Radiation Oncology, University of California Davis Medical Center, Sacramento, California, USA; and the ^cDivision of Radiology, University of California Davis Medical Center, Sacramento, California, USA.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

Manuscript received March 17, 2022; revised manuscript received June 17, 2022, accepted June 23, 2022.

ABBREVIATIONS AND ACRONYMS

4DCT = 4-dimensional computed tomograph

CMR = cardiac magnetic resonance

CSD = cardiac sympathetic denervation

ICD = implantable cardioverter-defibrillator

PTV = planning target volume

SBRT = stereotactic body radiation therapy

VT = ventricular tachycardia

tachycardia (VT) requiring antitachycardia pacing or shock (Figure 2). During a prior admission, coronary catheterization was performed that showed patent grafts not requiring intervention. Cardiac magnetic resonance (CMR) imaging demonstrated a large myocardial scar, consistent with prior infarct, involving the entire basal to apical inferior segments (Figure 3). Electrophysiology study identified a total of 5 separate hemodynamically well-tolerated, slow VTs in different areas of the inferior scar. Four were localized using entrainment activation mapping and successfully ablated (Figure 4). The

fifth, with a 500-ms cycle length, initially terminated but later recurred at 560 ms. Given morphology of the QRS complex and inability to ablate from endocardium, this VT was suspected to be epicardial in origin. Epicardial ablation was considered but would require a pericardial window, which was high risk because of prior coronary artery bypass graft and concern for bleeding from grafts caused by adhesions.

The patient underwent stellate ganglion block, which was well tolerated. He then underwent successful robotic sympathectomy of left T2 to T4. Right-sided sympathectomy was not performed because of large pleural plaque. He remained without VT for 72 hours and was discharged home 3 weeks before the current presentation.

DIFFERENTIAL DIAGNOSIS

Differential diagnosis for his wide-complex regular tachycardia included VT or supraventricular

FIGURE 2 Implantable Cardioverter-Defibrillator Interrogation Demonstrating Multiple Episodes of VT Requiring ATP or Shock

	VT	VF
Episodes	60	0
ATP Delivered	60	0
Shocks Delivered	3	0

ATP = antitachycardia pacing; VF = ventricular fibrillation; VT = ventricular tachycardia.

tachycardia with aberrancy. Given his history and prior electrocardiograms, his presentation was most consistent with VT.

INVESTIGATIONS

Electrolytes were within normal limits. Transthoracic echocardiography demonstrated stable global hypokinesia and akinesia of the inferior wall, with acute drop in ejection fraction from 35% to 20%.


MANAGEMENT

The patient's implantable cardioverter-defibrillator (ICD) was reprogrammed to have a lower VT detection zone with antitachycardia pacing therapy, which terminated his VT temporarily. Right heart catheterization demonstrated cardiogenic shock, with an index of 1.97 L/min/m² (Fick) and elevated filling pressures, which was managed with discontinuation of metoprolol, placement of an intra-aortic balloon

FIGURE 1 Electrocardiogram During Admission With Wide QRS Complex Monomorphic Tachycardia at a Rate of 110 Beats/Min




FIGURE 3 Late Gadolinium Enhancement Cardiac Magnetic Resonance Demonstrating Large Ischemic Scar With Corresponding Location in 17-Segment American Heart Association Cardiac Model




(A) Short axis of the mid left ventricle demonstrating ischemic subendocardial scar in segment 10. (B) Short axis of the left ventricle base demonstrating ischemic subendocardial scar in segments 3, 4, and 5. (C) Short axis of the left ventricle apex demonstrating ischemic subendocardial scar in segment 15. The **yellow arrows** show the ischemic scar.

FIGURE 4 Voltage Mapping Demonstrating Extensive Area of Scar



(A) Voltage map demonstrating extensive inferior scar (**red**) and healthy myocardium (**purple**) with adjacent border zones that were extensively ablated (**red tags**). **Blue tags** indicate areas of interest/fractionation. (B) Map of the patient's refractory source of ventricular tachycardia.



pump, and milrinone infusion. After diuresis, euvo-
lemia was confirmed with repeat right heart
catheterization. Telemetry demonstrated several epi-
sodes of VT despite multiple antiarrhythmic drug
therapies.


The patient was not a transplant candidate because
of his age. Because he continued to have intermittent
VT despite all available therapeutic options and car-
diomyopathy was rapidly progressing, it was decided
to attempt stereotactic body radiation therapy (SBRT)
for noninvasive cardiac radioablation. The off-label
nature of the therapy was discussed and, in con-
junction with radiation oncology, the decision
was made to proceed. Electroanatomic map images
were evaluated with 12-lead electrocardiogram cor-
relation of clinical VT. The source of VT was identified
as the mid to apical inferior wall, corresponding to the
location of scar on CMR.

For SBRT simulation, 10-phase respiratory gated
4-dimensional computerized tomography (4DCT)
was obtained with abdominal compression to limit
respiratory motion. CMR and the average intensity
projection from the 4DCT were fused in the treat-
ment planning system using rigid registration, and
segments for ablation were identified and con-
toured. Cardiac motion was assessed using the 10-
phase respiratory gated 4DCT to create an internal
target volume. An additional 5-mm planning target
volume (PTV) margin was added to account for daily
setup variability. The original target volume, inter-
nal target volume, and PTV resulted in 21.4, 29.9,
and 87.9 cm³, respectively (Figure 5). A total of
25 Gy in a single fraction was prescribed in accor-
dance with prior studies. The SBRT plan was
designed using 3 coplanar arcs using 6-MeV flat-
tening free-filter beams using RayStation treatment
planning software (RaySearch Laboratories), which
resulted in a highly conformal radiation plan with a
total delivery time of <4 minutes. Inverse plan
optimization was used to reduce the dose to healthy
myocardium, stomach, and bowel. The stomach was
constrained to a point maximum dose of <15 Gy
and <0.3 cm³ to 12 mL. The bowel was limited to a
point maximum dose of <15 Gy and, to meet normal
tissue constraints for the adjacent stomach, the PTV
coverage was 80%, receiving the prescription dose
of 25 Gy, with undercoverage of the portion of the
PTV directly abutting the stomach. Abdominal
compression was used throughout treatment, and
conebeam computed tomography was used for im-
age guidance immediately before treatment. ICD
interrogation postprocedure confirmed normal
function.

The patient tolerated the procedure without any
acute complications. For 3 days, he continued to have
nonsustained VT on telemetry. After 4 days, ven-
tricular ectopy burden significantly decreased.

DISCUSSION

Catheter ablation remains the mainstay treatment for
drug-refractory ventricular arrhythmias in patients
with structural heart disease; however, it is not
curative in many patients. Acute failure of VT abla-
tion can be seen in 10% of patients and long-term
recurrence in more than 50%, with epicardial origin
of VT identified as a major cause of ablation failure.¹
Epicardial ablation, percutaneously or via surgical
pericardial window, cannot always be performed in
patients with a history of cardiac surgery because of
the presence of pericardial adhesions and possible
damage to bypass grafts.



Autonomic modulation is an option for patients with persistent VT for whom pharmacologic and catheter ablation therapies have failed. Sympathectomy involves denervation of postganglionic sympathetic fibers from the stellate and T2 to T4 thoracic ganglia and is an effective treatment for refractory ventricular arrhythmia.² Cardiac sympathetic denervation (CSD) is included in the most recent American Heart Association/American College of Cardiology/Heart Rhythm Society guidelines as a reasonable therapy in selected patients with VT storm.³ Although CSD has been shown to reduce sustained VT and number of ICD shocks in approximately 50% of patients at a follow-up of 10 to 15 months, a cycle length of >400 ms and extensive scarring are associated with minimal response.^{2,4}

SBRT, commonly used to treat malignant tumors, uses highly focused radiation therapy and 3-dimensional imaging guidance. Cardiac side effects of radiation therapy have been observed after treatment of thoracic cancers, causing damage to arterial endothelium and premature coronary atherosclerosis, myocardial fibrosis, and conduction disturbances, usually 10 to 15 years after treatment and at doses of >40 Gy.⁵ Although myocyte cell death and subsequent fibrosis are expected years after exposure to chest radiation, decrease in VT has been seen nearly immediately after SBRT, suggesting additional mechanisms that are not yet entirely known.^{5,6} When used for the treatment of VT, radiation is usually delivered as single-fraction dose of 25 Gy, which is less than the required dosage

to induce late-stage cardiac fibrosis.⁶ As opposed to conventional catheter thermal ablation, where energy delivery to the epicardium or mid-myocardium may be limited by the depth of penetration, SBRT ablation targets transmural lesions. Histologic examination of explanted hearts of patients treated with SBRT did not demonstrate transmural fibrosis, further supporting the notion that a different mechanism is responsible for the reduction in VT seen at 25 Gy.⁶ Up-regulation of connexin 43 and cardiac sodium channel Na_v1.5 causing increased conduction velocity and QRS interval shortening have been suggested as the antiarrhythmic mechanism of cardiac SBRT at the treatment dose of 25 Gy.⁶

Noninvasive radioablation has been shown to be an effective and safe treatment for refractory VT, with markedly reduced VT burden in a series of 5 patients and prospective study of 17 patients.^{7,8} The effectiveness was noted after the initial blanking period to allow for inflammation from SBRT.⁷ This noninvasive procedure has been performed at varying levels of acuity, from the ambulatory setting to acutely ill patients on mechanical circulatory support.^{6,9} The most common side effects of SBRT are pericarditis and pneumonitis.⁸ The long-term effects are still not known because the first case series was published in 2017.⁷

Current recommendations for intractable ventricular arrhythmias in patients with reduced left ventricular function include consideration for heart transplantation and durable mechanical assist

devices or end-of-life conversations.¹⁰ SBRT represents an additional option for patients with refractory VT despite antiarrhythmic medications and who have undergone prior ablations. SBRT may be considered as a salvage therapy in patients with refractory VT; however, prospective trials and longer follow-up are still required. Because of its off-label use, current expert opinion recommends that use be restricted to patients with refractory VT and under the care of highly specialized interdisciplinary teams.⁹

FOLLOW-UP

At the 6-month follow-up, the patient was still free from VT recurrence and was without cardiac-related hospital admissions (Figure 6).

CONCLUSIONS

This case highlights the potential of SBRT as a salvage therapy in appropriately selected patients without other treatment options, with the benefit of radiation delivery to transmural lesions. After medical management, catheter ablation, and surgical left-sided CSD failed, focused cardiac radioablation was fast, painless, and effective in controlling VT. Careful long-term follow-up will be conducted.

ADDRESS FOR CORRESPONDENCE: Dr Hannah M. Bernstein, University of California Davis Medical Center, 4860 Y Street, Suite 2820, Sacramento, California 95817, USA. E-mail: hmb Bernstein@ucdavis.edu.

REFERENCES

1. Tokuda M, Kojodjojo P, Tung S, et al. Acute failure of catheter ablation for ventricular tachycardia due to structural heart disease: causes and significance. *J Am Heart Assoc.* 2013;2(3):e000072.
2. Vaseghi M, Barwad P, Malavassi Corrales FJ, et al. Cardiac sympathetic denervation for refractory ventricular arrhythmias. *J Am Coll Cardiol.* 2017;69(25):3070-3080. <https://doi.org/10.1016/j.jacc.2017.04.035>
3. Al-Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol.* 2018;72:e91-e220.
4. Murtaza G, Sharma SP, Akella K, et al. Role of cardiac sympathetic denervation in ventricular tachycardia: a meta-analysis. *Pacing Clin Electrophysiol.* 2020;43:828-837.
5. Curigliano G, Cardinale D, Dent S, et al. Cardiotoxicity of anticancer treatments: epidemiology, detection, and management. *CA Cancer J Clin.* 2016;66:309-325. <https://doi.org/10.3322/caac.21341>
6. Hohmann S, Hillmann HAK, Müller-Leisse J, et al. Stereotactic radioablation for ventricular tachycardia. *Herzschr Elektrophys.* 2022;33(1):49-54. <https://doi.org/10.1007/s00399-021-00830-y>
7. Cuculich PS, Schill MR, Kashani R, et al. Noninvasive cardiac radiation for ablation of ventricular tachycardia. *N Engl J Med.* 2017;377:2325-2336.
8. Robinson CG, Samson PP, Moore KM, et al. Phase I/II trial of electrophysiology-guided noninvasive cardiac radioablation for ventricular tachycardia. *Circulation.* 2018;139:313-321.
9. Krug D, Blanck O, Andratschke N, et al. Recommendations regarding cardiac stereotactic body radiotherapy for treatment refractory ventricular tachycardia. *Heart Rhythm.* 2021 18;(12):2137-2145.
10. Santangeli P, Rame JE, Birati EY, Marchlinski FE. Management of ventricular arrhythmias in patients with advanced heart failure. *J Am Coll Cardiol.* 2017;69:1842-1860.

KEY WORDS ablation, electrophysiology, stereotactic body radiation therapy, ventricular tachycardia