UCLA Proceedings of the UCLA Department of Medicine

Title

Acute Abdominal Pain in the Elderly

Permalink https://escholarship.org/uc/item/07z0m01v

Journal Proceedings of the UCLA Department of Medicine, 17(1)

Author Quinn, Roswell

Publication Date

2013-09-24

CLINICAL VIGNETTE

Acute Abdominal Pain in the Elderly

Roswell Quinn, MD PhD

Case Report

An 86-year old woman presented to the emergency room with dyspnea on exertion and fatigue. She had a history of congestive heart failure with preserved ejection fraction, atrial fibrillation on anticoagulation, hypertension, and prior transient ischemic attack. Home medications include metoprolol, warfarin, furosemide, and lisinopril. She frequently forgets her home medications and has not been taking her prescribed diuretic for over a year.

The patient reported 1 week of symptoms. She described being able to walk to the mailbox at baseline but was no longer able to do this and was unable to climb a flight of stairs. She became short of breath walking from room to room. Her exercise tolerance has declined over several years. The dyspnea is associated with dry cough, weakness, 5 pound weight gain, anorexia, and two loose stools. She had an episode of back pain located between her scapula 1 week ago that resolve spontaneously.

In the Emergency Room, the patient's vitals included: temperature 37.4, heart rate from 83 to 164, blood pressure 125/68, oxygen saturations 100% on room air with a respiratory rate of 16. Physical exam was notable for fine bibasilar crackles without increased work of breathing. JVP was elevated 9cm above the sternal angle. Heart sounds were normal with tachycardia and irregular rate, and there was trace-pitting edema of the bilateral lower extremities. Abdominal exam was benign. Laboratory testing included white blood cells 18K, hemoglobin 11.9, sodium 133, potassium 3.5, chloride 100, bicarbonate 20, BUN 32 creatinine 1.1, INR 2.6, BNP 638, troponin I negative. She received 1L on normal saline, 5mg of IV metoprolol times three, and underwent a CT angiogram of the chest that was negative for consolidation, pulmonary embolism, or effusions.

After admission to the internal medicine service, the patient received intravenous furosemide along with her home metoprolol, lisinopril, and warfarin. She was seen by infectious diseases and ACS work was negative. The morning following admission the patient reported to the team that her prior back pain radiated to the RUQ. Abdominal ultrasound revealed a perforated gallbladder. The patient's heart failure was medically optimized and an uncomplicated cholecystectomy was performed prior to discharge.

Discussion

Acute abdominal pain in older adults can present a diagnostic and therapeutic challenge. Physiologic changes with age can predispose individuals to abdominal pathology, such a biliary disease¹. Yet, older adults often have nonspecific symptomatology, present for evaluation later, have complex medical histories and coexisting disease, and can be poor surgical As a result, when compared to candidates. patients under 65 years old, those greater than 65 with acute abdominal pain have increased concurrent illnesses, longer hospital stays, more misdiagnosis, greater delays to surgery². This creates an unfortunate situation in which diagnostic accuracy is low, and mortality is high in this special population³.

Acute abdomen pain refers to new abdominal pain that is less than seven days old. The differential diagnosis of an acute abdomen remains broad in older adults. Along with common causes of an acute abdomen (cholecystitis, appendicitis, small bowel obstruction, pancreatitis, peptic ulcer disease, etc.) there are age specific considerations that require attention. These include conditions associated with age-related pathophysiology such as mesenteric ischemia, large bowel obstruction, diverticulosis, and aortic abdominal aneurysm. In addition, illnesses prevalent in the elderly - urinary tract infection, myocardial infection, or pneumonia - can present with abdominal pain.

Acute abdominal pain in the elderly can be challenging to diagnose. Patients frequently have atypical, seemingly benign, or even asymptomatic presentations. Research has this for bowel perforation, demonstrated cholecystitis, intra-abdominal appendicitis, infections, and peptic ulcer disease⁴⁻⁸. Morphine, so frequently used to treat abdominal pain in emergency rooms, can mask physical exam findings, specifically rigidity, in older adults with surgical abdomens⁹. One researcher has concluded that the physical exam could not reliably predict or exclude significant disease in older patients with acute abdominal pain¹⁰.

Initial diagnostic testing also fails to establish clear diagnoses in older patients with abdominal pain. White count, hemoglobin, liver function tests, and even lactate have not differentiated between surgical and non-surgical disease in inpatients when geriatric examined retrospectively¹¹. Unfortunately. newer diagnostic and surgical approaches have not improved outcomes in this population.¹² Misdiagnoses in this high-risk population have unfortunate outcomes. One small study found the mortality rate of patients over 80 years old with

KEY POINTS

Acute Abdominal Pain refers to new pain less than 7 days old.
In patients over 65 the differential diagnosis should be expanded to include age-related abdominal pathologies and common nonabdominal pathologies.
Older adults present later, have atypical presentations, and are more frequently misdiagnosed.
Delays to surgery, length of stay, morbidity and mortality are higher in older adults with acute abdominal pain. acute abdominal pain who were misdiagnosed was 59%¹³. A similar study found the mortality rate to be three times greater in misdiagnosed older adults when compared to their younger counterparts¹⁴.

The above differences in presentation and diagnosis of older adults with acute abdominal pain place this group at high risk for poor outcomes. When compared to individuals under 65 years old, older patients with abdominal pain can have double the time from admission to surgery³. Delays in presentation, diagnosis, and intervention -as seen in our case- confer increased morbidity. For example, a greater number of perforations have been seen in older cohorts with appendicitis¹⁵. When surgical interventions do occur earlier, outcomes remain poor. Emergent laparotomy in patients over 80 have high levels of morbidity (70% with life threatening post-operative complications) and mortality $(45\%)^{16}$. These unfortunate outcomes warrant a high level of suspicion and diagnostic acumen in adults greater than 65 years of age presenting with new abdominal pain that is less than last seven days in duration.

REFERENCES

- Lyon C, Clark DC. Diagnosis of acute abdominal pain in older patients. *Am Fam Physician*. 2006 Nov 1;74(9):1537-44. Review. PubMed PMID: 17111893.
- de Dombal FT. Acute abdominal pain in the elderly. J Clin Gastroenterol. 1994 Dec;19(4):331-5. Review. PubMed PMID: 7876517.
- Laurell H, Hansson LE, Gunnarsson U. Acute abdominal pain among elderly patients. *Gerontology*. 2006;52(6):339-44. Epub 2006 Aug 11. PubMed PMID:16905885.
- Brady P. Atypical presentation of postoperative faecal peritonitis in an elderly patient. *BMJ Case Rep.* 2013 Aug 9;2013. doi:pii: bcr2013009030. 10.1136/bcr-2013-009030. PubMed PMID: 23933858.
- Hilton D, Iman N, Burke GJ, Moore A, O'Mara G, Signorini D, Lyons D, Banerjee AK, Clinch D. Absence of abdominal pain in older persons with endoscopic ulcers: a prospective study. *Am J Gastroenterol.* 2001 Feb;96(2):380-4. PubMed PMID: 11232679.
- Elangovan S. Clinical and laboratory findings in acute appendicitis in the elderly. J Am Board Fam Pract. 1996 Mar-Apr;9(2):75-8. PubMed PMID: 8659268.
- Cooper GS, Shlaes DM, Salata RA. Intraabdominal infection: differences in presentation and outcome between younger patients and the elderly. *Clin Infect Dis.* 1994 Jul; 19(1):146-8. PubMed PMID: 7948517.
- Parker LJ, Vukov LF, Wollan PC. Emergency department evaluation of geriatric patients with acute cholecystitis. *Acad Emerg Med.* 1997 Jan;4(1):51-5. PubMed PMID: 9110012.
- Güngör F, Kartal M, Bektaş F, Söyüncü S, Yiğit O, Mesci A. Randomized controlled trial of morphine in

elderly patients with acute abdominal pain. *Ulus Travma Acil Cerrahi Derg*. 2012 Sep;18(5):397-404. doi: 10.5505/tjtes.2012.62534. PubMed PMID: 23188600.

- Marco CA, Schoenfeld CN, Keyl PM, Menkes ED, Doehring MC. Abdominal pain in geriatric emergency patients: variables associated with adverse outcomes. *Acad Emerg Med*. 1998 Dec;5(12):1163-8. PubMed PMID: 9864129.
- 11. **Parker JS, Vukov LF, Wollan PC**. Abdominal pain in the elderly: use of temperature and laboratory testing to screen for surgical disease. *Fam Med*. 1996 Mar;28(3):193-7. PubMed PMID: 8900552.
- Hui TT, Major KM, Avital I, Hiatt JR, Margulies DR. Outcome of elderly patients with appendicitis: effect of computed tomography and laparoscopy. *Arch Surg.* 2002 Sep;137(9):995-8; discussion 999-1000. PubMed PMID: 12215147.
- van Duren BH, Moghul M, Appleton SG, van Boxel GI. The acute abdomen. *BMJ*. 2013 Apr 24;346:f2549. doi: 10.1136/bmj.f2549. PubMed PMID: 23616034.
- Kizer KW, Vassar MJ. Emergency department diagnosis of abdominal disorders in the elderly. Am J Emerg Med. 1998 Jul;16(4):357-62. PubMed PMID: 9672450.
- Kraemer M, Franke C, Ohmann C, Yang Q; Acute Abdominal Pain Study Group. Acute appendicitis in late adulthood: incidence, presentation, and outcome. Results of a prospective multicenter acute abdominal pain study and a review of the literature. *Langenbecks Arch Surg.* 2000 Nov;385(7):470-81. Review. PubMed PMID: 11131250.
- Green G, Shaikh I, Fernandes R, Wegstapel H. Emergency laparotomy in octogenarians: A 5-year study of morbidity and mortality. World J Gastrointest Surg. 2013 Jul 27;5(7):216-21. doi: 10.4240/wjgs.v5.i7.216. PubMed PMID:23894689; PubMed Central PMCID: PMC3715657.

Submitted on September 24, 2013