

UCLA

UCLA Previously Published Works

Title

Genetic Basis for Clinical Response to CTLA-4 Blockade in Melanoma

Permalink

<https://escholarship.org/uc/item/0815417g>

Journal

New England Journal of Medicine, 371(23)

ISSN

0028-4793

Authors

Snyder, Alexandra
Makarov, Vladimir
Merghoub, Taha
[et al.](#)

Publication Date

2014-12-04

DOI

10.1056/nejmoa1406498

Peer reviewed

ORIGINAL ARTICLE

Genetic Basis for Clinical Response to CTLA-4 Blockade in Melanoma

Alexandra Snyder, M.D., Vladimir Makarov, M.D., Taha Merghoub, Ph.D., Jianda Yuan, M.D., Ph.D., Jesse M. Zaretsky, B.S., Alexis Desrichard, Ph.D., Logan A. Walsh, Ph.D., Michael A. Postow, M.D., Phillip Wong, Ph.D., Teresa S. Ho, B.S., Travis J. Hollmann, M.D., Ph.D., Cameron Bruggeman, M.A., Kasthuri Kannan, Ph.D., Yanyun Li, M.D., Ph.D., Ceyhan Elipenahli, B.S., Cailian Liu, M.D., Christopher T. Harbison, Ph.D., Lisu Wang, M.D., Antoni Ribas, M.D., Ph.D., Jedd D. Wolchok, M.D., Ph.D., and Timothy A. Chan, M.D., Ph.D.

ABSTRACT

BACKGROUND

Immune checkpoint inhibitors are effective cancer treatments, but molecular determinants of clinical benefit are unknown. Ipilimumab and tremelimumab are antibodies against cytotoxic T-lymphocyte antigen 4 (CTLA-4). Anti-CTLA-4 treatment prolongs overall survival in patients with melanoma. CTLA-4 blockade activates T cells and enables them to destroy tumor cells.

METHODS

We obtained tumor tissue from patients with melanoma who were treated with ipilimumab or tremelimumab. Whole-exome sequencing was performed on tumors and matched blood samples. Somatic mutations and candidate neoantigens generated from these mutations were characterized. Neoantigen peptides were tested for the ability to activate lymphocytes from ipilimumab-treated patients.

RESULTS

Malignant melanoma exomes from 64 patients treated with CTLA-4 blockade were characterized with the use of massively parallel sequencing. A discovery set consisted of 11 patients who derived a long-term clinical benefit and 14 patients who derived a minimal benefit or no benefit. Mutational load was associated with the degree of clinical benefit ($P=0.01$) but alone was not sufficient to predict benefit. Using genomewide somatic neopeptide analysis and patient-specific HLA typing, we identified candidate tumor neoantigens for each patient. We elucidated a neoantigen landscape that is specifically present in tumors with a strong response to CTLA-4 blockade. We validated this signature in a second set of 39 patients with melanoma who were treated with anti-CTLA-4 antibodies. Predicted neoantigens activated T cells from the patients treated with ipilimumab.

CONCLUSIONS

These findings define a genetic basis for benefit from CTLA-4 blockade in melanoma and provide a rationale for examining exomes of patients for whom anti-CTLA-4 agents are being considered. (Funded by the Frederick Adler Fund and others.)

From the Department of Medicine (A.S., T.M., M.A.P., J.D.W.), Human Oncology and Pathogenesis Program (A.S., V.M., A.D., L.A.W., K.K., T.A.C.), Swim across America–Ludwig Collaborative Research Laboratory (T.M., Y.L., C.E., C.L., J.D.W.), Department of Radiation Oncology (T.A.C.), Department of Pathology (T.J.H.), and Immunology Program, Ludwig Center for Cancer Immunotherapy (J.Y., P.W., T.S.H., J.D.W.), Memorial Sloan Kettering Cancer Center; Weill Cornell Medical College (A.S., M.A.P., J.D.W., T.A.C.); and Department of Mathematics, Columbia University (C.B.) — all in New York; the Department of Molecular and Medical Pharmacology (J.M.Z., A.R.) and the Department of Medicine, Division of Hematology–Oncology, Jonsson Comprehensive Cancer Center (A.R.), University of California, Los Angeles, Los Angeles; and Bristol-Myers Squibb, Princeton, NJ (C.T.H., L.W.). Address reprint requests to Dr. Chan at the Human Oncology and Pathogenesis Program, Memorial Sloan Kettering Cancer Center, 1275 York Ave., Box 20, New York, NY 10065, or at chant@mskcc.org; or to Dr. Wolchok at the Ludwig Center for Cancer Immunotherapy, Memorial Sloan Kettering Cancer Center, 1275 York Ave., New York, NY 10065, or at wolchokj@mskcc.org.

Drs. Snyder, Makarov, Merghoub, and Yuan and Drs. Wolchok and Chan contributed equally to this article.

This article was published on November 19, 2014, and updated on November 21, 2014, at NEJM.org.

N Engl J Med 2014;371:2189-99.
DOI: 10.1056/NEJMoa1406498

Copyright © 2014 Massachusetts Medical Society.

IMMUNE CHECKPOINT BLOCKADE HAS LED to durable antitumor effects in patients with metastatic melanoma, non–small-cell lung cancer, and other tumor types, but the factors determining whether a patient will have a response remain elusive.^{1,2} The fully human monoclonal antibodies ipilimumab and tremelimumab block cytotoxic T-lymphocyte antigen 4 (CTLA-4), resulting in T-cell activation. Some studies have established correlations between outcomes with ipilimumab and peripheral-blood lymphocyte count, markers of T-cell activation,³ an “inflammatory” microenvironment,^{4,5} and maintenance of high-frequency T-cell receptor clonotypes.⁶

The relationship among the genomic landscape of the tumor, the mutational load, and the benefit from treatment remains obscure. The immunogenicity resulting from nonsynonymous melanoma mutations has been shown in a mouse model,⁷ and the antigenic diversity of human melanoma tumors has been modeled in silico⁸ and in melanoma-specific CD8 T-cell responses after treatment with ipilimumab.⁹ Effector and helper T-cell function and regulatory T-cell depletion are necessary for the efficacy of CTLA-4 blockade,¹⁰ but there is not an association between a specific HLA type and a clinical benefit.¹¹ Melanomas have very high mutational burdens (0.5 to >100 mutations per megabase) as compared with other solid tumors.¹² Elegant studies have shown that somatic mutations can give rise to neoepitopes¹³ and that these may serve as neoantigens.^{14–16} We conducted a study to determine whether the genetic landscape of a tumor affects the clinical benefit provided by CTLA-4 blocking agents.

METHODS

SAMPLE ACQUISITION AND DNA PREPARATION

For the discovery set, we conducted whole-exome sequencing of DNA from tumors and matched normal blood from 25 ipilimumab-treated patients. A validation set included an additional 39 patients, of whom 5 were treated with tremelimumab. Primary tumor samples and matched normal peripheral-blood specimens were obtained after the patients had provided written informed consent. DNA was extracted, and exon capture was performed with the use of the SureSelect Human All Exon 50-Mb kit (Agilent Technologies). Enriched

exome libraries were sequenced on the HiSeq 2000 platform (Illumina) to provide a mean exome coverage of more than 100× (Memorial Sloan Kettering Cancer Center Genomics Core and Broad Institute).

IMMUNOGENICITY ANALYSIS OF SOMATIC MUTATIONS

We created a bioinformatic tool to translate all mutations in exomes and then evaluate binding with major histocompatibility complex (MHC) class I molecules. The neoantigen signature was generated from the nonamers containing four amino acid strings of peptides that are common to tumors from patients with a long-term benefit from therapy. Details are provided in the Supplementary Appendix, available with the full text of this article at NEJM.org.

INTRACELLULAR CYTOKINE STAINING

Candidate neoantigen peptides were synthesized (GenScript), cultured with autologous peripheral-blood mononuclear cells (PBMCs), and then analyzed by means of intracellular cytokine staining for interleukin-2, CD107a, macrophage inflammatory protein 1 β , tumor necrosis factor α , and interferon- γ on restimulation of cells with the candidate peptides.

STATISTICAL ANALYSIS

The Mann–Whitney test was used to compare mutational loads, and the log-rank test was used to compare Kaplan–Meier curves. The statistical methods used in the study are more fully described in the Supplementary Appendix.

RESULTS

MUTATIONAL LANDSCAPE OF MELANOMAS FROM THE STUDY PATIENTS


Baseline patient characteristics are shown in Table 1 (for more detailed information, see Tables S1 and S2 in the Supplementary Appendix). The study involved patients with and those without a long-term clinical benefit from therapy (CTLA-4 blockade alone or CTLA-4 blockade with resection of an isolated stable or nonresponding lesion). A long-term clinical benefit was defined by radiographic evidence of freedom from disease or evidence of a stable or decreased volume of disease for more than 6 months. Lack of a long-term ben-

Table 1. Clinical Characteristics of the Patients in the Discovery and Validation Sets, According to Clinical Benefit from Therapy.

Characteristic	Discovery Set		Validation Set	
	Long-Term Benefit (N=11)	Minimal or No Benefit (N=14)	Long-Term Benefit (N=25)	Minimal or No Benefit (N=14)
Age at start of treatment — yr				
Median	63	60	66	57
Range	39–70	48–79	33–90	18–74
Sex — no. of patients (%)				
Female	3 (27)	8 (57)	9 (36)	5 (36)
Male	8 (73)	6 (43)	16 (64)	9 (64)
Disease origin — no. of patients (%)				
Acral	0	3 (21)	1 (4)	1 (7)
Uveal	0	0	1 (4)	0
Cutaneous	10 (91)	8 (57)	15 (60)	11 (79)
Unknown primary	1 (9)	3 (21)	3 (12)	0
Not available	0	0	5 (20)	2 (14)
<i>BRAF</i> or <i>NRAS</i> mutation — no. of patients (%)				
No	1 (9)	6 (43)	17 (68)	11 (79)
Yes	10 (91)	8 (57)	8 (32)	3 (21)
Lactate dehydrogenase level at start of therapy — no. of patients (%)				
Normal	8 (73)	8 (57)	8 (32)	9 (64)
Above normal	2 (18)	5 (36)	3 (12)	3 (21)
Not available	1 (9)	1 (7)	14 (56)	2 (14)
Duration of response to therapy — wk				
Median	59	14	130	11
Range	42–361	11–23	64–376	3–29
Previous therapies — no.*				
Median	1	1	0	0
Range	0–3	0–2	0–2	0–3
Melanoma stage at time of diagnosis — no. of patients (%)				
IIIC	0	0	3 (12)	0
M1a	0	1 (7)	4 (16)	1 (7)
M1b	5 (45)	1 (7)	2 (8)	3 (21)
M1c	6 (55)	12 (86)	16 (64)	10 (71)
Overall survival — yr†				
Median	4.4	0.9	3.3	0.8
Range	2.0–6.9	0.4–2.7	1.6–7.2	0.2–2.1

* Previous therapies included interleukin-2 and cytotoxic chemotherapy.

† Overall survival was calculated from the date of the first dose of ipilimumab to the date of death or censoring of data.



efit was defined by tumor growth on every computed tomographic scan after the initiation of treatment (no benefit) or a clinical benefit lasting 6 months or less (minimal benefit). Representative scans are shown in Figure 1, and Figure S1 in the Supplementary Appendix.

To determine the genetic features associated with a sustained benefit from CTLA-4 blockade, we analyzed DNA in tumor and matched blood samples using whole-exome sequencing. In the discovery set, we generated 6.4 Gb of mapped sequence, with more than 99% of the target sequence covered to at least 10× depth and a mean exome coverage of 103× (Table S3 and Fig. S2 in

the Supplementary Appendix). The wide ranges of mutational burdens (Fig. 2A, and Table S3 in the Supplementary Appendix) and recurrent and driver mutations (Fig. S2C and S2D and Table S4 in the Supplementary Appendix) among samples were consistent with previously reported findings.¹⁷⁻¹⁹ The ratio of transitions to transversions (Fig. S2E in the Supplementary Appendix) and the frequency of nucleotide changes (Fig. S2F in the Supplementary Appendix) were similar in the discovery and validation sets.¹² No gene was universally mutated across patients with a sustained benefit.

ASSOCIATION BETWEEN MUTATIONAL BURDEN AND CLINICAL BENEFIT

We hypothesized that an increased mutational burden in metastatic melanoma samples would correlate with a benefit from CTLA-4 blockade. There was a significant difference in mutational load between patients with a long-term clinical benefit and those with a minimal benefit or no benefit, both in the discovery set ($P=0.01$ by the Mann-Whitney test) and in the validation set ($P=0.009$ by the Mann-Whitney test) (Fig. 2A, and Table S5 in the Supplementary Appendix). In the discovery set, a high mutational load was significantly correlated with improved overall survival ($P=0.04$ by the log-rank test) (Fig. 2B), and there was a trend toward improved survival in the validation set (Fig. S3A in the Supplementary Appendix). The latter set included eight patients with nonresponding tumors who otherwise had systemic disease control, which may confound the relationship between mutational load and survival. Further subdivision into four clinical categories was suggestive of a dose-response relationship in the discovery set (Fig. S3B in the Supplementary Appendix). These data indicate that a high mutational load correlates with a sustained clinical benefit from CTLA-4 blockade but that a high load alone is not sufficient to impart a clinical benefit, because there were tumors with a high mutational burden that did not respond to therapy.

SOMATIC NEOEPITOPES IN RESPONDING TUMORS AND EFFICACY OF CTLA-4 BLOCKADE


MHC class I presentation and cytotoxic T-cell recognition are required for ipilimumab activity.¹⁰ Because mutational load alone did not explain a clinical benefit from CTLA-4 blockade, we hypoth-

esized that the presence of specific tumor neoantigens might explain the varied therapeutic benefit. To identify these neoepitopes, we developed a bioinformatic pipeline incorporating prediction of MHC class I binding, modeling of T-cell receptor binding, patient-specific HLA type, and epitope-homology analysis (see the Methods section and Fig. S4 in the Supplementary Appendix).

We created a computational algorithm, called NAsseek, to translate all nonsynonymous missense mutations into mutant and nonmutant peptides (see the Methods section and Fig. S4 in the Supplementary Appendix). We examined whether a subgroup of somatic neoepitopes would alter the strength of peptide–MHC binding, using patient-specific HLA types (Table S3 in the Supplementary Appendix). We first compared the overall antigenicity trend of all mutant versus nonmutant peptides. In aggregate, the mutant peptides were predicted to bind MHC class I molecules with higher affinity than the corresponding nonmutant peptides (Fig. S5 in the Supplementary Appendix).

Using only peptides predicted to bind to MHC class I molecules (binding affinity, ≤ 500 nM), we searched for conserved stretches of amino acids shared by multiple tumors. We used standard methods of machine learning, hierarchical clustering, and signature derivation to identify consensus sequences (see the Methods section in the Supplementary Appendix).²⁰ We identified a number of tetrapeptide sequences that were shared by patients with a long-term clinical benefit but completely absent in patients with a minimal benefit or no benefit (Fig. 3A and 3B, and Table S6 in the Supplementary Appendix). It has been shown that short amino acid substrings comprise conserved regions across antigens recognized by a T-cell receptor.²¹ In these experiments, recognition of epitopes was driven by consensus tetrapeptides within the immunogenic peptides, and tetrapeptides within cross-reacting T-cell receptor epitopes were necessary and sufficient to drive T-cell proliferation, findings that are consistent with evidence that this polypeptide length can drive recognition by T-cell receptors.²² Tetrapeptides are used to model genome phylogeny because they occur relatively infrequently in proteins and typically reflect function.²³

We used the discovery set to generate a predictive signature from the candidate neoepitopes



(see the Methods section in the Supplementary Appendix). We found that the tetrapeptides common to each group (candidate neoepitopes) in-




Figure 3. Association of a Neopeptide Signature with a Clinical Benefit from CTLA-4 Blockade.

Candidate neopeptides were identified by means of mutational analysis, as described in the Methods section in the Supplementary Appendix. Panel A shows a heat map of candidate tetrapeptide neoantigens that were present in patients with a long-term clinical benefit but absent in patients with a minimal benefit or no benefit in the discovery set (comprising 25 patients). Each row represents a neopeptide; each column represents a patient. The vertical red line indicates the tetrapeptide signature associated with a response to blockade of cytotoxic T-lymphocyte antigen 4 (CTLA-4). The exact tetrapeptides, chromosomal loci, and non-mutant and mutant nonamers in which they occur are listed in Table S6 in the Supplementary Appendix. Panel B shows the same information for the validation set (comprising 39 patients). Panel C shows the Kaplan–Meier curves for overall survival in the discovery set for patients with the signature and those without the signature. Panel D shows the same data for the validation set.

cluded 101 shared exclusively among patients in the discovery set who had a long-term clinical benefit; this was also independently observed in the validation set (Fig. 3A and 3B, and Tables S6 and S7 in the Supplementary Appendix). This set of neopeptides defines a signature linked to a benefit from CTLA-4 blockade. Because of the size of our discovery set, we cannot exclude the possibility that additional biologically relevant epitopes exist and conversely that there are biologically relevant epitopes that were predicted bioinformatically but were not expressed or presented in patients with a minimal benefit or no benefit (Tables S7A and S7B in the Supplementary Appendix).

Shared tetrapeptide neopeptides did not simply result from a high mutational load. For example, in the discovery set, the patient with a minimal benefit or no benefit who had the greatest number of mutations (Patient SD7357, who had 1028 mutations) did not share any of the tetrapeptide signatures. This concept was illustrated again in the validation set, in which even tumors from patients with more than 1000 mutations (Patients NR9521 and NR4631) did not respond (Table S3 in the Supplementary Appendix). Simulation testing with five different models showed that the association between the neopeptide signature and a long-term clinical

benefit was highly significant and was unlikely to have resulted from chance alone ($P < 0.001$ for four methods and $P = 0.002$ for a fifth method) (Fig. S6 in the Supplementary Appendix). A high mutational load appeared to increase the probability, but not guarantee formation, of a neoepitope signature associated with a benefit. Consensus analysis revealed that the neoepitopes were not random. The frequencies of amino acids that made up the tetrapeptides in the group of patients with a long-term clinical benefit were different from those observed in the group with a minimal benefit or no benefit (Fig. S7A in the Supplementary Appendix).

The neoepitope signature derived from the discovery set correlated strongly with survival in both the discovery and validation sets ($P < 0.001$ for both comparisons by the log-rank test) (Fig. 3C and 3D). The correlation between mutational load and survival was not as strong (Fig. 2B, and Fig. S3A in the Supplementary Appendix).

The shared tetrapeptides were encoded by mutations in diverse genes across the genome (Fig. S7B and Table S6 in the Supplementary Appendix). Using RNA-sequencing data from the Cancer Genome Atlas, we confirmed that the genes harboring our somatic neoepitopes were widely expressed in melanoma (Table S8 in the Supplementary Appendix). In some cases, the amino acid change resulting from the somatic mutation led to a change in the tetrapeptide itself. In others, the mutant amino acid was separate from the tetrapeptide and altered MHC binding, as has been described previously.²⁴⁻²⁶

In addition, candidate neoepitopes common to both clinical groups were analyzed with the use of the Immune Epitope Database (www.iedb.org). This is the most comprehensive database of experimentally validated, published, and curated antigens, and it has been used to develop algorithms to identify antigens with high accuracy.¹⁴ The candidate neoepitopes common to patients with a long-term clinical benefit were homologous to many more viral and bacterial antigens in the database than were the neoepitopes common to patients with a minimal benefit or no benefit (Table S9 in the Supplementary Appendix). For example, the tetrapeptide substring BSSA was shared by patients with a long-term clinical benefit (Fig. 4A) and corresponds to the


precise antigenic portion of human cytomegalovirus immediate early epitope (MESSAKRKMDP-DNPD).²⁷ These data suggest that the neoepitopes in patients with strong clinical benefit from CTLA-4 blockade may resemble epitopes from pathogens that T cells are likely to recognize. The cross-reactive peptides defined by short peptide consensus sequences that were discovered by Birnbaum et al. with the use of an unbiased screen also had substantial homology to antigens in microbes.²¹ Although tantalizing, these observations will require further study to confirm.

Using a whole-exome sequencing approach, we characterized the predicted antigenic peptide space (see the Methods section in the Supplementary Appendix). As further validation of our study, we reidentified melanoma antigen recognized by T cells (MART-1, also known as MelanA), an experimentally validated melanocytic antigen (Fig. S8).²⁸ EKLS, which comprises the core amino acids of the MART-1 MHC class II epitope, was shared by patients with a long-term clinical benefit, and the phosphoserine moiety is critical for T-cell receptor recognition.²⁹ The frequency of leukocyte common antigen–positive cells and ratio of CD8–positive cells to FOXP3–positive cells were substantially different between patients with a long-term clinical benefit from ipilimumab and those with a minimal benefit or no benefit (Fig. S9 in the Supplementary Appendix).

IN VITRO VALIDATION OF PREDICTED IMMUNOGENIC PEPTIDES

Translation of next-generation sequencing into in vitro validation of peptide predictions has proven challenging, even in expert hands, with very low published validation rates.¹⁵ In vitro assays are hampered by the paucity of clinical samples, the sensitivity of preserved cells to the freeze–thaw process, the low frequency of anti-neoantigen T cells in clinical samples, and the very low sensitivity of T cells in vitro in the absence of the complex in vivo immunogenic microenvironment.

We attempted to optimize prediction by integrating multiple high-throughput approaches (Fig. S4 in the Supplementary Appendix). On the basis of our prediction algorithm, we generated pools of peptides and performed assays of T-cell activation for patients for whom we had suffi-



cient lymphocytes (see the Methods section in the Supplementary Appendix). Positives pools were observed for three of five patients (Fig. S10A, S10B, and S10C in the Supplementary Appendix). We identified the exact peptides for patients with adequate PBMCs. We found a polyfunctional T-cell response to the peptide TESPFEQHI in Patient CR9306 (Fig. S10D in the Supplementary Appendix) but not to its nonmutant counterpart, TKSPFEQHI. This response peaked at 60 weeks after the initiation of treatment (Fig. 4B). T-cell responses were absent in healthy donors (Fig. S10E in the Supplementary Appendix). TESPFEQHI had a predicted MHC class I affinity for B4402 of 472 nM, as compared with 18323 nM for TKSPFEQHI. ESPF is a common tetrapeptide found in the response signature and is a substrate (positions 176 through 179) of the hepatitis D virus large delta epitope p27 (PESPFA and ESPFAR).³⁰ TESPFEQHI results from a mutation in *FAM3C* (c.A577G;p.K193E), a gene highly expressed in melanoma (Table S8 in the Supplementary Appendix).

We also found that peptide GLEREGFTF elicited a polyfunctional T-cell response in Patient CR0095 (Fig. 4C, and Fig. S10F in the Supplementary Appendix), whereas nonmutant GLERGGFTF did not. This response peaked at 24 weeks after the initiation of treatment (Fig. 4C). GLEREGFTF arises from a mutation in *CSMD1* (c.G10337A;p.G3446E), which is also highly expressed in melanoma (Table S8 in the Supplementary Appendix), and the peptide has 80% homology to a known *Burkholderia pseudomallei* antigen (Immune Epitope Database Reference ID: 1027043). The lack of T-cell activation may not rule out a given neoantigen because in vitro assays are limited in sensitivity, as described above.

DISCUSSION

Anti-CTLA-4 and anti-programmed cell death 1 antibodies have resulted in long-term disease control in a subgroup of patients with melanoma.^{1,2} Here, we have illustrated the importance of tumor genetics in defining the basis of the clinical benefit from CTLA-4 blockade.

Our observations suggest a number of principles relevant to immunotherapy for cancer.

Although a high mutational load is associated with a benefit from immune checkpoint abrogation, this factor alone is not sufficient to impart a clinical benefit. Rather, there are somatic neoepitopes that are shared by patients with a prolonged benefit and are absent in those without a prolonged benefit. Owing to somatic mutations, a subset of proteins present in the tumor becomes recognized by the immune system as nonself, given their novelty in the tumor context.^{8,14,31,32} These concepts were formulated in the discovery set and confirmed in the validation set.

It is well known in the field of infectious diseases that an individual amino acid within a peptide can affect immunogenicity by altering peptide-MHC or peptide-T-cell receptor interactions.^{33,34} In cancers, the altered amino acid residue resulting from a single missense mutation can create a T-cell epitope from a previously self peptide.^{31,32,35} In the patients described here, altered amino acids resulting from tumor mutations caused the tumors to display somatic neoepitopes that elicited an antitumor response augmented by CTLA-4 blockade.

Our study has limitations. Although large for a genomic study (128 exomes), our sample size was limited, patients had received a variety of previous treatments, and tumor samples were obtained at various time points. Furthermore, although the panel of somatic neoepitopes (Fig. 3A and 3B, and Table S6 in the Supplementary Appendix) may constitute the most important ones, the in vivo relative immunologic contribution of each peptide is unclear. However, data showing that functionally important immunogenic epitopes persisted after treatment with expanded tumor-infiltrating lymphocytes suggest that the response to mutations may persist over time.¹⁶ Although the recapitulation of the neoantigen signature in the validation set suggests that this may provide a generally applicable tool for prediction of a benefit from immunotherapy, further studies will be needed to investigate the role of MHC class II molecules and the relative effects and characteristics of neoantigens in different cancers.

Our use of whole-exome sequencing to identify a genetic basis associated with a benefit from

CTLA-4 blockade provides proof of principle that tumor genomics can inform responses to immunotherapy. For the field of cancer genetics, these data suggest a need for an expanded definition of the previous categories of driver and passenger mutations. Our data show that exonic missense mutations in general confer increased MHC class I binding (Fig. S5A and S5B in the Supplementary Appendix) and confirm the hypothesis³⁶ that some mutations formerly categorized as passengers may in fact represent “immune determinants.”

Supported by grants from the Frederick Adler Fund, the National Institutes of Health, Swim across America, the Ludwig Trust, the Melanoma Research Alliance, the Stand Up to Cancer–Cancer Research Institute Immunotherapy Dream Team, the Hazen Polsky Foundation, and the STARR Cancer Consortium and by a Ruth L. Kirschstein National Research Service Award (T32CA009512, to Dr. Snyder). Bristol-Myers Squibb, the employer of two authors, did not provide funding for this study.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

We thank Martin Miller at Memorial Sloan Kettering Cancer Center (MSKCC) for his assistance with the NetMHC server, Agnes Viale and Kety Huberman at the MSKCC Genomics Core, Annamalai Selvakumar and Alice Yeh at the MSKCC HLA typing laboratory for their technical assistance, and John Khoury for assistance in chart review.

REFERENCES

- Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 2010;363:711-23. [Erratum, *N Engl J Med* 2010;363:1290.]
- Wolchok JD, Kluger H, Callahan MK, et al. Nivolumab plus ipilimumab in advanced melanoma. *N Engl J Med* 2013;369:122-33.
- Ku GY, Yuan J, Page DB, et al. Single-institution experience with ipilimumab in advanced melanoma patients in the compassionate use setting: lymphocyte count after 2 doses correlates with survival. *Cancer* 2010;116:1767-75.
- Ji RR, Chasalow SD, Wang L, et al. An immune-active tumor microenvironment favors clinical response to ipilimumab. *Cancer Immunol Immunother* 2012;61:1019-31.
- Gajewski TF, Louahed J, Brichard VG. Gene signature in melanoma associated with clinical activity: a potential clue to unlock cancer immunotherapy. *Cancer J* 2010;16:399-403.
- Cha E, Klinger M, Hou Y, et al. Improved survival with T cell clonotype stability after anti-CTLA-4 treatment in cancer patients. *Sci Transl Med* 2014;6:238ra70.
- Castle JC, Kreiter S, Diekmann J, et al. Exploiting the mutanome for tumor vaccination. *Cancer Res* 2012;72:1081-91.
- Srivastava N, Srivastava PK. Modeling the repertoire of true tumor-specific MHC I epitopes in a human tumor. *PLoS One* 2009;4(7):e6094.
- Kvistborg P, Philips D, Kelderman S, et al. Anti-CTLA-4 therapy broadens the melanoma-reactive CD8+ T cell response. *Sci Transl Med* 2014;6:254ra128.
- Peggs KS, Quezada SA, Chambers CA, Korman AJ, Allison JP. Blockade of CTLA-4 on both effector and regulatory T cell compartments contributes to the antitumor activity of anti-CTLA-4 antibodies. *J Exp Med* 2009;206:1717-25.
- Wolchok JD, Weber JS, Hamid O, et al. Ipilimumab efficacy and safety in patients with advanced melanoma: a retrospective analysis of HLA subtype from four trials. *Cancer Immunol* 2010;10:9.
- Alexandrov LB, Nik-Zainal S, Wedge DC, et al. Signatures of mutational processes in human cancer. *Nature* 2013;500:415-21.
- Segal NH, Parsons DW, Peggs KS, et al. Epitope landscape in breast and colorectal cancer. *Cancer Res* 2008;68:889-92.
- Matsushita H, Vesely MD, Koboldt DC, et al. Cancer exome analysis reveals a T-cell-dependent mechanism of cancer immunoeediting. *Nature* 2012;482:400-4.
- van Rooij N, van Buuren MM, Philips D, et al. Tumor exome analysis reveals neoantigen-specific T-cell reactivity in an ipilimumab-responsive melanoma. *J Clin Oncol* 2013;31(32):e439-e442.
- Tran E, Turcotte S, Gros A, et al. Cancer immunotherapy based on mutation-specific CD4+ T cells in a patient with epithelial cancer. *Science* 2014;344:641-5.
- Wei X, Walia V, Lin JC, et al. Exome sequencing identifies GRIN2A as frequently mutated in melanoma. *Nat Genet* 2011;43:442-6.
- Berger MF, Hodis E, Heffernan TP, et al. Melanoma genome sequencing reveals frequent PREX2 mutations. *Nature* 2012;485:502-6.
- Hodis E, Watson IR, Kryukov GV, et al. A landscape of driver mutations in melanoma. *Cell* 2012;150:251-63.
- Park YY, Park ES, Kim SB, et al. Development and validation of a prognostic gene-expression signature for lung adenocarcinoma. *PLoS One* 2012;7(9):e44225.
- Birnbaum ME, Mendoza JL, Sethi DK, et al. Deconstructing the peptide-MHC specificity of T cell recognition. *Cell* 2014;157:1073-87.
- Morita D, Yamamoto Y, Suzuki J, Mori N, Igarashi T, Sugita M. Molecular requirements for T cell recognition of N-myristoylated peptides derived from the simian immunodeficiency virus Nef protein. *J Virol* 2013;87:482-8.
- Stuart GW, Moffett K, Leader JJ. A comprehensive vertebrate phylogeny using vector representations of protein sequences from whole genomes. *Mol Biol Evol* 2002;19:554-62.
- Aleksic M, Dushek O, Zhang H, et al. Dependence of T cell antigen recognition on T cell receptor-peptide MHC confinement time. *Immunity* 2010;32:163-74.
- Insaiddoo FK, Borbulevych OY, Hosain M, Santhanagopalan SM, Baxter TK, Baker BM. Loss of T cell antigen recognition arising from changes in peptide and major histocompatibility complex protein flexibility: implications for vaccine design. *J Biol Chem* 2011;286:40163-73.
- Sliz P, Michielin O, Cerottini JC, et al. Crystal structures of two closely related but antigenically distinct HLA-A2/melanocyte-melanoma tumor-antigen peptide complexes. *J Immunol* 2001;167:3276-84.
- Lim JB, Kim HO, Jeong SH, et al. Identification of HLA-A*2402-restricted HCMV immediate early-1 (IE-1) epitopes as targets for CD8+ HCMV-specific cytotoxic T lymphocytes. *J Transl Med* 2009;7:72.
- Wong R, Lau R, Chang J, et al. Immune responses to a class II helper peptide epitope in patients with stage III/IV resected melanoma. *Clin Cancer Res* 2004;10:5004-13.
- Li Y, Depontieu FR, Sidney J, et al. Structural basis for the presentation of tumor-associated MHC class II-restricted phosphopeptides to CD4+ T cells. *J Mol Biol* 2010;399:596-603.
- Poisson F, Baillou F, Dubois F, Janvier B, Roingeard P, Goudeau A. Immune response to synthetic peptides of hepatitis delta antigen. *J Clin Microbiol* 1993;31:2343-9.
- Monach PA, Meredith SC, Siegel CT, Schreiber H. A unique tumor antigen produced by a single amino acid substitution. *Immunity* 1995;2:45-59.
- Dubey P, Hendrickson RC, Meredith SC, et al. The immunodominant antigen of an ultraviolet-induced regressor tumor is generated by a somatic point mutation

- in the DEAD box helicase p68. *J Exp Med* 1997;185:695-705.
33. Allen PM, Matsueda GR, Evans RJ, Dunbar JB Jr, Marshall GR, Unanue ER. Identification of the T-cell and Ia contact residues of a T-cell antigenic epitope. *Nature* 1987;327:713-5.
34. Anderson MW, Gorski J, Cutting edge: TCR contacts as anchors: effects on affinity and HLA-DM stability. *J Immunol* 2003;171:5683-7.
35. Noguchi Y, Chen YT, Old LJ. A mouse mutant p53 product recognized by CD4+ and CD8+ T cells. *Proc Natl Acad Sci U S A* 1994;91:3171-5.
36. Srivastava PK, Duan F. Harnessing the antigenic fingerprint of each individual cancer for immunotherapy of human cancer: genomics shows a new way and its challenges. *Cancer Immunol Immunother* 2013;62:967-74.

Copyright © 2014 Massachusetts Medical Society.