

UC San Diego

UC San Diego Previously Published Works

Title

Raising awareness about sleep disorders

Permalink

<https://escholarship.org/uc/item/08r5803f>

Journal

Lung India, 34(3)

ISSN

0970-2113

Authors

Jaiswal, Stuti J
Owens, Robert L
Malhotra, Atul

Publication Date

2017

DOI

10.4103/0970-2113.205331

Peer reviewed

Review Article

Raising awareness about sleep disorders

Stuti J Jaiswal^{1,2}, Robert L Owens³, Atul Malhotra³¹???, Scripps Translational Science Institute, ²Department of Internal Medicine, Scripps Clinic/Scripps Green Hospital, ³Division of Pulmonary, Critical Care and Sleep Medicine, University of California San Diego School of Medicine, La Jolla, CA 92037, USA

ABSTRACT

Sleep disorders affect a substantial number of people worldwide and may be increasing in prevalence. Unfortunately, many of these disorders can go unrecognized and untreated in clinical practice. Although physicians understand that good sleep is essential to patient health, approaching the issue is often glossed over in the primary care setting. Importantly, recent data also suggest that sleep disorders are becoming increasingly recognized in developing nations. Here, we review the pertinent features and treatment of some of the major sleep disorders, including obstructive sleep apnea, central sleep apnea, insomnia, and others, that affect patient health. We aim to promote clinician and public awareness of various sleep disorders with the intent of increasing rates of recognition and treatment.

KEY WORDS: Central sleep apnea, obstructive sleep apnea, sleep disorders

Address for correspondence: Dr. Stuti Jaiswal, Scripps Translational Science Institute, 3344 North Torrey Pines Court, La Jolla, CA 92037, USA.
E-mail: jaiswal.stuti@scrippshealth.org

INTRODUCTION

Clinicians frequently recommend a good night's sleep for their patients but rarely focus on the issue when seeing them in the clinical setting. Sleep problems may also be increasing in developing nations as a recent study suggested that almost 17% of patients surveyed across eight countries in Asia and Africa reported trouble with sleep.^[1] Literature over the past two decades has established a myriad of clinically important complications from various sleep disorders. Thus, a working knowledge of these disorders and their manifestations is imperative for optimal patient care. Here, we review some of the major sleep disorders with a goal of raising awareness about these conditions, eventually to stimulate further research and clinical focus in these areas.

SLEEP DEPRIVATION

Today's fast-paced lifestyle has led globally to an increase in sleep deprivation.^[2-4] Although the recommended

amount of sleep in a 24 h period is 7–8 h, recent data show that almost 30% of Americans are sleeping 6 h or less^[5] and other countries are also reporting a decline in sleep duration.^[6] The impact of this societal change is unclear, but some would argue that increasing sleep duration might have benefits for the population at large. Neurocognitive function decreases in a dose-dependent manner with chronic sleep deprivation,^[7] which can impair productivity at work and in daily functioning. In addition, epidemiological studies consistently associate shortened sleep duration with weight gain, risk of myocardial infarction, hypertension, and diabetes when compared to individuals with normal sleep duration.^[8-13] Of note, prolonged sleepers also have increased risk of mortality and cardiovascular complications when compared to 8 h per night sleepers although the mechanism remains unknown.^[12,13] Experimental studies have also shown important effects of imposing reduced sleep duration on normal sleepers resulting in impaired glucose metabolism, altered satiety hormone regulation,

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: ???

Access this article online	
Quick Response Code:	Website: www.lungindia.com
	DOI: ***

and autonomic dysfunction.^[14-18] The data currently fall short of providing rigorous evidence that extending sleep duration leads to important health benefits. Nonetheless, strong recommendations have been provided to educate the public regarding the health effects of sleep deprivation and the potential benefits of adequate sleep.^[19] The methods to achieve adequate sleep duration in the 7–8 h range depend on the underlying cause but include a safe place to sleep, decreased noise and screen-time before bed, lifestyle modifications, sleep hygiene recommendations, and prioritizing sleep.

OBSTRUCTIVE SLEEP APNEA

Perhaps the most clinically important sleep disorder is obstructive sleep apnea (OSA), which is defined by interruptions in breathing during sleep. The pathophysiology of the disorder relates to intermittent, partial, and/or complete collapse of the upper airway during sleep. OSA patients often snore and report daytime sleepiness although their clinical characteristics vary considerably. Obesity is a major risk factor for OSA although 30–40% of OSA patients are lean. In the US, roughly 6% of women and 13% of men have clinically important OSA defined by an apnea–hypopnea index >15/h.^[20] That is, individuals have a breathing problem during sleep at least once every 4 min. Worldwide, data are less abundant, but prevalence rates are similar, despite lower levels of obesity outside the US. For example, Heinzer *et al.* reported up to 50% of men in Switzerland had OSA with at least 15 breathing disturbances per hour of sleep.^[21] In India, some prevalence figures are reported, which suggest that roughly 7–9% of the middle-aged population suffer from this condition.^[22,23] Interestingly, although there are lower obesity rates reported in Asia than in the US, the prevalence of OSA is equivalent to that of Western nations,^[24,25] suggesting that the prevalence of OSA is likely to be higher for a given body mass index in Asian populations. Furthermore, as the obesity pandemic spreads, OSA prevalence figures in developing countries are likely to increase over time.

OSA has important consequences. With stoppages in breathing, individuals frequently have to wake up from sleep to breathe again, and thus sleep becomes very fragmented. This fragmentation leads to excessive daytime sleepiness, reduced quality of life, impaired neurocognitive function, increased risk of road traffic accidents, and other complications.^[26-29] In addition, with each breathing stoppage, gas exchange is disturbed due to pharyngeal collapse, and consequently, hypoxemia and hypercapnia can occur with associated catecholamine surges from arousal from sleep. These factors, along with others, likely underlie the substantial cardiovascular sequelae of OSA, which include increased risk of coronary artery disease, stroke, congestive heart failure (HF), and cardiac arrhythmias. In addition, OSA is well established as a risk factor for hypertension, based on animal models, human epidemiological studies, and rigorous randomized

controlled trials.^[30-33] Further details regarding the association of OSA and cardiovascular disease have been reviewed extensively elsewhere.^[34-36] Ongoing studies are examining the impact of intervention on OSA, but treatment of OSA is felt to improve cardiovascular risk although the extent of the improvement is still debated.

The diagnosis of OSA starts with a history and physical examination in those where clinical suspicion exists. In general, OSA diagnosis requires objective testing, with home sleep testing (HST) now emerging as a reasonable approach for the vast majority of patients.^[37] In-laboratory polysomnography (PSG) is historically the gold standard diagnostic test but is expensive and cumbersome and yields similar outcomes to simpler and cheaper tests. HST equipment varies in its complexity, but in general, simple devices (e.g., pulse oximeter, respiratory belt, and nasal pressure sensor) provide satisfactory diagnostic information to yield a diagnosis. In some cases, a repeat HST or in-laboratory testing may be required if a diagnosis remains elusive.

For therapy, continuous positive airway pressure (CPAP) remains the treatment of choice. Rigorous, randomized trials have shown that CPAP improves daytime symptoms and blood pressure.^[38-40] CPAP also likely reduces the risk of road traffic accidents and cardiovascular events although these data are still evolving. CPAP works well in those who tolerate it, but in those who are intolerant, further education and support are often required. CPAP adherence is likely improved with patient education, addressing nasal patency, optimizing the interface, and regular feedback/interactions. In those who truly cannot tolerate CPAP, alternative positive airway pressure (PAP) devices (e.g., bi-level PAP) may have value although the data are not compelling in clinical trials. Alternative therapies such as oral appliances and upper airway surgery have variable efficacy and have risk/expense associated with them. However, in their recent 2015 practice guidelines update, the American Academy of Sleep Medicine (AASM) recommends that treatment with custom oral appliance in patients who are intolerant of CPAP or who prefer alternate therapy as treatment with an oral appliance is preferred to no therapy.^[41] Conservative measures include weight loss^[42] through diet and exercise, the avoidance of alcohol,^[43,44] goal of 7–8 h of sleep at night, and maintenance of nasal patency. Positional therapy (e.g., avoidance of the supine posture) may have some value although, in general, it is hard to implement and monitor in the patient's home. Despite considerable progress in OSA, the majority of patients remains undiagnosed and untreated, speaking to the need for increased awareness and clinician suspicion for patients to receive optimal management.

CENTRAL SLEEP APNEA

Breathing centers in the medulla drive motor output to all of the various respiratory muscles including the diaphragm, muscles in the chest wall, and to the upper airways/tongue.

1 These breathing centers respond to blood PCO₂ and PO₂ 1
 2 levels, as well as other information from the periphery 2
 3 to regulate the rate and depth of inspiration. Affected 3
 4 individuals have inappropriate or absent responses to 4
 5 breathing stimuli, which leads to unstable ventilatory 5
 6 patterns during sleep. Clinically, central sleep apnea (CSA) 6
 7 is defined as cessation of airflow for at least 10 s in the 7
 8 absence of any inspiratory effort. The consequences and 8
 9 pathophysiology of CSA overlap to some degree with those 9
 10 of OSA, including increased sympathetic nervous system 10
 11 activation and arrhythmogenesis.^[45,46] In general, CSA can 11
 12 be divided into hypercapnic and nonhypercapnic types. 12

13
 14 Hypercapnic CSA can present in patients with strokes and 14
 15 brain lesions where direct neuronal injury occurs in the 15
 16 brain's chemoreceptive or respiratory rhythm generator 16
 17 regions. Patients with neuromuscular disorders such 17
 18 as myasthenia gravis or amyotrophic lateral sclerosis 18
 19 also fit into this category although the cause of their 19
 20 disease originates from neuromuscular dysfunction in 20
 21 ventilatory muscles in the periphery and not the central 21
 22 nervous system. Narcotic-induced CSA^[47,48] occurs 22
 23 in a dose-dependent response to central respiratory 23
 24 depression facilitated by actions on opioid receptors in 24
 25 the brainstem;^[49] patients receiving morphine equivalent 25
 26 daily dose of 200 mg or greater may have increased risk.^[50] 26

27 A recent study showed that 46% of patients on long-term 27
 28 opioid therapy demonstrated severe sleep disordered 28
 29 breathing with AHI >30/h and had a high risk of central 29
 30 apneas.^[47] Obesity hypoventilation syndrome (OHS), 30
 31 another form of hypercapnic CSA, is also clinically 31
 32 important given the global obesity epidemic although the 32
 33 exact pathophysiology of the disorder requires further 33
 34 elucidation. OHS occurs in a subset of obese patients; 34
 35 in general, elevated HCO₃⁻/PaCO₂ levels on laboratory 35
 36 work-up along with continued nocturnal desaturations 36
 37 despite CPAP therapy help in establishing the diagnosis. 37
 38 These patients usually require nighttime CPAP for 38
 39 treatment although bi-level PAP and new therapies such 39
 40 as intelligent volume-assured pressure support/averaged 40
 41 volume-assured pressure support are now commonly used. 41

42
 43 The most clinically relevant type of CSA is nonhypercapnic 43
 44 and presents in the context of HF patients with 44
 45 reduced ejection fraction, with an estimated 30–50% of 45
 46 patients affected.^[46,51] These patients develop a distinct, 46
 47 crescendo-decrescendo pattern of breathing termed 47
 48 Cheyne–Stokes respiration. Patients may have an 48
 49 exaggerated ventilatory response (hyperventilatory) to 49
 50 PCO₂ to the point where they lower the PCO₂ below the 50
 51 chemical apnea threshold, thereby causing apnea. Previous 51
 52 studies have examined the utility of CPAP therapy in HF 52
 53 patients with CSA, and although it may improve some 53
 54 outcomes in HF such as nocturnal oxygenation and 6-min 54
 55 walk time, it does not appear to improve mortality.^[52,53] 55
 56 Recently, a considerable amount of interest has arisen in 56
 57 the potential of adaptive (or auto) servo ventilation (ASV) 57
 58 in treating CSA in HF patients. ASV uses a noninvasive 58
 59 ventilator to measure patient ventilation and subsequently 59

adapt expiratory PAP and pressure support levels to 1
 eliminate apneas and hypopneas. **Although some studies** 2
have shown some short-term benefits and the reasons 3
behind this are currently unclear,^[54,55] a recent trial 4
of 1325 HF patients treated with ASV versus medical 5
therapy showed a significant increase in cardiovascular 6
and all-cause mortality at 12 months in the ASV group.^[56] 7
 Accordingly, the most recent AASM practice guidelines 8
 recommend that physicians stop prescribing ASV to treat 9
 CSA in patients with symptomatic HF and left ejection 10
 fraction <45%. Further studies are needed to determine 11
 the utility of ASV for short-term use and other outcome 12
 measures. In the meantime, the optimal therapy for 13
 Cheyne–Stokes respiration remains optimization of HF 14
 therapy, which can also reduce CSA severity. 15

16 INSOMNIA 17

18
 19 Insomnia remains one of the most common sleep 19
 20 complaints in the primary care clinic although the 20
 21 prevalence varies depending on the definition used and 21
 22 the population studied. Importantly, insomnia differs 22
 23 from sleep deprivation in that there is adequate time and 23
 24 opportunity to obtain sleep, but sleep is impaired and has 24
 25 negative consequences for the patient during the daytime. 25

26
 27 In the United States, the estimated prevalence of chronic 27
 28 insomnia ranges from 9% to 19%.^[57] Women report higher 28
 29 rates than men,^[58] and minority^[59] and elderly populations 29
 30 are also at an increased risk, with the latter reporting rates 30
 31 of approximately 50%.^[60,61] The International Classification 31
 32 of Sleep Disorders, Third Edition defines insomnia as 32
 33 difficulty with sleep initiation, sleep maintenance, or 33
 34 early sleep termination, a minimum of three nights per 34
 35 week for 3 months, with patients also reporting impaired 35
 36 daytime functioning.^[62] In general, most insomnia 36
 37 patients fall into the category of secondary insomnia, 37
 38 meaning that medication, a comorbid illness, exogenous 38
 39 substances (e.g., caffeine, nicotine, alcohol), another sleep 39
 40 disorder, or poor sleep environment are the root cause of 40
 41 symptoms. In contrast, primary insomnia occurs in the 41
 42 absence of these other factors and affects approximately 42
 43 25% of patients suffering from chronic insomnia and, in 43
 44 approximately, 5–10% of the US population in general.^[58,63] 44

45
 46 Insomnia reduces the quality of life^[64,65] and is also 46
 47 heavily associated with depression and anxiety although 47
 48 the directionality or causality of this relationship is not 48
 49 fully understood.^[66,67] A recent study examining the rates 49
 50 of accidents in over 5,000 patients with insomnia in 10 50
 51 countries showed that insomnia increased the number 51
 52 of motor vehicle and occupational accidents,^[68] which 52
 53 potentially places the lives of others at risk. 53

54
 55 Treatment for insomnia has frequently involved the use of 55
 56 pharmacologic interventions including benzodiazepines 56
 57 and nonbenzodiazepine sedative-hypnotic agents such as 57
 58 zolpidem, eszopiclone, and zaleplon. These interventions 58
 59 may have short-term efficacy, but long-term use is not 59

recommended due to issues with dependence, tolerance, as well as medication side effects.^[69-71] Interestingly, cognitive behavioral therapy (CBT) has been shown to improve sleep quality, as measured by sleep efficiency, time spent in slow-wave sleep, and total sleep time, when compared to pharmacologic intervention or placebo in older adults^[72] as well as younger patients.^[73] CBT including measures of sleep hygiene, sleep restriction, cognitive therapy, and relaxation may be one of the most effective treatments for insomnia.^[72] Internet-based behavioral therapies are now being used and are potentially “scalable” even in under-resourced areas.

RESTLESS LEGS SYNDROME AND PERIODIC LIMB MOVEMENTS

The prevalence of restless legs syndrome (RLS) varies depending on geographic location, but individuals in Asia experience lower rates of the disorder than individuals in North America and Europe, where disease rates have been reported as high as 30% in the latter two continents.^[74-76] RLS patients usually describe an uncomfortable sensation in the lower extremities, sometimes described as itching, tingling, or crawling, which leads to an irresistible urge to move that temporarily relieves the sensation. Symptoms are typically worse at rest compared to during activity and worse at night compared to daytime. The diagnosis requires exclusion of various mimics such as arthritis, vascular insufficiency, akathisia, and peripheral neuropathy. Patients can have milder disease, with symptoms occurring episodically with little to no sleep interruption or they may suffer from more severe disease, where episodes occur 3 or more times per week with severe interruption of nighttime sleep and presence of daytime symptoms. RLS can lead to insomnia due to an inability to fall asleep as patients often complain that their leg movements keep them awake.

Periodic limb movement (PLM) disorder involves repetitive episodes of stereotyped limb movements – usually, in the lower extremities occurring during sleep, and frequently occurs in conjunction with RLS; approximately 80–90% of RLS patients exhibit PLMs.^[77] PLMs are usually assessed during in-laboratory testing based on measurement of the anterior tibialis electromyogram. With the move toward HST, PLMs are becoming harder to assess and quantify. PLMs are frequently inconsequential but occasionally lead to sleep fragmentation.

The treatment of isolated PLMs is controversial, with minimal data showing improvement in important outcomes with therapy. On the other hand, RLS can respond to treatment with rigorous randomized controlled trials showing important outcomes.^[77] When treating patients with RLS, it is important to think about underlying causes, such as end-stage renal disease,^[78,79] low ferritin levels,^[80,81] and pregnancy.^[82] Patients with RLS related to low iron levels can respond to treatment with iron repletion in those with ferritin <50 mcg/L.^[83] In cases where

pharmacotherapy is required, pramipexole or ropinirole can be helpful although hypnotic agents, gabapentin, and physical massaging devices may all have value.^[77]

NARCOLEPSY

Our understanding of the neurobiology of sleep has progressed substantially over the last several decades, due largely in part to animal models that allow deeper understanding of the brain’s complexities in relationship to sleep. Animal models of narcolepsy show that certain hypothalamic lesions lead to a loss of muscle tone in response to emotional stimuli (cataplexy).^[84,85] Neurons in the posterior lateral hypothalamus secrete hypocretin (orexin), which is key in sustaining normal wakefulness and in preventing rapid eye movement (REM) activity from occurring at inappropriate times.^[86,87] Human and animal studies have implicated a loss of these neurons with reduced orexin secretion in narcolepsy.^[86,88] The classic tetrad for narcolepsy includes excessive daytime sleepiness, cataplexy, sleep paralysis, and hypnagogic hallucinations. People with narcolepsy often experience very fragmented sleep such that narcolepsy is often described as sleep encroaching on wakefulness and wakefulness encroaching on sleep. People with narcolepsy often go undiagnosed, and thus a high index of suspicion is required by clinicians to make the diagnosis. The diagnosis of narcolepsy is usually made by PSG with multiple sleep latency testing. The overnight sleep test is primarily to exclude other sleep disorders such as sleep apnea, PLMs. The daytime naps occur at 2-h intervals and are used to quantify sleep onset latency as well as propensity for REM sleep. People with objective evidence of sleepiness will have reduced sleep onset latency. People with multiple sleep onset REM episodes frequently have narcolepsy although occasional false positives can occur (e.g., depression, severe sleep apnea, and circadian misalignment). In patients where the diagnosis remains unclear, a lumbar puncture can occasionally be helpful to document low levels of hypocretin (orexin) in patients with narcolepsy plus cataplexy. In narcolepsy without cataplexy, cerebrospinal fluid orexin is typically normal, making lumbar puncture less helpful in these cases. The treatment of narcolepsy involves stimulants; modafinil is commonly used in the US although methylphenidate, dextroamphetamine, and other agents have utility.^[89] Nap therapy can also be helpful in some cases since people with narcolepsy frequently wake up feeling refreshed. Cataplexy on the other hand does not always require specific treatment depending on severity. Stimulus control can be helpful if certain emotional stimuli known to trigger cataplexy can be anticipated. If pharmacotherapy is required, certain antidepressant medications such as selective serotonin reuptake inhibitors or serotonin and norepinephrine reuptake inhibitors (SNRIs) can help suppress REM sleep and prevent cataplexy attacks; of note, antidepressant therapy is not currently recommended for treatment in narcolepsy alone.^[90] Sodium oxybate also has

a role as it helps consolidate sleep, can eliminate cataplexy, and improve daytime sleepiness; the combination of modafinil and sodium oxybate can be highly effective for afflicted individuals.^[91]

CONCLUSIONS

Our knowledge of the mechanisms and consequences of sleep disorders have become increasingly compelling over the last several decades and has now reached a point where we know that ignoring issues regarding sleep lead to poor clinical outcomes. Progress is being made in increasing awareness of sleep disorders, but further efforts are required to improve the situation, particularly in developing countries. The spread of Western culture, including technology, sleeping, and activity habits, as well as diet, suggest that sleep issues may only continue to increase in other countries. **Only with further efforts in advocacy, research, and education will major progress continue.** It is also important that clinicians utilize their knowledge and current research regarding sleep disorders to recognize these problems in the clinic and treat their patients accordingly.

Financial support and sponsorship

This research was funded in part by the NIH/NCATS flagship Clinical and Translational Science Award Grant (5KL2TR001112).

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Stranges S, Tigbe W, Gómez-Olivé FX, Thorogood M, Kandala NB. Sleep problems: An emerging global epidemic? Findings from the INDEPTH WHO-SAGE study among more than 40,000 older adults from 8 countries across Africa and Asia. *Sleep* 2012;35:1173-81.
2. Bonnet MH, Arand DL. We are chronically sleep deprived. *Sleep* 1995;18:908-11.
3. Goeger JA, Zijlstra FR, Dijk DJ. Sleep quantity, sleep difficulties and their perceived consequences in a representative sample of some 2000 British adults. *J Sleep Res* 2004;13:359-71.
4. Kronholm E, Partonen T, Laatikainen T, Peltonen M, Härmä M, Hublin C, et al. Trends in self-reported sleep duration and insomnia-related symptoms in Finland from 1972 to 2005: A comparative review and re-analysis of Finnish population samples. *J Sleep Res* 2008;17:54-62.
5. Krueger PM, Friedman EM. Sleep duration in the United States: A cross-sectional population-based study. *Am J Epidemiol* 2009;169:1052-63.
6. Bin YS, Marshall NS, Glozier N. Secular trends in adult sleep duration: A systematic review. *Sleep Med Rev* 2012;16:223-30.
7. Van Dongen HP, Maislin G, Mullington JM, Dinges DF. The cumulative cost of additional wakefulness: Dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep* 2003;26:117-26.
8. Gottlieb DJ, Redline S, Nieto FJ, Baldwin CM, Newman AB, Resnick HE, et al. Association of usual sleep duration with hypertension: The Sleep Heart Health Study. *Sleep* 2006;29:1009-14.
9. Gottlieb DJ, Punjabi NM, Newman AB, Resnick HE, Redline S, Baldwin CM, et al. Association of sleep time with diabetes mellitus and impaired glucose tolerance. *Arch Intern Med* 2005;165:863-7.
10. Newman AB, Enright PL, Manolio TA, Haponik EF, Wahl PW. Sleep disturbance, psychosocial correlates, and cardiovascular disease in 5201 older adults: The Cardiovascular Health Study. *J Am Geriatr Soc* 1997;45:1-7.

11. Gangwisch JE, Heymsfield SB, Boden-Albala B, Buijs RM, Kreier F, Pickering TG, et al. Short sleep duration as a risk factor for hypertension: Analyses of the first National Health and Nutrition Examination Survey. *Hypertension* 2006;47:833-9.
12. Cappuccio FP, Cooper D, D'Elia L, Strazzullo P, Miller MA. Sleep duration predicts cardiovascular outcomes: A systematic review and meta-analysis of prospective studies. *Eur Heart J* 2011;32:1484-92.
13. Ayas NT, White DP, Manson JE, Stampfer MJ, Speizer FE, Malhotra A, et al. A prospective study of sleep duration and coronary heart disease in women. *Arch Intern Med* 2003;163:205-9.
14. Spiegel K, Leproult R, Van Cauter E. Impact of sleep debt on metabolic and endocrine function. *Lancet* 1999;354:1435-9.
15. Spiegel K, Knutson K, Leproult R, Tasali E, Van Cauter E. Sleep loss: A novel risk factor for insulin resistance and Type 2 diabetes. *J Appl Physiol* (1985) 2005;99:2008-19.
16. Spiegel K, Tasali E, Penev P, Van Cauter E. Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med* 2004;141:846-50.
17. Taheri S, Lin L, Austin D, Young T, Mignot E. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med* 2004;1:e62.
18. Tochikubo O, Ikeda A, Miyajima E, Ishii M. Effects of insufficient sleep on blood pressure monitored by a new multi biomedical recorder. *Hypertension* 1996;27:1318-24.
19. Mukherjee S, Patel SR, Kales SN, Ayas NT, Strohl KP, Gozal D, et al. An official American thoracic society statement: The importance of healthy sleep. Recommendations and future priorities. *Am J Respir Crit Care Med* 2015;191:1450-8.
20. Peppard PE, Young T, Barnett JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol* 2013;177:1006-14.
21. Heinzer R, Vat S, Marques-Vidal P, Marti-Soler H, Andrieu D, Tobback N, et al. Prevalence of sleep-disordered breathing in the general population: The HypnoLaus study. *Lancet Respir Med* 2015;3:310-8.
22. Udawadia ZF, Doshi AV, Lonkar SG, Singh CI. Prevalence of sleep-disordered breathing and sleep apnea in middle-aged urban Indian men. *Am J Respir Crit Care Med* 2004;169:168-73.
23. Reddy EV, Kadiravan T, Mishra HK, Sreenivas V, Handa KK, Sinha S, et al. Prevalence and risk factors of obstructive sleep apnea among middle-aged urban Indians: A community-based study. *Sleep Med* 2009;10:913-8.
24. Lam B, Ip MS, Tench E, Ryan CF. Craniofacial profile in Asian and white subjects with obstructive sleep apnoea. *Thorax* 2005;60:504-10.
25. Lam B, Lam DC, Ip MS. Obstructive sleep apnoea in Asia. *Int J Tuberc Lung Dis* 2007;11:2-11.
26. Strohl KP, Brown DB, Collop N, George C, Grunstein R, Han F, et al. An official American thoracic society clinical practice guideline: Sleep apnea, sleepiness, and driving risk in noncommercial drivers. An update of a 1994 statement. *Am J Respir Crit Care Med* 2013;187:1259-66.
27. Kushida CA, Nichols DA, Holmes TH, Quan SF, Walsh JK, Gottlieb DJ, et al. Effects of continuous positive airway pressure on neurocognitive function in obstructive sleep apnea patients: The Apnea Positive Pressure Long-term Efficacy Study (APPLES). *Sleep* 2012;35:1593-602.
28. Johns MW. Daytime sleepiness, snoring, and obstructive sleep apnea. The Epworth Sleepiness Scale. *Chest* 1993;103:30-6.
29. Goncalves MA, Paiva T, Ramos E, Guilleminault C. Obstructive sleep apnea syndrome, sleepiness, and quality of life. *Chest* 2004;125:2091-6.
30. Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med* 2000;342:1378-84.
31. O'Connor GT, Caffo B, Newman AB, Quan SF, Rapoport DM, Redline S, et al. Prospective study of sleep-disordered breathing and hypertension: The Sleep Heart Health Study. *Am J Respir Crit Care Med* 2009;179:1159-64.
32. Marin JM, Agusti A, Villar I, Forner M, Nieto D, Carrizo SJ, et al. Association between treated and untreated obstructive sleep apnea and risk of hypertension. *JAMA* 2012;307:2169-76.
33. Walia HK, Li H, Rueschman M, Bhatt DL, Patel SR, Quan SF, et al. Association of severe obstructive sleep apnea and elevated blood pressure despite antihypertensive medication use. *J Clin Sleep Med* 2014;10:835-43.
34. Shamsuzzaman AS, Gersh BJ, Somers VK. Obstructive sleep apnea: Implications for cardiac and vascular disease. *JAMA* 2003;290:1906-14.

1 35. Baguet JP, Barone-Rochette G, Tamisier R, Levy P, Pépin JL. Mechanisms of cardiac dysfunction in obstructive sleep apnea. *Nat Rev Cardiol* 2012;9:679-88. 1

2 36. Hohl M, Linz B, Böhm M, Linz D. Obstructive sleep apnea and atrial arrhythmogenesis. *Curr Cardiol Rev* 2014;10:362-8. 2

3 37. Kuna ST, Gurubhagavatula I, Maislin G, Hin S, Hartwig KC, McCloskey S, et al. Noninferiority of functional outcome in ambulatory management of obstructive sleep apnea. *Am J Respir Crit Care Med* 2011;183:1238-44. 3

4 38. Pepperell JC, Ramdassingh-Dow S, Crosthwaite N, Mullins R, Jenkinson C, Stradling JR, et al. Ambulatory blood pressure after therapeutic and subtherapeutic nasal continuous positive airway pressure for obstructive sleep apnoea: A randomised parallel trial. *Lancet* 2002;359:204-10. 4

5 39. Jenkinson C, Davies RJ, Mullins R, Stradling JR. Comparison of therapeutic and subtherapeutic nasal continuous positive airway pressure for obstructive sleep apnoea: A randomised prospective parallel trial. *Lancet* 1999;353:2100-5. 5

6 40. Bakker JP, Edwards BA, Gautam SP, Montesi SB, Durán-Cantolla J, Aizpuru F, et al. Blood pressure improvement with continuous positive airway pressure is independent of obstructive sleep apnea severity. *J Clin Sleep Med* 2014;10:365-9. 6

7 41. Ramar K, Dort LC, Katz SG, Lettieri CJ, Harrod CG, Thomas SM, et al. Clinical practice guideline for the treatment of obstructive sleep apnea and snoring with oral appliance therapy: An update for 2015. *J Clin Sleep Med* 2015;11:773-827. 7

8 42. Chirinos JA, Gurubhagavatula I, Teff K, Rader DJ, Wadden TA, Townsend R, et al. CPAP, weight loss, or both for obstructive sleep apnea. *N Engl J Med* 2014;370:2265-75. 8

9 43. Scrima L, Broudy M, Nay KN, Cohn MA. Increased severity of obstructive sleep apnea after bedtime alcohol ingestion: Diagnostic potential and proposed mechanism of action. *Sleep* 1982;5:318-28. 9

10 44. Taasan VC, Block AJ, Boysen PG, Wynne JW. Alcohol increases sleep apnea and oxygen desaturation in asymptomatic men. *Am J Med* 1981;71:240-5. 10

11 45. Lanfranchi PA, Somers VK, Braghiroli A, Corra U, Eleuteri E, Giannuzzi P. Central sleep apnea in left ventricular dysfunction: Prevalence and implications for arrhythmic risk. *Circulation* 2003;107:727-32. 11

12 46. Javaheri S, Parker TJ, Liming JD, Corbett WS, Nishiyama H, Wexler L, et al. Sleep apnea in 81 ambulatory male patients with stable heart failure. Types and their prevalences, consequences, and presentations. *Circulation* 1998;97:2154-9. 12

13 47. Rose AR, Catcheside PG, McEvoy RD, Paul D, Kapur D, Peak E, et al. Sleep disordered breathing and chronic respiratory failure in patients with chronic pain on long term opioid therapy. *J Clin Sleep Med* 2014;10:847-52. 13

14 48. Walker JM, Farnley RJ, Rhondeau SM, Boyle KM, Valentine K, Cloward TV, et al. Chronic opioid use is a risk factor for the development of central sleep apnea and ataxic breathing. *J Clin Sleep Med* 2007;3:455-61. 14

15 49. Lalley PM. Mu-opioid receptor agonist effects on medullary respiratory neurons in the cat: Evidence for involvement in certain types of ventilatory disturbances. *Am J Physiol Regul Integr Comp Physiol* 2003;285:R1287-304. 15

16 50. Correa D, Farnley RJ, Chung F, Prasad A, Lam D, Wong J. Chronic opioid use and central sleep apnea: A review of the prevalence, mechanisms, and perioperative considerations. *Anesth Analg* 2015;120:1273-85. 16

17 51. Javaheri S. Sleep disorders in systolic heart failure: A prospective study of 100 male patients. The final report. *Int J Cardiol* 2006;106:21-8. 17

18 52. Bradley TD, Logan AG, Kimoff RJ, Sériès F, Morrison D, Ferguson K, et al. Continuous positive airway pressure for central sleep apnea and heart failure. *N Engl J Med* 2005;353:2025-33. 18

19 53. Sin DD, Logan AG, Fitzgerald FS, Liu PP, Bradley TD. Effects of continuous positive airway pressure on cardiovascular outcomes in heart failure patients with and without Cheyne-Stokes respiration. *Circulation* 2000;102:61-6. 19

20 54. Owada T, Yoshihisa A, Yamauchi H, Iwaya S, Suzuki S, Yamaki T, et al. Adaptive servoventilation improves cardiorenal function and prognosis in heart failure patients with chronic kidney disease and sleep-disordered breathing. *J Card Fail* 2013;19:225-32. 20

21 55. Yoshihisa A, Suzuki S, Yamaki T, Sugimoto K, Kunii H, Nakazato K, et al. Impact of adaptive servo-ventilation on cardiovascular function and prognosis in heart failure patients with preserved left ventricular ejection fraction and sleep-disordered breathing. *Eur J Heart Fail* 2013;15:543-50. 21

22 56. Cowie MR, Woehrl H, Wegscheider K, Angermann C, d'Ortho MP, Erdmann E, et al. Adaptive servo-ventilation for central sleep apnea in systolic heart failure. *N Engl J Med* 2015;373:1095-105. 22

23 57. Shochat T, Umphress J, Israel AG, Ancoli-Israel S. Insomnia in primary care patients. *Sleep* 1999;22 Suppl 2:S359-65. 23

24 58. Ohayon MM. Epidemiology of insomnia: What we know and what we still need to learn. *Sleep Med Rev* 2002;6:97-111. 24

25 59. Williams NJ, Grandne MA, Snipes A, Rogers A, Williams O, Airhihenbuwa C, et al. Racial/ethnic disparities in sleep health and health care: Importance of the sociocultural context. *Sleep Health* 2015;1:28-35. 25

26 60. Foley DJ, Monjan A, Simonsick EM, Wallace RB, Blazer DG. Incidence and remission of insomnia among elderly adults: An epidemiologic study of 6,800 persons over three years. *Sleep* 1999;22 Suppl 2:S366-72. 26

27 61. Foley DJ, Monjan AA, Brown SL, Simonsick EM, Wallace RB, Blazer DG. Sleep complaints among elderly persons: An epidemiologic study of three communities. *Sleep* 1995;18:425-32. 27

28 62. American Academy of Sleep Medicine. International classification of sleep disorders. 3rd ed. **???: American Academy of Sleep Disorders**; 2014. **AQ4** 28

29 63. Roth T, Roehrs T. Insomnia: Epidemiology, characteristics, and consequences. *Clin Cornerstone* 2003;5:5-15. 29

30 64. Zammitt GK, Weiner J, Damato N, Sillup GP, McMillan CA. Quality of life in people with insomnia. *Sleep* 1999;22 Suppl 2:S379-85. 30

31 65. Avidan AY. Insomnia in the geriatric patient. *Clin Cornerstone* 2003;5:51-60. 31

32 66. Ohayon MM, Roth T. Place of chronic insomnia in the course of depressive and anxiety disorders. *J Psychiatr Res* 2003;37:9-15. 32

33 67. Johnson EO, Roth T, Breslau N. The association of insomnia with anxiety disorders and depression: Exploration of the direction of risk. *J Psychiatr Res* 2006;40:700-8. 33

34 68. Léger D, Bayon V, Ohayon MM, Philip P, Ement P, Metlaine A, et al. Insomnia and accidents: Cross-sectional study (EQUINOX) on sleep-related home, work and car accidents in 5293 subjects with insomnia from 10 countries. *J Sleep Res* 2014;23:143-52. 34

35 69. Stranks EK, Crowe SF. The acute cognitive effects of zopiclone, zolpidem, zaleplon, and eszopiclone: A systematic review and meta-analysis. *J Clin Exp Neuropsychol* 2014;36:691-700. 35

36 70. Liappas IA, Malits PN, Dimopoulos NP, Gitsa OE, Liappas AI, Nikolaou ChK, et al. Zolpidem dependence case series: Possible neurobiological mechanisms and clinical management. *J Psychopharmacol* 2003;17:131-5. 36

37 71. Hajak G, Müller WE, Wittchen HU, Pittrow D, Kirch W. Abuse and dependence potential for the non-benzodiazepine hypnotics zolpidem and zopiclone: A review of case reports and epidemiological data. *Addiction* 2003;98:1371-8. 37

38 72. Sivertsen B, Omvik S, Pallesen S, Bjorvatn B, Havik OE, Kvale G, et al. Cognitive behavioral therapy vs zopiclone for treatment of chronic primary insomnia in older adults: A randomized controlled trial. *JAMA* 2006;295:2851-8. 38

39 73. Jacobs GD, Pace-Schott EF, Stickgold R, Otto MW. Cognitive behavior therapy and pharmacotherapy for insomnia: A randomized controlled trial and direct comparison. *Arch Intern Med* 2004;164:1888-96. 39

40 74. Koo BB. Restless legs syndrome: Relationship between prevalence and latitude. *Sleep Breath* 2012;16:1237-45. 40

41 75. Innes KE, Selfe TK, Agarwal P. Prevalence of restless legs syndrome in North American and Western European populations: A systematic review. *Sleep Med* 2011;12:623-34. 41

42 76. Ohayon MM, O'Hara R, Vitiello MV. Epidemiology of restless legs syndrome: A synthesis of the literature. *Sleep Med Rev* 2012;16:283-95. 42

43 77. Aurora RN, Kristo DA, Bista SR, Rowley JA, Zak RS, Casey KR, et al. The treatment of restless legs syndrome and periodic limb movement disorder in adults – An update for 2012: Practice parameters with an evidence-based systematic review and meta-analyses: An American Academy of Sleep Medicine Clinical Practice Guideline. *Sleep* 2012;35:1039-62. 43

44 78. Gigli GL, Adorati M, Dolso P, Piani A, Valente M, Brotini S, et al. Restless legs syndrome in end-stage renal disease. *Sleep Med* 2004;5:309-15. 44

45 79. Winkelman JW, Chertow GM, Lazarus JM. Restless legs syndrome in end-stage renal disease. *Am J Kidney Dis* 1996;28:372-8. 45

46 80. O'Keefe ST, Gavin K, Lavan JN. Iron status and restless legs syndrome in the elderly. *Age Ageing* 1994;23:200-3. 46

47 81. O'Keefe ST, Noel J, Lavan JN. Restless legs syndrome in the elderly. *Postgrad Med J* 1993;69:701-3. 47

48 82. Manconi M, Govoni V, De Vito A, Economou NT, Cesnik E, Casetta I, et al. Restless legs syndrome and pregnancy. *Neurology* 2004;63:1065-9. 48

49 83. Sun ER, Chen CA, Ho G, Earley CJ, Allen RP. Iron and the restless legs syndrome. *Sleep* 1998;21:371-7. 49

50 84. Riehl J, Nishino S, Cederberg R, Dement WC, Mignot E. Development 50

1	of cataplexy in genetically narcoleptic Dobermans. <i>Exp Neurol</i>	activity. <i>Proc Natl Acad Sci U S A</i> 1998;95:322-7.	1
2	1998;152:292-302.		2
3	85. Kushida CA, Baker TL, Dement WC. Electroencephalographic correlates	88. Thannickal TC, Moore RY, Nienhuis R, Ramanathan L, Gulyani S,	2
4	of cataplectic attacks in narcoleptic canines. <i>Electroencephalogr Clin</i>	Aldrich M, <i>et al.</i> Reduced number of hypocretin neurons in human	3
5	<i>Neurophysiol</i> 1985;61:61-70.	narcolepsy. <i>Neuron</i> 2000;27:469-74.	3
6	86. Chemelli RM, Willie JT, Sinton CM, Elmquist JK, Scammell T, Lee C,	89. Billiard M. Narcolepsy: Current treatment options and future approaches.	4
7	<i>et al.</i> Narcolepsy in orexin knockout mice: Molecular genetics of sleep	<i>Neuropsychiatr Dis Treat</i> 2008;4:557-66.	5
8	regulation. <i>Cell</i> 1999;98:437-51.	90. Vignatelli L, D'Alessandro R, Candelise L. Antidepressant drugs for	5
9	87. de Lecea L, Kilduff TS, Peyron C, Gao X, Foye PE, Danielson PE, <i>et al.</i>	narcolepsy. <i>Cochrane Database Syst Rev</i> 2008;???:CD003724.	6
10	The hypocretins: Hypothalamus-specific peptides with neuroexcitatory	91. Mayer G. The use of sodium oxybate to treat narcolepsy. <i>Expert Rev</i>	7
11		<i>Neurother</i> 2012;12:519-29.	8
12			8
13			9
14			10
15			11
16			12
17			13
18			14
19			15
20			16
21			17
22			18
23			19
24		Author Queries???	19
25		AQ1: Kindly provide department.	20
26		AQ2: Please check and confirm if the edit made to the	21
27		sentence retains the intended sense and amend if	22
28		necessary.	23
29		AQ3: Kindly review the sentence.	24
30		AQ4: Kindly provide publisher location and check the	25
31		publisher name.	26
32		AQ5: Kindly provide volume number.	27
33		AQ6: Please upload copyright form signed by all authors.	28
34			29
35			30
36			31
37			32
38			33
39			34
40			35
41			36
42			37
43			38
44			39
45			40
46			41
47			42
48			43
49			44
50			45
51			46
52			47
53			48
54			49
55			50
56			51
57			52
58			53
59			54

AQ5