

UC Irvine

UC Irvine Previously Published Works

Title

Interaction of CDKN2A and Sun Exposure in the Etiology of Melanoma in the General Population

Permalink

<https://escholarship.org/uc/item/0924m83v>

Journal

Journal of Investigative Dermatology, 131(12)

ISSN

0022-202X

Authors

Berwick, Marianne
Begg, Colin B
Armstrong, Bruce K
[et al.](#)

Publication Date

2011-12-01

DOI

10.1038/jid.2011.235

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Published in final edited form as:

J Invest Dermatol. 2011 December ; 131(12): 2500–2503. doi:10.1038/jid.2011.235.

Interaction of *CDKN2A* and Sun Exposure in the Etiology of Melanoma in the General Population

Marianne Berwick¹, Colin B. Begg², Bruce K. Armstrong³, Anne S. Reiner², Nancy E. Thomas⁴, Linda S. Cook¹, Irene Orlow², Anne Kricke³, Loraine D. Marrett⁵, Stephen B. Gruber⁶, Hoda Anton-Culver⁷, Robert C. Millikan⁴, Richard P. Gallagher⁸, Terry Dwyer⁹, Stefano Rosso¹⁰, Peter A. Kanetsky¹¹, and Julia Lee-Taylor¹² for the GEM Study Group

¹Department of Internal Medicine, University of New Mexico Cancer Center, University of New Mexico, Albuquerque, NM, USA

²Department of Epidemiology and Biostatistics, Memorial Sloan-Kettering Cancer Center, NY, USA

³Queen Elizabeth II Research Institute D02, University of Sydney, Sydney, New South Wales, Australia

⁴Lineberger Comprehensive Cancer Center, Chapel Hill, NC, USA

⁵Cancer Care Ontario, Toronto, Ontario, Canada

⁶Department of Epidemiology, University of Michigan, Ann Arbor, Michigan, USA

⁷Department of Epidemiology, University of California, Irvine, California, USA

⁸British Columbia Cancer Agency, Vancouver, British Columbia, Canada

⁹Menzies Centre, University of Tasmania, Hobart, Tasmania, Australia

¹⁰Piedmont Tumor Registry, Turin, Italy

¹¹Department of Epidemiology, University of Pennsylvania, Philadelphia, PA, USA

¹²Atmospheric Chemistry Division, National Center for Atmospheric Research, Boulder, CO, USA

TO THE EDITOR

A major goal in cancer prevention is to identify genetic and environmental risk factors and determine if they interact to increase risk. Melanoma is an excellent model because intermittent sun exposure is a well supported environmental risk factor for the development of melanoma and a genetic factor, *CDKN2A*, plays a major role in melanoma etiology. We have sequenced *CDKN2A*, the major familial melanoma gene, in 3,624 melanoma patients from nine centers in four countries (Berwick *et al.*, 2006, Orlow *et al.*, 2007), and we have

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Corresponding Author: Marianne Berwick, University of New Mexico, MSC07-4025, Albuquerque, NM 87131-0001, Telephone: 505-272-4369, Fax: 505-272-2570, mberwick@salud.unm.edu.

CONFLICT OF INTEREST

The authors state no conflict of interest.

identified the same patterns of sun exposure in risk for a second primary melanoma (Kricger *et al.*, 2007) as are found for incident, first primary melanomas (Gandini *et al.*, 2005). We therefore wanted to identify whether and how these two important risk factors – sun exposure and *CDKN2A* – might interact to increase melanoma risk.

We identified incident cases of melanoma from eight population-based registries in Australia (New South Wales and Tasmania), Italy (Piedmont area), Canada (British Columbia and Ontario), the United States (New Jersey, North Carolina, and Orange and San Diego Counties) and one hospital center in Michigan, which sees approximately 50% of the melanoma diagnosed in the state. The study design and details of the data collection have been previously published (Begg *et al.*, 2006). Briefly, single primary melanoma (SPM) controls were people diagnosed with an incident first invasive primary melanoma within a defined accrual period of 6 months during the year 2000, and multiple primary melanoma (MPM) cases were people diagnosed with an incident second- or higher-order invasive or *in situ* melanoma during a 3.5 year period from January 1, 2000. Inclusion of *in situ* cases was designed to avoid exclusion of people who could have been diagnosed with an invasive subsequent primary if the *in situ* lesion had not been removed. Participants gave informed consent, donated 4–6 buccal swabs or blood for DNA extraction, and completed questionnaires detailing demographics, phenotypic characteristics, family history of cancer and lifetime sun exposure behavior. The study protocol was approved by the Institutional Review Board of each participating institution. All study procedures adhered to the Helsinki guidelines. Sequencing was conducted for exons 1 α , 2 and 3 of the *CDKN2A* gene, which code for the p16 and or p14ARF proteins as previously described (Berwick *et al.*, 2006; Orlow *et al.* 2007).

Functional *CDKN2A* mutations (Orlow *et al.* 2007) were identified in 30 of 2,469 individuals with single primary melanomas (1.3%) and in 35 of 1,207 individuals with multiple primary melanoma (2.9%) (Berwick *et al.*, 2006). As we previously published, those with *CDKN2A* mutations are significantly younger than those with wildtype *CDKN2A*, more likely to have a family history of melanoma, more likely to have multiple melanomas and more nevi.

We used five measures of sun exposure previously reported by our group as important risk factors for second primary melanomas (Kricger *et al.*, 2007): (1) Ambient erythemal UVR (UVE) exposure at age 10 – a measure of early life sun exposure supported by migrant studies where the effect of ambient sun exposure is greatest in those who migrated before 10–15 years of age, (2) average annual hours on sunny holidays, (3) average annual hours of beach and waterside exposure, (4) lifetime painful or blistering burns, and (5) lifetime painful or blistering burns to the site of the melanoma. The latter four measures reflect an intermittent pattern of sun exposure, usually indicated by recreational exposure, the major form of sun exposure that has been associated with the development of melanoma (Armstrong and Kricger 1993; Gandini *et al.*, 2006). Odds ratios were calculated using logistic regression for unadjusted and adjusted stratified analyses, controlling for age, sex, study center, an age*sex interaction and ability to tan. Multiplicative interactions were assessed using cross product terms. All statistical analyses were undertaken using SAS Statistical Packages Version 9.2 (SAS, Inc, Cary, NC). All statistical tests were two-sided.

We observed no statistically significant multiplicative interactions between any measure of intermittent sun exposure and mutation of *CDKN2A* (Table 1). The stratified analyses show no statistically significant differences in effects of increasing sun exposure between mutation carriers and non-carriers, and the relative risk estimates are generally smaller than for the non-carriers.

These findings suggest that increasing sun exposure may add little to the increased risk of melanoma that is conferred by carriage of a *CDKN2A* mutation, and that those with a mutation are at high risk for the development of melanoma regardless of sun exposure. The adjusted relative risk of melanoma in carriers is 4.2 (95% CI 2.4, 7.7), as reported earlier in Berwick *et al.*, 2006.

Previous efforts to document a gene-environment interaction among *CDKN2A* carriers have reported similar findings. Consistent with the present results, Goldstein *et al.* (1998) did not find that sun exposure (measured as sunburns) was a significant risk factor for melanoma among families with *CDKN2A* mutations. Recently, Cust *et al.* (2011) reported that *CDKN2A* carriers appeared to have the same cumulative risk of melanoma regardless of ambient sun exposure thus suggesting that our observation of risk of second primary melanoma may be generalizable to all primary melanoma. Most studies of *CDKN2A* have focused on smaller populations (e.g., Nielson *et al.*, 2010) or on familial studies (such as Goldstein *et al.*, 1998; Bishop *et al.*, 2002; Cust *et al.*, 2011); GEM is the largest population-based series to sequence *CDKN2A* among individuals diagnosed with melanoma. Clearly, since the prevalence of *CDKN2A* mutations is very low, the ability to analyze the interaction between *CDKN2A* and solar exposure in the general population is necessarily limited, and inferences from such analyses are uncertain.

In summary, our study provides no evidence to suggest that the influences of *CDKN2A* mutational status and sun exposure on melanoma risk are related. In addition, we found little evidence that sun exposure increases the risk of melanoma in carriers, although our sample sizes are too small for a definitive conclusion on this issue. In the absence of further evidence people with *CDKN2A* mutations should receive at least the same sun protection advice as other people with similar phenotypic risk factors.

Acknowledgments

The study was conducted by the GEM Study Group: Marianne Berwick (PI, University of New Mexico), Memorial Sloan-Kettering Cancer Center, New York, NY, USA; Colin Begg (Co-PI), Irene Orlow (Co-Investigator), Urvi Mujumdar (Project Coordinator), Klaus Busam (Dermatopathologist), Pampa Roy (Laboratory Technician). *Study Centers:* The University of Sydney and The Cancer Council New South Wales, Sydney (Australia): Bruce Armstrong (PI), Anne Kricker (co-PI), Melisa Litchfield (Study Coordinator). Menzies Research Institute, University of Tasmania, Hobart (Australia): Terence Dwyer (PI, currently at the Murdoch Childrens Research Institute, Melbourne, Victoria), Paul Tucker (Dermatopathologist), Alison Venn (co-Investigator), Nicola Stephens (Study Coordinator). British Columbia Cancer Agency, Vancouver (Canada): Richard Gallagher (PI), Teresa Switzer (Coordinator). Cancer Care Ontario, Toronto (Canada): Loraine Marrett (PI), Elizabeth Theis (Co-Investigator), Lynn From (Dermatopathologist), Noori Chowdhury (Coordinator), Louise Vanasse (Coordinator). Centro per la Prevenzione Oncologia Torino, Piemonte (Italy): Stefano Rosso (PI), Roberto Zanetti (coPI), Carlotta Sacerdote (Coordinator). University of California, Irvine (USA): Hoda Anton-Culver (PI), Nancy Leighton (Coordinator). University of Michigan, Ann Arbor (USA): Stephen Gruber (PI), Joanne Jeter (Coordinator). New Jersey Department of Health and Senior Services, Trenton (USA): Judith Klotz (PI), Homer Wilcox (Co-PI), Helen Weiss (Coordinator). University of North Carolina, Chapel Hill (USA): Robert Millikan (PI), Nancy Thomas (Co-Investigator), Dianne Mattingly (Coordinator), Jon Player (Laboratory Technician). University of Pennsylvania,

Philadelphia, PA (USA): Timothy Rebbeck (PI), Peter Kanetsky (Co-Investigator), Amy Walker (Laboratory Manager), Saarene Panossian (Laboratory Technician). *Consultants*: Julia Lee Taylor and Sasha Madronich, National Centre for Atmospheric Research, Boulder, Colorado (USA).

This work is supported by the National Cancer Institute, awards CA 83180 and CA 112524 to MB. The National Center for Atmospheric Research is sponsored by the National Science Foundation.

Abbreviations

OR	odds ratio
CI	Confidence Interval
UVR	ultraviolet radiation
UVE	erythematous ultraviolet radiation
KJ/m²	kilojoules per square meter
MPM	multiple primary melanoma
SPM	single primary melanoma

References

1. Begg CB, Hummer AJ, Mujumdar U, et al. A design for cancer case-control studies using only incident cases: experience with the GEM study of melanoma. *Int J Epidemiol.* 2006; 35:756–64. [PubMed: 16556646]
2. Begg CB, Orlow I, Hummer AJ, et al. Lifetime risk of melanoma in CDKN2A mutation carriers in a population-based sample. *J Natl Cancer Inst.* 2005; 97:1507–15. [PubMed: 16234564]
3. Berwick M, Orlow I, Hummer AJ, et al. The prevalence of CDKN2A germ-line mutations and relative risk for cutaneous malignant melanoma: an international population-based study. *Cancer Epidemiol Biomarkers Prev.* 2006; 15:1520–5. [PubMed: 16896043]
4. Cust AE, Harlnad M, Makalic E, et al. Melanoma risk for CDKN2A mutation carriers who are relatives of population-based case carriers in Australia and the UK. *J Med Genet.* 2011; 48:266–72. [PubMed: 21325014]
5. Bishop DT, Demenais F, Goldstein AM, et al. Geographical variation in the penetrance of CDKN2A mutations for melanoma. *J Natl Cancer Inst.* 2002; 94:894–903. [PubMed: 12072543]
6. Gandini S, Sera F, Cattaruzza MS, et al. Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer.* 2005; 41:45–60. [PubMed: 15617990]
7. Goldstein AM, Falk RT, Fraser MC, et al. Sun-related risk factors for melanoma-prone families with CDKN2A mutations. *J Natl Cancer Inst.* 1998; 90:709–711. [PubMed: 9586669]
8. Kricger A, Armstrong BK, Goumas C, et al. Ambient UV, personal sun exposure and risk of multiple primary melanomas. *Cancer Causes Control.* 2007; 18:295–304. [PubMed: 17206532]
9. Nielsen K, Harbst K, Måsbäck A. Swedish CDKN2A mutation carriers do not present the atypical mole syndrome phenotype. *Melanoma Res.* 2010; 20:266–72. [PubMed: 20526219]
10. Orlow I, Begg CB, Cotignola J, et al. CDKN2A germline mutations in individuals with cutaneous malignant melanoma. *J Invest Dermatol.* 2007; 127:1234–43. [PubMed: 17218939]

Table 1

Analyses of the associations of multiple primary melanoma (MPM) with sun exposure measures stratified by CDKN2A mutational status.

Mutational Status	Exposure	SFM N=2469	MPM n=1207	OR _{crude} (95% CI)	* OR _{adjusted} (95% CI)	P ¹	P ²
UVE at age 10 (KJ/m ²)							
No mutation	303-912	1230	463	1.00	1.00		
	913-2084	1042	643	1.64 (1.42, 1.89)	1.26 (0.95, 1.67)		
	missing	167	66				
With mutation	303-912	17	15	1.00	1.0		
	913-2084	12	20	1.89 (0.69, 5.12)	1.18 (0.39, 3.54)	0.92	0.83
	missing	1	0				
Sunny Holiday – Average Hours per Year							
No mutation	0-19	1013	406	1.00	1.00		
	20-678	969	454	1.17 (0.99, 1.37)	1.33 (1.11, 1.59)		
	missing	457	312				
With mutation	0-19	12	11	1.00	1.00		
	20-678	14	16	1.25 (0.42, 3.69)	0.92 (0.28, 2.99)	0.55	0.24
	missing	4	8				
Beach and Waterside Activities – Average Hours per Year							
No mutation	0-24	1203	570	1.00	1.00		
	25-1857	1164	575	1.04 (0.91, 1.20)	1.20 (1.02, 1.41)		
	Missing	72	27				
With mutation	0-24	7	10	1.00	1.00		
	25-1857	23	25	0.76 (0.25 2.33)	0.87 (0.26, 2.85)	0.59	0.41
	Missing	0	0				
Lifetime Painful or Blistering Sunburns							
No mutation	0-10	1362	613	1.00	1.0		
	11-700	1023	533	1.16 (1.01, 1.33)	1.33 (1.13, 1.56)		
	missing	54	26				
With mutation	0-10	18	21	1.00	1.00		
	11-700	12	14	1.00 (0.37, 2.71)	0.99 (0.33, 2.91)	0.59	0.25
	missing	0	0				

Mutational Status	Exposure	SPM N=2469	MPM n=1207	OR _{crude} (95% CI)	* OR _{adjusted} (95% CI)	I ¹	p ²
Lifetime Painful or Blistering Burns to the Site of the Melanoma							
No mutation	None	1310	522	1.00	1.00		
	Any	945	544	1.45 (1.25, 1.67)	1.49 (1.27, 1.77)		
	missing	184	106				
With mutation	None	13	22	1.00	1.00		
	Any	11	13	0.69 (0.25, 1.94)	0.57 (0.19, 1.74)	0.09	0.09
	missing	6	0				

* Analyses are adjusted for age, sex, age-sex interaction, center and ability to tan.

¹ Multiplicative interaction using categories of sun exposure.

² Multiplicative interaction using continuous values of sun exposure.