

UC Berkeley

UC Berkeley Previously Published Works

Title

Thyroid Hormone Regulates the mRNA Expression of Small Heterodimer Partner through Liver Receptor Homolog-1

Permalink

<https://escholarship.org/uc/item/09z0p3rm>

Journal

Endocrinology and Metabolism, 30(4)

ISSN

2093-596X

Authors

Ahn, Hwa Young
Kim, Hwan Hee
Kim, Ye An
et al.

Publication Date

2015

DOI

10.3803/enm.2015.30.4.584

Peer reviewed



Thyroid Hormone Regulates the mRNA Expression of Small Heterodimer Partner through Liver Receptor Homolog-1

Hwa Young Ahn^{1,2}, Hwan Hee Kim¹, Ye An Kim¹, Min Kim¹, Jung Hun Ohn¹, Sung Soo Chung¹, Yoon-Kwang Lee³, Do Joon Park¹, Kyong Soo Park¹, David D. Moore³, Young Joo Park¹

¹Department of Internal Medicine, Seoul National University College of Medicine; ²Department of Internal Medicine, Chung-Ang University College of Medicine, Seoul, Korea; ³Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX, USA

Background: Expression of hepatic cholesterol 7 α -hydroxylase (CYP7A1) is negatively regulated by orphan nuclear receptor small heterodimer partner (SHP). In this study, we aimed to find whether thyroid hormone regulates SHP expression by modulating the transcriptional activities of liver receptor homolog-1 (LRH-1).

Methods: We injected thyroid hormone (triiodothyronine, T3) to C57BL/6J wild type. RNA was isolated from mouse liver and used for microarray analysis and quantitative real-time polymerase chain reaction (PCR). Human hepatoma cell and primary hepatocytes from mouse liver were used to confirm the effect of T3 *in vitro*. Promoter assay and electrophoretic mobility-shift assay (EMSA) were also performed using human hepatoma cell line.

Results: Initial microarray results indicated that SHP expression is markedly decreased in livers of T3 treated mice. We confirmed that T3 repressed SHP expression in the liver of mice as well as in mouse primary hepatocytes and human hepatoma cells by real-time PCR analysis. LRH-1 increased the promoter activity of SHP; however, this increased activity was markedly decreased after thyroid hormone receptor β /retinoid X receptor α /T3 administration. EMSA revealed that T3 inhibits specific LRH-1 DNA binding.

Conclusion: We found that thyroid hormone regulates the expression of SHP mRNA through interference with the transcription factor, LRH-1.

Keywords: Small heterodimer partner; Thyroid hormones; Cholesterol; Liver receptor homolog-1; Bile acids and salts

Received: 3 April 2015, Revised: 24 August 2015,

Accepted: 24 September 2015

Corresponding authors: Young Joo Park

Department of Internal Medicine, Seoul National University College of Medicine, 101 Daehak-ro, Jongno-gu, Seoul 03080, Korea

Tel: +82-2-2072-4183, Fax: +82-2-764-2199, E-mail: yjparkmd@snu.ac.kr

David D. Moore

Department of Molecular and Cellular Biology, Baylor College of Medicine, One Baylor Plaza, Houston, Texas 77030, USA

Tel: +1-713-798-3313, Fax: +1-713-798-3017, E-mail: moore@bcm.tmc.edu

Copyright © 2015 Korean Endocrine Society

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Small heterodimer partner (SHP) is an orphan nuclear receptor that, unlike other nuclear receptors, contains a putative ligand binding domain but lacks a classical DNA binding domain [1]. SHP mRNA is predominantly expressed in the liver, but it is also expressed in the adrenal gland, spleen, small intestine [2], and pancreas [3]. SHP has also been shown to interact with other nuclear receptors and transcription factors [4].

The expression of SHP is regulated by several other nuclear receptors. The orphan nuclear receptor, steroidogenic factor 1 (SF-1) and its liver homologue, liver receptor homolog-1 (LRH-1) transactivate the SHP promoter [5]. To date, at least 5 SF-1 binding sites have been identified in the promoter region of SHP [5], and LRH-1 is essential for basal SHP expression. When SHP protein expression is elevated by LRH-1, SHP forms a heterodimeric SHP/LRH-1 complex and this complex inactivates LRH-1 thus reducing SHP expression, which is an established auto-regulatory negative feedback loop for SHP [6]. Farnesoid X receptor (FXR) is another well-known inducer of SHP gene expression; an FXR binding site is located in the SHP promoter region [7]. In addition, liver X receptor α (LXR α) and hepatocyte nuclear factor 4 α (HNF4 α) directly regulate SHP promoter activity, and their respective binding sites have been previously identified [8,9]. Other nuclear receptors and inducers, such as estrogen receptor-related receptor γ and sterol regulatory element binding protein-1 (SREBP-1), have also been known to regulate SHP expression [4].

In the liver, SHP is known to perform several metabolic functions. It regulates bile acid synthesis and cholesterol metabolism by modulating cholesterol 7 α -hydroxylase (CYP7A1) expression through inhibition of LRH-1 and HNF4 α activity [10]. In addition, SHP regulates lipogenesis by inhibiting SREBP-1c expression via LXR [11], and serum triglyceride levels by repressing microsomal triglyceride transfer protein (MTP) expression by inhibiting LRH-1 binding to the MTP promoter [12].

Thyroid hormone is known to regulate cholesterol and bile acid metabolism, mainly through regulation of 3-hydroxy-3-methyl-glutaryl-CoA (HMG-CoA) reductase [13], low density lipoprotein (LDL) receptor [14], and CYP7A1 gene expression [15]. SHP has also been shown to play an important role in cholesterol metabolism [6] in the liver, mainly through transcriptional regulation of the CYP7A1 gene, which suggests a possible relationship or interaction between SHP and thyroid hormone. In our preliminary data using microarray analysis, we

found that SHP expression was decreased by thyroid hormone.

Thyroid hormone acts mainly through its nuclear receptors, thyroid hormone receptor (TR) α and β . TR β 1 is putatively expressed in the liver [16]. The functional TR complex consists of a heterodimer with retinoid X receptor (RXR). This complex then binds to a thyroid hormone response element (TRE) and regulates gene expression [17]. Although we observed that thyroid hormone inhibited SHP expression, we were unable to identify TR binding to a TRE in the SHP promoter. However, TR could function through interactions with other nuclear receptors, like peroxisome proliferator-activated receptor (PPAR) and LXR [16,18] at the level of transcription as well as by direct binding to TREs in the promoter region. Therefore, we speculated that TR β might regulate SHP expression by interacting with other transcription factors that regulate SHP promoter. To confirm this hypothesis, we performed this study to find whether thyroid hormone regulates SHP expression by modulating the transcriptional activities of LRH-1, which is required for basal SHP promoter activity in the liver.

METHODS

Animals

C57BL/6J wild type mice were used. Mice were housed in groups of 4 or 5 in plastic microisolator cages at 22°C with a 12-hour light/dark cycle. All animals were provided laboratory chow diet (Purina irradiated laboratory chow 38057, Purina Korea, Seoul, Korea) and water *ad libitum*. Animals were divided into 3 groups according to the duration of thyroid hormone treatment ($n=4$ or $n=5$ in each group). All experiments were performed in triplicate to confirm the results. Thyroid hormone, 3,5,3'-triiodothyronine (T3; Sigma Chemical Co., St. Louis, MO, USA) was prepared at a concentration of 1 mg/kg body weight in 20% DMSO and was administered via intraperitoneal injection. Six hours group: T3 treated group was administered T3 6 hours before sacrifice. Five days group: the T3 treated group was administered T3 once daily for 5 days. Animals were sacrificed 24 hours after the last T3 injection. All animals were sacrificed after fasting for 6 hours beginning at 6:00 AM. Mice were anesthetized by an intraperitoneal injection of Zoletil (Virvac, Carros, France). Blood was drawn by inferior vena cava puncture. The liver was quickly removed, frozen in liquid nitrogen, and then used for RNA extraction. All procedures were performed in accordance with the guidelines of the Seoul National University Bundang Hospital Animal Policy and Welfare Committee.

Cholesterol measurement

Serum was prepared from whole blood by centrifugation at $1,200 \times g$ for 15 minutes. Serum total cholesterol was measured with an enzymatic assay kit (Thermo DMA Inc., St. Louis, CO, USA) following the manufacturer's protocol.

Cell culture and transient transfection

Hepatocytes were isolated from wild type mice. Animals were anesthetized with Zoletil. The inferior vena cava was cannulated, and the portal vein was severed. Collagenase dissolved in Hanks' balanced salt solution (GIBCO Laboratories, Grand Island, NY, USA) was perfused through the liver at 37°C to disperse hepatocytes, and then the hepatocytes were further purified by Percoll (Sigma). The cells were seeded into collagen-coated 6-well plates at density of 1 to 2×10^6 cells/well in 2 mL of William's E medium supplemented with 10% fetal calf serum, penicillin (100 U/mL), streptomycin (0.1 mg/mL), insulin (10 mg/mL), and triamcinolone (1 nmol/L). We also used HepG2 and HEK293 cells. Cells were grown at 37°C in a humidified atmosphere of 5% $\text{CO}_2/95\%$ air. HepG2 cells were maintained in modified Eagle's medium plus 10% fetal bovine serum (FBS). T3 (100 nmol/L) was incubated with primary hepatocytes and HepG2 cells according to the response time (control 2, and 6 hours). After T3 treatment, RNA was isolated from the cells.

One day before transfection, confluent cells were trypsinized and plated into 6-well plates at a 1:4 ratio to allow the cells to reach 50% to 60% confluency at the time of transfection. Six hours before transfection, fresh medium containing 10% charcoal-stripped FBS was added to the cells. All transfections were performed using the calcium phosphate method. Luciferase values shown are the averages of triplicate samples. All plasmids used in this study were previously reported. Various 5' deletions of the human SHP promoter [5], the human SHP promoter [19], LRH-1 [19], TR β [20], and RXR α [21] were used.

RNA isolation and quantitative real-time polymerase chain reaction

Total RNA was isolated from frozen liver samples and cells using TRIzol Reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instruction. First-strand cDNA was synthesized from 1 μg of RNA using superscript II reverse transcriptase (Invitrogen). Quantitative real-time polymerase chain reaction (PCR) was performed using SYBR Green PCR master mix and an ABI prism 7500 Sequence Detection System (Ap-

plied Biosystems, Foster City, CA, USA). The PCR primers used for each gene are listed below (5' to 3'): mouse SHP-F (forward), TTG CAC CTG CAT CTC ACA GC; SHP-R (reverse), TCT TGG CTA GGA CAT CCA AG; glyceraldehyde 3-phosphate dehydrogenase (GAPDH)-F, TGT GTC CGT CGT GGATCT GA; GAPDH-R, CCT GCT TCA CCA CCT TCT TGA; human SHP-F, GGA CTT CCT TGG TTT GGA CA; SHP-R, CTC ATC CCA AGA AGG GAC AG; GAPDH-F, GAT CAT CAG CAA TGC CTC CT; GAPDH-R, TGT GGT CAT GAG TCC TTC CA. Relative mRNA expression was quantified using the comparative cycle threshold (Ct) method and expressed as $2^{-\Delta\Delta\text{Ct}}$.

Identification of differentially expressed genes

RNA was hybridized to Affymetrix GeneChip Human Gene 1.0 ST Arrays (Affymetrix, Santa Clara, CA, USA). Robust multichip analysis was used to normalize the expression data in the Expression Console software (Affymetrix) [22]. We calculated the log fold change of each probe in different conditions and independent samples. Student *t* test was performed to determine the significance of differential expression using GenePattern software [23].

Electrophoretic mobility-shift assay

Double-stranded oligonucleotides of the following sequences were used as probes: LRH-1 response element (LRE) in the human SHP promoter (LRE WT (-518 to -489), 5'-GAAGCA-GGGCCCAAGGTTAGGCAACAAG-3', LRE is underlined); LRE mutant competitor (LRE MT, 5'-GAAGCAGGGCCCttccaaAGCAACAAG-3'); consensus LRE (5'-GACTTCTGGAGTCAAGGTTGTTGGGCCATTC-3'). The probes were labeled with [α - ^{32}P] dATP using Klenow polymerase (Roche, Mannheim, Germany), and the labeled probe (35,000 cpm) was incubated with nuclear extracts of HepG2 cells treated with 100 nmol/L T3 in 10 mmol/L Hepes (pH 7.9) containing 50 mmol/L KCl, 0.1 mmol/L ethylenediaminetetraacetic acid, 0.25 mmol/L DTT, 0.1 mg/mL poly (dIdC), 0.01% nonidet P-40, and 10% glycerol at room temperature for 10 minutes. Competitors were added in 50-fold molar excess to the labeled probe.

Statistical analysis

Values are expressed as the mean \pm SD. The significance of differences between mean values was evaluated using Student *t* test. A $P < 0.05$ was considered statistically significant.

RESULTS

Thyroid hormone regulates the expression of genes involved in bile acid synthesis and cholesterol metabolism

Before conducting the microarray analyses, to confirm the effect of thyroid hormone on serum cholesterol, we measured serum cholesterol levels after intraperitoneal administration of T3 (1 mg/kg) to mice for 5 days. Serum total cholesterol levels decreased significantly after T3 injection (101.7 ± 12.0 mg/dL vs. 47.0 ± 9.7 mg/dL, $P < 0.05$). We profiled the transcriptional changes in mouse liver before and after 5 days of T3 treatment and examined the expression of genes related to the regulation of cholesterol and bile acid metabolism.

In microarray analyses (Table 1), genes related to cholesterol metabolism, HMG-CoA synthase and reductase was up-regulated by thyroid hormone. Among genes associated with high density lipoprotein cholesterol, apolipoprotein A-1 was up-regulated and scavenger receptor was down-regulated by thyroid hormone. However, the LDL receptor, the main receptor responsible for clearing cholesterol-laden LDL particles from the blood, was not significantly up-regulated by thyroid hormone, which is different from the results of previous studies [14,24].

Among the genes related to bile acid synthesis, CYP7A1 and mitochondrial sterol 27 hydroxylase (Cyp27a1) were up-regulated after T3 treatment, although the fold change of Cyp27a1 was small. In a previous study, thyroxine enhanced CYP7A1 expression but not Cyp27a1 expression in rat hepatocytes when added to dexamethasone-containing medium [25].

The expression of CYP7A1 and SHP was regulated by several transcription factors, such as FXR, LRH-1, HNF-4 α , and LXR α [4,26]. However, expression of these transcription factors did not significantly change in response to thyroid hormone in microarray analysis (Table 1). In contrast, the expression of SHP was markedly decreased by thyroid hormone; therefore, we assumed that thyroid hormone regulates SHP expression independent of other transcription factors, and might regulate A expression indirectly through changes in SHP expression.

To confirm this, we examined the changes in SHP mRNA level after T3 in mouse liver by real-time PCR. Level of serum T3 was significantly increased in 6-hour and 5-day group than control (231.4 ± 42.4 and 310.2 ± 65.8 ng/dL vs. 70.1 ± 20.7 ng/dL). As expected, the expression of SHP decreased after treatment with T3 for 5 days (Fig. 1A).

Table 1. Changes in Gene Expression after 5 Days of T3 Treatment in Mouse Liver

Gene symbol	Gene description	T3 treatment vs. control		
		Log fold change ^a	T-score	P value
Bile acid synthesis				
CYP7A1	Cytochrome P450 7A	1.008	3.477	0.025
CYP8B1	Sterol 12 α -hydroxylase	-0.364	-3.465	0.026
CYP27A1	Sterol 27-hydroxylase	0.500	3.784	0.020
Nuclear receptors				
SHP	Small heterodimer partner	-0.698	-7.061	0.002
FXR	Farnesoid X receptor	0.152	1.624	0.176
LXR α	Liver X receptor α	0.465	2.266	0.083
TR β	Thyroid hormone receptor β	0.119	1.423	0.225
HNF4 α	Hepatocyte nuclear receptor 4 α	-0.115	-1.955	0.118
LRH-1	Liver receptor homolog-1	-0.128	-1.623	0.177
Cholesterol metabolism				
HMG-CoA R	3-Hydroxy-3-methylglutaryl-coenzyme A reductase	1.516	3.935	0.018
HMG-CoA S1	3-Hydroxy-3-methylglutaryl-coenzyme A synthase 1	1.252	5.776	0.005
HMG-CoA S2	3-Hydroxy-3-methylglutaryl-coenzyme A synthase 2	0.074	0.699	0.529
SREBP1	Sterol regulatory element binding protein 1	-0.593	-2.543	0.062
VLDLR	Very low density lipoprotein receptor	-0.278	-1.301	0.262
SR-B1	Scavenger receptor	-0.792	-4.942	0.009
APOA1	Apolipoprotein A1	0.201	5.047	0.007
APOB	Apolipoprotein B	0.077	0.158	0.884
APOE	Apolipoprotein E	-0.025	-0.864	0.440
LDLR	Low density lipoprotein receptor	0.386	1.007	0.373

^aThe fold changes are shown in log scale.




Fig. 1. Effect of thyroid hormone (T3) treatment on small heterodimer partner (SHP) expression. Comparison of SHP expression in mouse liver (A, $n=4$ or $n=5$ in each group) after 6 hours or 5 days of thyroid hormone treatment (1 mg/g). Comparison of SHP expression in mouse primary hepatocytes (B) and human hepatoma cells (C) after 2 and 6 hours of T3 treatment (100 nmol/L). ^aSignificant differences compared to the control are $P < 0.05$.




Fig. 2. Effect of thyroid hormone (T3) on small heterodimer partner (SHP) promoter activity with or without thyroid hormone receptor (TR)/retinoid X receptor (RXR), and liver receptor homolog-1 (LRH-1). In each HepG2 cell sample, 300 ng of mouse SHP (A) or human SHP (B, C) promoter DNA was co-transfected with or without 75 ng of TR β and 75 ng of RXR α or 75 ng of LRH-1. Vehicle or T3 (100 nmol/L) was administered for 24 hours to determine the effect of thyroid hormone on SHP promoter expression. Luciferase activity was measured and normalized to β -galactosidase activity. Significant differences compared to the control are ^a $P < 0.05$ and ^b $P < 0.01$.

To confirm the specific regulatory effects of T3 on the expression of SHP, an *in vitro* study was performed using freshly isolated primary hepatocytes and human hepatoma HepG2 cells. Hepatocytes from C57BL/6L mice were isolated, and T3 (100 nmol/L) or vehicle was added to the culture media. As shown Fig. 1B, the expression of SHP decreased at 2 and 6 hours after T3 treatment. In HepG2 cells, T3 also decreased SHP expression after 6 hours of T3 treatment (Fig. 1C). Although the time sequence differed, T3 decreased the expression of SHP in both human and mouse; therefore, we evaluated the mechanism underlying the regulation of SHP by T3.

TR β /RXR α plus T3 decreases SHP promoter activity by interfering with LRH-1

We used full-length (~2 Kb) mouse and human SHP promoter constructs [5] to determine whether TR β affects the activity of the SHP promoter. We cotransfected HepG2 cells with SHP promoters along with an empty or TR β /RXR α expression vector and evaluated the effect of vehicle or T3 (100 nmol/L) administered over 24 hours on promoter activity. T3 treatment alone did not have a significant effect on mouse or human SHP promoter activity (Fig. 2A, B). When the SHP promoter was cotransfected with TR β /RXR α in the absence of T3, SHP promoter expression increased. However, this increased activity decreased significantly after administration of T3 (Fig. 2A, B). This result suggests that unliganded TR β /RXR α increases the activity of the SHP promoter and T3/TR β /RXR α repressed activity of SHP.

Chromatin immunoprecipitation and high throughput sequencing (ChIP-Seq) was used to define genome wide TR β binding sites in the liver [27]. In accord with the negative results of bioinformatics searches for conserved TREs in the human and mouse SHP promoters, no such sites were observed. Therefore, we presumed that the T3/TR β /RXR α complex might interact with other SHP promoter transcription factors. When we transfected the SHP promoter along with FXR, HNF4, or LRH-1, SHP promoter activity was increased (data not shown). Among these transcription factors, LRH-1 induced SHP promoter activity most strongly. LRH-1 is an abundant transcription factor in the liver; therefore, we checked for the interaction between LRH-1 and the T3/TR β /RXR α complex using the human SHP promoter construct (Fig. 2C). Interestingly, LRH-1 and unliganded TR β /RXR α markedly increased SHP promoter activity; however, when we added T3, promoter activity decreased significantly (Fig. 2C).

We then tested the functional interaction of TR and LRH-1




Fig. 3. Repression of expression from the 5' deletion small heterodimer partner (SHP) promoter by the thyroid hormone. The 200 ng of 5' human SHP promoter (full, -1490 and -175) was cotransfected with or without 100 ng of thyroid hormone receptor β (TR β) and 30 ng of retinoid X receptor α (RXR α) or 100 ng of liver receptor homolog-1 (LRH-1) in 293 cells. Thyroid hormone (T3; 100 nmol/L) was added. Luciferase activity was measured and normalized against renilla activity. NS, not significant. Significant differences compared to the control ^a $P < 0.01$.

using SHP promoter deletion constructs affecting LRH-1 binding sites [5]. To check the interaction, we used 3 fragments (-2080, -1490, and -175) of human SHP promoter in our study. As LRH-1 response element (LRE) in the human SHP promoter was located from -518 to -489, the activation of promoter by unliganded TR β /RXR α /LRH-1 was decreased in -175 fragments and when we added T3, overall activity of human SHP promoter by TR β /RXR α /LRH-1 was significantly reduced (Fig. 3).

Electrophoretic mobility-shift assays were performed with a double-stranded oligonucleotide containing the -518 human SHP promoter LRE to explore the mechanism of the T3 dependent inhibition of LRH-1 transactivation (Fig. 4). The affinity of LRH-1 binding decreased in nuclear extracts of human hepatoma cells treated with T3 for increasing times up to 24 hours (lanes 2 to 5). To verify the specific binding of LRH-1 to the probe, competition assays were also performed. An oligomer containing a mutated LRE sequence did not compete with the probe (lane 9), whereas an oligomer with the consensus LRE competed with the SHP LRE probe (lane 10).




Fig. 4. Interaction between thyroid hormone (T3) and liver receptor homolog-1 (LRH-1). Electrophoretic mobility-shift assay was performed using an oligomer containing the LRH-1 responsive element (LRE) in the human SHP promoter (-518) as a probe and nuclear extracts of human hepatoma cells treated with 100 nmol/L of T3 for increasing times as indicated. For competition assays, unlabeled oligomers containing the hLRH-1(-518) LRE (self; lane 8), an LRE mutant (lane 9), or a consensus LRE (lane 10) were used in a 50-fold molar excess. MT, mutant.

DISCUSSION

In this study, we demonstrated that liganded TR β suppressed the expression of SHP. Because there is no known TRE in SHP promoter, liganded TR β might decrease the expression of SHP by disturbing the transcriptional activity of LRH-1 to SHP. Therefore, our result could be one mechanism of regulation of CYP7A1 expression by thyroid hormone and TR β .

In fact, role of thyroid hormone to regulation of CYP7A1 was already studied in several previous studies. Murine CYP7A1 is highly inducible by thyroid hormone because there are TREs in the murine CYP7A1 promoter [28]. Therefore, thyroid hormone directly increases the expression of murine CYP7A1 and induces bile acid synthesis. In human, there are inconsistent results about regulation of CYP7A1 expression by thyroid hormone. Previously, human CYP7A1 expression was reduced by thyroid hormone in several studies [29,30]. Drover et al. [30] found that, unlike the CYP7A1 promoter site II TRE, the

sequence of the site III TRE is not conserved among different species, which might explain the divergent responses of human and murine CYP7A1 to thyroid hormone. In contrast, recent study showed that thyroid hormone increased expression of CYP7A1 similarly in human and mouse primary liver culture, human hepatocytes and mouse liver [31]. However, it is difficult to explain until now why several researchers announced different results about thyroid hormone and induction of CYP7A1. For this reason, the relationship between CYP7A1 and thyroid hormone it is not sufficient to explain the regulation of bile acid metabolism by thyroid hormone. However, in our study, SHP expression in the mouse (both *in vivo* and *in vitro*) and human hepatoma cells was identically decreased by thyroid hormone. Therefore, our results are important to determine the mechanism underlying bile acid metabolism regulation, as it is the first study to report repression of SHP expression by thyroid hormone.

Many potential transcription factors for SHP, such as LRH-1, HNF-4, FXR, LXR, and SREPB-1 [4] were identified in previous studies. Among these, LRH-1 induces strong transcriptional activity because at least 5 LRH-1 binding sites were identified in the SHP promoter [5].

The results of ChIP-Seq, TRE bioinformatics and direct cotransfection all indicate that TR β /RXR α does not regulate the SHP promoter directly. Among a number of transcription factors that are known to have such direct effects, LRH-1 has most binding sites and is essential for basal SHP promoter activity. In cotransfections of LRH-1 and TR β /RXR α , we observed a T3-dependent suppression of promoter activity. This effect was diminished in -175 fragments of human SHP promoter. We also observed a time-dependent decrease in specific LRH-1 DNA binding in T3 treated livers, but the basis for this effect remains to be determined.

Thyroid hormone can act as both a positive and negative regulator of gene expression. In positive regulation, TR binds to TREs located in the promoters of target genes. In the absence of T3, the unliganded TR/RXR complex with corepressors inhibits the transcriptional activity of target genes. Binding of T3 to TR induces structural changes and recruits coregulatory proteins, which enables transcriptional activation [17]. TRs also negatively regulate transcription with and without DNA binding. In contrast to positive regulation, when TR binds to specific negative TREs, liganded TR recruits a corepressor complex to repress transcriptional activity. Another model suggests that TR does not bind to DNA directly, but instead binds to another transcription factor via protein-protein interactions.

Then, liganded TR recruits a corepressor complex to inhibit transcriptional activity [32]. In contrast, unliganded TR binds to specific transcription factor and recruits a coactivator complex (CoA) leading to gene activation. The increased activity of SHP promoter by adding LRH-1 to unliganded TR β /RXR α as shown Fig. 2C could be explained by this mechanism, although it is not clearly known yet from our results.

In our study, we assumed that TR β did not bind to DNA directly, but interacted with another transcription factor, LRH-1. This interaction of TR with to other transcription factors has been demonstrated in previous studies; both TR β and LXR α heterodimerize with RXR α and compete for binding to DR4 elements in the CYP7A1 promoter [16]. In addition, PPAR selectively inhibits the transcriptional activity of TRs by competing for RXR and possibly non-RXR TR-auxiliary proteins [18]. Therefore, transcriptional regulation by TR through interaction with other transcriptional factors might be as important as regulation through classic TREs.

In conclusion, we found that thyroid hormone indirectly regulates the expression of SHP mRNA by interfering with the transcriptional activity of another transcription factor, LRH-1. This provides an additional mechanism to better understanding of the regulation of bile acid and cholesterol metabolism regulation by thyroid hormone.

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This study was supported by a grant from the Korean Health Technology R&D Project, the Ministry of Health and Welfare, Republic of Korea (A100589) and a grant from Seoul National University Bundang Hospital (03-2010-015). We thank Stephen D. Ayers from Methodist Hospital Research Institute for his assistance with data interpretation.

REFERENCES

- Seol W, Choi HS, Moore DD. An orphan nuclear hormone receptor that lacks a DNA binding domain and heterodimerizes with other receptors. *Science* 1996;272:1336-9.
- Lee HK, Lee YK, Park SH, Kim YS, Park SH, Lee JW, et al. Structure and expression of the orphan nuclear receptor SHP gene. *J Biol Chem* 1998;273:14398-402.
- Sanyal S, Kim JY, Kim HJ, Takeda J, Lee YK, Moore DD, et al. Differential regulation of the orphan nuclear receptor small heterodimer partner (SHP) gene promoter by orphan nuclear receptor ERR isoforms. *J Biol Chem* 2002;277:1739-48.
- Lee YS, Chanda D, Sim J, Park YY, Choi HS. Structure and function of the atypical orphan nuclear receptor small heterodimer partner. *Int Rev Cytol* 2007;261:117-58.
- Lee YK, Parker KL, Choi HS, Moore DD. Activation of the promoter of the orphan receptor SHP by orphan receptors that bind DNA as monomers. *J Biol Chem* 1999;274:20869-73.
- Lu TT, Makishima M, Repa JJ, Schoonjans K, Kerr TA, Auwerx J, et al. Molecular basis for feedback regulation of bile acid synthesis by nuclear receptors. *Mol Cell* 2000;6:507-15.
- Goodwin B, Jones SA, Price RR, Watson MA, McKee DD, Moore LB, et al. A regulatory cascade of the nuclear receptors FXR, SHP-1, and LRH-1 represses bile acid biosynthesis. *Mol Cell* 2000;6:517-26.
- Goodwin B, Watson MA, Kim H, Miao J, Kemper JK, Kliewer SA. Differential regulation of rat and human CYP7A1 by the nuclear oxysterol receptor liver X receptor- α . *Mol Endocrinol* 2003;17:386-94.
- Shih DQ, Screenan S, Munoz KN, Philipson L, Pontoglio M, Yaniv M, et al. Loss of HNF-1 α function in mice leads to abnormal expression of genes involved in pancreatic islet development and metabolism. *Diabetes* 2001;50:2472-80.
- Kir S, Zhang Y, Gerard RD, Kliewer SA, Mangelsdorf DJ. Nuclear receptors HNF4 α and LRH-1 cooperate in regulating CYP7A1 in vivo. *J Biol Chem* 2012;287:41334-41.
- Watanabe M, Houten SM, Wang L, Moschetta A, Mangelsdorf DJ, Heyman RA, et al. Bile acids lower triglyceride levels via a pathway involving FXR, SHP, and SREBP-1c. *J Clin Invest* 2004;113:1408-18.
- Huang J, Iqbal J, Saha PK, Liu J, Chan L, Hussain MM, et al. Molecular characterization of the role of orphan receptor small heterodimer partner in development of fatty liver. *Hepatology* 2007;46:147-57.
- Ness GC, Chambers CM. Feedback and hormonal regulation of hepatic 3-hydroxy-3-methylglutaryl coenzyme A reductase: the concept of cholesterol buffering capacity. *Proc Soc Exp Biol Med* 2000;224:8-19.
- Ness GC, Zhao Z. Thyroid hormone rapidly induces hepatic LDL receptor mRNA levels in hypophysectomized rats.

- Arch Biochem Biophys 1994;315:199-202.
15. Ness GC, Pendelton LC, Zhao Z. Thyroid hormone rapidly increases cholesterol 7 alpha-hydroxylase mRNA levels in hypophysectomized rats. *Biochim Biophys Acta* 1994;1214:229-33.
 16. Hashimoto K, Cohen RN, Yamada M, Markan KR, Monden T, Satoh T, et al. Cross-talk between thyroid hormone receptor and liver X receptor regulatory pathways is revealed in a thyroid hormone resistance mouse model. *J Biol Chem* 2006;281:295-302.
 17. Cheng SY, Leonard JL, Davis PJ. Molecular aspects of thyroid hormone actions. *Endocr Rev* 2010;31:139-70.
 18. Juge-Aubry CE, Gorla-Bajszczak A, Pernin A, Lemberger T, Wahli W, Burger AG, et al. Peroxisome proliferator-activated receptor mediates cross-talk with thyroid hormone receptor by competition for retinoid X receptor. Possible role of a leucine zipper-like heptad repeat. *J Biol Chem* 1995;270:18117-22.
 19. Lee YK, Moore DD. Dual mechanisms for repression of the monomeric orphan receptor liver receptor homologous protein-1 by the orphan small heterodimer partner. *J Biol Chem* 2002;277:2463-7.
 20. Park YJ, Lee EK, Lee YK, Park do J, Jang HC, Moore DD. Opposing regulation of cytochrome P450 expression by CAR and PXR in hypothyroid mice. *Toxicol Appl Pharmacol* 2012;263:131-7.
 21. Lee YK, Dell H, Dowhan DH, Hadzopoulou-Cladaras M, Moore DD. The orphan nuclear receptor SHP inhibits hepatocyte nuclear factor 4 and retinoid X receptor transactivation: two mechanisms for repression. *Mol Cell Biol* 2000;20:187-95.
 22. Irizarry RA, Bolstad BM, Collin F, Cope LM, Hobbs B, Speed TP. Summaries of Affymetrix GeneChip probe level data. *Nucleic Acids Res* 2003;31:e15.
 23. Reich M, Liefeld T, Gould J, Lerner J, Tamayo P, Mesirov JP. GenePattern 2.0. *Nat Genet* 2006;38:500-1.
 24. Lopez D, Abisambra Socarras JF, Bedi M, Ness GC. Activation of the hepatic LDL receptor promoter by thyroid hormone. *Biochim Biophys Acta* 2007;1771:1216-25.
 25. Stravitz RT, Vlahcevic ZR, Russell TL, Heizer ML, Avadhani NG, Hylemon PB. Regulation of sterol 27-hydroxylase and an alternative pathway of bile acid biosynthesis in primary cultures of rat hepatocytes. *J Steroid Biochem Mol Biol* 1996;57:337-47.
 26. Gilardi F, Mitro N, Godio C, Scotti E, Caruso D, Crestani M, et al. The pharmacological exploitation of cholesterol 7alpha-hydroxylase, the key enzyme in bile acid synthesis: from binding resins to chromatin remodelling to reduce plasma cholesterol. *Pharmacol Ther* 2007;116:449-72.
 27. Ayers S, Switnicki MP, Angajala A, Lammel J, Arumanayagam AS, Webb P. Genome-wide binding patterns of thyroid hormone receptor beta. *PLoS One* 2014;9:e81186.
 28. Shin DJ, Plateroti M, Samarut J, Osborne TF. Two uniquely arranged thyroid hormone response elements in the far upstream 5' flanking region confer direct thyroid hormone regulation to the murine cholesterol 7alpha hydroxylase gene. *Nucleic Acids Res* 2006;34:3853-61.
 29. Wang DP, Stroup D, Marrapodi M, Crestani M, Galli G, Chiang JY. Transcriptional regulation of the human cholesterol 7 alpha-hydroxylase gene (CYP7A) in HepG2 cells. *J Lipid Res* 1996;37:1831-41.
 30. Drover VA, Wong NC, Agellon LB. A distinct thyroid hormone response element mediates repression of the human cholesterol 7alpha-hydroxylase (CYP7A1) gene promoter. *Mol Endocrinol* 2002;16:14-23.
 31. Lammel Lindemann JA, Angajala A, Engler DA, Webb P, Ayers SD. Thyroid hormone induction of human cholesterol 7 alpha-hydroxylase (CYP7A1) in vitro. *Mol Cell Endocrinol* 2014;388:32-40.
 32. Weitzel JM. To bind or not to bind: how to down-regulate target genes by liganded thyroid hormone receptor? *Thyroid Res* 2008;1:4.