

UC San Diego

UC San Diego Previously Published Works

Title

Increased Fibrogenic Gene Expression in Multifidus Muscles of Patients With Chronic Versus Acute Lumbar Spine Pathology.

Permalink

<https://escholarship.org/uc/item/0bv651ph>

Journal

Spine, 45(4)

ISSN

0362-2436

Authors

Shahidi, Bahar
Fisch, Kathleen M
Gibbons, Michael C
[et al.](#)

Publication Date

2020-02-15

DOI

10.1097/brs.0000000000003243

Peer reviewed



Published in final edited form as:

Spine (Phila Pa 1976). 2020 February 15; 45(4): E189–E195. doi:10.1097/BRS.0000000000003243.

Increased Fibrogenic Gene Expression in Multifidus Muscles of Patients with Chronic Versus Acute Lumbar Spine Pathology

Bahar Shahidi, PT, PhD¹, Kathleen M. Fisch, PhD², Michael C. Gibbons, PhD³, Samuel R. Ward, PT, PhD^{1,3,4}

¹University of California San Diego Department of Orthopaedic Surgery, San Diego USA

²University of California San Diego, Center for Computational Biology & Bioinformatics, Department of Medicine, San Diego, USA

³University of California San Diego Department of Bioengineering, San Diego, USA

⁴University of California San Diego Department of Radiology, San Diego, USA

INTRODUCTION:

Low back pain is a common condition that affects 65–85% of the general population at some point in their lifetime(1). Although the causes of low back pain are complex and multifactorial, individuals with this condition demonstrate degenerative changes in the paraspinal muscles(2, 3). Most of these changes have been characterized on a whole-muscle level using imaging methods such as magnetic resonance imaging (MRI), or computed tomography (CT), and have described both atrophy of the paraspinal muscles in the form of reduced cross sectional area (CSA), as well as increased fatty infiltration throughout the muscles(3-6). On a microstructural scale, histological analyses of paraspinal muscle biopsies in both animal and human studies of lumbar spine degeneration also corroborate these findings(7, 8), with the addition of observations of muscle fiber degeneration(2, 7), fibrotic deposition(2, 9), inflammation(2, 10), and reduced vascularity(2, 11). It is not unusual to see more than 50% of the muscle compartment or sample replaced by fat and/or fibrotic tissue in these studies(2, 3), which is thought to have important implications regarding paraspinal muscle function in its role as a spinal stabilizer. Importantly, the presence of these muscle-specific changes has been associated with reduced patient function, higher levels of disability, higher likelihood of symptom recurrence, and poor post-surgical prognosis(8, 12-16).

These observations have led to additional questions about what is driving the tissue compositional changes of fat and fibrotic infiltration in this population. Some hypothesize that these compositional changes are a secondary consequence of disuse-related atrophy, in which non-muscle tissue fills in the area that used to be occupied by muscle. Alternatively, there is evidence that muscle cells actively degenerate as opposed to simple atrophy, a process that may be facilitated by inflammation or denervation(2, 17). As such, information

on the molecular pathways that drive tissue compositional change may provide key information on the mechanisms of muscle loss in this population, and ultimately guide more effective treatment strategies.

For example, with the clinical assumption that muscle-specific changes are driven by disuse-related atrophy, the primary approach towards mitigating these processes has been to reverse this using treatment such as targeted rehabilitation to facilitate muscle hypertrophy and ultimately restore strength and function. However, if the processes that are driving changes in muscle structure and function change and become more complex over the course of disease progression, then identifying and incorporating treatments that target these specific rate-limiting factors is important in restoring muscle function and ultimately improving clinical prognosis in this patient population.

One of the tools that has been used to investigate molecular drivers of chronic degenerative changes in tissues from a variety of musculoskeletal conditions is gene expression(18). As a tool, gene expression allows for an investigation of individual genes within functional pathways that are thought to contribute to a pathological phenotype. However, expression of a single gene in isolation often oversimplifies a complex process that likely involves several regulatory and feedback pathways. As such, investigations of single genes in the absence of their regulators often provides incomplete information regarding the impact of a single gene on a pathological phenotype. An additional challenge to performing gene expression in human muscle tissue is that profound tissue heterogeneity exists in samples, which can confound transcript abundance, making interpretation difficult (18, 19). Additionally, these tissue compositional changes are thought to progress with chronicity of disease, introducing an additional component of variability to human studies (2).

Currently, most literature utilizing gene expression as a tool for understanding muscle degenerative changes in the spine has utilized animal models such as rabbit and sheep(10, 20, 21), and only one other study exists investigating gene expression of paraspinal muscle in adults with lumbar spine pathology(22). In the aforementioned human study, gene expression from individuals with and without lumbar kyphosis was compared, and was limited to 3 genes primarily related to inflammation.

In light of the paucity of literature investigating molecular pathways of muscle degeneration over the course of human lumbar spine disease, the goal of this study was to investigate gene expression of functional groups of genes that were empirically consistent with the observed morphological phenotypes commonly observed in lumbar spine muscle degeneration. Therefore, multiple genes within the atrophic, myogenic, fibrogenic, adipogenic, and inflammatory pathways were compared between paraspinal muscle biopsies from individuals with acute and chronic lumbar spine pathology. We hypothesized that in atrophic, fibrogenic, and adipogenic pathways would be upregulated, and inflammatory and myogenic pathways would be downregulated in individuals with chronic pathology as compared to acute pathology. The clinical implication of these findings would help identify molecular pathways that are rate-limiting to recovery in this population in order to more appropriately prevent maladaptive changes in muscle as disease progresses.

METHODS:

This was a cross-sectional prospective observational study (Level I) of thirty-three muscle samples (N=8 acute, and 25 chronic) obtained from patients undergoing posterior approach lumbar spine surgery for degenerative lumbar spine pathology. Patients were categorized into the acute group if their duration of symptoms was less than 6 months, and they were categorized into the chronic group if their duration of symptoms was greater than 6 months. All patients underwent an informed consent process to obtain an intraoperative biopsy of the multifidus muscle in the lumbar spine and were included if they were undergoing surgery including decompressions, laminoforaminotomies, discectomies, or fusions. Patients with any diagnosed myopathy or systemic neurological condition were excluded. This study was performed in accordance with Declaration of Helsinki and under approval of the UC San Diego Institutional Review Board (IRB111647). Samples were obtained from a standardized location as previously described (2), split in half upon retrieval, and immediately flash frozen in liquid nitrogen cooled isopentane and stored at -80 degrees until processing. From one half of the sample, ten-micron sections were obtained from OCT-embedded frozen sample using a Leica (CM3050S, Buffalo Grove) cryostat. Gomori Trichrome stains were used to visualize gross muscle morphology and quantify tissue content using ImageJ (<http://imagej.nih.gov/ij>) as previously described(2). The other half of the tissue sample was homogenized and used for RNA extraction and subsequent gene expression analyses.

RNA isolation and quantitative PCR

For gene expression analysis, approximately 30–50 mg of the muscle biopsy was homogenized in bead tubes (Navy, NextAdvance) with TRIZOL (Ambion). RNeasy spin columns (Qiagen) were used to extract RNA using the manufacturer's protocol. One microgram of complimentary DNA (cDNA) was reverse transcribed with iScript cDNA Synthesis kits (Biorad). Quantitative PCR was carried out on custom plates on a BioRad CFX384 Touch qPCR analyzer for a panel of 42 genes associated with myogenic, atrophic, adipogenic, fibrogenic, and inflammatory pathways (Supplementary Table 1), with a cycle threshold determined using a SYBR green fluorophore. On-plate quality assessment was performed to assess gDNA contamination and RNA quality.

Statistical analysis

To determine the influence of tissue composition on RNA yield, concentration values (ng/ul) for yield were regressed against proportions of muscle, fat, and connective tissue from each biopsy sample. Raw cycle-threshold values (Ct values) were obtained from all samples and read into a qPCR expression set using the R Bioconductor package HTqPCR(23), and were quantile normalized to the mean Ct value to obtain gene expression values. A maximum Ct value of 39 was applied to all genes of interest to allow for statistical comparison, with lower values indicating higher expression in this method(23). Tissue composition was included in a linear regression model with normalized gene expression values to account for the potential influence of different tissue types on the genes of interest. Unsupervised hierarchical clustering using Euclidean distance was applied to the normalized expression values to determine the ability of gene expression patterns to differentiate between acute and chronic patient groups, and to identify potential sub-clustering within groups. Differential

expression values (delta-delta-Ct) were calculated with the limmaCtData wrapper in HTqPCR for the Bioconductor package limma using a moderated t-test. All raw p-values were adjusted for within-gene group multiple comparisons using the Benjamini & Hochberg method(24). Significance was set at an adjusted p-value threshold of $p < 0.05$, and trends were defined as adjusted p-values of < 0.08 .

RESULTS:

Tissue composition of the muscle biopsies obtained was heterogeneous, with muscle making up a mean (SD) of approximately 49.6 (18.0)%, adipose tissue making up 14.3 (12.3)%, and connective tissue (fibrosis) making up 21.2 (12.7)% of the tissue based on cross sectional analyses. There were no significant differences in tissue composition or RNA yield between acute and chronic biopsies (Table 1). Similarly, there was no relationship between tissue composition and gene expression values ($p > 0.134$) Unsupervised hierarchical clustering of normalized expression values resulted in two primary clusters that did not differentiate acute samples from chronic samples. There was no obvious sub-clustering segregating functional gene groups (Figure 1). There were no significant differences in gene expression patterns for genes within the atrophic ($p > 0.635$), adipogenic ($p > 0.317$), inflammatory ($p > 0.413$), or myogenic ($p > 0.320$) pathways, however, within the fibrosis gene group, CTGF was significantly upregulated ($p = 0.046$) in the chronic patients as compared to the acute patients. Similarly, the COL1A1 gene trended towards upregulation ($p = 0.061$), whereas MMP1 and MMP9 both trended towards downregulation in the chronic patients as compared to the acute patients ($p = 0.061$) (Figure 2).

DISCUSSION:

The broad aim of this study was to investigate the gene expression patterns for functional gene groups related to histomorphologic changes in muscle observed in acute and chronic lumbar spine pathology. Secondly, we wanted to ensure that these patterns were not confounded by tissue compositional heterogeneity within muscle samples, given that this is a common feature in this patient population, and has been shown to influence gene expression patterns in other musculoskeletal conditions(18). Although gene expression patterns for the majority of gene groups did not differentiate patients with acute versus chronic disease, several genes in the fibrosis category were significant or trended towards significance, suggesting that fibrosis remains an active biological process in the chronic disease state in contrast to adipogenesis, atrophy, or inflammation. Specifically, CTGF and COL1A1 were both upregulated in the chronic group as compared to the acute group, whereas MMP1 and MMP9 were downregulated. CTGF and COL1A1 are both related to activation and distribution of collagen networks. CTGF is a matricellular protein and is involved in extracellular matrix production (25, 26), and COL1A1 is a gene related to production of type I collagen. MMP1 and MMP9 are both matrix metalloproteinases, which are involved in the breakdown of extracellular matrix in normal physiological processes of tissue remodeling. MMP1 has the ability to cleave interstitial collagen types I, II, and II, and MMP9 is a gelatinase that degrades denatured collagen types IV, VII and X in many tissues(27). Together, the upregulation of collagen deposition and downregulation of

extracellular matrix breakdown is consistent with the fibrotic phenotype seen in chronically diseased paraspinal muscle.

Contrary to our hypothesis, we did not see any differential expression in the other gene pathways of atrophy, adipogenesis, myogenesis, or inflammation. These findings are interesting given that prior literature has suggested that atrophy and fatty infiltration are the primary phenotypic indications of declining muscle health in this population(5, 28-30). However, these observations have historically been informed by observations of decreased muscle cross sectional area and increased fatty deposition in magnetic resonance imaging (MRI)(5, 31, 32), where fibrosis or connective tissue would not be distinguishable from muscle in most clinically utilized imaging protocols (T1 or T2-weighted images), and therefore may not have been recognized as an obvious feature of disease. Together, along with prior literature, these findings suggest that the relationship between atrophy and fatty infiltration with chronicity could be overestimated, while the fibrogenic processes may have been underestimated. This may also indicate that the current treatment paradigm of reducing inflammation and reversing atrophy may have reduced efficacy in chronic stages of disease because these pathways are no longer active, as compared to fibrogenic processes.

The lack of differential expression in the inflammatory gene group is also in contradiction to the only other investigation of gene expression in the lumbar paraspinal muscle of individuals with lumbar spine pathology. However, the small number of genes studied, the small sample size, and the group of patients compared in this study may explain this discrepancy. In the aforementioned study, gene expression was compared between individuals with low back pain who had lumbar kyphosis (N=7) and those who had normal lumbar alignment (N=5). Their analysis was limited to $TNF\alpha$, and $IL6$ as inflammatory biomarkers, and $PGC-1\alpha$ as a marker of mitochondrial biogenesis. Although they reported roughly a two-fold increase in $TNF\alpha$ and a 4-fold increase in $IL6$ in the patient group with kyphotic posture as compared to the normal posture group, they reported p-values of 0.048 for both genes with no reported statistical corrections for multiple comparisons(22). Duration of symptoms was also not described in this study. These analytical strategies may have resulted in an overestimation of differential expression in comparison with the methods used in the current study, in which no significant differential expression was observed between acute and chronic patients. This also highlights the potential for overinterpretation of results when investigating a single gene as opposed to a cluster of genes within a pathway of interest.

The upregulation of fibrogenic genes has also been reported in muscle samples from patients with adolescent idiopathic scoliosis. A study investigating $TGF\beta$ gene expression between concave and convex sides of the scoliotic curve found that $TGF\beta$ was upregulated on the side of the concavity, which may suggest that chronic unloading plays a role in fibrogenic pathways in paraspinal muscle, although $TGF\beta$ is a gene that is implicated in multiple physiological processes (33). In a sheep model of intervertebral disc degeneration in which gene expression and histology were performed on the multifidus muscle pre- and post-injury, the multifidus muscle demonstrated no histologic signs of muscle atrophy, but exhibited increased connective tissue and fat proportions 6 months post injury as compared to pre-injury. In parallel, collagen-1 was upregulated after injury(9). Interestingly, gene

expression for inflammatory cytokines, such as $TNF\alpha$, and $IL1\beta$ were also upregulated in this study, whereas $TGF\beta$ was not significantly changed relative to baseline. Additionally, genes related to atrophy were also unchanged relative to baseline.

This study had some limitations. Primarily, our group definitions may have contributed to the lack of differential expression in the hypothesized pathways. In our study, we defined individuals with lumbar spine pathology of <6-month duration as being “acute”, and those with >6-month symptom duration as being “chronic”. However, clinical definitions of acute and chronic often use a more conservative threshold of 3 months due to the assumption that most musculoskeletal tissues (including muscle, bone, ligament, and tendon) should have had sufficient time to heal from an injury within this timeframe(34). However, it is reasonable that gene expression changes in muscle tissue may occur on an even shorter timeline given the high adaptive capacity of muscle relative to other musculoskeletal tissues(35-37). This is supported by the fact that even at the 6 -month timepoint, changes in tissue composition (high levels of fat and fibrosis) have already manifested, suggesting that the majority of gene expression in the pathways of interest have already occurred. Although we recognize this limitation, redefining our “acute” group with a more conservative timeline was not feasible due to the infrequency of surgically managed acute low back pain. Since standard of care for treatment of low back pain often includes weeks or months of conservative management in the form of pain medications, physical therapy, and injections, obtaining muscle samples from individuals undergoing surgical treatment with symptom durations of <3 months that have not been exposed to an acute trauma is rare. Given this limitation, the differential expression from these analyses may be more appropriately interpreted as a representation of processes associated with late-stage disease progression.

CONCLUSIONS:

This study is the first to evaluate a comprehensive set of atrophic, myogenic, fibrogenic, inflammatory, and adipogenic gene groups in paraspinal muscle of individuals with acute compared to chronic lumbar spine pathology. Our results demonstrate that an increase in fibrogenic gene expression is observed in individuals with late stage disease, whereas atrophic, adipogenic, and inflammatory programs are less predominant. These results provide important information in understanding muscle degeneration in human musculoskeletal disease and highlights the importance of studying functional groups of genes associated with phenotypes of interest. The lack of atrophic and inflammatory processes and upregulation of fibrogenic processes in the chronic stage of disease suggests that prevention or reversal of fibrogenesis may be a feasible treatment target for future research toward the goal of improved patient function and prognosis in this population.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

NIH R01HD088437 and TL1TR001443

The manuscript submitted does not contain information about medical device(s)/drug(s).

National Institutes of Health (grants RO1HD088437 and TL1TR001443) funds were received in support of this work.

No relevant financial activities outside the submitted work.

REFERENCES

1. Andersson GB. Epidemiological features of chronic low-back pain. *Lancet*. 1999;354(9178):581–5. doi: 10.1016/S0140-6736(99)01312-4. PubMed PMID: 10470716. [PubMed: 10470716]
2. Shahidi B, Hubbard JC, Gibbons MC, et al. Lumbar multifidus muscle degenerates in individuals with chronic degenerative lumbar spine pathology. *J Orthop Res*. 2017. Epub 2017/05/08. doi: 10.1002/jor.23597. PubMed PMID: 28480978.
3. Shahidi B, Parra CL, Berry DB, et al. Contribution of Lumbar Spine Pathology and age to Paraspinal Muscle Size and fatty Infiltration0Spine (Phila Pa 1976). 2016. doi: 10.1097/BRS.0000000000001848. PubMed PMID: 27517512.
4. Kader DF, Wardlaw D, Smith FW. Correlation between the MRI changes in the lumbar multifidus muscles and leg pain. *Clin Radiol*. 2000;55(2):145–9. doi: 10.1053/crad.1999.0340. PubMed PMID: 10657162. [PubMed: 10657162]
5. Kjaer P, Bendix T, Sorensen JS, et al. Are MRI-defined fat infiltrations in the multifidus muscles associated with low back pain? *BMC Med*. 2007;5:2. doi: 10.1186/1741-7015-5-2. PubMed PMID: 17254322; PMCID: PMC1796893. [PubMed: 17254322]
6. Parkkola R, Rytökoski U, Kormano M. Magnetic resonance imaging of the discs and trunk muscles in patients with chronic low back pain and healthy control subjects. *Spine (Phila Pa 1976)*. 1993;18(7):830–6. PubMed PMID: 8316880. [PubMed: 8316880]
7. Mannion AF, Weber BR, Dvorak J, et al.,. Fibre type characteristics of the lumbar paraspinal muscles in normal healthy subjects and in patients with low back pain. *J Orthop Res*. 1997;15(6):881–7. doi: 10.1002/jor.1100150614. PubMed PMID: 9497814. [PubMed: 9497814]
8. Ng JK, Richardson CA, Kippers V, et al. Relationship between muscle fiber composition and functional capacity of back muscles in healthy subjects and patients with back pain. *J Orthop Sports Phys Ther*. 1998;27(6):389–402. doi: 10.2519/jospt.1998.27.6.389. PubMed PMID: 9617724. [PubMed: 9617724]
9. Hodges PW, James G, Blomster L, et al. Multifidus muscle changes after back injury are characterized by structural remodeling of muscle, adipose and connective tissue, but not muscle atrophy: Molecular and morphological evidence. *Spine (Phila Pa 1976)*. 2015. doi: 10.1097/BRS.0000000000000972. PubMed PMID: 25943090.
10. Hodges PW, James G, Blomster L, et al. Can proinflammatory cytokine gene expression explain multifidus muscle fiber changes after an intervertebral disc lesion? *Spine (Phila Pa 1976)*. 2014;39(13):1010–7. doi: 10.1097/BRS.0000000000000318. PubMed PMID: 24718080. [PubMed: 24718080]
11. Hiepe P, Gussew A, Rzanny R, et al. Age-related structural and functional changes of low back muscles. *Exp Gerontol*. 2015;65:23–34. doi: 10.1016/j.exger.2015.02.016. PubMed PMID: 25735850. [PubMed: 25735850]
12. Airaksinen O, Herno A, Kaukanen E, et al. Density of lumbar muscles 4 years after decompressive spinal surgery. *Eur Spine J*. 1996;5(3):193–7. PubMed PMID: 8831123. [PubMed: 8831123]
13. Alaranta H, Tallroth K, Soukka A, et al. Fat content of lumbar extensor muscles and low back disability: a radiographic and clinical comparison. *J Spinal Disord*. 1993;6(2):137–40. PubMed PMID: 8504225. [PubMed: 8504225]
14. Barker KL, Shamley DR, Jackson D. Changes in the cross-sectional area of multifidus and psoas in patients with unilateral back pain: the relationship to pain and disability. *Spine (Phila Pa 1976)*. 2004;29(22):E515–9. PubMed PMID: 15543053. [PubMed: 15543053]
15. Demoulin C, Crielaard J-M, Vanderthommen M. Spinal muscle evaluation in healthy individuals and low-back-pain patients: a literature review. *Joint Bone Spine*. 2007;74(1):9–13. doi: 10.1016/j.jbspin.2006.02.013. [PubMed: 17174584]

16. Käser L, Mannion AF, Rhyner A, et al. Active therapy for chronic low back pain: part 2. Effects on paraspinal muscle cross-sectional area, fiber type size, and distribution. *Spine (Phila Pa 1976)*. 2001;26(8):909–19. PubMed PMID: 11317113. [PubMed: 11317113]
17. Mannion AF, Käser L, Weber E, et al. Influence of age and duration of symptoms on fibre type distribution and size of the back muscles in chronic low back pain patients. *Eur Spine J*. 2000;9(4): 273–81. PubMed PMID: 11261614; PMCID: PMC3611339. [PubMed: 11261614]
18. Gibbons MC, Fisch KM, Pichika R, et al. Heterogeneous muscle gene expression patterns in patients with massive rotator cuff tears. *PLoS One*. 2018;13(1):e0190439. Epub 2018/01/03. doi: 10.1371/journal.pone.0190439. PubMed PMID: 29293645; PMCID: PMC5749784. [PubMed: 29293645]
19. Rodriguez-Gonzalez FG, Mustafa DA, Mostert B, et al. The challenge of gene expression profiling in heterogeneous clinical samples. *Methods*. 2013;59(1):47–58. Epub 2012/06/02. doi: 10.1016/j.jymeth.2012.05.005. PubMed PMID: 22652627. [PubMed: 22652627]
20. James G, Blomster L, Hall L, et al. Mesenchymal Stem Cell Treatment of Intervertebral Disc Lesion Prevents Fatty Infiltration and Fibrosis of the Multifidus Muscle, but not Cytokine and Muscle Fiber Changes. *Spine (Phila Pa 1976)*. 2016;41(15):1208–17. Epub 2016/05/03. doi: 10.1097/BRS.0000000000001669. PubMed PMID: 27135642. [PubMed: 27135642]
21. Hodges PW, James G, Blomster L, et al. Multifidus Muscle Changes After Back Injury Are Characterized by Structural Remodeling of Muscle, Adipose and Connective Tissue, but Not Muscle Atrophy: Molecular and Morphological Evidence. *Spine (Phila Pa 1976)*. 2015;40(14): 1057–71. Epub 2015/05/07. doi: 10.1097/BRS.0000000000000972. PubMed PMID: 25943090. [PubMed: 25943090]
22. Kudo D, Miyakoshi N, Hongo M, et al. mRNA expressions of peroxisome proliferator-activated receptor gamma coactivator 1alpha, tumor necrosis factor-alpha, and interleukin-6 in paraspinal muscles of patients with lumbar kyphosis: a preliminary study. *Clin Interv Aging*. 2018;13:1633–8. Epub 2018/09/21. doi: 10.2147/CIA.S172952. PubMed PMID: 30233161; PMCID: PMC6135076. [PubMed: 30233161]
23. Dvinge H, Bertone P. HTqPCR: high-throughput analysis and visualization of quantitative real-time PCR data in R. *Bioinformatics*. 2009;25(24):3325–6. Epub 2009/10/08. doi: 10.1093/bioinformatics/btp578. PubMed PMID: 19808880; PMCID: PMC2788924. [PubMed: 19808880]
24. Reiner A, Yekutieli D, Benjamini Y. Identifying differentially expressed genes using false discovery rate controlling procedures. *Bioinformatics*. 2003;19(3):368–75. Epub 2003/02/14. PubMed PMID: 12584122. [PubMed: 12584122]
25. Morales MG, Cabello-Verrugio C, Santander C, et al. CTGF/CCN-2 over-expression can directly induce features of skeletal muscle dystrophy. *J Pathol*. 2011;225(4):490–501. Epub 2011/08/10. doi: 10.1002/path.2952. PubMed PMID: 21826667. [PubMed: 21826667]
26. Morales MG, Acuna MJ, Cabrera D, et al. The pro-fibrotic connective tissue growth factor (CTGF/CCN2) correlates with the number of necrotic-regenerative foci in dystrophic muscle. *J Cell Commun Signal*. 2018;12(1):413–21. Epub 2017/09/10. doi: 10.1007/s12079-017-0409-3. PubMed PMID: 28887614; PMCID: PMC5842176. [PubMed: 28887614]
27. Chen X, Li Y. Role of matrix metalloproteinases in skeletal muscle: migration, differentiation, regeneration and fibrosis. *Cell Adh Migr*. 2009;3(4):337–41. Epub 2009/08/12. PubMed PMID: 19667757; PMCID: PMC2802742. [PubMed: 19667757]
28. Fortin M, Macedo LG. Multifidus and paraspinal muscle group cross-sectional areas of patients with low back pain and control patients: a systematic review with a focus on blinding. *Phys Ther*. 2013;93(7):873–88. doi: 10.2522/ptj.20120457. PubMed PMID: 23504343; PMCID: PMC3704232. [PubMed: 23504343]
29. Freeman MD, Woodham MA, Woodham AW. The role of the lumbar multifidus in chronic low back pain: a review. *PM R*. 2010;2(2):142–6; quiz 1 p following 67. doi: 10.1016/j.pmrj.2009.11.006. PubMed PMID: 20193941. [PubMed: 20193941]
30. MacDonald DA, Moseley GL, Hodges PW. The lumbar multifidus: does the evidence support clinical beliefs? *Man Ther*. 2006;11(4):254–63. doi: 10.1016/j.math.2006.02.004. PubMed PMID: 16716640. [PubMed: 16716640]
31. Parkkola R, Korman M. Lumbar disc and back muscle degeneration on MRI: correlation to age and body mass. *J Spinal Disord*. 1992;5(1):86–92. PubMed PMID: 1571617. [PubMed: 1571617]

32. Wan Q, Lin C, Li X, et al. MRI assessment of paraspinal muscles in patients with acute and chronic unilateral low back pain. *Br J Radiol.* 2015;88(1053):20140546. doi: 10.1259/bjr.20140546. PubMed PMID: 26105517. [PubMed: 26105517]
33. Nowak R, Kwiecien M, Tkacz M, et al. Transforming growth factor-beta (TGF- beta) signaling in paravertebral muscles in juvenile and adolescent idiopathic scoliosis. *Biomed Res Int.* 2014;2014:594287. Epub 2014/10/15. doi: 10.1155/2014/594287. PubMed PMID: 25313366; PMCID: PMC4181945. [PubMed: 25313366]
34. Chou R Low back pain (chronic). *BMJ Clin Evid.* 2010;2010. Epub 2010/01/01. PubMed PMID: 21418678; PMCID: PMC3217809.
35. Chacon-Cabrera A, Lund-Palau H, Gea J, et al. Time-Course of Muscle Mass Loss, Damage, and Proteolysis in Gastrocnemius following Unloading and Reloading: Implications in Chronic Diseases. *PLoS One.* 2016;11(10):e0164951. Epub 2016/10/30. doi: 10.1371/journal.pone.0164951. PubMed PMID: 27792730; PMCID: PMC5085049. [PubMed: 27792730]
36. Bonaldo P, Sandri M. Cellular and molecular mechanisms of muscle atrophy. *Dis Model Mech.* 2013;6(1):25–39. doi: 10.1242/dmm.010389. PubMed PMID: 23268536; PMCID: PMC3529336. [PubMed: 23268536]
37. Ciciliot S, Schiaffino S. Regeneration of mammalian skeletal muscle. Basic mechanisms and clinical implications. *Curr Pharm Des.* 2010;16(8):906–14. PubMed PMID: 20041823. [PubMed: 20041823]




Figure 1. Hierarchical cluster analysis of multifidus muscle biopsies from acute and chronic patients. The patient group (Acute or Chronic) is indicated on the bottom axis of the heatmap, and gene abbreviations are indicated on the right side. Highly expressed genes are denoted by red coloring, and low-expressed genes are denoted by yellow coloring.




Figure 2. Delta-delta Ct ratios of expression between chronic and acute patient groups (acute group is the reference group). Bars with a solid outline indicate a significant comparison ($P < 0.05$), whereas hatched bars indicate a trend ($P < 0.08$). Solid bars with no outline are not significantly different.

Table 1.

Demographic and biopsy characteristics

	Acute	Chronic
Age (years)	62.5 (5.6)	61.1 (16.9)
Gender (M:F)	4:4	16:9
Fat (%)	16.3 (16.2)	14.2 (11.7)
Muscle (%)	46.2 (22.5)	50.9 (17.4)
Collagen (%)	21.2 (14.1)	20.1 (11.6)
RNA yield (ng/μl)	124.1 (57.6)	122.9 (55.3)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript