

UCSF

UC San Francisco Previously Published Works

Title

Acetaminophen inhibits cytochrome c redox cycling induced lipid peroxidation

Permalink

<https://escholarship.org/uc/item/0bw7065x>

Journal

Biochemical and Biophysical Research Communications, 423(2)

ISSN

0006-291X

Authors

Yin, Huiyong
Vergeade, Aurélie
Shi, Qiong
[et al.](#)

Publication Date

2012-06-01

DOI

10.1016/j.bbrc.2012.05.058

Peer reviewed

Published in final edited form as:

Biochem Biophys Res Commun. 2012 June 29; 423(2): 224–228. doi:10.1016/j.bbrc.2012.05.058.

Acetaminophen Inhibits Cytochrome C Redox Cycling Induced Lipid Peroxidation

Huiyong Yin^{a,b,c,*}, Aurélie Vergeade^a, Qiong Shi^a, William E. Zackert^a, Katherine C. Gruenberg^a, Magdalena Bokiej^a, Taneem Amin^a, Weizhen Ying^a, Tina S. Masterson^a, Sandra S. Zinkel^a, John A. Oates^{a,b}, Olivier Boutaud^{b,¶}, and L. Jackson Roberts II^{a,b,*},¶

^aDepartment of Medicine, Division of Clinical Pharmacology, Vanderbilt University School of Medicine, TN, 37232, USA

^bDepartment of Pharmacology, Division of Clinical Pharmacology, Vanderbilt University School of Medicine, TN, 37232, USA

^cLaboratory of Lipid Metabolism in Human Nutrition and Related Diseases, Institute for Nutritional Sciences, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, China

Abstract

Cytochrome (cyt) c can uncouple from the respiratory chain following mitochondrial stress and catalyze lipid peroxidation. Accumulating evidence shows that this phenomenon impairs mitochondrial respiratory function and also initiates the apoptotic cascade. Therefore, under certain conditions a pharmacological approach that can inhibit cyt c catalyzed lipid peroxidation may be beneficial. We recently showed that acetaminophen (ApAP) at normal pharmacologic concentrations can prevent hemoprotein-catalyzed lipid peroxidation *in vitro* and *in vivo* by reducing ferryl heme to its ferric state. We report here, for the first time, that ApAP inhibits cytochrome c-catalyzed oxidation of unsaturated free fatty acids and also the mitochondrial phospholipid, cardiolipin. Using isolated mitochondria, we also showed that ApAP inhibits cardiolipin oxidation induced by the pro-apoptotic protein, tBid. We found that the IC₅₀ of the inhibition of cardiolipin oxidation by ApAP is similar in both intact isolated mitochondria and cardiolipin liposomes, suggesting that ApAP penetrates well into the mitochondria. Together with our previous results, the findings presented herein suggest that ApAP is a pleiotropic inhibitor of peroxidase catalyzed lipid peroxidation. Our study also provides a potentially novel pharmacological approach for inhibiting the cascade of events that can result from redox cycling of cyt c.

Keywords

Acetaminophen; cytochrome c; lipid peroxidation; redox cycling

© 2012 Elsevier Inc. All rights reserved

Correspondence: L. Jackson Roberts, II, M.D. Departments of Pharmacology and Medicine, Division of Clinical Pharmacology, Vanderbilt University School of Medicine, Nashville, TN 37232-6602 Tel.: 615-322-3304; Fax: 615-322-470
jack.roberts@vanderbilt.edu.

*Current Address: Huiyong Yin, Institute for Nutritional Sciences, Shanghai Institute for Biological Sciences, Chinese Academy of Sciences, 294 Taiyuan Road, Shanghai, China 200031. hyyin@sibs.ac.cn.

¶Both authors contributed equally to the work

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

The normal function of cytochrome c (cyt c) is the transfer of electrons between the respiratory complexes III and IV in the mitochondria. However, cyt c also functions as a peroxidase. Acting as a peroxidase, cyt c catalyzes reduction of hydroperoxides, including H₂O₂, by a process that generates a radical that catalyzes lipid peroxidation [1–3].

This concept is derived from several observations starting in 1959 when Tappel and Zalkin were first to demonstrate that cytochrome c could catalyze peroxidation of unsaturated fatty acids [4]. This observation was later extended to the mitochondria when Scott and Hunter showed that generation of lipid peroxidation in the mitochondria was greatly dependent on cytochrome c [5], a finding extensively confirmed subsequently [6–14]. Cytochrome c catalyzed lipid peroxidation may be involved in many cellular processes ranging from phosphatidylserine externalization during apoptosis [15] to modification of mitochondrial function [16] and triggering cytochrome c release during apoptosis [15].

Acetaminophen (ApAP) is one of the most widely used analgesics. We and others have shown that it inhibits the prostaglandin H₂ synthase by reducing the protoporphyrin radical cation in the peroxidase active site of the enzyme, thereby blocking formation of the catalytic tyrosyl radical in the cyclooxygenase site [17,18]. We later showed that ApAP could similarly inhibit the peroxide-driven lipid peroxidation catalyzed by myoglobin and by hemoglobin [19] and presented the proof of concept that ApAP could be used *in vivo* to protect the kidney from oxidative damage following rhabdomyolysis [19].

Herein we describe for the first time the ability of ApAP to inhibit peroxidation of unsaturated fatty acids catalyzed by cytochrome c *in vitro* using free arachidonic acid and tetralinoleoyl cardiolipin. Importantly, we show that ApAP can prevent cardiolipin oxidation in isolated mitochondria following activation of apoptosis by tBid.

2. Materials and Methods

2.1 Reagents

Phospholipids, 1-palmitoyl-2-oleoyl-3-phosphatidylcholine (POPC), tetralinoleoyl cardiolipin (L₄CL), tetramyristeoyl cardiolipin (M₄CL) were purchased from Avanti Polar Lipids (Alabaster, AL) and used without further purification. All other chemicals were purchased from Sigma-Aldrich Chemical Company (Milwaukee, WI). HPLC quality solvents, such as methanol, water, 2-propanol, and acetonitrile were purchased from either Fisher Chemical (Phillipsburg, NJ) or EM Science (Gibbstown, NJ).

2.2 Inhibition of cyt c induced oxidation of arachidonic acid by ApAP

Oxidation of arachidonic acid (AA) by cyt c and H₂O₂ was performed based on minor modification of the protocol used for myoglobin [19]. Briefly, cyt c (50 μM) was mixed with [¹⁴C] AA (10 μM) and various concentrations of ApAP. The reaction was initiated with the addition of hydrogen peroxide (H₂O₂, 250 μM) and incubated at 37 °C for 3 h. Oxidation products were quantified as previously described [19]. Control experiments for each drug concentration were also performed in which cyt c was omitted. The radioactivity associated with products of oxidation of [¹⁴C] AA incubated without cyt c (background oxidation) was subtracted from each value obtained in presence of cyt c in the same conditions. The IC₅₀'s for ApAP were calculated using the logit method. Data are presented as the mean ± SEM.

2.3 Oxidation of L₄CL in liposome by cyt c and hydrogen peroxide

Oxidation of L₄CL in liposomes by cyt c and H₂O₂ was carried out based on a published protocol [20]. CL and POPC stored in chloroform were mixed in a glass vial and the solvent was removed by a flow of nitrogen. PBS (50 mM, pH 7.4) with 100 μM DTPA was added to achieve final concentrations of 50 μM CL and 200 μM POPC. Then, the lipid mixture was vortexed and sonicated for 1 min under nitrogen. Cyt c (5 μM final concentration) and various concentrations of ApAP ranging from 0 to 500 μM were added, and the reaction was initiated by adding H₂O₂ (100 μM final concentration). After 30 min at 37 °C, the reaction was stopped by adding 2 U/ml catalase. 0.75 % NaCl was added and the oxidation mixture was extracted with chloroform and methanol (2:1, v:v) containing 0.1 mM butyrate hydroxytoluene and 0.1 mM triphenylphosphine and 2.5 μg tetramyristeoylcardiolipin (M₄CL) as internal standard. The separated organic phase was evaporated, resuspended in methanol:acetonitrile:H₂O (60:20:20, v/v/v) and stored at -80°C until analysis by LC-MS as described below.

2.4 ApAP inhibits CL oxidation in isolated mitochondria

The efficacy of ApAP to inhibit CL oxidation in isolated mitochondria was determined as described. Mouse liver mitochondria were isolated as previously described [21]. 15 μg of mitochondria were pre-incubated with 0 to 400 μM ApAP for 10 min. Recombinant tBid (10 ng) was added to the mitochondria and the samples were incubated for 30 min at room temperature. Mitochondria were pelleted by centrifugation at 8000 rpm for 10 min at 4°C. The oxidation products of CL (OxCL) in the pellets were processed as described above for the liposomes and analyzed by LC-MS as described below.

2.5 Quantification of oxidation products of CL by Liquid Chromatography-Mass Spectrometry (LC-MS)

Analysis of oxidation products of CL by LC-MS was carried out as described previously [22]. The extracted lipid fraction was separated online by UPLC using a Waters Acquity UPLC system (Waters Corp., Milford, MA). Mass spectrometry analysis was performed on a Thermo Quantum Ultra triple quadrupole mass spectrometer (Thermo Scientific Inc., San Jose, CA, USA). The mass spectrometer was operated in the negative ion mode using selective reaction monitoring (SRM). Nitrogen was used as the sheath gas at 38 p.s.i. The capillary temperature was 350 °C. The spray voltage was 4.5 kV, and the tube lens voltage was 100 V. Data acquisition and analysis were performed using Xcalibur software, version 2.0. The molecular ion corresponding to the doubly charged monohydroxyL₄CL ([L₄CL-OH - 2H]²⁻; *m/z* = 731.6) was monitored as a major species of OxCL. The following transitions were monitored in SRM: M₄CL, *m/z* 619.6 to 227.2; L₄CL, *m/z* 723.6 to 279.2; CL-OH, *m/z* 731.6 to 279.2, and 731.6 to 295.2.

The area under the curve (AUC) for oxidized cardiolipins (OxCL), M₄CL and L₄CL were determined and the AUC of OxCL and L₄CL in each sample were normalized to the M₄CL AUC. The corrected values were used to calculate the ratio OxCL over total CL and were expressed as percent of total CL. Data are presented as the mean ± SEM.

3. Results

3.1 Acetaminophen inhibits cyt c redox cycling induced oxidation of arachidonic acid (AA)

We studied the ability of ApAP to inhibit peroxidation of unsaturated fatty acids catalyzed by cytochrome c *in vitro* using radiolabeled arachidonic acid. The experiments were performed in the presence of 250 μM H₂O₂ for 3 hours according to preliminary experiments showing that in these conditions all the substrate was not oxidized at the end of the reaction. Our results show that ApAP inhibits lipid peroxidation catalyzed by cyt c with

an IC_{50} of $91 \mu\text{M} \pm 47 \mu\text{M}$ ($n = 5$) (Figure 1). The IC_{50} of ApAP in these experimental conditions is well within its normal therapeutic range in humans (67–200 μM).

3.2 Acetaminophen prevents cyt c mediated cardiolipin oxidation in liposomes

Several studies have established that association of cyt c with several families of lipids enhances the ability of cyt c to catalyze lipid peroxidation [1,11,12,15]. In more recent work, evidence has emerged for a preponderant role of cardiolipin (CL), a class of lipid primarily associated with the mitochondria where cyt c is located. Tetralinoleoyl CL ($L_4\text{CL}$) is the major CL in mammalian cells and oxidation of $L_4\text{CL}$ leads to the formation of lipid hydroperoxides as the primary oxidation products [22]. We developed an LC-MS method to quantify the $L_4\text{CL}$ oxidation products. In this method, the CL-OOHs are reduced by triphenylphosphine and the resulting CLOHs are monitored by selective reaction monitoring. Tetramyristoyl CL ($M_4\text{CL}$) is used as an internal standard (Figure 2A). We found that ApAP potently inhibited cyt c catalyzed oxidation of $L_4\text{CL}$ (Figure 2B) with an IC_{50} of $22.8 \mu\text{M} \pm 7.6 \mu\text{M}$ ($n = 6$). It is noteworthy that we also found that the ability of ApAP to inhibit cyt c mediated CL oxidation is much greater than its ability to inhibit cyt c mediated oxidation of AA (Figure 1), which is consistent with the observation that binding to CL enhances cyt c ability to catalyze lipid peroxidation [11].

3.3 Acetaminophen inhibits cyt c mediated cardiolipin oxidation in mitochondria

Emerging evidence suggests that CL plays an active role in mitochondrial function, including the regulation of mitochondrial outer membrane (MOM), mobilization of cyt c, and anchoring of caspase-8 to mitochondria during death receptor-induced apoptosis [23]. Therefore, we investigated whether ApAP inhibits oxidation of CL in mitochondria isolated from mouse liver. We used tBid, a pro-apoptotic Bcl2 family member, to activate the mitochondria because of the evidence showing that tBid causes the remodeling of mitochondrial cristae and CL oxidation [24]. Our data shows that ApAP dose-dependently inhibits mitochondrial CL oxidation (Figure 3) with an IC_{50} of $32.5 \mu\text{M} \pm 5.5 \mu\text{M}$ ($n = 6$). Interestingly, the potency of ApAP to inhibit CL oxidation in isolated mitochondria is not significantly different from that obtained with liposomes (Mann Whitney, $p = 0.20$) (Figure 2B), suggesting that ApAP exhibits high penetrance into mitochondria.

4. Discussion

Recently we have shown that ApAP can prevent hemoprotein-catalyzed lipid peroxidation *in vitro* by reducing ferryl heme to its ferric state and that ApAP is highly protective against myoglobin redox cycling induced renal injury in an animal model of rhabdomyolysis-induced renal failure [19]. We have extended this discovery to the peroxidase activity of cyt c and obtained evidence indicating that ApAP is also capable of inhibiting cyt c redox cycling-induced lipid peroxidation.

As a heme protein, cyt c can function as a peroxidase. Its ability to catalyze oxidation of unsaturated fatty acids is well documented and it is thought to be contributing to several cellular dysfunctions following oxidative stress [1,4,5,15,16,20]. Using the same assay that we developed for myoglobin and hemoglobin [19], we showed that ApAP was able to inhibit lipid peroxidation catalyzed by H_2O_2 -activated cyt c *in vitro* at concentrations that are within the therapeutic range in humans.

Cyt c-dependent lipid peroxidation is known to be greatly increased when the protein associates with esterified lipids such as cardiolipin, the major lipid present in the mitochondria [11]. It is thought that binding of cyt c to CL unfolds the protein and facilitates the opening of the heme catalytic site to peroxides [25,26]. Partially unfolded cyt c in the

complex with CL exerts an almost 100-fold higher catalytic activity as a peroxidase in the presence of H₂O₂ than the native protein [27]. Using liposomes of cardiolipin, an established model for investigating this process, we show that ApAP inhibits oxidation of the polyunsaturated fatty acid-containing CL, L₄CL, with an IC₅₀ 4- fold lower than for free arachidonic acid.

CLs are predominantly located in the inner mitochondrial membranes and intimately interact with the electron transport chain (ETC) complexes involved in oxidative phosphorylation [28,29]. Furthermore, CL is also associated with members of the apoptotic machinery including cyt c, Bid, and caspase 8 [30,31]. Although ApAP is very efficient in inhibiting the prostaglandin synthases which are located in the endoplasmic reticulum, it has not been determined if ApAP also penetrates in the mitochondria. Using our mass spectrometric assay we investigated whether ApAP could inhibit CL oxidation in isolated mitochondria. The mitochondria were activated with tBid, as it was shown that this led to increased CL oxidation [24]. Our results clearly demonstrate that ApAP inhibits CL oxidation in isolated mitochondria with an IC₅₀ similar to that observed using L₄CL liposomes and purified cyt c. These data support the hypothesis that ApAP penetrates in the mitochondria and inhibits cyt c-dependent lipid peroxidation in response to tBid.

Our results may have broad potential implications as cyt c-catalyzed lipid peroxidation is involved in several biochemical processes. Oxidative stress is known to induce oxidation of phospholipids in the mitochondria by a cyt c-dependent mechanism [1–3,37] which has recently been shown to inhibit mitochondrial function and initiate the apoptotic response in a Bid-dependent fashion [38].

Cyt c also oxidizes phospholipids in the plasma membrane. An elegant work by Tyurina *et al.* suggested that cyt c binds to phosphatidylserine (PS) in the plasma membrane and catalyzes its peroxidation, leading to PS externalization [15], a trigger for phagocytic elimination of apoptotic cells [41,42].

A seminal discovery made by Kagan *et al.* suggests that CL peroxidation appears to be an early event preceding the release of cyt c and caspase activation [20]. The presence of four linoleic acid acyl chains in one molecule and the mitochondrial location make it susceptible to free radical oxidation [43]. The oxidation of CL is catalyzed by a peroxidase activity of the cyt c/CL complex. As a heme protein, cyt c can function as a peroxidase, albeit at a very low rate (about 1 M⁻¹S⁻¹) [44,45]. Native cyt c favors its normal function as an electron shuttle between complex III and IV of ETC. However, binding of CLs to cyt c unfolds the protein and paves the way for an opening of the heme catalytic site to hydrogen peroxides [25,26]. We provide evidence that ApAP inhibits CL oxidation induced by the peroxidase activity of cyt c/CL complex by inhibiting cyt c/CL peroxidase redox cycling.

As discussed above, cyt c redox cycling could have deleterious effects independent of caspase activation by catalyzing lipid peroxidation in the mitochondria, which leads to impaired respiratory function, or at the cell surface, which may eventually lead to cellular death. In this context, our results suggest that inhibition of cyt c catalyzed lipid peroxidation by ApAP could provide cytoprotection by preserving mitochondrial function as well as preventing activation of the phagocytic and caspase pathways.

In previous studies, we have demonstrated that ApAP is capable of inhibiting redox cycling of other hemoproteins, specifically myoglobin and hemoglobin [19]. ApAP also has been found to inhibit myeloperoxidase-induced lipid peroxidation [47]. Together with our results presented herein, these findings suggest that ApAP is a pleiotropic inhibitor of peroxidases, including cyt c. Although sustained inhibition of apoptosis by inhibiting cyt c catalyzed oxidation of cardiolipin could potentially allow damaged cells to survive and accumulate

additional mutations, leading to activation of oncogenes and uncontrolled cell proliferation (neoplasia), a pharmacologic approach that could temporarily prevent apoptotic cell death could have a profound beneficial impact in a variety of diseases such as myocardial infarction, stroke, and traumatic brain injury.

Acknowledgments

This work was supported by NIH grants GM15431, GM42056, ES13125 and HL088347. HY is partially supported by grants from the National Natural Science Foundation of China (31170809), National Key Basic Research Program of China (973 Program, #2012CB524900).

Abbreviations

| | |
|------------------------|--|
| AA | arachidonic acid |
| ApAP | acetaminophen |
| CL | cardiolipin |
| cyt c | cytochrome c |
| ETC | electron transport chain |
| HPLC | high performance liquid chromatography |
| L₄CL | tetralinoleoyl cardiolipin |
| LC-MS | liquid chromatography mass spectrometry |
| M₄CL | myristoyl CL |
| MOM | mitochondrial outer membrane |
| PGHS | prostaglandin H synthase |
| POPC | 1-palmitoyl-2-oleoyl-phosphatidylcholine |
| SRM | selective reaction monitoring |
| TLC | thin layer chromatography |
| UPLC | ultra pressure liquid chromatography. |

5. References

- [1]. Radi R, Turrens JF, Freeman BA. Cytochrome c-catalyzed membrane lipid peroxidation by hydrogen peroxide. *Arch. Biochem. Biophys.* 1991; 288:118–125. [PubMed: 1654818]
- [2]. Radi R, Sims S, Cassina A, Turrens JF. Roles of catalase and cytochrome c in hydroperoxide-dependent lipid peroxidation and chemiluminescence in rat heart and kidney mitochondria. *Free Radic. Biol. Med.* 1993; 15:653–659. [PubMed: 8138192]
- [3]. Radi R, Bush KM, Freeman BA. The role of cytochrome c and mitochondrial catalase in hydroperoxide-induced heart mitochondrial lipid peroxidation. *Arch. Biochem. Biophys.* 1993; 300:409–415. [PubMed: 8380970]
- [4]. Tappel AL, Zalkin H. Lipide peroxidation in isolated mitochondria. *Arch. Biochem. Biophys.* 1959; 80:326–333.
- [5]. Scott AA, Hunter FE Jr. Role of cytochrome C in glutathione induced swelling and lipid peroxidation in liver mitochondria. *Biochemical and biophysical research communications.* 1963; 11:461–465. [PubMed: 13992509]
- [6]. Kaschnitz RM, Hatefi Y. Lipid oxidation in biological membranes. Electron transfer proteins as initiators of lipid autoxidation. *Arch. Biochem. Biophys.* 1975; 171:292–304. [PubMed: 242270]
- [7]. Brown LR, Wuthrich K. A spin label study of lipid oxidation catalyzed by heme proteins. *Biochim. Biophys. Acta.* 1977; 464:356–369. [PubMed: 188473]

- [8]. Hamberg M. Decomposition of unsaturated fatty acid hydroperoxides by hemoglobin: Structures of major products of 13L-hydroperoxy-9,11-octadecadienoic acid. *Lipids*. 1975; 10:87–92. [PubMed: 1167926]
- [9]. Evans PJ, Akanmu D, Halliwell B. Promotion of oxidative damage to arachidonic acid and alpha 1-antiproteinase by anti-inflammatory drugs in the presence of the haem proteins myoglobin and cytochrome C. *Biochem. Pharmacol.* 1994; 48:2173–2179. [PubMed: 7811298]
- [10]. Iwase H, Sakurada K, Takatori T, Nagao M, Nijima H, Matsuda Y, Kobayashi M. Calcium ions potentiate lipoxygenase activity of cytochrome c at the physiological pH. *Biochemical and biophysical research communications*. 1998; 243:485–491. [PubMed: 9480835]
- [11]. Iwase H, Takatori T, Nijima H, Nagao M, Amano T, Iwate K, Matsuda Y, Nakajima M, Kobayashi M. Formation of leukotoxin (9,10-epoxy-12-octadecenoic acid) during the autoxidation of phospholipids promoted by hemoproteins. *Biochim. Biophys. Acta*. 1997; 1345:27–34. [PubMed: 9084498]
- [12]. Goni FM, Ondarroya M, Azpiazu I, Macarulla JM. Phospholipid oxidation catalyzed by cytochrome c in liposomes. *Biochim. Biophys. Acta*. 1985; 835:549–556. [PubMed: 2990565]
- [13]. Iwase H, Takatori T, Aono K, Iwate K, Takahashi M, Nakajima M, Nagao M. Establishment of a monoepoxide (leukotoxin and its isomer) producing system using a hydrogen peroxide-generating system. *Biochemical and biophysical research communications*. 1995; 216:483–488. [PubMed: 7488137]
- [14]. Hayakawa M, Ogawa T, Sugiyama S, Ozawa T. Hydroxyl radical and leukotoxin biosynthesis in neutrophil plasma membrane. *Biochemical and biophysical research communications*. 1989; 161:1077–1085. [PubMed: 2545194]
- [15]. Tyurina YY, Shvedova AA, Kawai K, Tyurin VA, Kommineni C, Quinn PJ, Schor NF, Fabisiak JP, Kagan VE. Phospholipid signaling in apoptosis: peroxidation and externalization of phosphatidylserine. *Toxicology*. 2000; 148:93–101. [PubMed: 10962127]
- [16]. Chen R, Feldstein AE, McIntyre TM. Suppression of mitochondrial function by oxidatively truncated phospholipids is reversible, aided by bid, and suppressed by Bcl-XL. *J. Biol. Chem.* 2009; 284:26297–26308. [PubMed: 19654426]
- [17]. Ouellet M, Percival MD. Mechanism of acetaminophen inhibition of cyclooxygenase isoforms. *Arch. Biochem. Biophys.* 2001; 387:273–280. [PubMed: 11370851]
- [18]. Boutaud O, Aronoff DM, Richardson JH, Marnett LJ, Oates JA. Determinants of the cellular specificity of acetaminophen as an inhibitor of prostaglandin H2 synthases. *Proc. Natl. Acad. Sci. U S A*. 2002; 99:7130–7135. [PubMed: 12011469]
- [19]. Boutaud O, Moore KP, Reeder BJ, Harry D, Howie AJ, Wang S, Carney CK, Masterson TS, Amin T, Wright DW, Wilson MT, Oates JA, Roberts LJ 2nd. Acetaminophen inhibits hemoprotein-catalyzed lipid peroxidation and attenuates rhabdomyolysis-induced renal failure. *Proc. Natl. Acad. Sci. U S A*. 2010; 107:2699–2704. [PubMed: 20133658]
- [20]. Kagan VE, Tyurin VA, Jiang J, Tyurina YY, Ritov VB, Amoscato AA, Osipov AN, Belikova NA, Kapralov AA, Kini V, Vlasova, Zhao Q, Zou M, Di P, Svistunenko DA, Kurnikov IV, Borisenko GG. Cytochrome c acts as a cardiolipin oxygenase required for release of proapoptotic factors. *Nat. Chem. Biol.* 2005; 1:223–232. [PubMed: 16408039]
- [21]. McBride HM, Silvius JR, Shore GC. Insertion of an uncharged polypeptide into the mitochondrial inner membrane does not require a trans-bilayer electrochemical potential: effects of positive charges. *Biochim. Biophys. Acta*. 1995; 1237:162–168. [PubMed: 7632709]
- [22]. Liu W, Porter NA, Schneider C, Brash AR, Yin H. Formation of 4-hydroxynonenal from cardiolipin oxidation: Intramolecular peroxy radical addition and decomposition. *Free Radical Biology and Medicine*. 2011; 50:166–178. [PubMed: 21047551]
- [23]. Montero J, Mari M, Colell A, Morales A, Basañez G, Garcia-Ruiz C, Fernández-Checa JC. Cholesterol and peroxidized cardiolipin in mitochondrial membrane properties, permeabilization and cell death. *Biochim. Biophys. Acta - Bioenergetics*. 2010; 1797:1217–1224.
- [24]. Scorrano L, Ashiya M, Buttle K, Weiler S, Oakes SA, Mannella CA, Korsmeyer SJ. A distinct pathway remodels mitochondrial cristae and mobilizes cytochrome c during apoptosis. *Dev. Cell*. 2002; 2:55–67. [PubMed: 11782314]

- [25]. Kagan VE, Bayir HA, Belikova NA, Kapralov O, Tyurina YY, Tyurin VA, Jiang J, Stoyanovsky DA, Wipf P, Kochanek PM, Greenberger JS, Pitt B, Shvedova AA, Borisenko G. Cytochrome c/cardiolipin relations in mitochondria: a kiss of death. *Free Radic. Biol. Med.* 2009; 46:1439–1453. [PubMed: 19285551]
- [26]. Hanske J, Toffey JR, Morenz AM, Bonilla AJ, Schiavoni KH, Pletneva EV. Conformational properties of cardiolipin-bound cytochrome c. *Proc. Natl. Acad. Sci. U S A.* 2012; 109:125–130. [PubMed: 22190488]
- [27]. Belikova NA, Tyurina YY, Borisenko G, Tyurin V, Samhan Arias AK, Yanamala N, Furtmuller PG, Klein-Seetharaman J, Obinger C, Kagan VE. Heterolytic reduction of fatty acid hydroperoxides by cytochrome c/cardiolipin complexes: antioxidant function in mitochondria. *J. Am. Chem. Soc.* 2009; 131:11288–11289. [PubMed: 19627079]
- [28]. Eble KS, Coleman WB, Hantgan RR, Cunningham CC. Tightly associated cardiolipin in the bovine heart mitochondrial ATP synthase as analyzed by ³¹P nuclear magnetic resonance spectroscopy. *J. Biol. Chem.* 1990; 265:19434–19440. [PubMed: 2147180]
- [29]. Fry M, Green DE. Cardiolipin requirement for electron transfer in complex I and III of the mitochondrial respiratory chain. *J. Biol. Chem.* 1981; 256:1874–1880. [PubMed: 6257690]
- [30]. Lutter M, Fang M, Luo X, Nishijima M, Xie X, Wang X. Cardiolipin provides specificity for targeting of tBid to mitochondria. *Nat. Cell Biol.* 2000; 2:754–761. [PubMed: 11025668]
- [31]. Schug ZT, Gottlieb E. Cardiolipin acts as a mitochondrial signalling platform to launch apoptosis. 2009; 1788:2022–2031.
- [32]. Jaques-Robinson KM, Golfetti R, Baliga SS, Hadzimichalis NM, Merrill GF. Acetaminophen is cardioprotective against H₂O₂-induced injury in vivo. *Exp. Biol. Med.* 2008; 233:1315–1322.
- [33]. Merrill GF, Merrill JH, Golfetti R, Jaques KM, Hadzimichalis NS, Baliga SS, Rork TH. Antiarrhythmic properties of acetaminophen in the dog. *Exp. Biol. Med.* 2007; 232:1245–1252.
- [34]. Merrill GF, Rork TH, Spiler NM, Golfetti R. Acetaminophen and myocardial infarction in dogs. *American Journal of Physiology - Heart Circulation Physiology.* 2004; 287:H1913–1920.
- [35]. Rork TH, Van Dyke K, Spiler NM, Merrill GF. Acetaminophen in the hypoxic and reoxygenated guinea pig myocardium. *Exp. Biol. Med.* 2004; 229:1154–1161.
- [36]. Nam TG, Nara SJ, Zagol-Ikapitte I, Cooper T, Valgimigli L, Oates JA, Porter NA, Boutaud O, Pratt DA. Pyridine and pyrimidine analogs of acetaminophen as inhibitors of lipid peroxidation and cyclooxygenase and lipoxygenase catalysis. *Org. Biomol. Chem.* 2009; 7:5103–5112. [PubMed: 20024105]
- [37]. Radi R, Beckman JS, Bush KM, Freeman BA. Peroxynitrite-induced membrane lipid peroxidation: the cytotoxic potential of superoxide and nitric oxide. *Arch. Biochem. Biophys.* 1991; 288:481–487. [PubMed: 1654835]
- [38]. McIntyre TM, Prescott SM, Stafforini DM. The emerging roles of PAF acetylhydrolase. *J. Lipid Res.* 2009; 50(Suppl):S255–259. [PubMed: 18838739]
- [39]. Circu ML, Aw TY. Reactive oxygen species, cellular redox systems, and apoptosis. *Free Radic. Biol. Med.* 2010; 48:749–762. [PubMed: 20045723]
- [40]. Ott M, Robertson JD, Gogvadze V, Zhivotovsky B, Orrenius S. Cytochrome c release from mitochondria proceeds by a two-step process. *Proc. Natl. Acad. Sci. U S A.* 2002; 99:1259–1263. [PubMed: 11818574]
- [41]. Fadok VA, Bratton DL, Frasch SC, Warner ML, Henson PM. The role of phosphatidylserine in recognition of apoptotic cells by phagocytes. *Cell Death Differ.* 1998; 5:551–562. [PubMed: 10200509]
- [42]. Fadok VA, Voelker DR, Campbell PA, Cohen JJ, Bratton DL, Henson PM. Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers specific recognition and removal by macrophages. *J. Immunol.* 1992; 148:2207–2216. [PubMed: 1545126]
- [43]. Yin H, Xu L, Porter NA. Free Radical Lipid Peroxidation: Mechanisms and Analysis. *Chem. Rev.* 2011; 111:5944–5972. [PubMed: 21861450]
- [44]. Florence TM. The degradation of cytochrome c by hydrogen peroxide. *J. Inorg. Biochem.* 1985; 23:131–141. [PubMed: 2983016]

- [45]. Radi R, Thomson L, Rubbo H, Prodanov E. Cytochrome c-catalyzed oxidation of organic molecules by hydrogen peroxide. *Arch. Biochem. Biophys.* 1991; 288:112–117. [PubMed: 1654817]
- [46]. Danial NN, Korsmeyer SJ. Cell death: critical control points. *Cell.* 2004; 116:205–219. [PubMed: 14744432]
- [47]. Chou T, Greenspan P. Effect of acetaminophen on the myeloperoxidase-hydrogen peroxide-nitrite mediated oxidation of LDL. *Biochim. Biophys. Acta.* 2002; 1581:57–63. [PubMed: 11960752]

Acetaminophen inhibits lipid peroxidation catalyzed by cytochrome c

Acetaminophen inhibits cardiolipin oxidation catalyzed by cytochrome c

Acetaminophen inhibits oxidation of mitochondrial cardiolipin induced by tBid




Figure 1. ApAP inhibits cyt c induced oxidation of arachidonic acid (AA)

[¹⁴C] AA was incubated with cyt c and various concentrations of ApAP as described in the methods section. The residual AA and the oxidized products were extracted and analyzed as described previously [19]. The oxidation is represented as the percentage of AA oxidized by cyt c at 3 hours versus the control in which no ApAP is added. Each data point represents the mean \pm S.E.M. (n = 4).




Figure 2. ApAP inhibits cyt c-catalyzed oxidation of cardiolipin

(A) Analysis of oxidation products of L₄CL by LC-MS. The following transitions were monitored: M₄CL, m/z 619.6 to 227.2; L₄CL, m/z 723.6 to 279.2; oxL₄CL, m/z 731.6 to 279.2 and 731.6 to 295.2. The structures of the doubly charged molecular ion [L₄CL-OH - 2H]²⁻ and of the monitored fragments are represented for OxL₄CL. M₄CL: tetramyristeoylcardiolipin. L₄CL: tetralinoleoylcardiolipin. OxL₄CL: monohydroxyL₄CL. (B) L₄CL liposomes were incubated with cyt c with various concentrations of ApAP and oxidized CL were quantified by LC/MS as described in the methods section. Oxidized CL are represented as % of total CL (mean ± S.E.M., n = 6)




Figure 3. ApAP inhibits CL oxidation in isolated mitochondria

Mouse liver isolated mitochondria were pretreated with various concentrations of ApAP for 1 h prior being activated by tBid. CL and its oxidized products were extracted and analyzed by LC/MS as described in the methods section. Oxidized CL is expressed as % of total CL (mean \pm S.E.M., n = 6)