

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

Quadrupole Moments of First Excited States in  $^{28}\text{Si}$ ,  $^{32}\text{S}$ , and  $^{40}\text{Ar}$

### Permalink

<https://escholarship.org/uc/item/0d8366pg>

### Authors

Nakai, K

Quebert, J L

Stephens, F S

et al.

### Publication Date

1970

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Submitted to Physical Review Letters

UCRL-19544  
Preprint

c. 2

LAWRENCE  
RADIATION LABORATORY

MAR 17 1970

LIBRARY AND  
DOCUMENTS SECTION

QUADRUPOLE MOMENTS OF FIRST EXCITED STATES IN  
 $^{28}\text{Si}$ ,  $^{32}\text{S}$ , AND  $^{40}\text{Ar}$

K. Nakai, J. L. Quebert, F. S. Stephens, and R. M. Diamond

January 1970

AEC Contract No. W-7405-eng-48

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

34  
**LAWRENCE RADIATION LABORATORY**  
**UNIVERSITY of CALIFORNIA BERKELEY**

UCRL-19544

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

QUADRUPOLE MOMENTS OF FIRST EXCITED STATES IN  $^{28}\text{Si}$ ,  $^{32}\text{S}$ , AND  $^{40}\text{Ar}^+$

K. Nakai<sup>\*</sup>, J. L. Quebert<sup>\*\*</sup>, F. S. Stephens, and R. M. Diamond

Lawrence Radiation Laboratory  
University of California  
Berkeley, California 94720

January 1970

ABSTRACT

Static quadrupole moments of the first excited states of  $^{28}\text{Si}$ ,  $^{32}\text{S}$ , and  $^{40}\text{Ar}$  have been measured using the reorientation effect in projectile Coulomb excitation. The results obtained are  $Q(^{28}\text{Si}, 2^+) = +0.11 \pm 0.05$  b,  $Q(^{32}\text{S}, 2^+) = -0.20 \pm 0.06$  b, and  $Q(^{40}\text{Ar}, 2^+) = +0.01 \pm 0.04$  b.

The reorientation effect<sup>1</sup> in projectile Coulomb excitation provides a sensitive way to measure quadrupole moments of light nuclei. This method was successfully applied to the measurement of the quadrupole moments of  $^{20}\text{Ne}$  and  $^{22}\text{Ne}$  ( $2^+$  states).<sup>2</sup> In the present work the first excited states in  $^{28}\text{Si}$ ,  $^{32}\text{S}$ , and  $^{40}\text{Ar}$  have been studied using beams from the Berkeley Hilac. These measurements indicate an oblate shape for  $^{28}\text{Si}$ , a prolate shape for  $^{32}\text{S}$ , and a spherical shape, within experimental error, for  $^{40}\text{Ar}$ .

The method used was almost exactly the same as in the Ne experiment.<sup>2</sup> The  $\gamma$ -ray yields from excited projectile and target nuclei at two projectile-scattering angles ( $160^\circ$  and  $90^\circ$ ) were measured simultaneously as particle- $\gamma$  coincidences between a NaI counter (7.5 cm  $\times$  7.5 cm) and a particle counter at each angle. A multidimensional-analysis program for the Hilac PDP-7 was used to store a  $\gamma$ -ray spectrum, a mixed particle spectrum from the two counters, and a time spectrum. In order to identify the particle counters in off-line analysis, the fast timing signals from one of the particle counters were delayed to produce two prompt peaks in the time spectra. The ratio of the excitation probability of the projectile to that of the target at each angle was deduced from the measured coincidence  $\gamma$ -ray spectra as:  $R^{160} = (N_P^{160}/N_T^{160}) \cdot (\epsilon_T/\epsilon_P)$ ,  $R^{90} = (N_P^{90}/N_T^{90}) \cdot (\epsilon_T/\epsilon_P)$ , where  $N$  is the integrated area of a photo-peak in the spectra, and  $\epsilon$  is the photo-peak efficiency of the NaI counter.

Superscripts indicate the scattering angle and subscripts distinguish between projectile (P) and target (T). A double ratio,  $R = (R^{160}/R^{90})$ , was also calculated which was less sensitive to many kinds of instrumental effects and to uncertainties in the  $B(E2)$  values of both projectile and target excitations. These ratios, after several corrections were made, were compared with calculated values using the deBoer-Winther Coulomb excitation program.<sup>3</sup>

This method is advantageous, because the excitation probability of the projectile is considerably more sensitive to its quadrupole moment than that of the target is to its moment. By measuring the target and projectile excitations simultaneously, several kinds of experimental uncertainties and instabilities cancel out, and by the proper choice of target nuclei, ambiguities involved in its excitation can be minimized. However, the estimation of the reorientation effect in the target excitation was more serious in the present case than in the Ne experiment, because the target nuclei were excited by heavier projectiles.

Since it was difficult to get low-energy beams of these projectiles at the Hilac, measurements have been made mainly with a  $^{206}\text{Pb}$  target. In the case of  $^{40}\text{Ar}$ , targets of  $^{120}\text{Sn}$  and  $^{130}\text{Te}$  were used as well as  $^{206}\text{Pb}$ . Because an accurate  $B(E2, 0^+ \rightarrow 2^+)$  value of  $^{206}\text{Pb}$  was needed, it was determined in separate experiments by a lifetime measurement of the  $2^+$  state using the Doppler-shift recoil-distance method.<sup>4</sup> The static quadrupole moment of the  $2^+$  state in  $^{206}\text{Pb}$  was assumed to be  $0.0 \pm 0.5 |Q_r|$ .<sup>5</sup>

The attenuation of the  $\gamma$ -ray angular distribution has been corrected using attenuation factors,  $G_2$  and  $G_4$ , determined in separate experiments. The interaction producing the attenuation was assumed to be magnetic. The correction to the  $\gamma$ -ray angular distribution and to the detector solid angle due to the motion of the  $\gamma$ -emitter was calculated by a numerical method. The higher velocities of projectile and recoiling target nuclei introduced larger corrections than was the case in the Ne experiment. However, the uncertainties due to these corrections are still only a small part of those in the final results.

Figure 1 shows the experimental results with best fits and calculated curves assuming  $Q = 0$  and  $\pm Q_r$ .<sup>5</sup> The best fits were obtained by least-square fittings to the experimental ratios. Because only two of the three ratios ( $R^{160}$ ,  $R^{90}$ ,  $R$ ) are independent, a correlated weight function was used in the fitting. Results of the fittings are summarized in Table I. Effects due to other low-lying  $0^+$ ,  $2^+$ ,  $4^+$ , and  $3^-$  states were estimated using experimental data<sup>6,7,8</sup> when available. No correction was applied unless the effect could be reliably calculated from experimental data, but additional uncertainty was given to the final values of  $B(E2, 0^+ \rightarrow 2^+)$  and  $Q$ . Corrections due to the effects of  $E4$  moments were also calculated using  $\beta_2$  and  $\beta_4$  from the (p,p') experiment<sup>9</sup> for  $^{28}\text{Si}$  and  $^{32}\text{S}$ , but for  $^{40}\text{Ar}$  only an estimated uncertainty was included. No correction was made for simultaneous excitation of projectile and target. The uncertainties introduced due to these effects were rather large, and could be appreciably reduced by further experimental information. The estimated corrections applied were small except for the case of  $^{32}\text{S}$ . The relatively large  $B(E2)$  values for the transitions between  $0_2^+$ ,  $2_2^+$ , and  $4_1^+$  states (3.78, 4.29, and 4.46 MeV) and the first excited state in  $^{32}\text{S}$  made a considerable change in  $Q(^{32}\text{S})$  ( $Q_{\text{uncorr}} = -0.14 \text{ b} \rightarrow Q_{\text{corr}} = -0.20 \text{ b}$ ).

The intrinsic quadrupole moments,  $Q_0$ ,<sup>10</sup> of the  $2^+$  states in the s-d shell nuclei deduced from the present projectile-excitation method are summarized in Fig. 2 together with values from other measurements.<sup>11-15</sup> The present result for  $^{28}\text{Si}$  agrees with a previous measurement.<sup>11</sup> The prolate shape ( $Q_0 > 0$ ) of  $^{32}\text{S}$  was somewhat unexpected; however, in this mass region calculations have shown that the minimum Hartree-Fock energies are very close for prolate and oblate shapes,<sup>16</sup> so that it does not seem unreasonable that

the addition of four nucleons can change the shape of a nucleus. In this regard, the different signs of the intrinsic quadrupole moments of  $^{28}\text{Si}(2^+)$  and  $^{27}\text{Al}(5/2^+)$  and of  $^{32}\text{S}(2^+)$  and  $^{33}\text{S}(3/2^+)$  are more striking. There are a number of possible reasons why the sign of  $Q_0$  for an odd-A nucleus might be different from that of a neighboring even-even one. Two rather obvious possibilities are: 1) if the state is a member of a  $K = 1/2$  band, the sign of  $Q_0$  will be opposite to that shown in the figure, and 2) if two shapes lie at nearly the same energy, the one preferred could be changed by the addition or removal of a particle from an orbital whose energy depends strongly on deformation. For all odd-A nuclei in this region, the sign of the quadrupole moment is correctly given by the position of the nucleus on the Nilsson single-particle diagram as determined by its nucleon number and the ground-state spin. The small static moment of  $^{40}\text{Ar}(2^+)$  (spherical shape) seems reasonable since this nucleus is quite near the doubly-closed shell.

#### ACKNOWLEDGMENTS

We are indebted to the Hilac crew for providing the heavy-ion beams. We wish to thank Drs. J. R. Leigh and K. H. Maier for their interest and discussions. Two of us (K.N. and J.L.Q.) are grateful to Professor I. Perlman for the opportunity to participate in this experiment at the Lawrence Radiation Laboratory.



## FOOTNOTES AND REFERENCES

<sup>†</sup> Work performed under the auspices of the U. S. Atomic Energy Commission.

\* On leave of absence from Osaka University, Osaka, Japan.

\*\* On leave of absence from CEN de Bordeaux-Gradignan, France, NATO fellow.

1. J. deBoer and J. Eichler, *Advances in Nuclear Physics*, Vol. 1, ed. by M. Baranger and E. Vogt, (Plenum Press Inc., New York, 1968).
2. K. Nakai, F. S. Stephens, and P. M. Diamond, submitted to *Nucl. Phys.*
3. A. Winther and J. deBoer, Coulomb Excitation, a collection of reprints, (Academic Press, New York, 1966) p. 303.
4. J. L. Quebert, K. Nakai, R. M. Diamond, and F. S. Stephens, to be published.
5.  $Q_r$  is the quadrupole moment calculated from the  $B(E2, 0^+ \rightarrow 2^+)$  value using the rigid rotor model.
6. S. J. Skorka, J. Hertel, and T. W. Retz-Schmidt, *Nuclear Data Tables* 2, 347 (1966).
7. G. T. Garvey, K. W. Jones, L. E. Carlson, A. G. Robertson, and D. F. H. Start, *Phys. Letters* 29B, 108 (1969); B. W. Sargent, J. R. Leslie, and J. H. Montague, *Contributions to the International Conference on Properties of Nuclear States*, Montreal, (1969); R. Lombard, P. Kossanyi-Demay, and G. R. Bishop, *Nucl. Phys.* 59, 398 (1964).
8. R. R. Johnson and R. J. Griffiths, *Nucl. Phys.* A117, 273 (1968).
9. R. de Swiniarski, C. Glashausser, D. L. Hendrie, J. Sherman, A. D. Bacher, and E. A. McClatchie, *Phys. Rev. Letters* 23, 317 (1969).
10.  $Q_0$  is deduced from the measured  $Q$  or  $B(E2, 0^+ \rightarrow 2^+)$  values in a model-dependent way using the formulae:

$$Q = \frac{3K^2 - I(I+1)}{(I+1)(2I+e)} Q_0, \quad \text{or} \quad B(E2, 0^+ \rightarrow 2^+) = \frac{5e^2}{16\pi} Q_0^2$$

11. O. Häusser, B. W. Hooton, D. Pelte, T. K. Alexander, and H. C. Evans, Phys. Rev. Letters 23, 320 (1969).
12. O. Häusser, B. W. Hooton, D. Pelte, T. K. Alexander, and H. C. Evans, Phys. Rev. Letters 22, 359 (1969).
13. D. Schwalm and B. Povh, Phys. Letters 29B, 103 (1969).
14. A. Bamberger, P. G. Bizzeti, and B. Povh, Phys. Rev. Letters 21, 1599 (1968).
15. V. S. Shirley, "Table of Nuclear Moments," in Hyperfine Structure and Nuclear Radiations, ed. by E. Matthias and D. A. Shirley, (North-Holland Publishing Co., Amsterdam, 1967) Appendix B; G. H. Fuller and V. W. Cohen, Nuclear Data Tables 5, 433 (1969).
16. G. Ripka, Advances in Nuclear Physics, Vol. 1, ed. by M. Baranger and E. Vogt, (Plenum Press Inc., New York, 1968).

#### FIGURE CAPTIONS

Fig. 1. Typical results of the experiments and analysis. The solid lines are the best fits (corresponding to an uncorrected  $Q$ ) and the dashed lines show the curves for  $Q = 0$  or  $\pm Q_r$ . The arrows indicate the "safe energy" ( $E_s$ ) defined in Ref. 1.

Fig. 2. Intrinsic quadrupole moments,  $Q_0^{10}$  in s-d shell nuclei. The circles indicate the intrinsic moments of first-excited  $2^+$  states determined from measured static quadrupole moments by: the present method (double circles), the method of the Chalk River group (closed circles), and that of the group at Heidelberg (open circles). The squares indicate the values calculated from measured  $B(E2, 0^+ \rightarrow 2^+)$  values. The intrinsic moments of odd-A

nuclei deduced from the spectroscopic quadrupole moment (assuming  $K = I$  except for  $^{19}\text{F}^*$  ( $K = 1/2, I = 5/2$ )) are shown by the diamonds.

Table I. Summary of the experimental results.


Target	Basis of calculations		Result (projectile)	
	$B(E2, \uparrow \text{target})$ $e^2 \times 10^{-50} \text{ cm}^4$	$Q(\text{target})$ (b)	$B(E2, 0^+ \rightarrow 2^+)$ $e^2 \times 10^{-50} \text{ cm}^4$	$Q(2^+)$ (b)
(a) $^{28}\text{Si}$ (1.78 MeV)				
$^{206}\text{Pb}$	$9.1 \pm 0.6^a$	$\begin{cases} +0.5  Q_r  \\ 0.0 \\ -0.5  Q_r  \end{cases}$	$3.3 \pm 0.4$ $3.3 \pm 0.4$ $3.3 \pm 0.4$	$0.126 \pm 0.045$ $0.110 \pm 0.045$ $0.098 \pm 0.045$
Summary	$9.1 \pm 0.6^a$	$(0.0 \pm 0.5  Q_r )^b$	$3.3 \pm 0.4$	$+0.11 \pm 0.05$
(b) $^{32}\text{S}$ (2.24 MeV)				
$^{206}\text{Pb}$	$9.1 \pm 0.6^a$	$\begin{cases} +0.5  Q_r  \\ 0.0 \\ -0.5  Q_r  \end{cases}$	$3.3 \pm 0.5$ $3.3 \pm 0.5$ $3.3 \pm 0.5$	$-0.192 \pm 0.055$ $-0.202 \pm 0.055$ $-0.215 \pm 0.055$
Summary	$9.1 \pm 0.6^a$	$(0.0 \pm 0.5  Q_r )^b$	$3.3 \pm 0.5$	$-0.20 \pm 0.06$
(c) $^{40}\text{Ar}$ (1.46 MeV)				
$^{120}\text{Sn}$	$\begin{cases} 23.0 \pm 1.2^c \\ 23.0 \pm 1.2^c \end{cases}$	$\begin{cases} +0.5  Q_r  \\ 0.0 \\ -0.5  Q_r  \end{cases}$ $(0.0 \pm 0.5  Q_r )^b$	$3.4 \pm 0.4$ $3.4 \pm 0.4$ <u><math>3.4 \pm 0.4</math></u>	$0.032 \pm 0.040$ $-0.021 \pm 0.040$ <u><math>-0.081 \pm 0.040</math></u>
$^{130}\text{Te}$	$30 \pm 3^d$	$-0.15 \pm 0.20^d$	$3.2 \pm 0.6$	$0.019 \pm 0.062$
$^{206}\text{Pb}$	$\begin{cases} 9.1 \pm 0.6^a \\ 9.1 \pm 0.6^a \end{cases}$	$\begin{cases} +0.5  Q_r  \\ 0.0 \\ -0.5  Q_r  \end{cases}$ $(0.0 \pm 0.5  Q_r )^b$	$3.0 \pm 0.5$ $3.0 \pm 0.5$ <u><math>3.0 \pm 0.5</math></u>	$0.052 \pm 0.072$ $0.040 \pm 0.072$ <u><math>0.022 \pm 0.072</math></u>
Summary	(Average)		$3.2 \pm 0.5$	$+0.01 \pm 0.04$

<sup>a</sup>Reference 4.

<sup>b</sup>Assumption:  $Q_r$  is the value calculated from the  $B(E2)$  using the rigid-rotor model.


<sup>c</sup>P. H. Stelson, F. K. McGowan, R. L. Robinson, W. T. Milner, and R. O. Sayer, Phys. Rev. 170, 1172 (1968).

<sup>d</sup>A. Christy, I. Hall, R. P. Harper, I. M. Nagib, and B. Wakefield, Contribution to the International Conference on Properties of Nuclear States, Montreal, (1969).



XBL 701-2138

Fig. 1



XBL 701-2137

Fig. 2

LEGAL NOTICE

*This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:*

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

*As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE RADIATION LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720