

eScholarship

Combinatorial Theory

Title

\(P\)-partitions with flags and back stable quasisymmetric functions

Permalink

<https://escholarship.org/uc/item/0db8n8q1>

Journal

Combinatorial Theory, 4(2)

ISSN

2766-1334

Authors

Nadeau, Philippe

Tewari, Vasu

Publication Date

2024

DOI

10.5070/C64264232

Supplemental Material

<https://escholarship.org/uc/item/0db8n8q1#supplemental>

Copyright Information

Copyright 2024 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at

<https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

P -PARTITIONS WITH FLAGS AND BACK STABLE QUASISYMMETRIC FUNCTIONS

Philippe Nadeau^{*1} and Vasu Tewari^{†2}

¹Universite Claude Bernard Lyon 1, CNRS, Ecole Centrale de Lyon, INSA Lyon, Université Jean Monnet, ICJ UMR5208,
69622 Villeurbanne, France
nadeau@math.univ-lyon1.fr

²Department of Mathematical and Computational Sciences, University of Toronto Mississauga, Mississauga, ON L5L 1C6,
Canada
vasu.tewari@utoronto.ca

Submitted: Mar 15, 2023; Accepted: Apr 11, 2024; Published: Sep 30, 2024

© The authors. Released under the CC BY license (International 4.0).

Abstract. Stanley's theory of (P, ω) -partitions is a standard tool in combinatorics. It can be extended to allow for the presence of a restriction, that is a given maximal value for partitions at each vertex of the poset, as was shown by Assaf and Bergeron. Here we present a variation on their approach, which applies more generally. The enumerative side of the theory is more naturally expressed in terms of back stable quasisymmetric functions. We study the space of such functions, following the work of Lam, Lee and Shimozono on back stable symmetric functions. As applications we describe a new basis for the ring of polynomials that we call forest polynomials. Additionally we give a signed multiplicity-free expansion for any monomial expressed in the basis of slide polynomials.

Keywords. P-partitions, quasisymmetric functions, slide polynomials

Mathematics Subject Classifications. 05E05, 06A07

Introduction

The study of (P, ω) -partitions was initiated by Stanley in his thesis [Sta72] wherein he established his *fundamental theorem* laying the groundwork. Since then (P, ω) -partitions, and several close cousins thereof, have proven to be very useful in combinatorics. We refer the reader to a recent survey by Gessel [Ges16] where the history is nicely recounted; see also [GR14, Section 5]. Several aspects of (P, ω) -partitions manifest themselves in the combinatorics of certain generating series $K_{(P, \omega)}$ which turn out to be quasisymmetric functions. A class that then emerges naturally is when P is a chain, in which case one obtains Gessel's fundamental quasisymmetric functions.

*P.N. is partially supported by French ANR grant ANR19-CE48-011-01 (COMBINÉ).

†V. T. acknowledges the support from Simons Collaboration Grant #855592.

An immediate, and pleasant, consequence of Stanley’s fundamental theorem is that $K_{(P,\omega)}$ expands *positively* in terms of fundamental quasisymmetric functions, i.e. the expansion involves nonnegative integer coefficients.

A polynomial analogue of the latter was introduced by Assaf and Searles [AS17] under the name slide polynomials. These were then realized by Assaf and Bergeron [AB20] in a manner akin to how fundamental quasisymmetric functions are defined via (P, ω) -partitions. To this end, the authors in *loc. cit.* introduced the notion of (P, ω, ρ) -partitions, which are (P, ω) -partitions in the traditional sense now subject to upper bound constraints imposed by the *restriction* ρ . The generating series $K_{(P,\omega)}$ from before turn into polynomials $K_{(P,\omega,\rho)}$. The natural analogue of Stanley’s fundamental theorem holds in this restricted setting as well, and Assaf–Bergeron establish that $K_{(P,\omega,\rho)}$ expands positively in terms of slide polynomials when ρ is AB-flag; see Section 2.2 for the precise definition.

The main purpose of this note is to introduce (P, Φ) -partitions, which are a mild generalization of those (P, ω, ρ) -partitions that satisfy the AB-flag condition. They form a class that is also easier to deal with, leading to simplified proofs. In particular, we find ourselves in the following desirable setting once more—an analogue of Stanley’s fundamental theorem holds and this allows us to describe the set of (P, Φ) -partitions as a disjoint union of (L, Φ) -partition sets as L varies over the set of linear extensions of P . This then allows us to express $K_{(P,\Phi)}$ as a sum of the $K_{(L,\Phi)}$, each of which either equals a slide polynomial or 0. As a particular instance of $K_{(P,\Phi)}$ we introduce forest polynomials, which are studied in greater detail in the context of the permutahedral variety in [NT24].

It turns out that the *back stable* setting is a particularly convenient place to view these results. By allowing our (P, Φ) -partitions to take values in the negative integers, the polynomials $K_{(P,\Phi)}$ become generating series $\overleftarrow{K}_{(P,\Phi)}$ in the alphabet x_i where $i < N$ for some $N \in \mathbb{Z}$. The $\overleftarrow{K}_{(P,\Phi)}$ naturally live inside the space of *back stable quasisymmetric functions* \overleftarrow{QR} , which we define and study in Section 4. In particular we establish that the back stable limits of slide polynomials form a basis for \overleftarrow{QR} .

Finally we give an explicit description expressing a monomial in the basis of slide polynomials. It turns out that the coefficients that arise belong to $\{0, \pm 1\}$. In other words, this expansion is *signed multiplicity-free*.

1. Combinatorial preliminaries

Given a sequence $\mathbf{c} = (c_i)_{i \in \mathbb{Z}}$ of nonnegative integers, we define its *support* $\text{Supp}(\mathbf{c})$ to be the set of indices i such that $c_i > 0$. We call \mathbf{c} an \mathbb{N} -vector if $\text{Supp}(\mathbf{c})$ is finite. If $\text{Supp}(\mathbf{c}) \subseteq [n] := \{1, \dots, n\}$ for some positive integer n , we occasionally write $\mathbf{c} = (c_1, \dots, c_n)$ and refer to \mathbf{c} as a *weak composition*. For any \mathbb{N} -vector \mathbf{c} , we let $|\mathbf{c}| := \sum_i c_i$ denote its *weight*. The finite sequence of positive integers obtained by omitting all 0s from \mathbf{c} is called a *strong composition*. Henceforth, by composition, we shall always mean strong composition. If $\alpha = (\alpha_1, \dots, \alpha_{\ell(\alpha)})$ is a composition of weight $r \geq 0$, then we denote this by $\alpha \vDash r$. Here $\ell(\alpha)$ is the *length* of α . The unique composition of weight and length both 0 will be denoted by \emptyset . We will need two operations on compositions— *concatenation* and *near-concatenation*. Both take compositions $\alpha = (\alpha_1, \dots, \alpha_p)$

and $\beta = (\beta_1, \dots, \beta_q)$ as input; the former produces $\alpha \cdot \beta = (\alpha_1, \dots, \alpha_p, \beta_1, \dots, \beta_q)$ whereas the latter produces $(\alpha_1, \dots, \alpha_p + \beta_1, \beta_2, \dots, \beta_q)$.

When writing our \mathbb{N} -vectors $\mathbf{c} = (c_i)_{i \in \mathbb{Z}}$ we use a vertical bar to distinguish the ‘left half’ (\dots, c_{-1}, c_0) and the ‘right half’ (c_1, c_2, \dots) . For example $\mathbf{c} = (\dots, 0, 2|0, 1, 2, 0, 2, 0, \dots)$ has $c_0 = 2$ separated from $c_1 = 0$ by a vertical bar. We will be particularly interested in \mathbb{N} -vectors supported on the set of positive integers \mathbb{Z}_+ , in which case we will omit the vertical bar and the zeros left of it.

We denote the set of words in an alphabet \mathcal{A} by \mathcal{A}^* . We denote the *length* of any word $w \in \mathcal{A}^*$ by $\ell(w)$. We let ϵ denote the *empty word*, i.e. the unique word of length 0. A subset of \mathcal{A}^* which we care about is $\text{Inj}(\mathcal{A})$, the set of *injective words*, i.e. words with all distinct letters.

We now proceed to define the ordered alphabet that is relevant for our purposes. It is obtained by ‘augmenting’ \mathbb{Z} .

Definition 1.1. Let $\overline{\mathbb{Z}}$ be the ordered alphabet with letters $i^{[j]}$ where $i \in \mathbb{Z}$ and $j \in \mathbb{Z}_+$. We have a linear order on $\mathbb{Z} \sqcup \overline{\mathbb{Z}}$ given by $i < i^{[1]} < i^{[2]} < i^{[3]} < \dots < i + 1$ for all i . We define the *value* of $i^{[j]}$ by $\text{val}(i^{[j]}) = i$.

The order is the lexicographic order on $\mathbb{Z} \times \mathbb{Z}_+$, but the notation will serve to highlight the prevalent role of the first factor.

There is a natural way to go from \mathbb{Z}^* to $\text{Inj}(\overline{\mathbb{Z}})$: given w in \mathbb{Z}^* , one associates a word $W = \text{stan}(w)$ by labeling the occurrences of the same letter i in w from left to right by $i^{[1]}, i^{[2]}, \dots$. For instance $w = 1221625$ gets mapped to $W = 1^{[1]}2^{[1]}2^{[2]}1^{[2]}6^{[1]}2^{[3]}5^{[1]}$. This process is injective, and will be used as our natural embedding

$$\text{stan} : \mathbb{Z}^* \hookrightarrow \text{Inj}(\overline{\mathbb{Z}}). \tag{1.1}$$

2. Revisiting Stanley and Assaf–Bergeron

We first recall the setting and main results of the celebrated theory of (P, ω) -partitions due mainly to Stanley. We then explain how this theory extends in the presence of a *restriction* by recalling pertinent results of Assaf and Bergeron.

In this section (P, \leq_P) is a finite poset. We will interchangeably use P to denote both the poset as well as the set underlying the poset. We denote the cover relation by \prec_P .

Definition 2.1. A P -partition is a function $f : P \rightarrow \mathbb{Z}_+$ such that $f(u) \geq f(v)$ whenever $u \prec_P v$.

2.1. Stanley’s theory of (P, ω) -partitions.

The starting point is to fix in addition a bijective labeling $\omega : P \rightarrow \{1, \dots, \#P\}$. The pair (P, ω) then forms a labeled poset. A (P, ω) -partition is a P -partition f such that $f(u) > f(v)$ whenever $u \prec_P v$ and $\omega(u) > \omega(v)$. Let $\text{Part}(P, \omega)$ denote the set of (P, ω) -partitions.

A *linear extension* L of P is a linear ordering of P extending \leq_P . Thus a linear extension is a totally ordered set with P as its underlying set. Let $\text{Lin}(P)$ be the set of linear extensions of P . For example, the (P_0, ω_0) -partitions for the example illustrated in Figure 2.1 are the functions $f : P_0 \rightarrow \mathbb{Z}_+$ that satisfy $f(a) \geq f(b)$ and $f(c) > f(b)$. Throughout this note, we depict strict inequalities using dashed edges in Hasse diagrams.




Figure 2.1: A labeled poset and its two linear extensions.

Theorem 2.2 (Stanley’s fundamental theorem [Sta72, Theorem 6.2]). *We have*

$$\text{Part}(P, \omega) = \bigsqcup_{L \in \text{Lin}(P)} \text{Part}(L, \omega).$$

For (P_0, ω_0) in Figure 2.1, the two linear extensions are shown together with their induced labeling on the right. Theorem 2.2 says that (P_0, ω_0) -partitions are the functions f satisfying either $f(a) \geq f(c) > f(b)$ or $f(c) > f(a) \geq f(b)$, as can be directly checked.

Now for any (P, ω) one can consider the generating function

$$K_{(P, \omega)} = \sum_f \prod_{u \in P} x_{f(u)} \quad (2.1)$$

where the sum is over the set of all (P, ω) -partitions. Then Stanley’s theorem has the following expansion as an immediate corollary:

$$K_{(P, \omega)} = \sum_{L \in \text{Lin}(P)} K_{(L, \omega)}. \quad (2.2)$$

What makes this particularly useful is that the series $K_{(L, \omega)}$ are precisely the fundamental quasisymmetric functions introduced by Gessel [Ges84], as we now detail.

Recall that a series f in the variables x_i for $i \in I$, with I an interval in \mathbb{Z} , is *quasisymmetric* if for any composition $\alpha = (\alpha_1, \dots, \alpha_k)$ and any subsets $\{i_1 < \dots < i_k\}$ and $\{j_1 < \dots < j_k\}$ of I , the coefficients of $x_{i_1}^{\alpha_1} \dots x_{i_k}^{\alpha_k}$ and $x_{j_1}^{\alpha_1} \dots x_{j_k}^{\alpha_k}$ in f are the same.

Let \mathbf{x}_+ denote the set of *positively-indexed variables* $\{x_i \mid i \in \mathbb{Z}_+\}$. Given a subset S of $\{1, \dots, r-1\}$, the *fundamental quasisymmetric function* $F_{r, S} \in \mathbb{Q}[\mathbf{x}_+]$ is defined by

$$F_{r, S} = \sum_{\substack{1 \leq i_1 \leq \dots \leq i_r \\ i_j < i_{j+1} \text{ if } j \in S}} x_{i_1} \dots x_{i_r}. \quad (2.3)$$

For instance $F_{3, \{1\}} = \sum_{1 \leq i_1 < i_2 \leq i_3} x_{i_1} x_{i_2} x_{i_3}$ and $F_{3, \{2\}} = \sum_{1 \leq i_1 \leq i_2 < i_3} x_{i_1} x_{i_2} x_{i_3}$. The $F_{r, S}$ for $r \geq 0$ and S a subset of $\{1, \dots, r-1\}$ form a basis of the space of quasisymmetric functions in \mathbf{x}_+ . Let $\alpha = (\alpha_1, \dots, \alpha_\ell) \vDash r$ be the composition corresponding to $S \subseteq \{1, \dots, r-1\}$ under the folklore correspondence given by $S = \{\alpha_1, \alpha_1 + \alpha_2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{\ell-1}\}$. We will then use freely the notation F_α to denote $F_{r, S}$. For instance $F_{(1,2)} = F_{3, \{1\}}$ and $F_{(2,1)} = F_{3, \{2\}}$.

Now if L is a chain $v_1 \prec_L \cdots \prec_L v_r$ with a labeling ω , define

$$\text{Des}(\omega) = \{i \in [r - 1] \mid \omega(v_i) > \omega(v_{i+1})\}.$$

If we denote the composition of r corresponding to $\text{Des}(\omega)$ by α_ω , then it is easily verified that

$$K_{(L,\omega)} = F_{\text{rev}(\alpha_\omega)}.$$

Here rev reverses its input. This shows that the expansion (2.2) expresses any $K_{(P,\omega)}$ positively in the basis of fundamental quasisymmetric functions. For the poset in Figure 2.1, we get the expansion $K_{(P_0,\omega_0)} = F_{(2,1)} + F_{(1,2)}$.

2.2. Restricted partitions

We now want to constrain P -partitions to be dominated by certain fixed values at each element of P . Let (P, ω) be a labeled poset.

Definition 2.3. Fix $\rho : P \rightarrow \mathbb{Z}_+$ a map, called a *restriction*. We define (P, ω, ρ) -partitions as those (P, ω) -partitions f that satisfy $f(u) \leq \rho(u)$ for all $u \in P$.

Let $\text{Part}(P, \omega, \rho)$ be the set of all (P, ω, ρ) -partitions. Assaf and Bergeron [AB20] have already considered this extension. We recall some of their results (see Remark 2.6 for easy comparison). First they note that Theorem 2.2 adapts immediately:

Theorem 2.4 ([AB20, Theorem 3.14]). *For any restriction ρ , we have*

$$\text{Part}(P, \omega, \rho) = \bigsqcup_{L \in \text{Lin}(P)} \text{Part}(L, \omega, \rho).$$

So, for instance, if we impose the restriction $\rho(a) = 3, \rho(b) = \rho(c) = 2$ for the poset P_0 in Figure 2.1, then $\text{Part}(P_0, \omega, \rho)$ comprises exactly two functions: $f(a) = f(c) = 1, f(b) = 2$ and $f(c) = 1, f(a) = f(b) = 2$. The former comes from the linear extension in the middle and the latter from that on the right.

Definition 2.3 leads to a restricted version of (2.1) allowing us to define generating functions $K_{(P,\omega,\rho)}$, which are now polynomials, in the obvious manner. Theorem 2.4 then naturally gives a restricted version of (2.2) for $K_{(P,\omega,\rho)}$.

Now the issue here is that, when L is a linear order, the family of functions $K_{(L,\omega,\rho)}$ for varying ω and ρ is too large, and is in particular not free. There is however a very natural subfamily that forms a basis of the space of all polynomials, namely the *slide polynomials* [AS17] to which we will come back later. A remaining issue is that, for general restrictions ρ , the expansion of $K_{(P,\omega,\rho)}$ is not necessarily positive in the slide basis, cf. [AB20, Example 3.12].

Assaf and Bergeron are thus led to consider a restricted set of restrictions ρ . These are defined as follows: Say that the restriction ρ is an *AB-flag* for (P, ω) if the following conditions hold (cf. [AB20, Definition 4.1]).

- (AB1) If $u \prec_P v$, then $\rho(u) \geq \rho(v)$;
- (AB2) If $u \prec_P v$ and $\rho(u) > \rho(v)$, then $\omega(u) > \omega(v)$.

Their main result is then the following:

Theorem 2.5. *If ρ is an AB-flag for (P, ω) , the polynomial $K_{(P, \omega, \rho)}$ expands in the slide basis with nonnegative integral coefficients.*

We postpone the definition of slide polynomials to the next section, preferring to define them via a seemingly different perspective than their original definition [AS17, Definition 3.6].

In order to prove Theorem 2.5, Assaf and Bergeron modify ρ to have it satisfy in addition a certain “well-labeled” restriction; see [AB20, Section 4]. This modification is needed because the AB-flag property alone does not transfer to linear extensions as can be easily seen. As Assaf and Bergeron demonstrate in [AB20, Proposition 4.3], the set of flagged (P, ρ) -partitions remains unchanged in spite of these modifications. The proof of [AB20, Proposition 4.3] has a minor gap. To assist the reader following their proof, we note some key aspects and record the fix. They employ the fact that if one takes the Hasse diagram of any (P, ω) and removes all strict edges, then the resulting connected components inherit an ordering from P . Note that they start with an AB-flag but at this stage this plays no role since a constant AB-flag puts no restriction on ω . As the example in Figure 2.2 shows, we can in fact get two connected components that are not ordered. That being noted, the removal of strict edges does indeed give a preorder on the connected components. Now if one considers the induced order, then the AB-flag property forces all elements of these supercomponents to have the same ρ -values, and the remainder of their proof goes through.




Figure 2.2: Counterexample to an argument in the proof of [AB20, Proposition 4.3].

In the next section, we will find a simpler condition on restrictions, which at the same time is more general than AB-flags and for which (positivity of) the slide expansion will follow immediately.

Remark 2.6. Our setup differs slightly from [AB20]. First, our P -partitions are order-reversing instead of order-preserving. To go between the two formulations, one simply has to go from a poset (P, \leq_P) to its opposite (P, \geq_P) . Also, the function ω is implicit in their work, since they consider the set underlying P to be $\{1, \dots, \#P\}$ already. This allows for compact notation, but is not well adapted to having different labelings on a given abstract poset. Furthermore, the forthcoming notion of (P, Φ) -partitions would be cumbersome to use in this setting.

3. (P, Φ) -partitions

We now define our notion of (P, ω) -partitions with restrictions. Recall the ordered alphabet $\overline{\mathbb{Z}_+}$ from Section 1, given by letters $i^{[j]}$ with $i, j \in \mathbb{Z}_+$.

Definition 3.1 (Labeled flag Φ). Let P be a poset. A *labeled flag* on P is an *injective* function $\Phi : P \rightarrow \overline{\mathbb{Z}_+}$.

Definition 3.2 ((P, Φ) -partitions). Let (P, Φ) be a poset with a labeled flag. A (P, Φ) -*partition* is a function $f : P \rightarrow \mathbb{Z}_+$ such that for any $u, v \in P$:

- $f(u) \geq f(v)$ if $u \prec_P v$;
- $f(u) > f(v)$ if $u \prec_P v$ and $\Phi(u) > \Phi(v)$;
- $f(u) \leq \text{val}(\Phi(u))$.

We denote the set of all (P, Φ) -partitions by $\text{Part}(P, \Phi)$. An example is given in Figure 3.1. On the left is the Hasse diagram of the *diamond* poset with the labeled flag in blue. On the right are the five (P, Φ) -partitions. The e -labels on the covers may be momentarily ignored.




Figure 3.1: A poset with labeled flag (P, Φ) and its five (P, Φ) -partitions.

Let us note immediately that the notion of (P, Φ) -partitions is simply a concise way to encode a class of (P, ω, ρ) -partitions with a certain condition on ρ . Namely, say that ρ is an *LF-flag* if it satisfies

(LF) For any $u, v \in P$, $\rho(u) > \rho(v)$ implies $\omega(u) > \omega(v)$.

Proposition 3.3. Fix a poset P . Then the following hold.

- (1) If Φ is a labeled flag, then $\text{Part}(P, \Phi) = \text{Part}(P, \omega, \rho)$ for some ω, ρ satisfying (LF).
- (2) Conversely, for any ω, ρ satisfying (LF), there exists a labeled flag Φ such that $\text{Part}(P, \Phi) = \text{Part}(P, \omega, \rho)$.

Proof. For (1), define $\omega = \omega(\Phi)$ as the unique bijection $P \rightarrow \{1, \dots, \#P\}$ that satisfies $\omega(u) < \omega(v)$ if and only if $\Phi(u) < \Phi(v)$. Define also $\rho = \rho(\Phi)$ by $\rho(u) = \text{val}(\Phi(u))$.

We first claim that ρ is LF-flag. Indeed, we have $\rho(u) > \rho(v)$ if and only if $\text{val}(\Phi(u)) > \text{val}(\Phi(v))$, which in turn implies that $\Phi(u) > \Phi(v)$. By the definition of ω this immediately yields $\omega(u) > \omega(v)$.

We now claim that $\text{Part}(P, \Phi) = \text{Part}(P, \omega, \rho)$. Say $f \in \text{Part}(P, \Phi)$. Then we know that $f(u) \leq \text{val}(\Phi(u)) = \rho(u)$ for all $u \in P$. Furthermore, $u \prec_P v$ implies $f(u) \geq f(v)$. Finally $u \prec_P v$ and $\Phi(u) > \Phi(v)$ implies $f(u) > f(v)$. Now note that $\Phi(u) > \Phi(v) \iff \omega(u) > \omega(v)$. It follows that $f \in \text{Part}(P, \omega, \rho)$. The inclusion $\text{Part}(P, \omega, \rho) \subseteq \text{Part}(P, \Phi)$ is essentially the same proof.

For the converse in (2), given ρ an LF-flag define $\Phi(u) = \rho(u)^{[\omega(u)]}$. Like before we need to show that $\text{Part}(P, \omega, \rho) = \text{Part}(P, \Phi)$, and only the second condition in Definition 3.2 is not trivial. It follows however readily from the fact that $\Phi(u) > \Phi(v)$ is here equivalent to $\omega(u) > \omega(v)$ because of the condition (LF). \square

3.1. Comparison with AB-flags

The notion of (P, Φ) -partitions extends the notion of AB-flagged partitions defined in the previous section:

Proposition 3.4. *Let ρ be an AB-flag for (P, ω) . Define a labeled flag on P by $\Phi(u) = \rho(u)^{[\omega(u)]}$. Then*

$$\text{Part}(P, \omega, \rho) = \text{Part}(P, \Phi).$$

Proof. Elements of both sets are P -partitions that are bounded by ρ . It remains to show that the strictness conditions are imposed for the same cover relations. Thus the claim we have to prove is that if $u \prec_P v$, then $\Phi(u) > \Phi(v)$ if and only if $\omega(u) > \omega(v)$.

Pick then $u, v \in P$ such that $u \prec_P v$. Now $\Phi(u) > \Phi(v)$ is equivalent to either $\rho(u) > \rho(v)$ or $(\rho(u) = \rho(v) \text{ and } \omega(u) > \omega(v))$ by definition of the order on $\overline{\mathbb{Z}_+}$. In the latter case, we are already done. In the former case, condition (AB2) implies $\omega(u) > \omega(v)$. The converse also holds since we have necessarily $\rho(u) \geq \rho(v)$ by (AB1). \square

The next example shows that in fact, the notion of (P, Φ) -partitions is strictly more general than that of AB-flagged partitions.

Example 3.5. Consider (P, Φ) given by the labeled Hasse diagram on the left of Figure 3.1. Then $\text{Part}(P, \Phi)$ consists of the five partitions shown on the right.

Let us consider possible choices of (ω, ρ) such that $\text{Part}(P, \omega, \rho) = \{f_1, f_2, f_3, f_4, f_5\}$ (Note that we know by Proposition 3.3 that such a choice is always possible). The cover relation e_1 is necessarily strict, otherwise the partition f with constant value 1 would be valid. The other cover relations are necessarily weak since there is a f_i with equality for each of them. This imposes a unique choice for ω , namely $\omega(v_i) = i$ for $i = 1, 2, 3, 4$. As for ρ , we must clearly have $\rho(v_2) = 3$ and $\rho(v_3) \geq 3$. Then $\rho(v_1) = 1$ is also imposed, since any other choice would make it possible to switch the value of v_1 in f_1 from 1 to 2, resulting in a partition f not in the allowed f_i 's.

Now assume in addition that ρ is an AB-flag. By (AB1) the remaining choices for ρ are $\rho(v_4) = 1$ and $\rho(v_3) = 3$. This gives us $\rho(v_3) > \rho(v_4)$ while $\omega(v_3) < \omega(v_4)$, contradicting (AB2). This shows that there are no ω, ρ with ρ an AB-flag such that $\text{Part}(P, \Phi) = \text{Part}(P, \omega, \rho)$.

3.2. Slide polynomials

Since (P, Φ) -partitions form a certain class of (P, ω, ρ) -partitions, we have the immediate corollary of Theorem 2.4:

Corollary 3.6. *Let (P, Φ) be a poset with a labeled flag. We have*

$$\text{Part}(P, \Phi) = \bigsqcup_{L \in \text{Lin}(P)} \text{Part}(L, \Phi).$$

We introduce the generating polynomial

$$K_{(P, \Phi)} = \sum_{f \in \text{Part}(P, \Phi)} \prod_{u \in P} x_{f(u)}. \tag{3.1}$$

Corollary 3.6 gives:

$$K_{(P, \Phi)} = \sum_{L \in \text{Lin}(P)} K_{(L, \Phi)}. \tag{3.2}$$

We will now see that all the (nonzero) $K_{(L, \Phi)}$ are *slide polynomials* which we introduce next. Write $L = v_1 \prec v_2 \prec \dots \prec v_r$. Any (L, Φ) -partition f can be encoded in a sequence (i_1, \dots, i_r) with $i_j = f(v_j)$, while (L, Φ) is encoded in the injective word $W = a_1 \dots a_r \in \text{Inj}(\overline{\mathbb{Z}_+})$ by simply setting $a_i = \Phi(v_i)$. The corresponding generating function $K_{(L, \Phi)}$ is thus given by the explicit series

$$\mathfrak{F}(W) := K_{(L, \Phi)} = \sum_{\substack{i_1 \geq i_2 \geq \dots \geq i_r > 0 \\ i_j > i_{j+1} \text{ if } a_j > a_{j+1} \\ i_j \leq \text{val}(a_j)}} x_{i_1} x_{i_2} \dots x_{i_r}. \tag{3.3}$$

Example 3.7. Consider $W = 3^{[1]}3^{[2]}1^{[1]}$. Then (3.3) says

$$\mathfrak{F}(W) = \sum_{i_1 \geq i_2 \geq i_3 > 0} x_{i_1} x_{i_2} x_{i_3}$$

where $3 \geq i_1 \geq i_2 > i_3$ and $i_3 \leq 1$. Thus $\mathfrak{F}(W) = x_3^2 x_1 + x_3 x_2 x_1 + x_2^2 x_1$.

Note that replacing the $3^{[2]}$ in the middle by any letter in $\overline{\mathbb{Z}_+}$ larger than $3^{[2]}$ does not alter $\mathfrak{F}(W)$. Finally note that it can be the case that the sum defining $\mathfrak{F}(W)$ is empty. For instance, if $W = 1^{[2]}1^{[1]}$, then $\mathfrak{F}(W) = 0$. This ‘anomaly’ will be fixed when we work in the back stable setting.

Given a weak composition \mathbf{c} with positive support, consider the following word $W_{\mathbf{c}}$: for each $i > 0$, let $X_i = i^{[1]}i^{[2]} \dots i^{[c_i]}$. Then $W_{\mathbf{c}} = X_{i_1} X_{i_2} \dots X_{i_m}$ where $i_1 > i_2 > \dots > i_m$ are the indices i such that $c_i > 0$. For instance if $\mathbf{c} = (0, 2, 4, 0, 1, 0, \dots)$ then $W_{\mathbf{c}} = 5^{[1]}3^{[1]}3^{[2]}3^{[3]}3^{[4]}2^{[1]}2^{[2]}$.

Definition 3.8 ([AB20]). The *slide polynomial* \mathfrak{F}_c is defined as $\mathfrak{F}(W_c)$.

Proposition 3.9. For any $W \in \text{Inj}(\overline{\mathbb{Z}_+})$, the polynomial $\mathfrak{F}(W)$ is either zero or is equal to \mathfrak{F}_c for a unique c .

Proof. The uniqueness of c holds because slide polynomials form a basis of $\mathbb{Q}[\mathbf{x}_+]$. This is the content [AS17, Theorem 3.9], and follows readily from a triangular change of basis with monomials.

Now given W , we must construct a word W_c such that $\mathfrak{F}(W) = \mathfrak{F}(W_c)$. By scanning W from left to right, let us show one can define such a word in an algorithmic fashion. One has to find a standard word W_c with the same descents as W , which moreover gives the same upper bounds. We now explain the construction.

We first construct a word U in \mathbb{Z}^* . If W is empty, then so is U . If $W \in \overline{\mathbb{Z}_+}$, then $U = \text{val}(W)$. Otherwise, say $W = w_1 \dots w_r$ for $r \geq 2$, with w_i pairwise distinct letters in $\overline{\mathbb{Z}_+}$. Suppose we have scanned letters w_1 through w_k for $1 \leq k < r$ and constructed a word $u_1 \dots u_k$. If $w_{k+1} < w_k$, then $u_{k+1} := \min(\text{val}(w_{k+1}), u_k - 1)$. If $w_{k+1} > w_k$, then $u_{k+1} := u_k$. Repeating this we get $U = u_1 \dots u_r \in \mathbb{Z}^*$, a word with nonincreasing letters.

If any letter u_i is nonpositive, then $\mathfrak{F}(W) = 0$. Otherwise $\text{stan}(U) = W_c$ for a unique \mathbb{N} -vector c with positive support, and by induction one checks that W_c satisfies $\mathfrak{F}(W) = \mathfrak{F}(W_c)$ as wanted. \square

We refer to the resulting W_c as $\text{RS}(W)$. For instance, let $W = 5^{[1]}6^{[5]}8^{[3]}3^{[2]}3^{[1]}1^{[2]}2^{[1]}3^{[3]}$. Then $U = 55532111$ which in turn means that $\text{RS}(W) = 5^{[1]}5^{[2]}5^{[3]}3^{[1]}2^{[1]}1^{[1]}1^{[2]}1^{[3]}$. This equals W_c for $c = (\dots, 0|3, 1, 1, 0, 3, 0, \dots)$. Note that if the last letter in W is replaced by say $2^{[3]}$, then one gets $\mathfrak{F}(W) = 0$. This is because $W_c = \text{RS}(W)$ satisfies $\text{Supp}(c) \not\subseteq \mathbb{Z}_+$.

The combinatorics explained in the proof are already presented by Reiner and Shimozono in [RS95, Lemma 8]: the minor differences are that the authors in *loc. cit.* work from right to left, and work with \mathbb{Z}_+ instead of $\overline{\mathbb{Z}_+}$.

3.3. Forest polynomials

We close this section by introducing a novel family of polynomials that forms the core of the work [NT24] that links the quotient of $\mathbb{Q}[x_1, \dots, x_n]$ by the ideal of positive degree quasisymmetric polynomials in x_1, \dots, x_n to the Schubert class expansion of the cohomology class of the permutahedral variety Perm_n . We lay some foundations for that work here.




Figure 3.2: An indexed forest with its flag.

Let S be a finite set of integers. It decomposes uniquely as $S = I_1 \sqcup \dots \sqcup I_k$, where each I_j is a maximal subset of consecutive integers in S . An *indexed forest* F on S is the data of a rooted binary tree T_j with $|I_j|$ internal nodes for any $j \in [k]$. Its size $|F|$ is the cardinality of S , and we represent the latter by unit intervals of the integer line. See Figure 3.2 for an indexed forest (ignoring the labels in red for now) with $|F| = 6$. It comprises three trees supported on the intervals $[2, 4]$, $[7]$, and $[11, 12]$ from left to right.

Let us denote the internal nodes (i.e. everything but the leaves) in F by $\text{IN}(F)$. We will treat an indexed forest F as a Hasse diagram of a poset (also called F) with underlying set $\text{IN}(F)$. *What roles do the intervals play?* They determine a labeled flag Φ_F as follows: for each left leaf lf , let m be the label of the unit interval to its right. Let v_1, \dots, v_k be the inner nodes on the left branch ending at lf , from bottom to top. Then define $\Phi_F(v_i) := m^{[i]}$ for $i = 1, \dots, k$. In Figure 3.2 the left leaves have labels 2, 4, 7 and 11, completely determining Φ_F as shown in red.

Definition 3.10. The *forest polynomial* $\mathfrak{P}_F \in \mathbb{Q}[\mathbf{x}_+]$ is defined as

$$\mathfrak{P}_F = K_{(F, \Phi_F)}.$$

Explicitly, \mathfrak{P}_F is the sum of monomials $\prod_{v \in \text{IN}(F)} x_{f(v)}$ over all labelings $f : \text{IN}(F) \rightarrow \mathbb{Z}_+$ satisfying $f(v) \leq \text{val}(\Phi_F(v))$ for all v , and that are weakly increasing down left edges and strictly increasing down right edges.

Thus \mathfrak{P}_F is the sum of monomials over ‘semi-standard’ increasing labelings of the internal nodes of the indexed forest F . For example, given the indexed forest F in Figure 3.3 one has

$$\mathfrak{P}_F = \sum_{\substack{2 \geq a \geq b \\ 4 \geq c > b}} x_a x_b x_c = x_2^2 x_4 + x_1 x_2 x_4 + x_1^2 x_4 + x_2^2 x_3 + x_1 x_2 x_3 + x_1^2 x_3 + x_1^2 x_2 + x_1 x_2^2.$$




Figure 3.3: An indexed forest F with $c(F) = (0, 2, 0, 1)$.

We can obtain the expansion of \mathfrak{P}_F into slide polynomials thanks to (3.2). Note that a linear extension of the poset F corresponds to a *decreasing forest*, i.e. a bijective labeling $\text{IN}(F) \rightarrow \{1, \dots, |F|\}$ that is decreasing down edges. Let $\text{Dec}(F)$ be the set of such decreasing labelings, and for any such labeling ℓ we note $W(\ell)$ the injective word read from the flag Φ_F . We thus get

$$\mathfrak{P}_F = \sum_{\ell \in \text{Dec}(F)} \mathfrak{F}(W(\ell)). \tag{3.4}$$

Returning to our running example we get two decreasing labelings for the F in Figure 3.3, which in turn imply the slide expansion

$$\mathfrak{P}_F = \mathfrak{F}(2^{[1]} 4^{[1]} 2^{[2]}) + \mathfrak{F}(4^{[1]} 2^{[1]} 2^{[2]}) = \mathfrak{F}_{12} + \mathfrak{F}_{0201}.$$

By appealing to a simple bijection between \mathbb{N} -vectors and indexed forests we can show that the set of forest polynomials is a basis for the polynomial ring $\mathbb{Z}[\mathbf{x}_+]$. Indeed, let $c(F)$ be the weak composition given by $c_i(F)$ being the number of nodes on the left branch leading to the leaf just left of the label i . It is easily seen that the correspondence $F \mapsto c(F)$ is a bijection between indexed forests and \mathbb{N} -vectors supported on \mathbb{Z}_+ . In fact $\mathbf{x}^{c(F)}$ is the revlex leading monomial in \mathfrak{P}_F , which in turn suffices to show the following:

Theorem 3.11. *The set of forest polynomials forms a basis for $\mathbb{Z}[\mathbf{x}_+]$.*

4. Back stable version

We shall see that the theory of (P, Φ) -partitions (and (P, ω, ρ) -partitions more generally) becomes nicer when we allow P -partitions to take nonpositive values. In the expansion (3.2) of $K_{(P, \Phi)}$, some terms on the right-hand side can be zero. In fact $K_{(P, \Phi)}$ itself may be zero, and this phenomenon leads to some technical investigation in [AB20]. Removing the lower bound for (P, Φ) -partitions ensures that no term will cancel, leading to a more pleasant theory.

The corresponding generating functions then belong to a certain class of series. We describe some structural theory of these series, borrowing notation and drawing motivation from [LLS21]. We also note that some of the series we consider arose naturally in previous work; see [NT21, Proposition 8.9] and [TWZ22, Section 3].

4.1. Back stable (P, Φ) -partitions

We now consider (P, Φ) -partitions f with the added liberty that f can now take values in \mathbb{Z} instead of \mathbb{Z}_+ .

Definition 4.1 (Back stable (P, Φ) -partitions). Given a poset P , let Φ be an injective function $\Phi : P \rightarrow \overline{\mathbb{Z}}$. A *back stable* (P, Φ) -partition is a function $f : P \rightarrow \mathbb{Z}$ such that for any $u, v \in P$:

- $f(u) \geq f(v)$ if $u \prec_P v$;
- $f(u) > f(v)$ if $u \prec_P v$ and $\Phi(u) > \Phi(v)$;
- $f(u) \leq \text{val}(\Phi(u))$.

Denote by $\overleftarrow{K}_{(P, \Phi)}$ the corresponding generating function. Note that this is now a homogeneous series of degree $\#P$ in $(x_i)_{i \in \mathbb{Z}}$ such that only a finite number of x_i for $i > 0$ occur. Then Stanley's fundamental theorem, in the form of Corollary 3.6, holds with no change. For L a chain, identified as a word $W \in \text{Inj}(\overline{\mathbb{Z}})$ as in Section 3.2, let $\overleftarrow{\mathfrak{F}}(W) = \overleftarrow{K}_{(L, \Phi)}$. Then we have

$$\overleftarrow{K}_{(P, \Phi)} = \sum_{L \in \text{Lin}(P)} \overleftarrow{\mathfrak{F}}(W). \quad (4.1)$$

The series $\overleftarrow{K}_{(P, \Phi)}$ are never zero. In particular, none of the series $\overleftarrow{\mathfrak{F}}(W)$ vanishes in (4.1), so the right-hand side is an actual summation over all linear extensions. Another advantage is

that $\overleftarrow{K}_{(P,\Phi)}$ lives inside the vector space of back stable quasisymmetric functions, introduced in the next subsection. This space comes equipped with several linear maps, which allow for recovering both the classical story of (P, ω) -partitions and quasisymmetric functions as well as the polynomial story involving (P, Φ) -partitions developed here.

4.2. Back stable quasisymmetric functions

We work with the set of variables $\mathbf{x} := \{x_i \mid i \in \mathbb{Z}\}$. Given $b \in \mathbb{Z}$, define $\mathbf{x}_{\leq b} := \{x_i \mid i \leq b\}$. In the case $b = 0$, we set $\mathbf{x}_- := \mathbf{x}_{\leq 0}$. Let $\mathbf{x}_+ := \{x_i \mid i \geq 1\}$. Let $\text{QSym}(\mathbf{x}_{\leq b})$ be the \mathbb{Q} -algebra of quasisymmetric functions in the ordered alphabet $\mathbf{x}_{\leq b}$. The algebra $\text{QSym}(\mathbf{x}_-)$ will be denoted by QSym .

Let $R \subseteq \mathbb{Q}[\mathbf{x}]$ denote the set of bounded degree formal power series f in \mathbf{x} with the property that there exists an $N \in \mathbb{Z}$ such that no x_i appears in f for $i > N$.

Definition 4.2. Let $f \in R$. We say that f is *back quasisymmetric* if there exists a $b \in \mathbb{Z}$ such that for any sequence of positive integers (a_1, \dots, a_k) and any monomial m in $\mathbf{x}_{>b}$, the coefficient of $x_{i_1}^{a_1} \cdots x_{i_k}^{a_k} m$ in f equals that of $x_{j_1}^{a_1} \cdots x_{j_k}^{a_k} m$ whenever $i_1 < \cdots < i_k \leq b$ and $j_1 < \cdots < j_k \leq b$.

Equivalently, f is back quasisymmetric if there exists a $b \in \mathbb{Z}$ such that $f \in \text{QSym}(\mathbf{x}_{\leq b}) \otimes \mathbb{Q}[x_{b+1}, x_{b+2}, \dots]$.

Remark 4.3. For the reader looking for something in the spirit of [LLS21], the following alternative definition should suffice. Call $f \in R$ back quasisymmetric if there exists a $b \in \mathbb{Z}$ such that $\sigma_i(f) = f$ for all $i \leq b$ where σ_i acts via Hivert’s *quasisymmetrizing action* [Hiv00, Section 3].

Let \overleftarrow{QR} denote the space of back quasisymmetric functions. Clearly, the space \overleftarrow{R} of back symmetric functions [LLS21] is a subset of \overleftarrow{QR} . We have the following analogue of [LLS21, Proposition 3.1].

Proposition 4.4. *We have that*

$$\overleftarrow{QR} = \text{QSym} \otimes \mathbb{Q}[\mathbf{x}].$$

Proof. Both QSym and $\mathbb{Q}[\mathbf{x}]$ are contained in \overleftarrow{QR} . Let us show that their abstract tensor product naturally embeds naturally in \overleftarrow{QR} . We need to prove that for any linear relation in \overleftarrow{QR}

$$\sum_{\alpha, \mathbf{c}} u_{\alpha, \mathbf{c}} F_{\alpha}(\mathbf{x}_-) \mathbf{x}^{\mathbf{c}} = 0, \tag{4.2}$$

the coefficients $u_{\alpha, \mathbf{c}} \in \mathbb{Q}$ are all zero. Here α and \mathbf{c} run over compositions and \mathbb{N} -vectors respectively. Now fix an \mathbb{N} -vector \mathbf{d} . After dividing the relation in (4.2) by $\mathbf{x}^{\mathbf{d}}$, we get

$$\sum_{\alpha} u_{\alpha, \mathbf{d}} F_{\alpha}(\mathbf{x}_-) = - \sum_{\alpha, \mathbf{c} \neq \mathbf{d}} u_{\alpha, \mathbf{c}} F_{\alpha}(\mathbf{x}_-) \mathbf{x}^{\mathbf{c}-\mathbf{d}}. \tag{4.3}$$

Now we apply the shift $x_i \mapsto x_{i+b}$ for b a nonnegative integer, and then set $x_i = 0$ for $i \leq 0$. The expression $F_{\alpha}(\mathbf{x}_-) \mathbf{x}^{\mathbf{c}-\mathbf{d}}$ becomes $F_{\alpha}(x_1, \dots, x_b) \prod_i x_{i+b}^{c_i-d_i}$. As b goes to infinity, this quantity

goes to zero since $c \neq d$. The limit here is the usual one for series, where for any fixed monomial the coefficients eventually stabilize.

The relation in (4.3) in the limit then gives

$$\sum_{\alpha} u_{\alpha, \mathbf{d}} F_{\alpha}(\mathbf{x}_+) = 0. \quad (4.4)$$

We can now conclude that $u_{\alpha, \mathbf{d}} = 0$ for all α by linear independence of fundamental quasisymmetric functions. Since \mathbf{d} was arbitrary, we get the desired result.

Having thus shown $\overleftarrow{QR} \supseteq \text{QSym} \otimes \mathbb{Q}[\mathbf{x}]$, we proceed to prove the reverse inclusion. Let $f \in \overleftarrow{QR}$, so that there exists $b \in \mathbb{Z}$ such that $f \in \text{QSym}(\mathbf{x}_{\leq b}) \otimes \mathbb{Q}[x_{b+1}, x_{b+2}, \dots]$. By linearity, it is enough to assume that $f = F_{\alpha}(\mathbf{x}_{\leq b})P$ with $P \in \mathbb{Q}[x_{b+1}, x_{b+2}, \dots]$. Say $b \geq 1$. Write the fundamental quasisymmetric function $F_{\alpha}(\mathbf{x}_{\leq b})$ as $F_{\alpha}(\mathbf{x}_- + x_1 + \dots + x_b)$. We know (cf. for instance [AFNT15, Section 2.3]) that this expands as

$$F_{\alpha}(\mathbf{x}_{\leq b}) = \sum_{\beta \odot \gamma = \alpha \text{ or } \beta \cdot \gamma = \alpha} F_{\beta}(\mathbf{x}_-) F_{\gamma}(x_1, \dots, x_b). \quad (4.5)$$

Now suppose $b \leq -1$. Then $F_{\alpha}(\mathbf{x}_{\leq b}) = F_{\alpha}(\mathbf{x}_- - x_0 - \dots - x_{b+1})$. This time we know that

$$F_{\alpha}(\mathbf{x}_{\leq b}) = \sum_{\beta \odot \gamma = \alpha \text{ or } \beta \cdot \gamma = \alpha} (-1)^{|\gamma|} F_{\beta}(\mathbf{x}_-) F_{\gamma}(x_{b+1}, \dots, x_0). \quad (4.6)$$

In both cases, this shows that $f = F_{\alpha}(\mathbf{x}_{\leq b})P$ is an element of $\text{QSym} \otimes \mathbb{Q}[\mathbf{x}]$ as wanted. \square

4.3. Some maps defined on back stable quasisymmetric functions

As in [LLS21, Section 3.4], we consider the evaluation map $\eta_0 : \mathbb{Q}[\mathbf{x}] \rightarrow \mathbb{Q}$ obtained by setting $x_i = 0$ for all $i \in \mathbb{Z}$. In other words, it picks the constant term in a polynomial. It induces the map $1 \otimes \eta_0$ on $\text{QSym} \otimes \mathbb{Q}[\mathbf{x}]$: it essentially picks out the term in QSym and forgets the polynomial part. Following Lam–Lee–Shimozono, we will abuse notation and refer to this induced map on \overleftarrow{QR} by η_0 as well.

Let $\gamma : \overleftarrow{QR} \rightarrow \overleftarrow{QR}$ be the map shifting variables $x_i \mapsto x_{i+1}$ for $i \in \mathbb{Z}$ [LLS21, Section 3.3]. Finally let $\pi_+ : \overleftarrow{QR} \rightarrow \mathbb{Q}[\mathbf{x}_+]$ be the *truncation map* obtained by setting $x_i = 0$ for $i \leq 0$; see proof of [LLS21, Proposition 3.18]. Note that all of these maps are algebra morphisms, and the previous two were already employed in the proof of Proposition 4.4.

Proposition 4.5. *For any $f \in \overleftarrow{QR}$,*

$$\eta_0(f)(\mathbf{x}_+) = \lim_{b \rightarrow \infty} \pi_+(\gamma^b(f)).$$

Note that $\eta_0(f)$ lives in $\text{QSym}(\mathbf{x}_-)$. The notation $\eta_0(f)(\mathbf{x}_+)$ means that we now write it in the variables \mathbf{x}_+ using the natural isomorphism between $\text{QSym}(\mathbf{x}_-)$ and $\text{QSym}(\mathbf{x}_+)$. The limit is the usual one for series, already used in the proof of Proposition 4.4: the coefficients of any fixed monomial eventually stabilize.

Proof. By linearity it is enough to prove it for $f = g\mathbf{x}^c$ with $g \in \mathbb{Q}\text{Sym}$, with $\mathbf{x}^c = x_{i_1}^{a_1} \cdots x_{i_k}^{a_k}$. Then $\eta_0(f) = g$ if $\mathbf{x}^c = 1$ and 0 otherwise. On the other hand, for b large enough $\pi^+(\gamma^b(f)) = g(x_1, \dots, x_b)x_{i_1+b}^{a_1} \cdots x_{i_k+b}^{a_k}$. This has limit 0 if any of the a_i is nonzero, and $g(\mathbf{x}_+)$ otherwise as wanted. \square

4.4. Back stable slides

We now discuss the *back stable slides* $\overleftarrow{\mathfrak{F}}_c := \overleftarrow{\mathfrak{F}}(W_c)$ beginning with establishing that they belong in \overleftarrow{QR} . In particular, this would show via (4.1) that $\overleftarrow{K}_{(P,\Phi)} \in \overleftarrow{QR}$ as well. To this end, we will need a result from [TWZ22] that we now recall for the reader's convenience.

Let \mathbf{c} be an \mathbb{N} -vector. We let $\text{flat}(\mathbf{c})$ denote the sequence formed by the positive entries in \mathbf{c} . Call a decomposition $\mathbf{c} = \mathbf{d} + \mathbf{e}$ where addition is component-wise *good* if either $\text{flat}(\mathbf{c}) = \text{flat}(\mathbf{d}) \cdot \text{flat}(\mathbf{e})$ or $\text{flat}(\mathbf{c}) = \text{flat}(\mathbf{d}) \odot \text{flat}(\mathbf{e})$ holds. Then [TWZ22, Lemma 3.5] in a special case states that:

Lemma 4.6. *Let \mathbf{c} be an \mathbb{N} -vector such that $\text{Supp}(\mathbf{c}) \subseteq \mathbb{Z}_+$. Then $\overleftarrow{\mathfrak{F}}_c$ has the following expansion in $\mathbb{Q}\text{Sym} \otimes \mathbb{Q}[\mathbf{x}_+]$:*

$$\overleftarrow{\mathfrak{F}}_c = \sum_{\text{good } \mathbf{c}=\mathbf{d}+\mathbf{e}} F_{\text{flat}(\mathbf{d})}(\mathbf{x}_-) \overleftarrow{\mathfrak{F}}_{\mathbf{e}}(\mathbf{x}_+).$$

Example 4.7. Let $\mathbf{c} = (0, 2, 0, 2, 0, 0, \dots)$. Then $\text{flat}(\mathbf{c}) = (2, 2)$ and one can easily check that we have the following five decompositions for $(2, 2)$: $\emptyset \cdot (2, 2)$, $(1) \odot (1, 2)$, $(2) \cdot (2)$, $(2, 1) \odot (1)$, $(2, 2) \cdot \emptyset$. These in turn translate to five good decompositions and we obtain

$$\overleftarrow{\mathfrak{F}}_{0202} = \overleftarrow{\mathfrak{F}}_{0202}(\mathbf{x}_+) + F_1(\mathbf{x}_-) \overleftarrow{\mathfrak{F}}_{0102}(\mathbf{x}_+) + F_2(\mathbf{x}_-) \overleftarrow{\mathfrak{F}}_{0002}(\mathbf{x}_+) + F_{21}(\mathbf{x}_-) \overleftarrow{\mathfrak{F}}_{0001}(\mathbf{x}_+) + F_{22}(\mathbf{x}_-).$$

Proposition 4.8. *For any \mathbb{N} -vector \mathbf{c} , we have $\overleftarrow{\mathfrak{F}}_c \in \overleftarrow{QR}$. As a consequence, $\overleftarrow{K}_{(P,\Phi)} \in \overleftarrow{QR}$ for any poset P with labeled flag Φ .*

Proof. It suffices to assume $\text{Supp}(\mathbf{c}) \subseteq \mathbb{Z}_+$. Indeed, abuse notation and define $\gamma(\mathbf{c})$ to be the \mathbb{N} -vector obtained by shifting \mathbf{c} once to the right. Then it is clear that for $i \in \mathbb{Z}$ that

$$\gamma^i \left(\overleftarrow{\mathfrak{F}}_c \right) = \overleftarrow{\mathfrak{F}}_{\gamma^i(\mathbf{c})}.$$

Now note that \overleftarrow{QR} is closed under shifting.

When $\text{Supp}(\mathbf{c}) \subseteq \mathbb{Z}_+$, then Lemma 4.6 immediately implies that $\overleftarrow{\mathfrak{F}}_c \in \overleftarrow{QR}$. As for showing $\overleftarrow{K}_{(P,\Phi)} \in \overleftarrow{QR}$, note that it is a sum of back stable slides as in (4.1). \square

The next result describes the distinguished role played by the back stable slides in \overleftarrow{QR} . It is the analogue of [LLS21, Theorem 3.5].

Theorem 4.9. *The back stable slides $\overleftarrow{\mathfrak{F}}_c$ for \mathbf{c} an \mathbb{N} -vector form a \mathbb{Q} -basis of \overleftarrow{QR} .*

Proof. The linear independence holds for the same reason as in [LLS21]. Indeed the revlex leading monomial in $\overleftarrow{\mathfrak{F}}_{\mathbf{c}}$ is $\mathbf{x}^{\mathbf{c}}$. We now show that the $\overleftarrow{\mathfrak{F}}_{\mathbf{c}}$ are spanning.

Pick $f \in \overleftarrow{QR}$. Without loss of generality assume that $f \in \mathbb{Q}\text{Sym} \otimes \mathbb{Q}[\mathbf{x}_+]$: indeed, like before, this follows from the fact that \overleftarrow{QR} is closed under shifting.

We know that slide polynomials form a basis for $\mathbb{Q}[\mathbf{x}_+]$, so we may assume that f has the form $f = F_{\alpha}(\mathbf{x}_-)\overleftarrow{\mathfrak{F}}_{(c_1, \dots, c_k)}$. If $c_i = 0$ for all $1 \leq i \leq k$, then we are done. Otherwise let $\mathbf{d} = (\dots, 0, \alpha|c_1, \dots, c_k, 0, \dots)$. By [TWZ22, Lemma 3.5] (essentially the statement in Lemma 4.6) the difference $f - \overleftarrow{\mathfrak{F}}_{\mathbf{d}}$ is a sum of $F_{\beta}(\mathbf{x}_-)\overleftarrow{\mathfrak{F}}_{(a_1, \dots, a_k)}$ where $a_1 + \dots + a_k < c_1 + \dots + c_k$. Induction implies the claim. \square

Remark 4.10. As the reader may expect at this stage, by allowing our indexed forests to be supported on \mathbb{Z} rather than \mathbb{Z}_+ , we may easily define *back stable forest polynomials*. The resulting family of polynomials $\overleftarrow{\mathfrak{P}}_F \in \overleftarrow{QR}$ then expands as a sum of back stable slide polynomials, one for each element in $\text{Dec}(F)$.

The lemma next saying that slides and fundamental quasisymmetric functions simultaneously inhabit $\overleftarrow{\mathfrak{F}}_{\mathbf{c}}$ is straightforward.

Lemma 4.11. *We have $\eta_0(\overleftarrow{\mathfrak{F}}_{\mathbf{c}}) = F_{\text{flat}(\mathbf{c})}(\mathbf{x}_-)$. Additionally*

$$\pi_+(\overleftarrow{\mathfrak{F}}_{\mathbf{c}}) = \begin{cases} \overleftarrow{\mathfrak{F}}_{\mathbf{c}} & \text{if } \text{Supp}(\mathbf{c}) \subseteq \mathbb{Z}_+; \\ 0 & \text{otherwise.} \end{cases}$$

Our next result follows from Stanley's theory again. It gives a *shuffle rule* for multiplying back stable slide polynomials. Let \mathbf{c} and \mathbf{d} be \mathbb{N} -vectors. We have $W_{\mathbf{c}}$ and $W_{\mathbf{d}}$ defined as usual. Define the set of shuffles $W_{\mathbf{c}} \sqcup W_{\mathbf{d}}$ where we replace every instance of $i^{[j]}$ in $W_{\mathbf{d}}$ by $i^{[j+c_i]}$. This shift ensures, amongst other things, that the set of shuffles comprises injective words.

Proposition 4.12. *Given \mathbb{N} -vectors \mathbf{c} and \mathbf{d} we have*

$$\overleftarrow{\mathfrak{F}}_{\mathbf{c}} \cdot \overleftarrow{\mathfrak{F}}_{\mathbf{d}} = \sum_{L \in W_{\mathbf{c}} \sqcup W_{\mathbf{d}}} \overleftarrow{\mathfrak{F}}(L).$$

Note that each summand on the right-hand side is a back stable slide polynomial. Hitting the expansion with η_0 gives us the usual shuffle product for $F_{\text{flat}(\mathbf{c})} \cdot F_{\text{flat}(\mathbf{d})}$, while applying π_+ recovers the rule for slides.

Example 4.13. Let $\mathbf{c} = (0, 1, 0, 2)$ and $\mathbf{d} = (0, 1)$. Then $W_{\mathbf{c}} = 4^{[1]}4^{[2]}2^{[1]}$ and $W_{\mathbf{d}} = 2^{[1]}$. We have that

$$W_{\mathbf{c}} \sqcup W_{\mathbf{d}} = \{4^{[1]}4^{[2]}2^{[1]}2^{[2]}, 4^{[1]}4^{[2]}2^{[2]}2^{[1]}, 4^{[1]}2^{[2]}4^{[2]}2^{[1]}, 2^{[2]}4^{[1]}4^{[2]}2^{[1]}\}$$

Consider the $L \in W_{\mathbf{c}} \sqcup W_{\mathbf{d}}$ from left to right. The corresponding $\overleftarrow{\mathfrak{F}}(L)$ are $\overleftarrow{\mathfrak{F}}_{(0,2,0,2)}$, $\overleftarrow{\mathfrak{F}}_{(1,1,0,2)}$, $\overleftarrow{\mathfrak{F}}_{(1,2,0,1)}$, and $\overleftarrow{\mathfrak{F}}_{(1,3)}$ respectively. So we infer that

$$\overleftarrow{\mathfrak{F}}_{0102} \cdot \overleftarrow{\mathfrak{F}}_{01} = \overleftarrow{\mathfrak{F}}_{0202} + \overleftarrow{\mathfrak{F}}_{1102} + \overleftarrow{\mathfrak{F}}_{1201} + \overleftarrow{\mathfrak{F}}_{13}.$$

5. The inverse slide Kostka matrix

We showed in Theorem 4.9 that back stable slide polynomials form a basis of \overleftarrow{QR} . In view of Proposition 4.4, another basis is given by the $F_\alpha(\mathbf{x}_-)\mathbf{x}^c$ for any composition α and \mathbb{N} -vector c . In this section we give an explicit change of basis from the first basis to the second. Since we know how to multiply back stable slide polynomials by Proposition 4.12, it is enough to treat separately the case of $F_\alpha(\mathbf{x}_-)$ and \mathbf{x}^c . Now $F_\alpha(\mathbf{x}_-)$ is already equal to the back stable slide $\overleftarrow{\mathfrak{F}}_c$ with $c = (\dots, 0, \alpha|0, 0, \dots)$, so it remains to express a monomial \mathbf{x}^c in terms of back stable slide polynomials.

Recall that we define $\overleftarrow{\mathfrak{F}}_c$ as $\overleftarrow{\mathfrak{F}}(W_c)$, where W_c is a special word attached to c . These words are precisely the standardizations of *nonincreasing* words in \mathbb{Z}^* , that is words where letters decrease weakly from left to right. We will use letters C, D, E for nonincreasing words, to recall that they correspond bijectively to \mathbb{N} -vectors c, d, e , but we will not use the latter. We will also write simply $\mathfrak{F}(C)$ and $\overleftarrow{\mathfrak{F}}(C)$ as no confusion is caused.

Let \mathcal{P} denote the set of all nonincreasing words. Fix $C = C_1C_2 \cdots C_m \in \mathcal{P}$ with $C_i \in \mathbb{Z}$. By grouping equal values one may write $C = M_1^{m_1}M_2^{m_2} \cdots M_t^{m_t}$ where $m_i > 0$ and $M_i > M_{i+1}$ for $1 \leq i < t$. Fix $i \in \{1, \dots, t\}$, and let $x_0 = M_i$ and $M_{t+1} = -\infty$ by convention. Now let

$$B_i = \{x_1 \cdots x_{m_i} \in \mathcal{P} \mid x_{j+1} \in \{x_j, x_j - 1\} \text{ for } 0 \leq j \leq m_i - 1, \text{ and } x_{m_i} > M_{i+1}\}. \tag{5.1}$$

This given, define the following set that is crucial for us:

$$\mathbb{B}_C = \{X^1 \cdots X^t \mid X^i \in B_i \text{ for } 1 \leq i \leq t\}.$$

Note that $\mathbb{B}_C \subseteq \mathcal{P}$ by construction. Elements $D \in \mathbb{B}_C$ are completely characterized by the set $S_C(D)$ of indices j such that $x_{j+1} = x_j - 1$ in (5.1). In particular, this allows us to identify \mathbb{B}_C with a distinguished subset of the set of sequences (S_1, \dots, S_t) where each $S_i \subseteq \{0, \dots, m_i - 1\}$. Such sequences in turn may naturally be identified as subsets of $\{0, 1, \dots, m_1 + \cdots + m_t - 1\}$. In fact, as we shall soon see, this aforementioned association has even further structure; we have an isomorphism between appropriate posets.

Example 5.1. Let $C = 442 = 4^2 2^1$. Then B_1 and B_2 equal $\{44, 43, 33\}$ and $\{2, 1\}$ respectively. We thus have

$$\mathbb{B}_C = \{442, 332, 432, 441, 331, 431\}.$$

Recording the $S_C(D)$ as D varies over \mathbb{B}_C gives us the following subsets of $\{0, 1, 2\}$:

$$\{\emptyset, \{0\}, \{1\}, \{2\}, \{0, 2\}, \{1, 2\}\}.$$

Note that these subsets give a lower order ideal in the Boolean lattice on subsets of $\{0, 1, 2\}$.

We are now ready to state the main result in this subsection. It expresses the monomial $\mathbf{x}(C) := x_{C_1} \cdots x_{C_m}$ in terms of back stable slides. In particular, the expansion is *signed multiplicity free*.

Theorem 5.2. *The back stable slide expansion of $\mathbf{x}(C)$ is given by*

$$\mathbf{x}(C) = \sum_{D \in \mathbb{B}_C} (-1)^{|S_C(D)|} \overleftarrow{\mathfrak{F}}(D). \quad (5.2)$$

Recall that all words are nonincreasing in this section, so that all $\overleftarrow{\mathfrak{F}}(D)$ can be written in the form $\overleftarrow{\mathfrak{F}}_d$ immediately. We illustrate the theorem with an example.

Example 5.3. Let $C = 442$, so that $C = W_c$ with $c = (0, 1, 0, 2, 0, \dots)$. We already computed \mathbb{B}_C in Example 5.1. Theorem 5.2 then states:

$$\begin{aligned} x_2 x_4^2 &= \overleftarrow{\mathfrak{F}}(442) - \overleftarrow{\mathfrak{F}}(332) - \overleftarrow{\mathfrak{F}}(432) - \overleftarrow{\mathfrak{F}}(441) + \overleftarrow{\mathfrak{F}}(331) + \overleftarrow{\mathfrak{F}}(431) \\ &= \overleftarrow{\mathfrak{F}}_{0102} - \overleftarrow{\mathfrak{F}}_{012} - \overleftarrow{\mathfrak{F}}_{0111} - \overleftarrow{\mathfrak{F}}_{1002} + \overleftarrow{\mathfrak{F}}_{102} + \overleftarrow{\mathfrak{F}}_{1011}. \end{aligned}$$

We need some preparation before presenting the proof of Theorem 5.2. We will appeal to poset-theoretic terminology freely; the reader is referred to [Sta97, Chapter 3] for any undefined jargon.

Definition 5.4. Let $\mathcal{P}_m \subseteq \mathcal{P}$ be the set of all nonincreasing words of length m . For $C, D \in \mathcal{P}$, define $D \leq_m C$ if and only if $D_i \leq C_i$ for all i and $D_i > D_{i+1}$ whenever $C_i > C_{i+1}$.

This makes it clear that (\mathcal{P}_m, \leq_m) is a poset. In fact it is *locally finite*, thus we have the existence of a Möbius function μ . We recall that it is defined on all (D, C) with $D \leq_m C$ by $\mu(C, C) = 1$ for all C , and whenever $D <_m C$,

$$\sum_{D \leq_m E \leq_m C} \mu(E, C) = 0. \quad (5.3)$$

We illustrate a convex subset of \mathcal{P}_3 in Figure 5.1.

Toward describing the Möbius function of \mathcal{P}_m , we will first show that it is a lattice, that is, any two elements have a *join* (least upper bound) and a *meet* (greatest lower bound). It is helpful to consider an example of a join of two elements, as that will guide the construction that follows.

Consider $C = 555322$ and $D = 664421$. Any common upper bound must be component-wise greater than both C and D . This leads one to propose $E' = 665422$. Now observe that $E'_2 > E'_3$ yet $C_2 = C_3$, and $E'_3 > E'_4$ but $D_3 = D_4$. Therefore, whilst E' is component-wise greater than both C and D , we do not have $C \leq_m E'$ and $D \leq_m E'$. There is an easy fix: increment E'_3 and E'_4 appropriately so that the resulting word does not have strict descents in positions 2 and 3. Indeed, check that $E = 666622$ meets these criteria and therefore $E = C \vee D$.

Proposition 5.5. *For any $m \geq 1$, \mathcal{P}_m is a lattice.*

Proof. Let C, D be two elements of \mathcal{P}_m . We will first prove the existence of their join by generalizing the construction above. Define $E' \in \mathcal{P}_m$ by $E'_i = \max(C_i, D_i)$. Let J be the set consisting of 1 and all $i \in \{2, \dots, m\}$ such that $C_{i-1} > C_i$ and $D_{i-1} > D_i$. So, for instance, in the example leading to this proposition, we would have $J = \{1, 5\}$. Now define $E \in \mathcal{P}_m$ by setting $E_i = E'_{j_k}$ where j_k the maximal element of J such that $j_k \leq i$.




Figure 5.1: Hasse diagram of the convex subset of \mathcal{P}_3 formed by all words below 444 and with positive letters.

We claim that E is the join of C and D . Let us first check that E is an upper bound for C and D . Indeed we have $E_i \geq E'_i$ and $E'_i \geq C_i, D_i$, so E is component-wise greater than C and D . Second, $E_{i-1} > E_i$ implies that $i \in J$ by construction, which by definition entails $C_{i-1} > C_i$ and $D_{i-1} > D_i$. Thus we infer that $E \geq_m C, D$.

Now let G satisfy $G \geq_m C, D$. One thus has clearly that $G_i \geq E'_i$ for all i , and since $i \notin J$ implies that $G_{i-1} = G_i$, it follows that in fact $G_i \geq E_i$ for all i . Now if $G_{i-1} > G_i$, one necessarily has $i \in J$. We have thus $E_i = E'_i = \max(C_i, D_i)$ and $E_{i-1} \geq E'_{i-1} = \max(C_{i-1}, D_{i-1}) > \max(C_i, D_i) = E_i$ where the last inequality follows from $C_{i-1} > C_i$ and $D_{i-1} > D_i$. This shows that $G \geq_m E$, which completes the proof that any two elements have a join.

We now need to show that any two elements have a meet. This could be done explicitly as above. We will rather adapt the abstract argument of [Sta97, Proposition 3.3.1] used there in the case of a finite bounded poset.

Let $C, D \in \mathcal{P}_m$. Note first that C, D always have a common lower bound L : for instance, let k be the minimal value of all letters occurring in C and D , and pick $L = k(k-1) \cdots (k-m+1)$. Consider the set \mathcal{X} of all elements of \mathcal{P}_m that lie below C and D and above L . Now \mathcal{X} is finite and contains L , so by the first half of the proof we can define the join $M := \bigvee \mathcal{X}$, i.e. the join of all elements in \mathcal{X} .

By construction $M \leq_m C, D$. We claim that it is in fact the meet of C and D . Indeed, let M' be any element below C and D . Then $M' \leq_m M' \vee L$, and this join is in \mathcal{X} . It follows that $M' \leq_m M$ as wanted. We have thus shown that any two elements admit a join and a meet, and so \mathcal{P}_m is a lattice. \square

We now compute the Möbius function explicitly:

Proposition 5.6. *Let $C, D \in \mathcal{P}_m$ with $D \leq_m C$. We have*

$$\mu(D, C) = \begin{cases} (-1)^{|S_C(D)|} & \text{if } D \in \mathbb{B}_C \\ 0 & \text{if } D \notin \mathbb{B}_C. \end{cases}$$

Proof. We will apply the crosscut theorem [Sta97, Corollary 3.9.4], which we can since \mathcal{P}_m is a lattice by Proposition 5.5. We shall need an explicit description for the set \mathcal{G}_C of coatoms of C in \mathcal{P}_m . We will use $C = 555322 \in \mathcal{P}_6$ as a running example.

Write $C = M_1^{m_1} M_2^{m_2} \cdots M_t^{m_t}$ where $m_i > 0$ and $M_i > M_{i+1}$ for $1 \leq i < t$, as when we defined \mathbb{B}_C earlier. Let i such that $M_i > M_{i+1} + 1$, or $i = t$; we will call such an index *good*. For $j \in \{0, \dots, m_i - 1\}$, define the word C_i^j as the word C where $M_i^{m_i}$ is replaced with $M_i^j (M_i - 1)^{m_i - j}$. Note that, in comparison to C , the word C_i^j has an additional strict inequality if $j > 0$. For a good index i , we let $\mathcal{G}_{C,i} := \{C_i^j \mid 0 \leq j \leq m_i - 1\}$. For $C = 555322 = 5^3 3^1 2^2$, the good indices are 1 and 3, and we have $\mathcal{G}_{C,1} = \{444322, 544322, 554322\}$, $\mathcal{G}_{C,3} = \{555311, 555321\}$. Then as an immediate consequence of the definition of \leq_m we have

$$\mathcal{G}_C = \bigcup_{i \text{ good}} \mathcal{G}_{C,i}.$$

Now we let

$$\mathbb{G}(D, C) := \{J \subseteq \mathcal{G}_C \mid \bigwedge J = D\}.$$

The crosscut theorem then says

$$\mu(D, C) = \sum_{J \in \mathbb{G}(D, C)} (-1)^{|J|}. \quad (5.4)$$

If $\mathbb{G}(D, C)$ is empty, then we have $\mu(D, C) = 0$.

Assume now that $\mathbb{G}(D, C)$ is nonempty, and write $D = \bigwedge J$ for some $J \subseteq \mathcal{G}_C$. Observe the crucial fact that

$$(*) \quad D \in \mathbb{B}_C \iff J \text{ satisfies } |J \cap \mathcal{G}_{C,i}| < M_i - M_{i+1} \text{ for all good } i < t.$$

In our running example, this says $|J \cap \{554322, 544322, 444322\}| < 5 - 3 = 2$. The observation (*) hinges on the fact that the cardinality of the intersection $|J \cap \mathcal{G}_{C,i}|$ forces strict descents in the corresponding positions in the appropriate substring in $\bigwedge J = D$, and that the entries in this substring must be strictly greater than M_{i+1} for the resulting word to belong to \mathbb{B}_C .

If J satisfies (*), then it is unique and consists precisely of the elements C_i^j that have a strict descent in $S_C(D)$. Therefore $|J| = |S_C(D)|$, and it follows from (5.4) that $\mu(D, C)$ is equal to $(-1)^{|S_C(D)|}$ as desired. In fact, \mathbb{B}_C is naturally identified via $D \mapsto S_C(D)$ to a lower ideal of a boolean lattice –as hinted in Example 5.1– and we recover indeed the classical Möbius function of the latter.

Finally we consider the case where J is such that $D = \bigwedge J \notin \mathbb{B}_C$. By (*), there exists a good index $i < t$ such that $|J \cap \mathcal{G}_{C,i}| \geq M_i - M_{i+1}$. Pick the minimal such i , and consider $i_0 > i$ the next good index. Note that i_0 must exist as t is always a good index by definition. Then define $J' = J \Delta \{C_{i_0}^0\}$ where Δ is the symmetric difference. In our running example, an element not in \mathbb{B}_C is given by $D = 433210 = \bigwedge J$ where $J = \{544322, 444322, 555321\}$. We have $i_0 = 3$ and thus $J' = J \cup \{C_{i_0}^0 = 555311\}$. The map χ_C defined by $J \mapsto J'$ is clearly an involution on \mathcal{G}_C . Moreover, one checks that $\bigwedge J' = \bigwedge J$, so χ_C restricts to an involution on $\mathbb{G}(D, C)$. It follows that the sum in (5.4) is zero, which completes the proof. \square

We conclude by establishing the claimed monomial-to-slide expansion.

Proof of Theorem 5.2. Let $C \in \mathcal{P}_m$. By definition we have

$$\overleftarrow{\mathfrak{F}}(C) = \sum_{D \leq_m C} \mathbf{x}(D). \tag{5.5}$$

We want to apply Möbius inversion to (5.5), but the sum on the right is infinite. We thus restrict to the polynomial case temporarily:

$$\mathfrak{F}(C) = \sum_{D \leq_m C, D > 0} \mathbf{x}(D). \tag{5.6}$$

Here we take $D > 0$ to mean that all components of D are strictly positive. By Möbius inversion we get:

$$\mathbf{x}(C) = \sum_{D \leq_m C, D > 0} \mu(D, C) \mathfrak{F}(D). \tag{5.7}$$

Note that this is indeed the same Möbius function of the full poset \mathcal{P}_m : imposing $D > 0$ gives us an upper ideal, which is a convex subset, thereby preserving the Möbius values by (5.3). By Proposition 5.6, we obtain:

$$\mathbf{x}(C) = \sum_{D \in \mathbb{B}_C, D > 0} (-1)^{|S_C(D)|} \mathfrak{F}(D). \tag{5.8}$$

This gives us the polynomial case.

Now for $i > 0$, apply (5.8) to $\gamma^i(C) = (C_1 + i) \cdots (C_m + i)$, and then shift the resulting expansion by $\gamma^{-i} : x_k \mapsto x_{k-i}$. By passing to the limit in the result, we get (5.2) (for $C > 0$) since $\overleftarrow{\mathfrak{F}}(C)$ is the limit of $\gamma^{-i} \mathfrak{F}(\gamma^i(C))$ when i tends to infinity. For general C , one has $\gamma^i(C) > 0$ for i large enough, and then one concludes using $\overleftarrow{\mathfrak{F}}(C) = \gamma^{-i} \overleftarrow{\mathfrak{F}}(\gamma^i(C))$. \square

Acknowledgements

We are grateful to Darij Grinberg for helpful comments on both content and exposition. We are also grateful to the referees for their careful reading and helpful suggestions.

References

[AB20] S. Assaf and N. Bergeron. Flagged (\mathcal{P}, ρ) -partitions. *European J. Combin.*, 86:103085, 17, 2020. doi:10.1016/j.ejc.2020.103085.

[AFNT15] J.-C. Aval, V. Féray, J.-C. Novelli, and J.-Y. Thibon. Quasi-symmetric functions as polynomial functions on Young diagrams. *J. Algebraic Combin.*, 41(3):669–706, 2015. doi:10.1007/s10801-014-0549-y.

- [AS17] S. Assaf and D. Searles. Schubert polynomials, slide polynomials, Stanley symmetric functions and quasi-Yamanouchi pipe dreams. *Adv. Math.*, 306:89–122, 2017. doi:10.1016/j.aim.2016.10.015.
- [Ges84] I. M. Gessel. Multipartite P -partitions and inner products of skew Schur functions. In *Combinatorics and algebra (Boulder, Colo., 1983)*, volume 34 of *Contemp. Math.*, pages 289–317. Amer. Math. Soc., Providence, RI, 1984.
- [Ges16] I. M. Gessel. A historical survey of P -partitions. In *The mathematical legacy of Richard P. Stanley*, pages 169–188. Amer. Math. Soc., Providence, RI, 2016. doi:10.1090/mbk/100/10.
- [GR14] D. Grinberg and V. Reiner. Hopf Algebras in Combinatorics. 2014. arXiv:1409.8356.
- [Hiv00] F. Hivert. Hecke algebras, difference operators, and quasi-symmetric functions. *Adv. Math.*, 155(2):181–238, 2000. doi:10.1006/aima.1999.1901.
- [LLS21] T. Lam, S. J. Lee, and M. Shimozono. Back stable Schubert calculus. *Compos. Math.*, 157(5):883–962, 2021. doi:10.1112/S0010437X21007028.
- [NT21] P. Nadeau and V. Tewari. A q -analogue of an algebra of Klyachko and Macdonald’s reduced word identity, 2021. arXiv:2106.03828.
- [NT24] P. Nadeau and V. Tewari. Forest polynomials and the class of the permutahedral variety. *Adv. Math.*, 453:Paper No. 109834, 33, 2024. URL: <https://doi-org.myaccess.library.utoronto.ca/10.1016/j.aim.2024.109834>, doi:10.1016/j.aim.2024.109834.
- [RS95] V. Reiner and M. Shimozono. Key polynomials and a flagged Littlewood-Richardson rule. *J. Combin. Theory Ser. A*, 70(1):107–143, 1995. doi:10.1016/0097-3165(95)90083-7.
- [Sta72] R. P. Stanley. *Ordered structures and partitions*. Memoirs of the American Mathematical Society, No. 119. American Mathematical Society, Providence, R.I., 1972.
- [Sta97] R. P. Stanley. *Enumerative combinatorics. Vol. 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original.
- [TWZ22] V. Tewari, A. T. Wilson, and P. B. Zhang. Chromatic nonsymmetric polynomials of Dyck graphs are slide-positive. *Proc. Amer. Math. Soc.*, 150(5):1873–1888, 2022. doi:10.1090/proc/15664.