

Lawrence Berkeley National Laboratory

Recent Work

Title

SOME FIRST RESULTS FROM THE MARK II AT SPEAR

Permalink

<https://escholarship.org/uc/item/0g0052xd>

Author

Abrams, G.S.

Publication Date

1979-06-01

UC 34 C+H

LBL-9320
SLAC-PUB-2350



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA, BERKELEY, CA

Physics, Computer Science & Mathematics Division

Presented at the International Conference on High Energy Physics, Geneva, Switzerland, June 27 - July 4, 1979

SOME FIRST RESULTS FROM THE MARK II AT SPEAR

Stanford Linear Accelerator Center, Lawrence Berkeley Laboratory and Department of Physics, University of California, Berkeley


June 1979

RECEIVED
LAWRENCE
BERKELEY LABORATORY
Sep 17
AUG 28 1979

LIBRARY AND DOCUMENTS SECTION

For Reference

Not to be taken from this room



Prepared for the U. S. Department of Energy under Contract W-7405-ENG-48

34d

1/19/81 12:30, 1

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

SOME FIRST RESULTS FROM THE MARK II AT SPEAR

G.S. Abrams, M.S. Alam, C.A. Blocker, A.M. Boyarski, M. Breidenbach, C.H. Broll, D.L. Burke, W.C. Carithers, W. Chinowsky, M.W. Coles, S. Cooper, B. Couchman, W.E. Dieterle, J.B. Dillon, J. Dorenbosch, J.M. Dorfan, M.W. Eaton, G.J. Feldman, H.G. Fischer, M.E.B. Franklin, G. Gidal, G. Goldhaber, G. Hanson, K.G. Hayes, T. Himel, D.G. Hitlin, R.J. Hollebeek, W.R. Innes, J.A. Jaros, P. Jenni, A.D. Johnson, J.A. Kadyk, A.J. Lankford, R.R. Larsen, M.J. Longo, D. Lüke, V. Lüth, J.F. Martin, R.E. Millikan, M.E. Nelson, C.Y. Pang, J.F. Patrick, M.L. Perl, B. Richter, J.J. Russell, D.L. Scharre, R.H. Schindler, R.F. Schwitters, S.R. Shannon, J.L. Siegrist, J. Strait, H. Taureg, V.I. Telnov, M. Tonutti, G.H. Trilling, E.N. Vella, R.A. Vidal, I. Videau, J.M. Weiss, H. Zacccone.

Presented by J.M. Weiss[†]

[†]Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

and

Lawrence Berkeley Laboratory and Department of Physics
University of California, Berkeley, California 94720

ABSTRACT

Preliminary results are given from the Mark II experiment at SPEAR on radiative decays of the $\psi'(3684)$ and on inclusive baryon production from 3.67 to 7.4 GeV center-of-mass energy. A 90% confidence level upper limit of 0.12% is given for $BR[\psi' \rightarrow \gamma\eta_c'(3455)] \times BR[\eta_c'(3455 \rightarrow \gamma\psi)]$.

1. INTRODUCTION

This paper presents some first results from the Mark II experiment at the Stanford Linear Accelerator Center e^+e^- storage ring facility SPEAR. These include results on:

- i) radiative decays of the $\psi'(3684)$;
- ii) inclusive baryon production in e^+e^- annihilation.

A number of other topics are discussed by G. Gidal elsewhere in these proceedings¹⁾.

2. THE MARK II DETECTOR

A schematic diagram of the Mark II detector is shown in Fig. 1. Particles originating in the intersection region pass through a thin stainless steel vacuum pipe, a cylindrical scintillation pipe counter, 16 layers of cylindrical drift chambers²⁾, and a layer of 48 time-of-flight (TOF) scintillation counters.

They then may penetrate the 1.3 radiation length solenoidal coil and enter one of 8 lead-liquid argon (LA) shower counters³⁾ (14 r.l.) which surround the inner detector. In addition, a muon detection system uses proportional tubes interspersed among the split magnet flux returns shown in Fig. 1 and additional iron absorbers on the top and sides which are not shown in the figure. Finally, both endcap regions of the detector are covered by additional shower detectors.

Performance features may be summarized here by:

- i) 200 μ average spatial resolution for the drift chambers which provide highly efficient tracking for $p > 100$ MeV/c over 75% of 4π ster. and some coverage out to 85% of 4π ster.

* Work supported by the Department of Energy under contract number DE-AC03-76SF00515.




Fig. 1. Schematic view of the Mark II detector.

- ii) 300 psec. time-of-flight resolution for hadrons (270 psec. for Bhabha events) over 75% of 4π ster. giving 1σ π/K separation at 1.3 GeV/c and 1σ K/p separation at 2.0 GeV/c.
- iii) $\delta E/E = .12/\sqrt{E}$ (E in GeV) energy resolution for photons and electrons from the liquid argon shower counters which cover 73% of 4π ster. The efficiency of these devices has been measured using the photons in fully-constrained $\psi \rightarrow \pi^+ \pi^- \pi^0$ and $\pi^+ \pi^- \pi^+ \pi^- \pi^0$ events and agrees well with detailed Monte Carlo calculations. This efficiency rises from about 30% at $E_\gamma = .15$ GeV to approximately 85% at $E_\gamma = .30$ GeV and 95% above .55 GeV.

3. RADIATIVE DECAYS OF THE $\psi'(3684)$

Events of the form



are selected by the following criteria:

- i) exactly 2 tracks of opposite charge from the primary vertex with an invariant mass between 2.8 and 3.4 GeV ($p_e < 1.75$ GeV/c).
- ii) At least 2 photons found in the liquid argon, each separated from the nearest charged track by at least 0.2 m. ($E_\gamma > .1$ GeV). More than 2 photons are allowed due to the possibility of spurious "photons" generated from random preamplifier noise.

These events are then fit to the hypothesis (1) using the program SQUAW with 5 constraints. Figure 2(a) shows the $\gamma\gamma$ mass spectrum for events satisfying (1) with a $\chi^2 < 15$. The events in the prominent η peak at .548 GeV coming from $\psi' \rightarrow \eta\psi$ are then eliminated by requiring $m_{\gamma\gamma} < .530$ GeV, leaving events which are candidates for the desired cascade decay



The high mass $\gamma\psi$ combinations are plotted in Fig. 2(b) and show two clear peaks corresponding to the $\chi(3505)$ and $\chi(3550)$. In addition, there is a smooth background consistent in both size and shape with that expected from $\psi' \rightarrow \pi^0\pi^0\psi$. While there is no clear peak corresponding to the $\chi(3410)$, the events near that mass are consistent with the small branching ratio previously observed for that state^{4,5,6,7}). On the other hand, there is no evidence in these data for a peak at 3455 MeV which has been previously reported⁴) as $\chi(3455)$ and which has been suggested as a candidate for the pseudoscalar η'_c . Table 1 presents preliminary branching ratios obtained using our knowledge of the detector acceptance.




Fig. 2(a). $\gamma\gamma$ mass spectrum for events satisfying $\psi' \rightarrow \gamma\gamma\psi$

2(b). High mass $\gamma\psi$ combinations.

Table 1

Summary of Branching Ratios $\psi' \rightarrow \gamma\psi$ *

	Mark II %	Mark I ⁴⁾ %	DESY-Heidelberg ⁷⁾ %
η	$3.8 \pm .5$	$4.3 \pm .8$	$3.6 \pm .5$
$\chi(3550)$	$1.2 \pm .3$	$1.0 \pm .6$	$1.0 \pm .2$
$\chi(3505)$	$2.5 \pm .3$	$2.4 \pm .8$	$2.5 \pm .4$
$\chi(3455)$	$<.12^{**}$	$.8 \pm .4$	$<.25$
$\chi(3410)$	$.08 \pm .08$	$.2 \pm .2$	$.14 \pm .09$

* $BR[\psi' \rightarrow \gamma\chi] \times BR[\chi \rightarrow \gamma\psi]$

** 90% confidence level

4. INCLUSIVE BARYON PRODUCTION IN e^+e^- ANNIHILATIONS

The inclusive production of \bar{p} , Λ and $\bar{\Lambda}$ has been studied with data concentrated in the center-of-mass energy range 4.5-6.0 GeV covering the region of a previously reported rise in $R(p+\bar{p})$ ⁸⁾. Multi-hadron events with at least 3 detected tracks are selected. The time-of-flight system is used to identify p and \bar{p} to 2.0 GeV/c using a weight method which is helpful above 1.3 GeV/c. However, only the \bar{p} results are used here due to beam-gas backgrounds. The Λ and $\bar{\Lambda}$ are observed by their $p\pi^-$ and $\bar{p}\pi^+$ decay modes with an rms mass resolution of about 4 MeV.

Preliminary results, corrected for acceptance, are presented in Fig. 3 as a ratio of the inclusive cross-section to the μ -pair production cross-section. For the proton case, $R(\bar{p}+p) = 2\sigma_{\bar{p}}/\sigma_{\mu\mu}$. All errors shown are statistical only and do not include an estimated systematic uncertainty of $\pm 30\%$, believed largely energy independent⁹⁾.

The measurements are consistent with previous experiments^{8,10)}. They show in detail, however, that the rise in the inclusive baryon production in e^+e^- annihilation is smooth and occurs principally between 4.6 and 5.2 GeV center-of-mass energy.

This work is supported by the U.S. Department of Energy under contract number DE-AC03-76SF00515.




Fig. 3(a). Inclusive \bar{p} production vs. c.m. energy.
3(b). Inclusive Λ and $\bar{\Lambda}$ production vs. c.m. energy.

* * *

REFERENCES

- 1) G. Gidal, Proc. of Internat. Conf. on High Energy Physics, Geneva, Switzerland, 27 June - 4 July, 1979.
- 2) W. Davies-White et al., SLAC Preprint SLAC-PUB-2181 (1978), to be published in Nucl. Instrum. Methods.
- 3) G.S. Abrams et al., IEEE Trans. on Nucl. Sci. NS-25, 309 (1978).
- 4) J.S. Whitaker et al., Phys. Rev. Lett. 37, 1596 (1976).
- 5) W. Braunschweig et al., Phys. Lett. 57B, 407 (1975).
- 6) V. Blobel, Proc. of the XIIth Rencontre de Moriond, Flaine, France, March 1977.
- 7) W. Bartel et al., DESY Report DESY 78/49 (1978).
- 8) M. Piccolo et al., Phys. Rev. Lett. 39, 1503 (1977).
- 9) This uncertainty is dominated by model dependence of the acceptance calculation which is still at a preliminary stage.
- 10) R. Brandelik et al., Nucl. Phys. B148, 189 (1979).

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720