

# UC San Diego

## UC San Diego Previously Published Works

### Title

Diversity and distribution of the bmp gene cluster and its Polybrominated products in the genus Pseudoalteromonas

### Permalink

<https://escholarship.org/uc/item/0jr9934m>

### Journal

Environmental Microbiology, 21(5)

### ISSN

1462-2912

### Authors

Busch, Julia  
Agarwal, Vinayak  
Schorn, Michelle  
[et al.](#)

### Publication Date

2019-05-01

### DOI

10.1111/1462-2920.14532

Peer reviewed

# Diversity and distribution of the *bmp* gene cluster and its Polybrominated products in the genus *Pseudoalteromonas*

Julia Busch,<sup>1,2</sup> Vinayak Agarwal,<sup>1,2†</sup> Michelle Schorn,<sup>1,2</sup>  
Henrique Machado,<sup>1,3</sup> Bradley S. Moore,<sup>1,2,4</sup>  
Greg W. Rouse,<sup>5</sup> Lone Gram<sup>3</sup> and Paul R. Jensen<sup>1,2\*</sup>

<sup>1</sup>Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA.

<sup>2</sup>Scripps Center for Oceans and Human Health, Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA.

<sup>3</sup>Department of Biotechnology and Biomedicine, Technical University of Denmark, Søtofts Plads bldg. 221, DK-2800, Kgs. Lyngby, Denmark.

<sup>4</sup>Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California San Diego, La Jolla, CA, USA.

<sup>5</sup>Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA.

## Summary

The production of pentabromopseudilin and related brominated compounds by *Pseudoalteromonas* spp. has recently been linked to the *bmp* biosynthetic gene cluster. This study explored the distribution and evolutionary history of this gene cluster in the genus *Pseudoalteromonas*. A phylogeny of the genus revealed numerous clades that do not contain type strains, suggesting considerable species level diversity has yet to be described. Comparative genomics revealed four distinct versions of the gene cluster distributed among 19 of the 101 *Pseudoalteromonas* genomes examined. These were largely localized to the least inclusive clades containing the *Pseudoalteromonas luteoviolacea* and *Pseudoalteromonas phenolica* type strains and show clear evidence of gene and gene cluster loss in certain lineages. *Bmp* gene phylogeny is largely congruent with the *Pseudoalteromonas* species phylogeny, suggesting

vertical inheritance within the genus. However, the gene cluster is found in three different genomic environments suggesting either chromosomal rearrangement or multiple acquisition events. *Bmp* conservation within certain lineages suggests the encoded products are highly relevant to the ecology of these bacteria.

## Introduction

The structure of the bromine rich antibiotic pentabromopseudilin was the first marine microbial natural product described (Lovell, 1966). It was isolated from an obligate marine bacterium identified at the time as *Pseudomonas* sp. (Burkholder *et al.*, 1966) but was likely a member of the genus now described as *Pseudoalteromonas*. Subsequently, pentabromopseudilin has been reported from multiple *Pseudoalteromonas* spp. (Isnansetyo and Kamei, 2003; Fehér *et al.*, 2010; Vynne *et al.*, 2011; Vynne *et al.*, 2012; Whalen *et al.*, 2015; El Gamal *et al.*, 2016a, 2016b; Agarwal *et al.*, 2017) and a single strain (MMB-1) of *Marinomonas mediterranea* (Agarwal *et al.*, 2014). These two genera of *Gammaproteobacteria* belong to different Orders, raising the possibility that the associated biosynthetic gene cluster (BGC) was exchanged by horizontal gene transfer. Biosynthetically related brominated natural products (BNPs) have also been reported from diverse organisms ranging from sponge-associated cyanobacteria to seabirds (Agarwal *et al.*, 2017). In addition to pentabromopseudilin, *Pseudoalteromonas* strains produce brominated phenol and pyrrole monomers and dimers as well as polybrominated diphenyl ethers (PBDEs). One brominated pyrrole monomer, tetra-bromopyrrole, appears to play an important role in the settlement and development of coral larvae (Tebben *et al.*, 2011; Sneed *et al.*, 2014). PBDEs are also of particular interest due to their negative implications for human health via endocrine disruption and links to neurodevelopmental deficits and cancer (Hallgren *et al.*, 2001; Zhou *et al.*, 2002; Fernie *et al.*, 2005; Richardson *et al.*, 2008).

Starting in the late 1920s, PBDEs were produced industrially and used as flame-retardants in a wide variety of products such as electronics and textiles (Johnson-Restrepo and Kannan, 2009). While anthropogenic PBDEs can be

Received 27 October, 2018; revised 10 January, 2019; accepted 12 January, 2019. \*For correspondence. E-mail pjensen@ucsd.edu; Tel. 858-534-7322; Fax 858-534-1318.

†Present address: Georgia Institute of Technology School of Chemistry & Biochemistry 901 Atlantic Drive Atlanta, GA 30332-0400, USA.

introduced into the ocean via surface runoff, there is increasing evidence that these compounds are also produced naturally in the marine environment (Reddy *et al.*, 2004; Teuten and Reddy, 2007). Due to their lipophilicity, these molecules bioaccumulate in the blubber and tissues of marine mammals, thus suggesting that a human diet rich in seafood could lead to increased PBDE exposure (Venkateswaran and Dohmoto, 2000; Malmvärn *et al.*, 2005; Teuten *et al.*, 2005).

The recent identification of the *bmp* gene cluster as the biosynthetic origin of pentabromopseudilin and related BNPs in *M. mediterranea* MMB-1 and select *Pseudoalteromonas* strains (Agarwal *et al.*, 2014; El Gamal *et al.*, 2016a) provides a unique opportunity to more broadly assess the biosynthesis of these compounds. The *bmp* cluster encodes the production of polybrominated phenol and pyrrole monomers (2–4), the phenolic homodimeric antibiotic bromophene (5), PBDEs (6), polybrominated biphenyls (7) and the brominated phenol/pyrrole heterodimer pentabromopseudilin (1) (Fig. 1). The gene cluster encodes an ACP-TE di-domain protein (Bmp1) and two flavin-dependent brominases (Bmp2 and Bmp5) that catalyse the bromination of pyrrole and phenol monomers respectively (2–4). Bmp3, 4 and 8 play roles in bromopyrrole monomer synthesis and Bmp6 is involved in the biosynthesis of bromophene


and PBDEs (6–7). Bmp7 catalyses the oxidative coupling of bromophenol and bromopyrrole monomers in the presence of Bmp9, Bmp10 and NADH.

Previous studies of BNP production in *Pseudoalteromonas* spp. focused on a few specific strains (Isnansetyo and Kamei, 2003; Fehér *et al.*, 2010). As a result, we lack a broader understanding of the diversity and distribution of the gene cluster within the genus and how this relates to compound production. Using metabolomics and a multilocus phylogeny generated from 144 *Pseudoalteromonas* strains, we mapped the distribution of *bmp* genes and pentabromopseudilin production. The results suggest a complex evolutionary history for the gene cluster and provide clues as to how this biosynthetic capacity evolved within the genus.

## Results


### *Pseudoalteromonas* species phylogeny

A *Pseudoalteromonas* species phylogeny that includes type strains for all 42 named species encompasses considerable diversity (Fig. 2). The primary bifurcation distinguishes the *Pseudoalteromonas spongiae* and *Pseudoalteromonas piratica* type strains from the remainder of the genus, which can be delineated into three major clades. Clade 3 contains the




**Fig. 1.** The *bmp* gene cluster and its products.

A. Five versions of the *bmp* gene cluster have been identified: four in *Pseudoalteromonas* spp. (A, C-E) and one in *M. mediterranea* MMB-1 (B). Hash marks indicate an ~150 bp insertion in *bmp1* (version C). The small molecule products detected from each version are indicated. B. Bmp functional roles in the biosynthesis of brominated natural products. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**Fig. 2.** *Pseudoalteromonas* species phylogeny and *bmp* gene cluster distribution. A maximum likelihood phylogeny generated from concatenated 16S, *gyrB*, *pyrH*, *recA* and *rpoD* nucleotide sequences. The three least inclusive clades containing the *bmp* cluster are highlighted in color. Lettered brackets (A, C, D and E) indicate the version of the *bmp* cluster. The 42 type strains are indicated in bold italics and the 17 strains isolated as part of this study by CN prefixes followed by country of origin (USA or Fiji) and source (SW = seawater, PT = plankton tow, GA = green alga, RA = red alga, BA = brown alga). Squares indicate *bmp* distribution (filled = present, empty = absent) and *bmp* version (fully filled = version A, half-filled = partial BCG). Method of detection (GM = genome sequence or PCR) indicated. Filled triangles indicate the detection of pentabromopseudolin, half-filled triangles indicate the detection of tetrabromopseudolin and empty triangles indicate that no compounds associated with the *bmp* gene cluster were detected. Roman numerals (as designated in Fig. 3) indicate 17 candidate new species identified based on coherence between 95% ANI groupings and the species phylogeny. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**Fig. 3.** ANI analysis. Average nucleotide identity of 91 *Pseudoalteromonas* genome sequences. The vertical dashed line specifies 95% ANI and the grey circles designate ANI species groupings. Strain names are followed by clade designations (from Fig. 2) and the 28 ANI groups (95%) that lack type strains indicated by Roman numerals (I–XXVIII). Coloured blocks correspond to sub-clade designations (from Fig. 2). The distribution of the 10 genes in the *bmp* gene cluster are displayed on the right with the version of the gene cluster (from Fig. 1) indicated on the right. Hash marks indicate an ~150 bp insertion in *bmp1*. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

*Pseudoalteromonas ulvae*, *Pseudoalteromonas tunicata* and *Pseudoalteromonas denitrificans* type strains while the remainder of the type strains fall within clades 1 and 2. Numerous clades at varying depths in the phylogeny do not contain a type strain, suggesting some species level diversity has yet to be described. To further assess the potential for taxonomic novelty, average nucleotide identity (ANI) was calculated among all publicly available genome sequences ( $N = 91$ ). At the suggested >95% ANI cut-off for species designations (Goris *et al.*, 2007), 42 species level groups were resolved of which 14 contained a type strain (Fig. 3). Two of these (*Pseudoalteromonas elyakovii*/*Pseudoalteromonas piscicida* and *Pseudoalteromonas flavipulchra*/*Pseudoalteromonas distincta*) contain more than one type strain suggesting these lineages have been over-described. The 28 ANI groups that lacked a type strain were assigned Roman numerals (I–XXVIII, Fig. 3). The 26 type strains for which genome sequences were not available likely correspond to many of these ANI groups, however at least two can be considered unnamed based on this metric. To further address taxonomic novelty, we searched the

species tree for clades that lacked a type strain and were coherent with a 95% ANI group. Using this approach, we identified 17 candidate new species, which are indicated in Fig. 2 by the Roman numeral assigned to their ANI group. Of the 17 strains isolated as part of this study (CN designations, Fig. 2), seven originated from the United States and 10 from Fiji. All of the Fiji strains were isolated from macroalgae, and a majority clade with *Pseudoalteromonas luteoviolacea* while four clade with one of the new ANI species (XXVI). Given that the processing and isolation efforts were not consistent among sample types and locations, no correlations can be drawn between these variables and the species phylogeny.

*Distribution of the bmp gene cluster among Pseudoalteromonas spp.*

We next mapped the distribution of the *bmp* gene cluster onto the *Pseudoalteromonas* species phylogeny. Of the 101 *Pseudoalteromonas* genomes mined (91 published, 10 unpublished), 19 contain some version of this gene

cluster and all of these fall within clade 2 (Fig. 2). Three least inclusive sub-clades containing the *bmp* BGC can be identified: one containing *P. luteoviolacea*, one containing *P. phenolica*, and one containing five named species. In cases where genome sequences were not available, PCR was used to test for the presence of the *bmp* BGC. The 12 strains that were PCR positive and sequence verified for the *bmp* cluster all fall within the least inclusive clade containing *P. luteoviolacea*. *Pseudoalteromonas luteoviolacea* and *P. phenolica* are the only named species whose type strains contain the BGC. Six of the candidate ANI species also contain the BGC (sp. III, XXII, XXIII, XXV, XXVI and XXVII).

A comparative analysis of the *bmp* gene cluster revealed four versions (A, C–E) among the *Pseudoalteromonas* strains and a fifth version (B) in *M. mediterranea* strain MMB-1 (Fig. 1). Version A contains 10 genes and has been experimentally linked to the production of pentabromopseudilin and related BNPs (Agarwal *et al.*, 2014). It was observed in 25 of the clade 2 strains including the *P. phenolica* and *P. luteoviolacea* type strains and five of the candidate ANI species (Fig. 2). Version B was only observed in a single *M. mediterranea* strain (MMB-1) and has one additional gene annotated as a putative permease. Version C has an ~150 bp insertion in *bmp1* suggesting it may be non-functional. This insertion was sequence verified by PCR in the two strains (NCIMB 1942 and NCIMB 2035) in which it was observed. The genome sequences for PS5 (ANI sp. IV) and A757 (ANI sp. III) contain a partial *bmp* cluster that lacks *bmp5*–8 (version D) (Fig. 3). These are the only two strains with any form of the BGC in the least inclusive clade containing 2ta6 and *P. peptidolytica* (Fig. 2). Strains S4047 and S4054 (both ANI sp. XXIII) possess version E of the gene cluster, which lacks *bmp3*–7 (Fig. 3). This truncated version, along with a pair of strains that lack the entire gene cluster (H33 and H33-S), are part of a well-supported sub-clade that is dominated by version A (Fig. 2), suggesting they result from gene loss events. Strains isolated as part of this study and for which genomes were not available were assigned version A if sequence verified PCR products from both *bmp2* and *bmp5* were detected. They were not assigned version B, since this has only been detected in a single strain of *M. mediterranea*, or versions C–E based on the results of the metabolomics analyses (below).

#### Pentabromopseudilin production


*Pseudoalteromonas* strains possessing different versions of the *bmp* gene cluster were analysed for pentabromopseudilin (1) production (Supporting Information Fig. S1). Pentabromopseudilin is a heterodimer comprised of brominated pyrrole and phenol monomers. All 10 genes in version A of the

gene cluster are required for the production of this compound in *Pseudoalteromonas* spp. (Agarwal *et al.*, 2014). Therefore, any *Pseudoalteromonas* strains that produce this compound can be expected to contain *bmp* version A, since version B has not been observed in the genus. Of the 25 *Pseudoalteromonas* strains containing *bmp* version A, 24 were confirmed for pentabromopseudilin production (Fig. 2) thus matching metabolic predictions. The 25th strain (H11) was not available for testing. H33 and H33-S are the only strains in clade 2 that are missing the entire *bmp* gene cluster and, as expected, no BNPs were detected (Vynne *et al.*, 2012). We also failed to detect brominated compounds from the strains that possess *bmp* version C (NCIMB 1942 and NCIMB 2035). The 150 bp insertion in the ACP-TE encoding *bmp1* gene distinguishes version C from A and may explain the lack of BNP production in these strains. Tetrabromopyrrole (2) was the only BNP detected from strains with *bmp* version D, which is consistent with the expected phenotype given that genes encoding production of the polybrominated phenol monomer (*bmp5*–6), the bromopyrrole homodimer (*bmp7*) and tribromopyrrole via the debromination of tetrabromopyrrole (*bmp8*) are missing (Fig. 1B). Additionally, BNPs were not detected from the two strains containing version E of the BGC (S4047 and S4054), which lacks *bmp3*–7. These results are consistent with the functional characterization of the gene cluster, as the products of the missing genes are required to synthesize brominated pyrrole and phenol monomers.

#### Evolutionary history of the *bmp* gene cluster

A concatenated phylogeny of the *bmp* genes derived from all 19 *Pseudoalteromonas* genome sequences that contained some version of the BGC was compared with a species tree generated for the same strains (Fig. 4). While no outgroup was used in the *bmp* phylogeny, the sequences from *M. mediterranea* strain MMB-1 were included and the tree was rooted consistently with the species tree (Fig. 4A). With the exception of the position of strains MMB-1 and S3663, the two trees are largely congruent. The *bmp* phylogeny shows that the sister clades containing *P. luteoviolacea* and A757 and the more distantly related *P. phenolica* clade are consistent with the species tree, supporting a single acquisition of the *bmp* cluster in the genus. The relationship between the *bmp* BGCs observed in *M. mediterranea* strain MMB-1 and *Pseudoalteromonas* spp. supports the hypothesis that it was acquired by horizontal gene transfer.

To further assess if a single acquisition event could account for the distribution of the *bmp* gene cluster in the genus *Pseudoalteromonas*, we explored its genomic environment, expecting it to be similar among strains. Pairwise comparisons revealed high levels of sequence identity in and around the *bmp* BGC for strains within the large clade containing CPMOR-1 and -2 indicating positional conservation in this lineage (Fig. 5A). These results



**Fig. 4.** *Bmp* gene phylogeny.

A. Concatenated maximum likelihood phylogeny of *bmp* gene sequences.

B. Concatenated maximum likelihood phylogenies of 16S, *gyrB*, *pyrH*, *recA* and *rpoD* sequences derived from the 19 *Pseudoalteromonas* and one *M. mediterranea* strain (MMB-1) that contain the *bmp* gene cluster. The *bmp* gene tree is rooted based on the species phylogeny while the species tree is rooted with MMB-1. Type strains are indicated by species names.


further support the hypothesis that strain S4054 experienced partial BGC loss while strain NCIMB 1942 experienced BGC modification resulting in *bmp* versions C and E respectively. The detection of a remnant of *bmp7* in strain S4054 provides additional support for the gene loss hypothesis (Supporting Information Fig. S2), while it can be inferred that the complete BGC was lost in strain H33. Outside of this clade, the lack of sequence identity between strains A757 and H33 suggests that the BGC is in a different genomic environment in strain A757 and the related strain PS5, the only two strains that possess version D of the BGC. When more distantly related strains are compared (HI1 and *P. phenolica*, Fig. 5B), the lack of sequence identity outside the BGC is more difficult to interpret, but could also indicate different genomic environments. A comparison of strain HI1, which contains *bmp* version A, with strain A757, which contains version D, reveals little sequence identity outside of the BGC (Fig. 5C), again suggesting different genomic environments with version D representing a partial loss of the cluster. While congruence between the *bmp* gene phylogeny and the *Pseudoalteromonas* species phylogeny suggests a single acquisition event for the BGC, the genomic environments in which it occurs could be

explained by either multiple acquisition events or intragenomic rearrangements. Regardless of which explanation is correct, there is clear evidence within sub-clades of the *Pseudoalteromonas* phylogeny that the gene cluster has been maintained, modified or lost following a model of vertical inheritance (Fig. 6).

## Discussion

Characterization of the *bmp* BGC revealed it accounts for a diverse set of compounds including the potent antibiotic pentabromopseudilin (Burkholder *et al.*, 1966), the coral larval settlement cue tetrabromopyrrole (El Gamal *et al.*, 2016a) and PBDEs whose industrial production and use as flame-retardants have been banned in some countries due to human health implications (Johnson-Restrepo and Kannan, 2009). These compounds have primarily been reported from members of the genus *Pseudoalteromonas*, yet despite their ecological and environmental significance, compound production and the distribution of the associated biosynthetic genes have not been investigated.

The results presented here indicate a narrow distribution and complex evolutionary history for the *bmp* gene cluster that includes gene modifications and deletions that affect



**Fig. 5.** Genomic environment of the *bmp* gene cluster.

A. The species phylogeny is contrasted with conservation of the *bmp* gene cluster (red) and surrounding genomic environments (grey). Strain names are followed by the version of the gene cluster (A, C, D or E) and candidate ANI species identifiers. Opacity indicates percent sequence identity with anything below 50% not shown.

B. A high level of sequence identity is observed in the *bmp* gene cluster between two distantly related strains (HI1 and *P. phenolica*) but not in the surrounding genomic environment.


C. Strain A757 (*bmp* version D) reveals a partial loss of the *bmp* gene cluster. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

compound production. While congruence between the species and *bmp* phylogenies supports a single acquisition event (Fig. 4), this model requires eight complete or partial gene cluster losses to explain the observed distributions. Given that the gene cluster was observed in different genomic environments in some strains, the possibility of multiple acquisition events cannot be ruled out. However, different environments could also be accounted for by intragenomic rearrangements, as has previously been shown for the genus *Salinispora* (Letzel *et al.*, 2017). While the evolutionary history of the *bmp* gene cluster remains unresolved at

the genus level, there are three points from which vertical inheritance can be inferred with confidence (Fig. 6). Even within these sub-lineages, there is clear evidence of gene cluster loss or modification, attesting to its complex evolutionary history.

There is ample evidence that the gene clusters associated with natural product biosynthesis are exchanged by horizontal gene transfer. Despite this, there was a strong correlation between taxonomy, *bmp* composition and compound production within the genus *Pseudoalteromonas*. For example, all *P. phenolica* strains analysed maintained





**Fig. 6.** Evolutionary history of the *bmp* gene cluster in *Pseudoalteromonas*.

A. Inferred ancestral gene clusters are mapped onto a condensed species phylogeny (animated from Fig. 2). The distribution of each gene (coloured lines) is indicated based on presence or absence in that lineage (truncated or missing lines = absence).

B. Versions of the gene cluster (A, C, D or E) observed in relation to the tree with evolutionary events indicated as maintenance, loss or modification. Losses include complete or partial deletion of the gene cluster. An insertion in *bmp1* is the only modification observed (indicated with a star). Percent sequence identities between homologues indicated. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

the same version of the BGC and produced the same natural products. The least inclusive clade that includes *P. luteoviolacea* is also highly conserved for these features, with the exception of a few loss events. The conservation of *bmp* within certain clades suggests that the products encoded provide an important selective advantage and add to growing evidence that natural products can be useful taxonomic markers for bacteria (Hoffmann *et al.*, 2018).

During the course of this study, gene clusters related to *bmp* were detected in sponge cyanobacterial symbionts (Agarwal *et al.*, 2017). While these gene clusters have yet to be fully characterized, they include *bmp5-7* homologues, which encode the production of PBDEs, but not the genes required to make brominated pyrrole molecules

or pentabromopseudilin. These results suggest that an even more complex evolutionary history remains to be resolved for the *bmp* gene cluster.

Remarkable new insight is being gained into the evolutionary processes that drive natural product biosynthesis (Medema *et al.*, 2014; Ruzzini and Clardy, 2016; Lind *et al.*, 2017). These studies have helped reveal the dynamic processes of gene gain, loss and degradation in BGCs (Letzel *et al.*, 2017) and how subtle changes in regulatory genes can have substantial effects on both gene expression and compound production (Amos *et al.*, 2017). This study adds to the growing number of examples in which the evolutionary history of a BGC provides insight into the processes that establish lineage specific chemical diversity (Freel *et al.*, 2011).

While multiple versions of the *bmp* cluster were observed, selection appears to favour the maintenance of version A, which encodes the biosynthesis of the potent antibiotic pentabromopseudilin (1). In cases where the BGC was lost, it may be complemented by the products of another BGC, as was shown with siderophore biosynthesis in marine actinomycetes (Bruns *et al.*, 2017). Alternatively, relaxed selective pressures associated with some habitats may lead to BGC loss. Regardless of the factors driving loss events, their occurrence within well-supported lineages demonstrates the unpredictable nature of microbial natural product discovery and presents a caveat to their applications in chemotaxonomy. The bioinformatic analyses performed here provided a powerful predictor of chemical phenotypes, as there was a perfect correlation between the version of the BGC detected and BNP production among the strains analysed. Continued access to more genome sequences we will undoubtedly expand our understanding of the evolutionary history of the *bmp* gene cluster in *Pseudoalteromonas* and other taxa in which it may be observed.

## Experimental procedures

### Bacterial isolation

Seawater samples were collected at Scripps Institution of Oceanography (SIO), plankton tows (20 µm mesh size) were performed from the SIO pier, and algal samples were collected by SCUBA from Catalina Island, CA and the Coral Coast, Fiji (Supporting Information Table S1). The algae, which were identified as red, green or brown based on visual appearance, were gently shaken to remove excess water before a small piece of tissue was added to 10 ml of 0.2 µm-filtered seawater. The algal samples were mixed and diluted 1:10, 1:50 and 1:100 in sterile seawater, and 100 µl of each dilution spread onto Difco 2216 media with 15 g/l agar added. Seawater and plankton tow samples were diluted 1:2, 1:10 and 1:50 and similarly plated. Bacterial colonies were isolated based on morphology, specifically selecting for dark purple and other pigmentation that is characteristic of *Pseudoalteromonas* spp. Strains were identified at the genus level based on 16S rRNA sequence analysis (see below) and glycerol stocks frozen (−80 °C) in Difco 2216 broth with 25% glycerol.

### Genome mining, primer design and PCR

Published genome sequences are available for 16 of 42 *Pseudoalteromonas* type strains while *gyrB* and 16S sequences are available for an additional 12 (Supporting Information Table S2). Our analysis also included an additional 75 published *Pseudoalteromonas* genomes (Supporting Information Table S3), 10 unpublished genomes and 17 strains isolated as part of this study. Genome

sequences were acquired from the NCBI and JGI public databases or sequenced at SIO. They were uploaded and annotated in RAST (Aziz *et al.*, 2008) and mined for 16S, *gyrB*, *pyrH*, *recA* and *rpoD* based on annotations. *Bmp1-10* gene sequences were identified using the internal BLAST search tool in RAST with gene sequences from *P. luteoviolacea* strain 2ta16 as queries. Ten unpublished *Pseudoalteromonas* genomes sequenced at the Technical University of Denmark were similarly mined. PCR was used to generate sequences for strains isolated as part of this study but for which genome sequences were not available. Degenerate PCR primers were designed based on the translated and MUSCLE aligned *gyrB*, *pyrH*, *recA*, *rpoD* and *bmp* sequences (Edgar *et al.*, 2004). The primers were validated across different *Pseudoalteromonas* spp. using Primacal software and manual analyses (Supporting Information Table S4) (Gadberry *et al.*, 2005). All of the mined and amplified sequences have been deposited in NCBI (MK421604-MK421630; MK469519-MK468590).

For genomic DNA extractions, strains were cultured overnight at room temperature in 10 ml of marine broth (Difco 2216) with shaking at 230 rpm. Cells were pelleted (16,000g, 2 min) from 1 ml of culture, re-suspended in 600 µl of Nuclei Lysis Solution (Promega), and incubated at 80 °C for 5 min after which 200 µl of Protein Precipitate Solution (Promega) was added. The samples were vortexed at high speed for 20 s, incubated on ice for 5 min, centrifuged (16,000g, 10 min), and the supernatant transferred into 600 µl of isopropanol. The samples were mixed by gentle inversion, centrifuged (13,000 rpm, 10 min), and the DNA pellet washed once with 70% ethanol, dried and rehydrated in water. PCR thermocycling conditions were as follows: 5 min of initial denaturation at 95 °C, 30 cycles of denaturation at 95 °C for 1 min, annealing for 1 min at 52–55 °C, extension at 72 °C for 1 min, and a final extension at 72 °C for 5 min. PCR products were sequenced using Sanger methods by Eton Biosciences (<https://www.etonbio.com/>) and trimmed for quality before analysis.

### Phylogenetic analyses

Nucleotide sequences used to establish the species phylogeny were aligned using MAFFT (Katoch *et al.*, 2002) then concatenated. Maximum likelihood analysis of the partitioned data was run using RAxML with 100 rapid bootstrap replicates and the GTR + G model (Stamatakis, 2014). To assess the phylogeny of the *bmp* cluster, the MUSCLE alignments of the *bmp1*, 2, 9 and 10 gene nucleotide sequences were concatenated and a maximum likelihood analysis performed using RAxML with 100 rapid bootstrap replicates and GTR + G model (Edgar *et al.*, 2004; Stamatakis, 2014). All trees were visualized using FigTree v1.4.2.

*Pentabromopseudilin analysis*

Bacterial cultures were grown at room temperature with agitation (200 rpm) for 48 h in 50 ml of Difco 2216 media with 1 g/l of KBr added. The cultures were extracted twice with 50 ml ethyl acetate and the organic fractions combined and dried *in vacuo*. The dried extracts were dissolved in 500 µl of MeOH and analysed by LC/MS using previously described methods (Agarwal *et al.*, 2014). Pentabromopseudilin production was verified in comparison to an authentic standard or based on previously reported results.

*Comparative genomics*

ANI was determined for 91 *Pseudoalteromonas* genomes using custom scripts available at [https://github.com/juanu/ANI\\_analysis](https://github.com/juanu/ANI_analysis). A distance matrix was generated from all pairwise comparisons and ANI divergence (100 – ANI) was calculated. Pairwise comparisons between contigs to assess the genomic environment of the *bmp* BGC were performed using the Artemis Comparison Tool v13.0.0 (Carver *et al.*, 2005).

**Acknowledgements**

This work was jointly supported by the US National Science Foundation (OCE-1313747) and the US National Institute of Environmental Health Sciences (P01-ES021921) through the Oceans and Human Health program. We acknowledge N. Patin for providing strains and the National Institutes of Health award U19-TW007401-01 for access to samples from Fiji. The authors declare no conflict of interest.

**References**

Agarwal, V., El Gamal, A., Yamanaka, K., Poth, D., Kersten, R.D., Schorn, M., *et al.* (2014) Biosynthesis of polybrominated aromatic organic compounds by marine bacteria. *Nat Chem Biol* **10**: 640–647. <https://doi.org/10.1038/nchembio.1564>.

Agarwal, V., Blanton, J.M., Podell, S., Taton, A., Schorn, M. A., Busch, J., *et al.* (2017) Metagenomic discovery of polybrominated diphenyl ether biosynthesis by marine sponges. *Nat Chem Biol* **13**: 537–543. <https://doi.org/10.1038/nature15540>. Genetic.

Amos, G.C.A., Awakawa, T., Tuttle, R.N., Letzel, A.-C., Kim, M. C., Kudo, Y., *et al.* (2017) Comparative transcriptomics as a guide to natural product discovery and biosynthetic gene cluster functionality. *Proc Natl Acad Sci* **114**: E11121–E11130. <https://doi.org/10.1073/pnas.1714381115>.

Aziz, R.K., Bartels, D., Best, A., DeJongh, M., Disz, T., Edwards, R.A., *et al.* (2008) The RAST server: rapid annotations using subsystems technology. *BMC Genomics* **9**: 1–15. <https://doi.org/10.1186/1471-2164-9-75>.

Bruns, H., Crüsemann, M., Letzel, A.-C., Alanjary, M., McInerney, J.O., Jensen, P.R., *et al.* (2017) Function-related

replacement of bacterial siderophore pathways. *ISME J* **12**: 320–329. <https://doi.org/10.1038/ismej.2017.137>.

Burkholder, P.R., Pfister, R.M., and Leitz, F.H. (1966) Production of a pyrrole antibiotic by a marine bacterium. *Appl Microbiol* **14**: 649–653.

Carver, T.J., Rutherford, K.M., Berriman, M., Rajandream, M. A., Barrell, B.G., and Parkhill, J. (2005) ACT: the Artemis comparison tool. *Bioinformatics* **21**: 3422–3423. <https://doi.org/10.1093/bioinformatics/bti553>.

Edgar, R.C., Drive, R.M., and Valley, M. (2004) MUSCLE : Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**: 1792–1797. <https://doi.org/10.1093/nar/gkh340>.

El Gamal, A., Agarwal, V., Diethelm, S., Rahman, I., Schorn, M.A., Sneed, J.M., *et al.* (2016a) Biosynthesis of coral settlement cue tetrabromopyrrole in marine bacteria by a uniquely adapted brominase–thioesterase enzyme pair. *Proc Natl Acad Sci* **113**: 3797–3802. <https://doi.org/10.1073/pnas.1519695113>.

El Gamal, A., Agarwal, V., Rahman, I., and Moore, B.S. (2016b) Enzymatic reductive dehalogenation controls the biosynthesis of marine bacterial pyrroles. *J Am Chem Soc* **138**: 13167–13170. <https://doi.org/10.1021/jacs.6b08512>.

Fehér, D., Barlow, R., McAtee, J., and Hemscheidt, T.K. (2010) Highly brominated antimicrobial metabolites from a marine *Pseudoalteromonas* sp. *J Nat Prod* **73**: 1963–1966. <https://doi.org/10.1021/np100506z>.

Fernie, K.J., Shutt, J.L., Mayne, G., Hoffman, D., Letcher, R. J., Drouillard, K.G., and Ritchie, I.J. (2005) Exposure to polybrominated diphenyl ethers (PBDEs): changes in thyroid, vitamin A, glutathione homeostasis, and oxidative stress in American kestrels (*Falco sparverius*). *Toxicol Sci* **88**: 375–383. <https://doi.org/10.1093/toxsci/kfi295>.

Freel, K.C., Nam, S.J., Fenical, W., and Jensen, P.R. (2011) Evolution of secondary metabolite genes in three closely related marine Actinomycete species. *Appl Environ Microbiol* **77**: 7261–7270. <https://doi.org/10.1128/AEM.05943-11>.

Gadberry, M.D., Malcomber, S.T., Doust, A.N., and Kellogg, E. A. (2005) Primaclade - a flexible tool to find conserved PCR primers across multiple species. *Bioinformatics* **21**: 1263–1264. <https://doi.org/10.1093/bioinformatics/bti134>.

Goris, J., Konstantinidis, K.T., Klappenbach, J.A., Coenye, T., Vandamme, P., and Tiedje, J.M. (2007) DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* **57**: 81–91. <https://doi.org/10.1099/ijs.0.64483-0>.

Hallgren, S., Sinjari, T., Håkansson, H., and Darnerud, P. (2001) Effects of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) on thyroid hormone and vitamin A levels in rats and mice. *Arch Toxicol* **75**: 200–208. <https://doi.org/10.1007/s002040000208>.

Hoffmann, T., Krug, D., Bozkurt, N., Duddela, S., Jansen, R., Garcia, R., *et al.* (2018) Correlating chemical diversity with taxonomic distance for discovery of natural products in Myxobacteria. *Nat Commun* **9**: 1–10. <https://doi.org/10.1038/s41467-018-03184-1>.

Isnansetyo, A., and Kamei, Y. (2003) MC21-A, a bactericidal antibiotic produced by a new marine bacterium, *Pseudoalteromonas Phenolica* sp. Nov. O-BC30T, against methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* **47**: 480–488. <https://doi.org/10.1128/AAC.47.2.480>.

- Johnson-Restrepo, B., and Kannan, K. (2009) An assessment of sources and pathways of human exposure to polybrominated diphenyl ethers in the United States. *Chemosphere* **76**: 542–548. <https://doi.org/10.1016/j.chemosphere.2009.02.068>.
- Katoh, K., Misawa, K., Kuma, K.-i., and Miyata, T. (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast fourier transform. *Nucleic Acids Res* **30**: 3059–3066.
- Letzel, A.C., Li, J., Amos, G.C.A., Millán-Aguíñaga, N., Ginigini, J., Abdelmohsen, U.R., et al. (2017) Genomic insights into specialized metabolism in the marine actinomyete *Salinispora*. *Environ Microbiol* **19**: 3660–3673. <https://doi.org/10.1111/1462-2920.13867>.
- Lind, A.L., Wisecaver, J.H., Lameiras, C., Wiemann, P., Palmer, J.M., Keller, N.P., et al. (2017) Drivers of genetic diversity in secondary metabolic gene clusters within a fungal species. *PLoS Biol* **15**: 1–26. <https://doi.org/10.1371/journal.pbio.2003583>.
- Lovell, F.M. (1966) The structure of a bromine-rich marine antibiotic. *Journal of The American Chemical Society* **88**: 4510–4511.
- Malmväm, A., Marsh, G., Kautsky, L., Athanasiadou, M., Bergman, A., and Asplund, L. (2005) Hydroxylated and methoxylated brominated diphenyl ethers in the red algae *Ceramium tenuicorne* and blue mussels from the Baltic Sea. *Environ Sci Technol* **39**: 2990–2997.
- Medema, M.H., Cimermančić, P., Sali, A., Takano, E., and Fischbach, M.A. (2014) A systematic computational analysis of biosynthetic gene cluster evolution: lessons for engineering biosynthesis. *PLoS Comput Biol* **10**: e1004016. <https://doi.org/10.1371/journal.pcbi.1004016>.
- Reddy, C.M., Xu, L., O'Neil, G.W., Nelson, R.K., Eglinton, T. I., Faulkner, D.J., et al. (2004) Radiocarbon evidence for a naturally produced, bioaccumulating halogenated organic compound. *Environ Sci Technol* **38**: 1992–1997.
- Richardson, V.M., Staskal, D.F., Ross, D.G., Diliberto, J.J., DeVito, M.J., and Birnbaum, L.S. (2008) Possible mechanisms of thyroid hormone disruption in mice by BDE 47, a major polybrominated diphenyl ether congener. *Toxicol Appl Pharmacol* **226**: 244–250. <https://doi.org/10.1016/j.taap.2007.09.015>.
- Ruzzini, A.C., and Clardy, J. (2016) Gene flow and molecular innovation in bacteria. *Curr Biol* **26**: R859–R864. <https://doi.org/10.1016/j.cub.2016.08.004>.
- Sneed, J.M., Sharp, K.H., Ritchie, K.B., and Paul, V.J. (2014) The chemical cue tetrabromopyrrole from a biofilm bacterium induces settlement of multiple Caribbean corals. *Proc R Soc B Biol Sci* **281**: 1–9. <https://doi.org/10.1098/rspb.2013.3086>.
- Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>.
- Tebben, J., Tapiolas, D.M., Motti, C.A., Abrego, D., Negri, A.P., Blackall, L.L., et al. (2011) Induction of larval metamorphosis of the coral *Acropora millepora* by tetrabromopyrrole isolated from a *Pseudoalteromonas* bacterium. *PLoS One* **6**: 1–8. <https://doi.org/10.1371/journal.pone.0019082>.
- Teuten, E.L., and Reddy, C.M. (2007) Halogenated organic compounds in archived whale oil: a pre-industrial record. *Environ Pollut* **145**: 668–671. <https://doi.org/10.1016/j.envpol.2006.08.022>.
- Teuten, E.L., Xu, L., and Reddy, C.M. (2005) Two abundant bioaccumulated halogenated compounds are natural products. *Science* **307**: 917–920. <https://doi.org/10.1126/science.1106882>.
- Venkateswaran, K., and Dohmoto, N. (2000) *Pseudoalteromonas peptidolytica* sp. Nov., a novel marine mussel-thread-degrading bacterium isolated from the sea of Japan. *Int J Syst Evol Microbiol* **50**: 565–574.
- Vynne, N.G., Månsson, M., Nielsen, K.F., and Gram, L. (2011) Bioactivity, chemical profiling, and 16S rRNA-based phylogeny of *Pseudoalteromonas* strains collected on a global research cruise. *Marine Biotechnol* **13**: 1062–1073. <https://doi.org/10.1007/s10126-011-9369-4>.
- Vynne, N.G., Mansson, M., and Gram, L. (2012) Gene sequence based clustering assists in dereplication of *Pseudoalteromonas luteoviolacea* strains with identical inhibitory activity and antibiotic production. *Mar Drugs* **10**: 1729–1740. <https://doi.org/10.3390/md10081729>.
- Whalen, K.E., Poulson-Ellestad, K.L., Deering, R.W., Rowley, D.C., and Mincer, T.J. (2015) Enhancement of antibiotic activity against multidrug-resistant bacteria by the efflux pump inhibitor 3,4-dibromopyrrole-2,5-dione isolated from a *Pseudoalteromonas* sp. *J Nat Prod* **78**: 402–412. <https://doi.org/10.1021/np500775e>.
- Zhou, T., Taylor, M.M., DeVito, M.J., and Crofton, K.M. (2002) Developmental exposure to brominated diphenyl ethers results in thyroid hormone disruption. *Toxicol Sci* **116**: 105–116.

## Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

**Figure S1** Detection of pentabromopseudilin by LCMS. Strains isolated as part of this study were screened for pentabromopseudilin production. (A) Extracted ion chromatograms for the mass of pentabromopseudilin (553.67 g/mol) from a strain with version A of the BGC (CNY-791, top) and one that lacks the BGC (CNY-943, middle). Pentabromopseudilin standard (bottom). All strains containing *bmp* version A produced pentabromopseudilin with an absolute intensity of at least  $1.4 \times 10^6$  (B) Isotopic pattern of a pentabromopseudilin standard and the compound detected from strain CNY-791 (C).

**Fig. S2.** Evidence for partial loss of the *bmp* cluster. A region immediately downstream of *bmp2* in strain S4054 (version E) has high sequence identity to a portion of *bmp7* in strain HI1 (version A), supporting gene loss in S4054.

**Table S1.** Source of *Pseudoalteromonas* strains isolated as part of this study.

**Table S2.** Type strains without published genome sequences.

**Table S3.** *Pseudoalteromonas* genomes used in this study. Type strains indicated in bold. Roman numerals (I–XXVIII) correspond to the 28 ANI groups (95%) that lack type strains (from Fig. 3). Asterisks (\*) indicate 95% ANI groups where type strain genome sequences were unavailable.

**Table S4.** PCR primer sequences.