

Lawrence Berkeley National Laboratory

Recent Work

Title

Relative Yield Functions Pi Negative Mesons from Carbon

Permalink

<https://escholarship.org/uc/item/0pq300nd>

Authors

Jones, Stanley
White, Stephen

Publication Date

1949-12-07

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

UNCLASSIFIED

RELATIVE YIELD FUNCTIONS FOR π^- MESONS FROM CARBON

Stanley B. Jones and R. Stephen White

December 7, 1949

Berkeley, California

-2-

INSTALLATIONNo. of Copies

Argonne National Laboratory	8
Armed Forces Special Weapons Project	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute	1
Brookhaven National Laboratory	8
Bureau of Medicine and Surgery	1
Bureau of Ships	1
Carbide & Carbon Chemicals Corp. (K-25)	4
Carbide & Carbon Chemicals Corp. (Y-12)	4
Chicago Operations Office	1
Cleveland Area Office	1
Columbia University (Dunning)	1
Columbia University (Failla)	1
Dow Chemical Company	1
General Electric Company, Richland	6
Hanford Operations Office	1
Idaho Operations Office	1
Iowa State College	2
Kansas City	1
Kellex Corporation	2
Knolls Atomic Power Laboratory	4
Los Alamos	3
Mallinckrodt Chemical Works	1
Massachusetts Institute of Technology (Gaudin)	1
Massachusetts Institute of Technology (Kaufmann)	1
Mound Laboratory	3
National Advisory Committee for Aeronautics	2
National Bureau of Standards	2
Naval Radiological Defense Laboratory	2
NEPA Project	2
New Brunswick Laboratory	1
New York Operations Office	5
North American Aviation, Inc.	1
Oak Ridge National Laboratory	8
Patent Advisor, Washington	1
Rand Corporation	1
Sandia Base	1
Sylvania Electric Products, Inc.	1
Technical Information Branch, ORE	15
U. S. Public Health Service	1
UCLA Medical Research Laboratory (Warren)	1
University of California Radiation Laboratory	5
University of Rochester	2
University of Washington	1
Western Reserve University (Friedell)	2
Westinghouse Electric Corp.	4
Total	119

Information Division
Radiation Laboratory
Univ. of California
Berkeley, California

-3-

ABSTRACT

Relative Yield Functions for π^- Mesons From Carbon

Stanley B. Jones and R. Stephen White

December 7, 1949

This paper described determinations of the relative yield curves for π^- mesons generated by alpha-particle and proton beams striking carbon targets. The mesons counted had energies of 2-10 Mev and angles to the beam direction up to about 45° .

-4-

RELATIVE YIELD FUNCTIONS FOR π^- MESONS FROM CARBON

Stanley B. Jones and R. Stephen White

Radiation Laboratory, Department of Physics
University of California
Berkeley, California

December 7, 1949

The artificial production of mesons has made possible the determination of meson yield as a function of the energy of the bombarding particle. This paper reports such determinations for π^- mesons generated by the Berkeley 184-inch cyclotron with both proton and alpha-particle beams incident upon a carbon target with the restriction that only mesons of energies 2-10 Mev and of angle to the beam up to about 45° were counted. A preliminary report of this study has been published in abstract form.⁽¹⁾

¹S. B. Jones and R. S. White, Phys. Rev. 75, 1468 (1949); Phys. Rev. 76, 588 (1949).

The beam energies were selected by placing the targets at predetermined distances from the center of the cyclotron. As indicated in Fig. 1, the π mesons produced at the target move approximately in circular orbits in the magnetic field. In particular, the π^- mesons circulate in the counterclockwise direction as illustrated. Some of the low energy ones are slowed down and stopped in stacked photographic plates placed from four inches to seven inches from the leading edge of the target as shown in Fig. 2. These plates are then examined for meson tracks over an area corresponding to meson energies of 2-10 Mev by means of a microscope with a traveling stage.

The bombardments were made using the carbon targets illustrated schematically in Fig. 2. The target dimensions in the plane perpendicular to the beam

-5-

direction are 1 inch by 3 inches. The thickness for the alpha-particle beam was 1/16 inch with an exposure time of 20 minutes and for the proton beam, 1/32 inch with an exposure time of 30 seconds. Most of the mesons originate from the small target area, measuring 1 inch along the leading edge by 1/8 inch perpendicular to the edge, which first intercepts the beam, since this region is most intensely bombarded by the cyclotron beam. This was found by monitoring the target with thin polystyrene.

The integral of the beam-current was obtained by observing the relative amounts of C^{11} formed in the $C^{12}(\alpha, n)C^{11}$ and $C^{12}(p, pn)C^{11}$ reactions for the alpha-particle and proton bombardments, respectively, through measurement of the positron activity of C^{11} which has a half-life of 20.5 minutes. A correction was made for the decay of the C^{11} activity in the target during the bombardment by alpha-particles. A correction for the variation of the cross section for $C^{12}(\alpha, n)C^{11}(2)$ and $C^{12}(p, pn)C^{11}(3)$ with energy was also included.

²R. L. Thornton and R. W. Senseman, Phys. Rev. 72, 872 (1947)

³L. Aamodt, V. Peterson and R. Phillips, to be published.

At each beam energy, a single plate from the center of a stack of at least five plates was selected for study. Each plate was scanned to a depth of 2 mm. from the leading edge for a measured distance along the edge. Nearly identical areas were scanned on the plates for the various beam energies and meson density over the scanned and adjacent areas did not change rapidly. The number of mesons counted were corrected for the integrated beam current, plate thickness, and area scanned.

The apparatus was selective for π^- mesons. However, positive mesons might possibly have entered the plates and it is certain that some μ^- mesons also have

-6-

entered the plates. In order to include only π^- mesons, we counted only σ (star-forming) mesons. Since few, if any, μ^- mesons form stars^(4,5), the

⁴W. Y. Chang, Rev. Mod. Phys. 21, 166 (1949).

⁵Goldschmidt, Clermont, King, Muirhead and Ritson, Proc. Phys. Soc. 61, 183 (1948).

total number of π^- mesons in the energy interval examined will then be proportional to the number of σ mesons since a fixed fraction of π^- mesons form stars⁽⁶⁾.

⁶F. L. Adelman and S. B. Jones, Phys. Rev. 75, 1468 (1949).

The data are given in Table I.

Table I

Beam Energy	No. σ Mesons Observed	Relative Volume Scanned	Relative Integrated Beam Current	Corrected Relative Yield
<u>I. Alpha Particle Beam</u>				
390	241	1.00	1.00	100 percent
340	109	.740	1.80	34
305	24	.797	1.87	7
265	4	.806	3.85	0.5
<u>II. Proton Beam</u>				
345	75	1.00	1.00	100 percent
305	126	1.08	3.30	47
270	46	.794	3.57	22
235	19	.944	3.46	8
200	3	1.12	3.76	1
165	0	1.23	1.97	0

-7-

Plots of the data in the tables are given in Figs. 3 and 4.

Dr. W. H. Barkas, by the method of energy and momentum balance, has calculated the absolute minimum threshold for meson production using the most favorable reactions⁽⁷⁾. These are indicated in the figures.

⁷W. H. Barkas, Phys. Rev. 75, 1109 (1949)

It should be emphasized that this study does not give the relative total cross section for π^- meson production but only the relative yields in a particular meson energy interval. V. Peterson⁽⁸⁾ measured an absolute differential cross

⁸V. Peterson, Private communication.

section for π^- mesons produced by 390 Mev alpha-particles on a 1/4 inch carbon target. For π^- mesons of energy 2-5 Mev and of angles with the beam direction up to about 45° he found a value of $3.0 \pm 0.8 \times 10^{-32} \text{ cm}^2 \text{ Mev}^{-1} \text{ steradian}^{-1}$ per carbon nucleus. Richman and Wilcox⁽⁹⁾ measured a value of $2.1 \pm .6 \times 10^{-28} \text{ cm}^2$

⁹C. Richman and H. Wilcox, Bulletin Am. Phys. Soc., Stanford Meeting, Dec. 29, 1949)

steradian^{-1} per carbon nucleus at 90 degrees to the beam direction for the combined cross section for both π^- mesons and π^+ mesons produced by 345 Mev protons on carbon.

The authors are indebted to Professor Ernest O. Lawrence for his continued interest in this work. We also wish to thank Professor R. L. Thornton and Drs. Eugene Gardner, C. M. G. Lattes, and Hugh Bradner for much helpful discussion. We are very grateful to James Vale and the cyclotron crew for making the bombardments. This paper is based on work performed under the auspices of the Atomic Energy Commission.




Fig. 1

Plan view of cyclotron (meson trajectories enlarged).

0 1 2 3
FEET




Fig. 2

Schematic diagram of plate holder.

0 1 2 3
INCHES




Fig. 3
Meson excitation curve for protons, normalized to 100 at 345 Mev with a meson energy interval of 2-10 Mev.




Fig. 4

Meson excitation curve for alpha-particles normalized to 100 at 390 Mev with a meson energy interval of 2-10 Mev.