

## **UC Irvine**

### **Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health**

#### **Title**

A Food Impaction in Crohn's Ileal Stricture

#### **Permalink**

<https://escholarship.org/uc/item/0q917421>

#### **Journal**

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 11(1)

#### **ISSN**

1936-900X

#### **Authors**

Hokama, Akira  
Kishimoto, Kazuto  
Ihama, Yasushi  
[et al.](#)

#### **Publication Date**

2010

#### **Copyright Information**

Copyright 2010 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed

## A Food Impaction in Crohn's Ileal Stricture

Akira Hokama, MD, PhD\*  
Kazuto Kishimoto  
Yasushi Ihama  
Hiroschi Chinen  
Fukunori Kinjo  
Jiro Fujita

First Department of Internal Medicine, University of the Ryukyus, Okinawa, Japan

Supervising Section Editor: Sean Henderson, MD

Submission history: Submitted August 11, 2009; Revision Received August 28, 2009; Accepted September 14, 2009

Reprints available through open access at [http://escholarship.org/uc/uciem\\_westjem](http://escholarship.org/uc/uciem_westjem)

[West J Emerg Med. 2010; 11(1):106.]



**Figure.** Computed tomography scan showing a circumferential food obstruction (arrow) in an ileal stricture with wall thickening.

A 47-year-old man with a four-year history of ileo-colonic Crohn's disease (CD) presented with abdominal pain and vomiting. On examination, his temperature was 38.2°C and a distended and generalized tender abdomen without peritoneal signs was noted. Laboratory tests revealed an increased white blood cell count of 10,000/mm<sup>3</sup> and a C-reactive protein of 6.2 mg/dl. A plain abdominal radiograph revealed air-fluid levels in the small intestine. A computed tomography (CT) scan disclosed a circumferential obstruction (arrow) in an ileal stricture with wall thickening (Figure 1). When questioned, the patient stated that his previous dinner consisted of dried cuttlefish and vegetable roots, which might be difficult to digest. Gastrointestinal decompression with nasogastric tube and antibiotic treatment resolved his symptoms. The patient remained well and was discharged with maintenance infliximab treatment.

This case is an important reminder that CD patients are potentially at a greater risk of bowel obstruction with less digestible food. In other rare instances, bowel obstruction by

food, fruit pits, phytobezoar or foreign bodies has actually led physicians to the diagnosis of CD.<sup>1-3</sup> In these cases, a CT scan has been the key in assessing the bowel obstruction, providing a clear identification of the anatomical site to help identify the cause.

---

*Address for Correspondence:* Akira Hokama, MD, PhD. First Department of Internal Medicine, University of the Ryukyus, 207 Uehara, Nishihara, Okinawa 903-0215, Japan. Email: [hokama-a@med.u-ryukyu.ac.jp](mailto:hokama-a@med.u-ryukyu.ac.jp)

---

### REFERENCES

1. Kaufman D, Lazinger M, Fogel S, et al. Fruit pit obstruction leading to the diagnosis of Crohn's disease. *Am J Surg.* 2001; 182:530.
2. Shedda S, Frizelle F. Images in clinical medicine. Crohn's disease and an olive. *N Engl J Med.* 2006; 355:e22.
3. Slim R, Chemaly M, Yaghi C, et al. Silent disease revealed by a fruit-ileal obstruction. *Gut.* 2006; 55:181-90.