

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

SEAECH FOR CORE-EXCITED STATES IN N15

### Permalink

<https://escholarship.org/uc/item/0rc5g799>

### Authors

Harvey, B.G.  
Meriwether, J.R.  
Nercy, A. Bussiere de  
et al.

### Publication Date

1964-04-01

**University of California**  
**Ernest O. Lawrence**  
**Radiation Laboratory**

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

SEARCH FOR CORE-EXCITED STATES IN N<sup>15</sup>

**Berkeley, California**

## DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

B. G. Harvey, J. R. Meriwether, and A. Bussiere de Nercy<sup>†</sup>

Univ. of Calif., Lawrence Radiation Laboratory, Berkeley

and

D. J. Horen

U.S. Naval Radiological Defense Laboratory

#### ABSTRACT

Differential cross-sections for the scattering of 40.2-MeV alpha particles by  $N^{15}$  and  $O^{16}$  have been measured. The elastic cross section for  $N^{15}$  contains a deep minimum at about 88 degrees in the center of mass system, analogous to that previously reported for the scattering of 41-MeV alpha particles by  $O^{16}$ . Alpha groups which correspond to excitation of levels in  $N^{15}$  at 5.276, 6.328, 7.16, 7.31, and 7.57 MeV (as well as others) have been observed, and their angular distributions determined. Preliminary analysis of the data indicates that the levels at 5.276 and 7.57 MeV are formed by the coupling of the odd  $p_{1/2}$  nucleon to a core excitation whose description is similar to that which gives rise to the 6.13-MeV level (3-) in  $O^{16}$ . Analogously, the 7.31-MeV level in  $N^{15}$  appears to be formed by the coupling of the  $p_{1/2}$  nucleon to a core excitation whose description is similar to that which gives rise to the 7.12-MeV level (1-) in  $O^{16}$ .

Proceeding Paris Conference  
on Nuclear Physics  
July 2-8, 1964

UCRL-11378

UNIVERSITY OF CALIFORNIA  
Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

SEARCH FOR CORE-EXCITED STATES IN  $N^{15}$

B. G. Harvey, J. R. Meriwether, A. Bussiere de Nercy  
and D. J. Horen

April 1964

B. G. Harvey, J. R. Meriwether, and A. Bussiere de Nercy  
Univ. of Calif., Lawrence Radiation Laboratory, Berkeley

and

D. J. Horen

U.S. Naval Radiological Defense Laboratory

The interpretation of the level structures of odd-A nuclei in terms of the weak coupling of the odd nucleon to collective excitations of the even-even core has been the subject of a number of theoretical and experimental investigations. In order to test the ability of this model to interpret some of the levels in light nuclei, we have studied the scattering of 40.2-MeV alpha particles by  $N^{15}$  and  $O^{16}$ .  $N^{15}$  was chosen because 1) it might be considered as having one hole in a doubly closed shell, 2) it has a ground state spin of  $1/2$ , and 3) with the exception of the two levels at 5.276 and 5.304 MeV, the energy spacings of the first few levels are comparable or greater than the energy resolution obtainable with our gas target system. In addition, previous investigators have suggested that the 6.13-MeV ( $3^-$ ) level in  $O^{16}$  can be described in collective terms. Hence, if the core excitation model is valid for  $N^{15}$ , one would expect to observe two states with spins and parities  $5/2^+$  and  $7/2^+$  whose "center of gravity" lies at an energy of about 6.13 MeV. The two levels in  $N^{15}$  at 5.276 MeV ( $5/2^+$ ) and 7.57 MeV ( $7/2^+$  probable) appeared as likely candidates.

Gaseous targets of nitrogen enriched to 99.7 per cent in  $N^{15}$  and natural oxygen, contained in a 7.6 cm diameter cell at a pressure of about 10 cm Hg, were bombarded with 40.2 MeV alpha particles from the Berkeley 224-cm variable energy cyclotron. A lithium-drifted silicon detector was used to measure angular distributions of the scattered

particles out to center of mass angles of about  $105^\circ$  and  $70^\circ$  for  $N^{15}$  and  $O^{16}$ , respectively. The overall resolution, including the angular resolution of the gas target system, varied from about 100 to 250 keV.

The angular distributions for excitation of the levels in  $N^{15}$  at energies of 5.276 (5/2+), 7.31 (3/2+), and 7.57 (7/2+) MeV are shown in Fig. 1. For comparison, the distributions for excitation of the levels in  $O^{16}$  at energies of 6.13 (3-) and 7.12 (1-) MeV are also shown. The deep minimum in the elastic distribution for  $N^{15}$  is similar to that reported by Yavin and Farwell [1] for the scattering of 41-MeV alpha particles by  $O^{16}$ . The angular distributions for excitation of the 5.276-MeV (5/2+) and 7.57-MeV (7/2+) levels in  $N^{15}$  are in phase with the elastic group, as well as that which leads to excitation of the 3- level in  $O^{16}$  at an energy of 6.13 MeV. Except for an as yet unexplained systematic error,<sup>†</sup> the sum of the differential cross sections for excitation of these two states in  $N^{15}$  is equal to that for excitation of the 6.13-MeV level in  $O^{16}$ . These results are indicative that the 5.276- and 7.57-MeV levels in  $N^{15}$  arise from the coupling of the odd  $p_{1/2}$  nucleon to a core excitation whose description is similar to that which gives rise to the 6.13-MeV level in  $O^{16}$ .

It is interesting to note that the angular distribution for excitation of the 7.31-MeV level (3/2+) in  $N^{15}$  is similar to that for excitation of the 7.12-MeV (1-) level in  $O^{16}$ . In addition, the differential cross section (after being increased by 20 per cent<sup>‡</sup>) is about two-thirds that for exciting the 7.12-MeV level in  $O^{16}$ . Hence, it is tempting to consider the 7.31-MeV level in  $N^{15}$  as one member of a doublet which arises from the coupling of the odd  $p_{1/2}$  nucleon to a 1- core excitation.

A more detailed discussion of these results will be presented elsewhere.

## NOTES AND REFERENCES

† On leave from Laboratoire Joliot-Curie, ORSAY, FRANCE.

\* Our measured  $N^{15}$  elastic cross section was about 20 per cent lower than that for  $O^{16}$ .

[1] A. I. Yavin and G. W. Farwell, Nucl. Phys. 12, 1 (1959).



## FIGURE CAPTION

Fig. 1. Angular distributions for scattering of 40.2-MeV alpha particles which lead to excitation of levels in  $N^{15}$  at energies of 5.27 (5/2+), 7.31 (3/2+) and 7.57 (7/2+) MeV, and in  $O^{16}$  at energies of 6.13 (3-) and 7.12 (1-) MeV. The absolute cross sections for the  $N^{15}$  data are probably 20 per cent too low. (See footnote to text.)




Fig. 1.

MUB-3167

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

