

Lawrence Berkeley National Laboratory

Recent Work

Title

POMERANCHUK REPULSION AND RESONANCE NARROWING

Permalink

<https://escholarship.org/uc/item/0s20g64g>

Author

Chew, Geoffrey F.

Publication Date

1965-06-30

University of California

Ernest O. Lawrence
Radiation Laboratory

POMERANCHUK REPULSION AND RESONANCE NARROWING

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

POMERANCHUK REPULSION AND RESONANCE NARROWING

Geoffrey F. Chew

June 30, 1965

POMERANCHUK REPULSION AND RESONANCE NARROWING*

Geoffrey F. Chew

Department of Physics and Lawrence Radiation Laboratory
University of California, Berkeley, California

June 30, 1965

ABSTRACT

The contribution of the P (Pomeranchuk) and P' trajectories to the generalized two-particle (low energy) potential is shown to be repulsive and effectively of long range. A rough expression for the P potential is given in terms of the high-energy total cross section and associated diffraction peak. It is argued that Pomeranchuk repulsion represents the many-particle channels that dominate high energies and that have an important narrowing effect on resonance widths even though these channels are closed in the low-energy resonance region.

I. INTRODUCTION

In a recent paper there were discussed certain consequences of employing Regge poles rather than fixed J poles as the source of two-particle generalized potentials.¹ An important omission in that paper was an estimate of potentials arising from trajectories for which the first physical J value fails to have an associated pole of the S matrix. Two well-established trajectories are of this type, the so-called P (Pomeranchuk) and P' trajectories where the first associated particles have $J = 2$,² whereas the first physical angular momentum value is $J = 0$. The purpose of the present paper is to show that the $J = 0$ components of the P and P' Regge potentials are repulsive and effectively of long range. They may constitute the major bootstrap component, so far overlooked, that tends to make resonances narrow.

Bootstrap calculations of low-baryon-number particles on the basis of fixed-spin potentials have always yielded larger widths than experimentally observed.³ It is well known from the dynamics of particles with large baryon number (classical nuclear physics) that the proliferation of many-body channels, open at high energies, systematically narrows the widths of low-energy resonances for which these channels are closed. No estimates have heretofore been given of this effect for particles of low baryon number, but the Reggeized strip approximation⁴ includes the high-energy inelastic effect and therefore should manifest the narrowing tendency.

In the new form of strip approximation the generalized two-body potential is represented as a sum over contributions from the leading Regge trajectories of crossed reactions. Reference 1 shows that when the leading physical J value on the trajectory has an associated physical particle, one may associate the potential in the conventional manner with "exchange" of this particle, although there is an important form factor which reduces the strength and extends the range--relative to a fixed-spin (elementary) particle potential. A small part of the P and P' potentials may be associated in such a sense with exchange of the $J = 2$ $f(1250)$ and $f'(1525)$ particles, but the major component belongs to $J = 0$ -- where no particles exist. We suggest that physically this latter component represents the aforementioned dynamical effect of many-particle channels, closed inside the strip where the potential is to be employed, but open above the strip boundary.

Why is such an identification plausible? First of all, the P and P' trajectories account for most of the total cross section in the high-energy region where multiple production dominates.⁵ Second, as we shall see, the $J = 0$ component of the P and P' potentials is always repulsive and of a range--corresponding to the forward peaks of high energy diffraction scattering--that is relatively long. When such a long-range repulsion is added to a shorter range attraction from "ordinary" particle-exchange, one has the dynamical situation favorable to narrow resonances.⁶

II. THE $J = 0$ COMPONENT OF THE POMERANCHUK POTENTIAL

In Reference 1 it was explained that inside the s strip one may make a Legendre polynomial expansion in z_t of the s -reaction potential associated with the i th Regge pole communicating with the t reaction. Since the Pomernanchuk trajectory is of even signature, we have

$$V_P^S(t,s) = \sum_{J \text{ even}} (2J + 1) V_J^P(t) P_J(z_t) \quad , \quad (\text{II:1})$$

where

$$z_t(s,t) = \frac{s + q_a^2(t) + q_b^2(t)}{2q_a(t)q_b(t)} \quad , \quad (\text{II:2})$$

$$q_a^2(t) = \frac{t}{4} - m_a^2 \quad , \quad q_b^2(t) = \frac{t}{4} - m_b^2 \quad , \quad (\text{II:3})$$

if the s reaction connects channels with particle masses m_a and m_b . It should suffice for our qualitative discussion here, as it did in Reference 1, to employ the Khuri-Jones formula for $V_J^P(t)$:

$$V_J^P(t) = \beta_P [q_a(t)q_b(t)]^{\alpha_P(t)} \frac{\gamma_P(t)}{J - \alpha_P(t)} e^{-[J - \alpha_P(t)]\xi_1(t)} \quad , \quad (\text{II:4})$$

where β_P is a crossing matrix element (always positive for the Pomernanchuk pole), $\gamma_P(t)$ is the reduced residue (also positive

near $t = 0$), and $\alpha_p(t)$ is the Pomeranchuk trajectory. The function $\xi_1(t)$ is given by

$$\xi_1(t) = \log [z_1(t) + (z_1^2(t) - 1)^{1/2}] \quad , \quad (\text{II:5})$$

where $z_1(t) = z(s_1, t)$, s_1 being the strip width, that is, the lowest energy at which the imaginary part of the full amplitude can be approximated by the imaginary part of the potential. It appears experimentally that $s_1 \gtrsim 4 \text{ GeV}^2$.

The qualitative discussion of Reference 1 may be applied to $V_{J=2}^P(t)$, associating this force component with $f(1250)$ exchange, although the damping here with respect to elementary particle exchange is severe. Our rough estimate would give a reduction at $t = 0$ by a factor $\approx e^{-2.7}$, almost one order of magnitude, so the $J = 2$ component of the Pomeranchuk potential is relatively minor, although attractive (positive). The $J = 0$ component, on the other hand, is, for $|t| \ll s_1$,

$$V_{J=0}^P(t) = -\beta_P \alpha_P^{-1}(t) \gamma_P(t) s_1^{\alpha_P(t)} \quad , \quad (\text{II:6})$$

strongly repulsive. The result for P' is similar. One may usefully compare (II:6) to the high-energy limit of the imaginary part of the amplitude--which is the same as the high-energy limit of the imaginary part of the Pomeranchuk potential:



$$\text{Im } V_P^s(t,s) \xrightarrow{s \rightarrow \infty} \beta_P \left[\frac{\sqrt{\pi}}{2} (2\alpha_P(t)+1) \frac{\Gamma(\alpha_P(t) + 1/2)}{\Gamma(\alpha_P(t) + 1)} \right] \gamma_P(t) s^{\alpha_P(t)} \quad (\text{II:7})$$

Observe that for s not enormously larger than s_1 the t dependence of the two forms is similar. Thus the "shape" of the Pomeranchuk potential is essentially that of the high-energy diffraction peak. Using the optical theorem,

$$\sigma^{\text{tot}}(s) \underset{s \gg s_1}{\approx} \frac{16\pi}{s} \text{Im } V_P^s(t=0, s) \quad (\text{II:8})$$

together with the fact that $\alpha_P(0) = 1$, we may establish the normalization to be

$$V_{J=0}^P(t=0) \approx -\frac{s_1}{24\pi^2} \sigma^{\text{tot}}(\infty) \quad (\text{II:9})$$

Had we used the Chew-Jones expression for the Regge formula^{1,4} rather than the Khuri-Jones expression, we should have found in (II:9) a coefficient $-s_1/16\pi^2$, corresponding to a slightly different significance for the parameter s_1 . Since actual dynamical calculations are more likely to be based on the Chew-Jones expression, we shall use this latter normalization in what follows. (The arguments to be made here are only qualitative, so a factor of 3/2 is of no consequence.)

III. AN APPARENT CONTRADICTION

Estimating the t -discontinuity (or imaginary part) of (II:6), one finds it negative in the region between the 2π threshold and the mass squared of $f(1250)$.⁸ Since the t discontinuity of any t -reaction partial-wave elastic amplitude must be positive, a doubt arises about the correctness of (II:6).

In fact, Chew and Teplitz⁹ proposed a technique for evaluation of the potential which precludes a negative result for the potential carrying the vacuum quantum numbers. The reasoning of these authors, however, depended on the neglect of double spectral functions throughout the "corner" regions where both variables (s and t) are inside their respective strips. This is equivalent to assuming that inside the t strip the entire t discontinuity is contained in the potential for the s reaction.

Such is, of course, not strictly the case, and if one asks where (in t) the discontinuity of (II:6) becomes large, one sees that it is in the region where $\text{Im } \alpha_p(t)$ is large, that is, the upper portion of the t strip above the mass squared of $f(1250)$. In view of the relatively narrow width of the f we can be sure that $\text{Im } \alpha_p(t)$ remains small for $t < m_f^2$.¹⁰ Now, in the upper portion of the t strip (inside the s strip) there may be substantial components of the Mandelstam double spectral function arising from iteration of lower t components in the potential. This double spectral function contributes to the total t discontinuity but is excluded (by definition)

from the potential. Were the double spectral function sufficiently large it could produce the required positive sign for the complete t discontinuity, even though the potential (II:6) may be negative.

Towards the lower edge of the t strip the potential must dominate the t discontinuity, so (II:6) cannot there be a good approximation to the complete (vacuum-like) potential. Here the procedure recommended by Chew and Teplitz seems appropriate in order to include the effect of secondary trajectories and "background."

Notice that our conjectured mechanism for removing the contradiction between (II:6) and the positive-definiteness requirement, through the double spectral function, implies the inadequacy of approximating the left-hand discontinuities in an N/D calculation by the discontinuities of the potential. This is perhaps not surprising if one recalls that this latter approximation has been especially deficient in handling strongly repulsive forces.¹¹

A lesser paradox is the circumstance that the "range" of the Pomeranchuk potential (II:6), as measured by its logarithmic derivative at $t = 0$, is longer than would be given by a dispersion-relation estimate based on the region of $t(>m_p^2)$ where the imaginary part becomes large. For pion-pion scattering, as an example, the inverse logarithmic derivative with respect to t of the diffraction amplitude (II:7) at $t = 0$ and $s \approx s_1$ is $\approx 0.5 \text{ GeV}^2$,¹² while $m_p^2 = 1.6 \text{ GeV}^2$. The explanation here is that the imaginary part of $V_{J=0}^P(t)$ oscillates when the imaginary part of $\alpha_p(t)$ becomes large, leading to cancellations in the dispersion integral, so the dependence on t near $t = 0$ may

be steeper than given by the elementary estimate, which tacitly assumes an absence of cancellations. This circumstance means that Pomeranchuk repulsion even while behaving dynamically like a long-range force, does not correspond to a "nearby" left-hand singularity in partial-wave amplitudes. It is a superposition of distant singularities on both right (outside the strip) and left, in which the oscillatory character of the discontinuity is an essential feature. To represent such an effect in N/D models by a few phenomenological poles on the left is probably hopeless.

IV. ESTIMATE OF THE IMPORTANCE OF POMERANCHUK REPULSION

Let us now examine for a much studied example, the $I = 1$ $\pi\pi$ channel, the relative importance of the potentials associated with the P and ρ trajectories, the latter being the only one usually considered for this system.

In our previous paper¹ we have roughly estimated the ρ potential as

$$V_{\pi\pi, I=1}^{\rho}(t) = 3(1 + s/2q_t^2)V_1^{\rho}(t) \quad , \quad (IV:1)$$

where, for $|t| \ll s_1$,

$$V_1^{\rho}(t) \approx q_t^2 \frac{4\Gamma_{\rho}/m_{\rho}}{m_{\rho}^2 - t} e^{-2\alpha'_{\rho}(m_{\rho}^2 - t)} \quad , \quad (IV:2)$$

the effective crossing matrix element here being equal to $1/2$.¹³

The potential is attractive, to be compared to our estimate above of the repulsive Pomeranchuk potential:

$$V_{\pi\pi, I=1}^P(t) \approx -\frac{s_1}{8\pi^2} \sigma_{\pi\pi}^{\text{tot}}(\infty) \frac{\text{Im } A_{\pi\pi}(s_1, t)}{\text{Im } A_{\pi\pi}(s_1, 0)} \quad . \quad (IV:3)$$

Although the detailed shape of the high-energy $\pi\pi$ forward diffraction amplitude is not known, it should suffice here to represent it by a simple exponential of the above-mentioned width 0.5 GeV^2 .

The value of $\sigma^{\text{tot}}(0)$ is taken as 10 mb ,¹² leading to

$$V_{\pi\pi, I=1}^P(t) \approx -0.3 s_1 e^{2t}, \quad (\text{IV:4})$$

where s_1 and t are to be evaluated in units of GeV^2 . For the ρ potential, using a width $\Gamma_\rho = 110 \text{ MeV}$, a mass $m_\rho = 0.77 \text{ GeV}$, and a trajectory slope $\alpha'_\rho = 0.5 m_\rho^{-2}$, we have

$$V_{\pi\pi, I=1}^\rho(t) \approx 1.1 \left(\frac{t}{4} - m_\pi^2 + \frac{s}{2} \right) \frac{e^{1.7t}}{1-1.7t}. \quad (\text{IV:5})$$

Comparing (IV:4) and (IV:5), one should notice two points: (a) The t dependence of the two potentials is not very different, but the ρ potential has a major component increasing linearly with s , while the Pomeranchuk potential is independent of s . (b) In the lower half of the strip, where $s < s_1/2$, the Pomeranchuk repulsion is entirely comparable in magnitude to the ρ attraction.

The s -increasing aspect of the ρ potential means that in N/D dynamics this component, acting like a very short-range attractive force, tends to dominate the denominator function and thus to control the existence and location of resonances in the amplitude. On the other hand, the width of a resonance (resonances are expected to occur in the lower half of the strip) is proportional to the numerator function at the resonance energy--which is sensitive to the value of the potential in this low-energy region (the "long-range force").

Thus a drastic reduction of the potential in the resonance region should lead to an important resonance narrowing effect.

It has already been remarked that with such a strong repulsion one may not employ the N/D device of replacing left-hand partial-wave cuts by the cuts of the potential. It will be necessary to perform at least a few steps of the Mandelstam iteration in order to achieve a believable dynamical result. The results of such calculations will, one hopes, be reported at a later time.

V. CONCLUSION

The presence of Pomeranchuk repulsion in all two-particle channels may explain why resonance widths have so uniformly been overestimated in non-Reggeized bootstrap calculations. At the same time, certain aspects of the qualitative estimates heretofore given of the attractive forces essential to forming bound states and resonances are not invalidated by Reggeization. There remains a correlation with the concept of particle exchange, and the sign (attraction or repulsion) generally survives. We can understand in this way the success of crude bootstrap arguments that use crossing matrices and almost nothing more. The estimates given in this paper and in Reference 1 indicate, however, that to achieve even semiquantitative accuracy in the dynamics it will be necessary to employ Regge potentials together with the Mandelstam iteration or the equivalent thereto.

ACKNOWLEDGMENTS

I am deeply indebted to Dr. P. D. B. Collins and Dr. V. L. Teplitz for intensive discussions of the ideas described here. Their numerical calculations provided an invaluable stimulant.

REFERENCES

- * Work done under auspices of the U.S. Atomic Energy Commission.
1. G. F. Chew, Approximation of Regge Potentials Through Form Factors (Lawrence Radiation Laboratory Report UCRL-16101, May 1965, to be published in the Supplement to the Progress of Theoretical Physics.
 2. Presumed to be the $f(1250)$ and the $f'(1525)$, the latter reported by Barnes, et al., Phys. Rev. Letters (1965).
 3. See, for example, J. R. Fulco, G. L. Shaw, and D. Y. Wong, Phys. Rev. 137, B1242(1965).
 4. G. F. Chew, Phys. Rev. 129, 2363 (1963); G. F. Chew and C. E. Jones, Phys. Rev. 135, B208 (1964).
 5. See R. Phillips and W. Rarita (Lawrence Radiation Laboratory Report UCRL-16033, April 1965), submitted to Phys. Rev., for a recent review of all high-energy πN and KN experiments and for additional references.
 6. One may say that the system becomes "trapped" inside the repulsive barrier and takes a long time to find its way out. A long lifetime means a narrow width.
 7. This result is confirmed by numerical calculations of Collins and Teplitz based on the Chew-Jones potential (private communication, Berkeley, (1965)).
 8. The essential point is that $\text{Im } \alpha_p(t)$ is positive.
 9. G. F. Chew and V. L. Teplitz, Phys. Rev. 137, B139 (1965).
 10. For $t = m_f^2$, $\text{Im } \alpha_p = \Gamma_f m_f \left(\frac{d \text{Re } \alpha_p}{dt} \right)_{t = m_f^2}$, and the trajectory

slope appears to be less than 1 GeV^{-2} . Thus for $\Gamma_f \approx 100 \text{ MeV}$,

$\text{Im } \alpha_p \lesssim 0.1$ at $t = m_f^2$.

11. J. G. Bjorken and A. Goldberg, Nuovo Cimento 16, 539 (1960).
12. G. F. Chew and V. L. Teplitz, Phys. Rev. 136, B1154 (1964).
13. In the $\pi\pi$ problem, both crossed reactions contain the poles in question, so the total potential treated in this section is twice that from the t reaction above.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

