

Lawrence Berkeley National Laboratory

Recent Work

Title

EXPERIMENTAL COMPARISON BETWEEN ON- AND OFF-MASS-SHELL INELASTICITIES IN π -p SCATTERING. II

Permalink

<https://escholarship.org/uc/item/0vn3s9x3>

Authors

Colton, Eugene
Schlein, Peter E.
Ma, Z. Ming
et al.

Publication Date

1970-06-01

c.2

EXPERIMENTAL COMPARISON BETWEEN ON- AND
OFF-MASS-SHELL INELASTICITIES IN π^-p SCATTERING. II

RECEIVED
LAWRENCE
RADIATION LABORATORY

JUN 25 1970

LIBRARY AND
DOCUMENTS SECTION

Eugene Colton, Peter E. Schlein
Z. Ming Ma and Gerald A. Smith

June 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-19771
247

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

EXPERIMENTAL COMPARISON BETWEEN ON- AND OFF-MASS-SHELL

INELASTICITIES IN π^-p SCATTERING. II *

Eugene Colton

Lawrence Radiation Laboratory
University of California
Berkeley, California 94720

and

Peter E. Schlein⁺CERN
Geneva, Switzerland

and

Z. Ming Ma and Gerald A. Smith

Physics Department
Michigan State University
East Lansing, Michigan 48823

ABSTRACT

It is shown that the experimental cross-section ratio of the reactions $pp \rightarrow \Delta^{++}$ (neutrals) and $pp \rightarrow \Delta^{++}(\pi^-p)$ (at 6.6 GeV/c) for low momentum transfer to the Δ^{++} ($|t| < 0.2 \text{ GeV}^2$) is approximately the same as the cross-section ratio $\sigma(\pi^-p \rightarrow \text{neutrals})/\sigma(\pi^-p \rightarrow \pi^-p)$ for π^-p effective mass less than 1.56 GeV. The disagreement observed between the ratios for masses greater than 1.56 GeV can be explained by increased values of t_{\min} and non- Δ^{++} background in the pp reactions. These results imply that pion exchange dominates $pp \rightarrow \Delta^{++}$ (neutrals) for low momentum transfer to the Δ^{++} .

In a recent analysis of the reactions

$$pp \rightarrow \Delta^{++}(\pi^- p) \quad (1a)$$

$$\Delta^{++}(\pi^- \pi^+ n) \quad (1b)$$

$$\Delta^{++}(\pi^- \pi^0 p) \quad (1c)$$

at $6.6 \text{ GeV}/c^1$ we showed that the relative yields of the bracketed systems at low momentum transfer ($|t_{p\Delta}| < 0.2 \text{ GeV}^2$) as a function of effective mass of that system are very nearly the same as the relative cross sections of the three on-mass-shell $\pi^- p$ reactions

$$\pi^- p \rightarrow \pi^- p \quad (2a)$$

$$\pi^- p \rightarrow \pi^- \pi^+ n \quad (2b)$$

$$\pi^- p \rightarrow \pi^- \pi^0 p \quad (2c)$$

This agreement suggests that the bracketed systems in reactions (1a)-(1c) result from the scattering of an incident proton on a virtual pion from the other proton when it is in the virtual state $p \rightarrow \Delta^{++} \pi^-$. It furthermore suggests that the off-mass-shell corrections to the pole equation descriptions of the three reactions (1a)-(1c) are very similar. Otherwise the physical region comparison of the three reactions could not agree so well.

In a subsequent analysis of the reactions²

$$pp \rightarrow n(\pi^+ p) \quad (3a)$$

$$\rightarrow n(\pi^+ p \pi^+ \pi^-) \quad (3b)$$

we compared the relative yields of the bracketed systems in reactions (3a) and (3b) and the relative cross sections for

$$\pi^+ p \rightarrow \pi^+ p \quad (4a)$$

$$\pi^+ p \rightarrow \pi^+ p \pi^+ \pi^- \quad (4b)$$

In this case the comparison showed that the experimental ratios [of (3b)

to (3a)] systematically exceeded the π^+p on-shell cross section ratios by $\sim 25 - 40\%$. However, the difference can be explained solely by a $\Delta^-(1238)$ background contribution to reaction (3b).

We present in this note a further comparison (also at 6.6 GeV/c) which is a followup to the analysis of Ref. 1. The relative yields of the low momentum transfer components ($|t| < 0.2 \text{ GeV}^2$) of the reactions

$$pp \rightarrow \Delta^{++}(\text{neutrals}) \quad (5a)$$

$$pp \rightarrow \Delta^{++}(\pi^- p) \quad (5b)$$

are compared to the relative cross sections for

$$\pi^- p \rightarrow \text{neutrals} \quad (6a)$$

$$\pi^- p \rightarrow \pi^- p \quad (6b)$$

The data of reaction (5a) consists of a 2.34 $\mu\text{b}/\text{event}$ sample of 4026 unfitted two-prong events with $M_{p\pi^+} < 1.5 \text{ GeV}$ and forward neutrals in the c.m. For these events the ionization confidence level is required to favor a $p\pi^+$ interpretation instead of pp for the two outgoing charged tracks. The data for reaction (5b), as before,¹ consists of a 0.36 $\mu\text{b}/\text{event}$ sample of 7514 $pp \rightarrow p\pi^+\pi^-$ events. The data for reactions (5a) and (5b) are obtained by applying the simultaneous cuts³

$$\begin{aligned} 1.14 < M_{p\pi^+} < 1.30 \text{ GeV} \\ |t| < 0.2 \text{ GeV}^2 \end{aligned} \quad (7)$$

We take $|t|$ to be the lower of the two possible momentum transfers squared from the beam or target proton to the Δ^{++} and restrict the discussion henceforth to the events passing the cuts (7).

The effective mass spectra of the bracketed systems of reactions (5a) and (5b) are presented in Figs. 1(a) and 1(b), respectively. The smooth curve drawn in Fig. 1(a) represents an estimate of the background due to the reaction $pp \rightarrow p\pi^+n$. The intercept at 1166 MeV is the point above which no $pp \rightarrow p\pi^+n$ constrained fits are observed. In Fig. 2 we show the ratio of the cross sections of reactions (5a) to (5b) as a

function of mass of the bracketed systems (M_x). The ratios have been corrected for $pp \rightarrow p\pi^+n$ background in reaction (5a) for $M_x < 1.166$ GeV. The cross-hatched band shows the known on-mass-shell ratio of reactions (6a) to (6b).⁴ The experimental and on-mass-shell ratios agree quite well below 1560 MeV except for the first point from 1080 - 1120 MeV (which may just be due to poor statistics or an overestimate in $p\pi^+n$ background). Above 1560 MeV the experimental ratios exceed the on-shell ratios by a factor of two or three.⁵ This disagreement at large M_x may just be due to the use of events with generally larger values of $|t|$ (because t_{\min} increases with M_x) or simply non- Δ^{++} background in the peripheral data of reaction (5a).

We have looked at the $p\pi^+$ mass spectra in reactions (5a) and (5b) for several regions of the mass of the bracketed systems (X) in order to see if the difference in ratios for large M_x is due to non- Δ^{++} background in the data of reaction (5a). For $M_x < 1500$ MeV the non- Δ^{++} background appears to be small. However, the phase space available to the $p\pi^+$ mass for large M_x is so small ($\max M_{p\pi^+} < 1.4$ GeV) that one cannot judge the percentage of non- Δ^{++} events in this sample. Assuming that non- Δ^{++} background, if present, is weakest in the central region of the Δ^{++} we have again calculated the experimental cross-section ratios of reactions (5a) to (5b) using peripheral ($|t| < 0.2$ GeV²) data with a narrower Δ^{++} cut ($1.18 < M_{p\pi^+} < 1.26$ GeV). These ratios are identical to those displayed in Fig. 2 for $M_x < 1.56$ GeV except for the larger errors (less events are utilized in this case). The ratios for the 5 M_x bins with $M_x > 1.56$ GeV are represented by the points with dashed errors in Fig. 2. For $M_x > 1.56$ GeV the ratios

calculated using a narrow Δ^{++} cut are in better agreement with the on-shell ratios than those experimental ratios obtained with data satisfying both restrictions of Eq. (7).

Experimental determinations of on-shell inelastic cross sections ($x\pi \rightarrow y, y \neq x\pi$) in processes of the type $xp \rightarrow yn$ or $xp \rightarrow y\Delta^{++}$ can be obtained by extrapolation of the appropriate Chew-Low distribution to the pion exchange pole. Our results show, as before,^{1,2} that the form of the extrapolation function used should have approximately the same form used for elastic $x\pi$ scattering.


FOOTNOTES AND REFERENCES

- * Supported in part by the United States Atomic Energy Commission (LRL, UCLA) and the National Science Foundation (MSU).
- + John Simon Guggenheim fellow, on leave from University of California, Los Angeles (1969-1970).
- 1. Eugene Colton, Peter E. Schlein, Eugene Gellert, and Gerald A. Smith, *Phys. Rev. Letters* 21, 1548 (1968).
- 2. Eugene Colton and Peter E. Schlein, *Phys. Rev.* D,1, 373 (1970).
- 3. In the case of the $pp\pi^+\pi^-$ events, if both $M_{p\pi^+}$ combinations fall within the Δ^{++} band and both have $|t| < 0.2 \text{ GeV}^2$ we use that combination with the smallest $|t|$. This procedure tends to distort the $M_{p\pi^-}$ distribution slightly by depleting combinations at intermediate and high $p\pi^-$ masses and correspondingly adding the other low t combinations to the low end of the $M_{p\pi^-}$ spectrum.
- 4. The π^-p elastic cross sections were constructed from the CERN phase shifts, see, e.g., A. Donnachie, R. G. Kirsopp, and C. Lovelace, CERN TH 838, Addendum; 1967 (unpublished). For a summary of $\pi^-p \rightarrow$ neutrals cross sections see, e.g., C. B. Chiu, R. D. Eandi, A. C. Helmholtz, R. W. Kenney, B. J. Moyer, J. A. Poirier, W. B. Richards, R. J. Cence, V. Z. Peterson, N. K. Sehgal, and V. J. Stenger, *Phys. Rev.* 156, 1415 (1967).
- 5. W. E. Ellis, R. R. Kinsey, T. W. Morris, and R. S. Panvini, BNL 14426, 1970 (unpublished). These authors also report that their experimental ratios exceed the on-mass-shell ratios by roughly a factor of two (see Fig. 5(a)).

FIGURE CAPTIONS


Fig. 1. Effective mass distributions of X in $pp \rightarrow \Delta^{++} X$ for $|t| < 0.2$ GeV^2 at 6.6 GeV/c. The Δ^{++} selection is 1.14 - 1.30 GeV;
(a) X = neutrals. (b) X = $p\pi^-$.

Fig. 2. Cross-section ratio $\sigma[pp \rightarrow \Delta^{++}(\text{neutrals})]/\sigma[pp \rightarrow \Delta^{++}(p\pi^-)]$ plotted as a function of the mass of the non- Δ^{++} final state system, for momentum transfer to the Δ^{++} $|t| < 0.2$ GeV^2 . The dashed points in the uppermost five mass bins represent the ratios calculated using $1.18 < M_{p\pi^+} < 1.26$ GeV for the Δ^{++} selection. The cross-hatched band is the known (Ref. 4) on-shell ratio $\sigma(\pi^-p \rightarrow \text{neutrals})/\sigma(\pi^-p \rightarrow \pi^-p)$.



XBL 704-2730

Fig. 1



XBL704-2729

Fig. 2

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720