

Lawrence Berkeley National Laboratory

Recent Work

Title

Experimental and Theoretical Characterization of Multifragment Events

Permalink

<https://escholarship.org/uc/item/0x813865>

Authors

Moretto, L.G.
Roussel-Chomaz, P.
Colonna, M.
et al.

Publication Date

1991-06-01



Lawrence Berkeley Laboratory


UNIVERSITY OF CALIFORNIA

Presented at the 7th Adriatic International Conference on Nuclear Physics: Heavy Ion Physics—Today and Tomorrow, Island of Brioni, Yugoslavia, May 27–June 1, 1991, and to be published in the Proceedings

Experimental and Theoretical Characterization of Multifragment Events

L.G. Moretto, P. Roussel-Chomaz, M. Colonna, N. Colonna, and G.J. Wozniak

June 1991



LOAN COPY |
 Circulates |
 for 4 weeks | Bldg. 50 Library.
 Copy 2
 LBL-30931

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Experimental and Theoretical Characterization of Multifragment Events

L. Moretto, P. Roussel-Chomaz, M. Colonna,
N. Colonna, and G.J. Wozniak

Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

June 1991

This report has been reproduced directly from the best available copy.

This work was supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of Nuclear Physics, of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098

Experimental and Theoretical Characterization of Multifragment Events

L.G. Moretto, P. Roussel-Chomaz^{a)}, M. Colonna^{b)}, N. Colonna, and G.J. Wozniak

Nuclear Science Division, Lawrence Berkeley Laboratory

1 Cyclotron Road, Berkeley, CA94720

Abstract

Multifragment events are shown to be associated with specific sources characterized by their mass and excitation energy through the incomplete fusion model. Excitation functions for the different multifragment decay channels are found to be almost independent of the system and the incident energy. Preliminary comparisons of the data with dynamical calculations followed by statistical decay calculations are discussed.

Introduction

At intermediate energies ($E/A \leq 50$ MeV), two sources of complex fragments have been identified [1-4]: a fast, non-equilibrium source which produces light fragments at forward angles in normal kinematics and an equilibrium component, originating from the statistical decay of a compound nucleus via binary processes which cover the entire range of mass asymmetry in the exit channel. These compound nuclei are formed in complete or incomplete fusion reactions, depending on the incident energy and on the mass asymmetry of the system.

Within the incomplete fusion picture, it is possible to correlate the mass and excitation energy of the product nucleus with the degree of fusion by means of the source velocity. In reverse kinematics, for large impact parameters the nuclei formed are slightly heavier than the projectile and move at slightly lower velocity. As the impact parameter decreases, the projectile picks up more and more mass from the target, the velocity of the compound system decreases and its excitation energy increases. This correlation, clearly observed for the 18 MeV/u $^{139}\text{La} + ^{64}\text{Ni}$ reaction [5], made it possible to study, at one incident energy, the decay properties of hot nuclei over a large excitation energy range.

We have extended this method to ternary, quaternary, etc... events[6]. We have also compared our experimental results to some very preliminary results obtained by coupling a Landau-Vlasov type calculation [7] describing the dynamical stage of the collision with a statistical binary decay code [8,9] used to describe the deexcitation process. These results may help us to understand the role played by the dynamics and by the statistics in these collisions.

^{a)} Permanent address: CEN Saclay, 91191 Gif sur Yvette Cedex France

^{b)} Present address: Universita di Catania, 95129 Catania, Italy

Experimental results

Multifragment emission has been studied in the reactions induced by ^{139}La beams on different targets: ^{12}C , ^{27}Al , ^{40}Ca , ^{51}V , $^{\text{nat}}\text{Cu}$, and ^{139}La . Because of the reverse kinematics, even the heaviest fragments have velocities large enough to be identified easily with simple ΔE - E telescopes. Furthermore the reaction products are focused in a narrow cone around the beam direction; therefore the detection efficiency is good even with a detection system of modest size.

As an overview of the reaction mechanism, Fig. 1 presents for the 2-body events, the correlation between the velocities (normalized to the beam velocity) reconstructed for the source of the fragments by the relation $v_s = \sum m_i v_i / \sum m_i$ and the total charge detected, for six different energies ranging from 18 to 55 MeV/u and four entrance channel asymmetries. The data corresponding to ^{129}Xe beams have been shifted by 3 Z-units to make the comparison easier ($\Delta Z_{\text{La-Xe}}=3$).

The first row corresponds to the most asymmetric system $\text{La/Xe} + ^{12}\text{C}$, which has relatively low available energies in the center of mass system, and presents a very simple pattern. At 18 MeV/u, the source velocity distribution peaks at the value expected for complete fusion, which corresponds to the solid line, and the total charge detected is the total charge of the system. In this case, complete fusion has occurred and only neutrons have been evaporated. When the incident energy increases, the distributions move to higher source velocities and lower total detected charge. The same description holds for the ^{27}Al target. The only difference is that due to the higher excitation energies that can be reached, the evaporation is more extensive and the detected charge is less than that of the primary compound nucleus, even at 18 MeV/u.

The pattern observed for the heavier targets is more complicated. At 18 MeV/u we observe a very nice illustration of the transition from complete fusion ($Z_1 + Z_2 \approx Z_P + Z_T$) to incomplete fusion, with a ridge line going to lower secondary charge when the source velocity increases, which is again what is expected for incomplete fusion in reverse kinematics. As the incident energy increases, and the excitation energy available in the reaction increases, the pattern shifts towards lower Z values because of the evaporation process. At 35 MeV/u the pattern observed is upright, indicating that the system lost by evaporation as many nucleons as it had gained from the fusion with the target. The patterns observed for ^{51}V and $^{\text{nat}}\text{Cu}$ above 35 MeV/u are quite different, with a ridge line going to lower secondary detected charge when the source velocity decreases, which is the opposite to what was observed at lower incident energy.

We shall now consider n-fold events, i.e. events where n fragments are detected in coincidence, with n = 2, 3, 4 and even 5 at 55 MeV/u. Fig. 2 presents the Z distributions for n-fold events for all the systems measured at 40 MeV/u. For the ^{12}C target a narrow peak is observed, but as the mass of target increases, this peak broadens and shifts to lower detected charge. These effects arise from the larger range of mass transfers and from the increase of light particle evaporation due to the larger range of excitation energies available.




Fig. 1: Linear contour plots of the source velocity versus total detected charge for 2-fold coincidence events, for 6 incident energies and 4 different entrance channel asymmetries. The beam energy and the target are indicated in the first row and column, respectively. The total available energy in the c.m. system is indicated in the lower right of each frame. The horizontal lines and the vertical arrows indicate the complete fusion velocity for each system and the projectile charge, respectively.

The tail at low total detected charges also increases with the mass of the target, and this is related to the increase of higher n-fold events where only 2 fragments are detected. The same Z_{total} distribution plotted for 3-fold and 4-fold events presents a peak centered at approximately the same value, but with a reduced tail to low Z_{total} , indicating that most of these events are complete.

In the following we will restrict ourselves to events where the total measured charge is higher than 30, in order to exclude those events where one fragment is clearly missing and to avoid biasing our kinematical reconstructions. Fig. 3 presents the source velocity distributions obtained at 40 MeV/u for all the targets and for the different fragment multiplicities. The observed peak broadens significantly when the mass of the target increases. This width has two different origins: incomplete fusion processes and the broadening from evaporation. This contribution has been estimated with the statistical code Gemini[8, 9]. In the case of the ^{12}C target, the width can be explained almost entirely by light particle evaporation, but in the case of the heavier targets, a wide range of excitation energies contributes effectively to complex fragment emission. For a given target, the requirement of larger multiplicity of complex fragments selects out events with lower source velocities and therefore higher excitation energies. The same trend has been observed with Ne+Au at 60 MeV/u [10].




Fig. 2: Total detected charge for some of the systems measured at 40 MeV/u as a function of the multiplicity in the exit channel.

To study the behavior of hot nuclear systems as their excitation energy increases, excitation functions for the binary, ternary, etc. decay channels have been constructed. More precisely, Fig. 4 presents the evolution of the proportion of n -fold events with respect to the total number of coincidence events as a function of the excitation energy inferred from the source velocity through the incomplete fusion model, for four incident energies. The picture obtained is quite striking. First, at all energies the 3-fold and 4-fold event probabilities (and 5-fold at 55 MeV/u) increase significantly up to excitation energies as high as 8 MeV/u. This energy dependence is a good indication that the relation between E^* and the source velocity is valid and also confirms that the width of the source velocity distribution is only partly due to light particle evaporation (if it were only particle evaporation, the excitation functions would be flat). Second, the increase observed in these excitation functions is essentially the same, for all systems, and at all bombarding energies, suggesting that the branching ratios for binary, ternary, quaternary decays depend mainly upon excitation energy (and angular momentum) and not upon the dynamics of the reaction.

To conclude on these excitation functions, their independence with respect to target-projectile combination and to incident energy suggests a competition between the different decay channels independent of the entrance channel and therefore supports the idea of an intermediate system whose decay properties are mainly determined by its excitation energy and angular momentum.




Fig. 3: Same as Fig. 2 for the source velocities expressed as the ratio of the source to the beam velocity.

La + X , ZTOT>30




Fig. 4: Proportion of 2-3-4-5 fold events as a function of the excitation energy per nucleon for the systems studied at 4 different energies.

Boltzmann-Nordheim-Vlasov calculations

Kinetic equations, like the Landau-Vlasov, Boltzmann-Uehling-Uhlenbeck or Boltzmann-Nordheim-Vlasov equations, have been widely used to simulate the evolution of heavy-ion collisions in the intermediate energy range[11-16]. So far, these transport theories have not been able to reproduce the yields of fragments in the mass range between the projectile and target[16]. On the other hand, statistical approaches, successfully used at low energy, where complex fragments are assumed to be produced in the decay of compound nuclei formed in fusion or incomplete fusion reactions, take minimal account of entrance channel effects and cannot reproduce the non-equilibrium features observed at intermediate energy[17].

The two approaches described above are complementary and can be combined in a framework which incorporates both dynamical evolution and statistical decay. More specifically, we will show that the experimental charge distributions as well as the main features of the coincident events can be reproduced by coupling a dynamical approach describing the formation of the primary fragments, to a statistical stage where these excited primary fragments can undergo any division from particle evaporation to symmetric fission.

The dynamic stage of the collision was simulated by solving the Boltzmann-Nordheim-Vlasov (BNV) equation. The resulting average trajectory in phase space, is followed up to a time called for the sake of convenience, the "relaxation" time. We have chosen the relaxation time as the time when the slope of the emitted nucleon mean energy curve changes, indicating the transition from preequilibrium emission to evaporation from a more equilibrated source.

At the relaxation time, a clustering procedure[15] is used to calculate, for each impact parameter, the primary fragment observables: mass, charge, velocity, angle, excitation energy and angular momentum. In this procedure a cluster is formed by the test particles that satisfy the relation $|r_i - r_j| < D$, where D is set to the minimum value that reproduces the mass of the target and projectile at $t = 0$ ($D = 1.5$ fm). With this method we obtain the most probable distribution of primary fragments with all their properties at the relaxation time as a function of impact parameter.




Fig.5: Comparison of the experimental and calculated charge distribution for the system La + Al at 55 MeV/u. No normalization factor has been applied to the simulation. The statistical errors on experimental data are smaller than the diamonds and the errors related to the extraction procedure are smaller than 20% except above $Z = 35$ where the cross section can be underestimated by up to 60% due to the reduced angular coverage for the very heavy fragments.

Finally, these sources are allowed to undergo sequential binary decays. The deexcitation process has been simulated with the statistical decay code GEMINI[8]. In this code, all decay channels are considered, from light particle emission to symmetric division.

The model described above has been applied to the reaction $^{139}\text{La} + ^{27}\text{Al}$ at 55 MeV/u[18]. For the most central impact parameters ($b = 1, 2$ fm), the BNV calculations predict the occurrence of fusion accompanied by preequilibrium emission. For $b = 3$ fm, the mechanism observed can be characterized as fast fission. Finally the more peripheral reactions lead to a process whose features are very reminiscent of deep inelastic collisions as they are observed at low incident energy. At each impact parameter, preequilibrium emission of light particles is observed. Fig. 5 shows the cross section as a function of atomic number Z for the system under study. Good agreement is observed between the data and the simulation.




Fig. 6 : Comparison of the experimental (solid line) and calculated (dashed line) total charge (left-hand column) and source velocity (right-hand column) distributions for different fragment multiplicities in the case of $^{139}\text{La} + ^{27}\text{Al}$ at 55 MeV/u. The spectra have been normalized to the same maximum.

More severe constraints are imposed by the coincidence events. Coincident events have been separated according to the detected fragment multiplicity. A n -fold event is defined as an event where n fragments with $Z \geq 4$ are detected. Table 1 compares the different branching ratios for the binary, ternary and quaternary channels obtained from the data and the calculation after filtering through the detector acceptance. In both cases, 2-fold events represent the bulk of the detected fragments, and the branching ratios for 3-fold events agree within 20%.

The experimental total charge and source velocity distributions with the corresponding theoretical quantities obtained from the calculations after filtering through our detection efficiency, are shown in Fig. 6. The overall agreement is satisfactory. The peak position in the Z_{tot} distribution for the 2-fold events is well reproduced. For the 3-fold events, the peak position obtained in the calculation is somewhat larger than the data. The shift observed in the Z_{tot} peak position may also be due to an underestimate of the excitation energy deposited in the primary sources. The source velocity distributions (shown Fig. 6) are gated for $Z_{\text{tot}} \geq 30$, in order to reduce to a reasonable level the contamination arising from incompletely detected events and to avoid biasing our kinematical reconstructions. For all types of events, the position of the calculated peak is in good agreement with the data, but the width is underestimated for 3 and 4-fold events.




Fig. 7: Comparison of the linear contour Dalitz plots obtained from the data and from the simulation before and after filtering through the detection efficiency, for the system $^{139}\text{La} + ^{27}\text{Al}$ at 55 MeV/u.

The 3-body decay features are best shown by means of Dalitz plots. In the Dalitz plots presented here, a gate has been applied to the total detected charge ($Z_{\text{tot}} > 30$). Fig. 7 compares the data with the results of the "BNV+Gemini" calculation before and after filtering. It should be stressed that at the end of the dynamical stage the Dalitz plot would be empty, since only either an incomplete fusion residue or a target and projectile remnant are produced. After the deexcitation stage, the simulated Dalitz plot is very similar to the data, with a predominance of events with one heavy and two light fragments. The relative abundance of these events located in the corners of the triangle is considerably reduced after filtering through our detection efficiency, because the heavy fragments are strongly forward peaked and very often do not pass through the filter.

Acknowledgements

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the US Department of Energy under contract DE-AC03-76SF00098.

Table 1 : Comparison of the experimental and calculated proportions of multifragment events. A gate has been set on the total detected charge ($Z_{\text{tot}} \geq 30$).

	2-fold/Total	3-fold/Total	4-fold/Total
Data	0.869	0.122	0.0088
Simulation	0.855	0.144	0.0017

References

- [1] R.J. Charity et al., *Nucl. Phys.* A511, 59 (1990)
- [2] M. Fatyga et al., *Phys. Rev. Lett.* 58, 2527 (1987)
- [3] B. Borderie et al., *Phys. Lett.* B205, 26 (1988)
- [4] E. Plagnol et al., *Phys. Lett.* B221, 11 (1989)
- [5] N. Colonna et al., *Phys. Rev. Lett.* 62, 1833 (1989)
- [6] Y. Blumenfeld et al., *Phys. Rev. Lett.* 66, 576 (1991)
- [7] A. Bonasera et al., *Phys. Lett.* B244, 169 (1990)
- [8] R.J. Charity et al., *Nucl. Phys.* A483, 371 (1988)
- [9] R.J. Charity et al., *Nucl. Phys.* A476, 516 (1988)
- [10] R. Bougault et al., *Phys. Lett.* B232, 291 (1989)
- [11] Ch. Gregoire et al., *Nucl. Phys.* A465, 315 (1987)
- [12] H. Stöcker and W. Greiner, *Phys. Rep.* 137 (1986) 277
- [13] G.F. Bertsch and S. Das Gupta, *Phys. Rep.* 160 (1988) 190 and refs therein
- [14] A. Bonasera, G.F. Burgio, M. Di Toro, *Phys. Lett. B* 221 (1989) 233
A. Bonasera, G. Russo, H.H. Wolter, *Phys. Lett. B* 246 (1990) 337
- [15] A. Bonasera et al., *Phys. Lett. B* 244 (1990) 169
- [16] A. Adorno et al., *Nucl. Phys.* A in press
- [17] D.R. Bowman et al., *Nucl. Phys.* A523 (1991) 386
- [18] P. Roussel-Chomaz et al., in preparation

LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
INFORMATION RESOURCES DEPARTMENT
BERKELEY, CALIFORNIA 94720