

UC Davis

UC Davis Previously Published Works

Title

Heterotrophic nitrogen fixation in oligotrophic tropical marshes: changes after phosphorus addition

Permalink

<https://escholarship.org/uc/item/0xk5x7sm>

Journal

Hydrobiologia: The International Journal of Aquatic Sciences, 627(1)

ISSN

1573-5117

Authors

Černá, Barbora
Rejmánková, Eliška
Snyder, Jenise M.
[et al.](#)

Publication Date

2009-07-01

DOI

10.1007/s10750-009-9715-y

Peer reviewed

Heterotrophic nitrogen fixation in oligotrophic tropical marshes: changes after phosphorus addition

Barbora Černá · Eliška Rejmánková ·
Jenise M. Snyder · Hana Šantrůčková

Received: 24 April 2008 / Revised: 8 January 2009 / Accepted: 14 January 2009 / Published online: 13 February 2009
© The Author(s) 2009. This article is published with open access at Springerlink.com

Abstract In order to determine the impact of nutrient enrichment on phosphorus (P) limited wetlands, we established experimental P additions in marshes throughout northern Belize. P significantly increased macrophyte primary production, which led to the rapid elimination of cyanobacterial mats. The replacement of cyanobacterial mats by macrophytes constrained autotrophic nitrogen (N) fixation, increased the quantity, and changed the quality of organic matter input to the sediments. We predicted that the activity of sediment heterotrophic N fixers will be impacted by these alterations in carbon input. We used the acetylene reduction technique to measure potential (glucose amended) nitrogenase activity (NA) in sediments from controls and treatment plots that have been P enriched for four years and dominated either by *Eleocharis cellulosa*, or *Typha domingensis* for two years. NA in P-enriched plots was 2–3 orders of magnitude higher than NA in controls. NA was positively correlated with the soil reactive P, both total organic and microbial

carbon, live root biomass, and total phospholipid fatty acids (PLFA) as an indicator of active microbial biomass. It was negatively correlated with the concentration of ammonium-N. Path analysis revealed that the indirect effect of P on NA through the root biomass was more important than the direct effect of P. NA of the upper sediment layer was consistently higher in *Eleocharis* than in *Typha* dominated plots, despite the higher litter input by *Typha*. We feel that the higher levels of lignin and phenolics occurring in *Typha* litter, relative to *Eleocharis*, constrained NA in *Typha* plots.

Keywords Heterotrophic nitrogen fixation · *Typha* · *Eleocharis* · Cyanobacteria · Phosphorus · Salinity

Introduction

Biological nitrogen (N) fixation contributes “new” N to the majority of ecosystems, but despite its importance in ecosystem functioning, ecological controls of N fixation are relatively poorly understood (Vitousek et al., 2002). Research on biological N fixation and its role in maintaining soil N fertility in wetland ecosystems has focused on rice fields (Roger & Ladha, 1992; Roger, 1995) and coastal wetlands such as salt marshes and mangroves (Pelegri & Twilley, 1998; Bagwell & Lovell, 2000; Nielsen et al., 2001; Tyler et al., 2003; Moseman, 2007). However, such research is rather limited in freshwater marshes (Bristow, 1973;

Handling editor: Luis Mauricio Bini

B. Černá · H. Šantrůčková
Faculty of Science, University of South Bohemia,
Branišovská 31, České Budějovice 370 05,
Czech Republic

E. Rejmánková (✉) · J. M. Snyder
University of California Davis, One Shields Ave.,
Davis, CA 95616, USA
e-mail: erejmankova@ucdavis.edu

Tjepkema & Evans, 1976; Inglett et al., 2004; Scott et al., 2005).

Herbaceous wetland ecosystems generally lack symbiosis-involving legumes and thus the most important N fixers are N-fixing cyanobacteria and sediment heterotrophic bacteria (Howarth et al., 1988a). Cyanobacterial N fixation has been described for a variety of freshwater and salt marshes, as well as mangroves (Paerl et al., 1994; Rejmánková & Komárková, 2000; Tyler et al., 2003; Inglett et al., 2004; Steppe & Paerl, 2005; Scott et al., 2007; Vargas & Novelo, 2007). The contribution of cyanobacterial N fixation to the ecosystem N economy is relatively high, contributing on average 1–10 (and up to 76) g N m⁻² y⁻¹ (Howarth et al., 1988a).

Heterotrophic N fixation has been reported to be closely associated with plant roots (Boyle & Patriquin, 1981; Bergholz et al., 2001; Nielsen et al., 2001; Moseman, 2007), but the complex interaction between plant roots, root exudates, and microorganisms are poorly understood (Coleman, 2008). Most importantly, though, plants are known to provide a source of available carbon to sediment diazotrophs (Dakora & Drake, 2000; Bürgmann et al., 2005). Heterotrophic N fixation varies greatly in wetland sediment, and the rates are generally about an order of magnitude lower than in cyanobacterial mats (Howarth et al., 1988a). Although the N fixed by diazotrophs may not be immediately available for plant uptake, plants may still benefit from the asymbiotic fixation over time, as nitrogen becomes available through organic N depolymerization and mineralization (Bürgmann et al., 2005).

Across ecosystems, N fixation is controlled by a variety of abiotic (bottom-up) and biotic (top-down) factors (Vitousek et al., 2002). In wetlands, these factors include light, redox potential, salinity, carbon substrate availability, inorganic N and P concentrations, Fe and Mo, availability and grazers (Capone & Kiene, 1988; Howarth et al., 1999). In saline environments, bacteria have to cope with ionic stress, which is balanced by intracellular osmoprotective compounds, many of which are sugars (e.g., disaccharide trehalose, Welsh & Herbert, 1999) or contain N (amino acids and their derivatives such as glycine betaine, Csonka, 1989; Galinski & Trüper, 1994). N fixation thus provides an important source of N in these, often N-limited, environments (Casselman et al., 1981; Whiting & Morris, 1986).

Ecosystem functioning of tropical herbaceous wetlands is generally poorly known in spite of their ecological significance. Wetlands of our study system in northern Belize are situated in a flat karstic area with a calcareous marl bedrock and range in size from small <1 ha marshes to large shallow inland lagoons (>100 km²). Essentially, all of these wetlands are dominated by benthic cyanobacterial mats with scattered macrophytes, mainly *Eleocharis* spp., and they are strongly phosphorus (P) limited (Rejmánková et al., 1995; Rejmánková, 2001). No N limitation has been detected in any of the reported studies.

Cyanobacterial mats are characterized by high primary production and relatively high rates of N fixation (Rejmánková & Komárková, 2000, 2005). Trophic status of some of these wetlands has been changing recently due to increasing input of nutrients from fertilizer runoff from the sugar cane fields and other crops (Johnson & Rejmánková, 2005).

Using these wetlands as a model system, we have established a field manipulative experiment to obtain a mechanistic explanation for an ecosystem level response to increased nutrient input along a salinity gradient. The central hypothesis was: The macrophyte/microphyte competition following P addition will constrain cyanobacterial N fixation, which eventually will lead to N limitation. We have confirmed that P addition promotes the growth of macrophytes (*Eleocharis* spp. and *Typha domingensis*), which rapidly reduces cyanobacterial mats (Rejmánková & Houdková, 2006). The competitive replacement of cyanobacterial mats by macrophytes following P addition limits N fixation by cyanobacteria and changes the quantity and quality of organic matter input to the sediments. There is a significant difference in the amount and quality of litter produced by the two macrophytes (litter production on average 800 and 1700 g m⁻² y⁻¹ for *Eleocharis* and *Typha*, respectively; Rejmánková et al., 2008). We expect that the activity of sediment microorganisms, and consequently the N and P mineralization processes will be strongly impacted by the quality of input of organic carbon through litter and root exudates. We also expect that, due to processing litter with high C/N ratio, soil microorganisms will require extra N to produce new microbial biomass. In situations where the pool of available N is limited, most of N will be immobilized by microbes. Whether this will eventually lead to competition between

plants and microorganisms for N, and lead to plant N limitation, will depend on the availability of organic matter, and, consequently, heterotrophic N fixation.

In this study, we report on the differences in the sediment heterotrophic N fixation among controls and P-enriched plots dominated either by *Eleocharis cellulosa* or *Typha domingensis*, and attempt to explain these differences in the context of abiotic and biotic factors. We hypothesize that: (i) elimination of cyanobacterial mats in P-enriched plots shifts the microbial activity from autotrophs to heterotrophs and increases the microbial activity in sediments, mainly in the rhizosphere; (ii) in *Typha* dominated plots, the large quantity of litter and larger amount of C-rich root exudates will result in higher demand by sediment microorganisms for N and will lead to higher heterotrophic N fixation than in *Eleocharis* dominated plots; (iii) in higher salinity marshes, the microorganisms may need to invest more energy into osmotic regulation than to the enzyme production, which will result in lower heterotrophic N fixation.

Methods

Study site

Our study sites are located in lowlands of northern Belize, Central America within a 50-km radius of 18°9'58" N and 88°31'28" W. This part of the Yucatan Peninsula is an uplifted marine platform composed of a 2–3 km thick sequence of cretaceous and Tertiary limestone, dolomite, and gypsum (Weidie, 1985). Marsh hydrology is closely linked to the ground water system, and water levels are controlled primarily by regional precipitation. The majority of wetlands in the study area remain flooded or saturated year round. Most of the coastal plain contains a freshwater lens many meters thick floating on seawater. The porous limestone aquifer allows inland seawater intrusion, and, as a result, the mixing of fresh and marine waters can occur. Furthermore, the “fresh” ground waters are nearly saturated with carbonate and sulfate, derived from the dissolution of platform rocks. The net result of these factors is that there are large differences in ion content (sulfate, bicarbonate, and chloride) and conductivity of inland wetlands, which vary by several orders of magnitude (0.2–7 mS). Sediments in low conductivity marshes

are usually formed by peaty clays, while high conductivity marshes have predominantly marly sediments.

Main primary producers in these systems are several species of emergent macrophytes (*Eleocharis cellulosa*, *E. interstincta*, *Cladium jamaicense*, and *Typha domingensis*) and speciose communities of microphytes represented mostly by cyanobacteria (Rejmánková et al., 2004).

Treatment plot history

Fifteen marshes of diverse salinities, all dominated by sparse macrophytes (*Eleocharis* spp.) and cyanobacterial mats, have been studied as a part of a project aimed at the assessment of ecosystem response to nutrient addition along a salinity gradient (Rejmánková et al., 2008). Four 10 × 10 m plots were established in each marsh in August 2001; one represented a control, the remaining three received N, P and N&P addition in August 2001, August 2002, and March 2005, N was added as ammonium nitrate and P as triple super phosphate in amounts corresponding to 20 and 10 g m⁻² y⁻¹, respectively. In March 2003, one individual of *Typha domingensis* was planted in each plot. In the majority of the controls and N addition plots, the *Typha* plant did not survive. In P-enriched plots, however, they have been growing vigorously and outcompeting *Eleocharis*. In January 2005, the P addition plots in 6 marshes were manipulated to be half-dominated by *Eleocharis* and half dominated by *Typha*. Two of these 6 marshes, one from the low and one from the high salinity categories, were selected to measure the impact of P addition and macrophyte dominance on the heterotrophic N fixation (Fig. 1).

Soil and water sampling and analyses

Samples were collected from controls, *Eleocharis* dominated P addition plots, and *Typha* dominated P addition plots in August 2005 (beginning of wet season) and March 2007 (dry season), respectively (Fig. 1). The plots were always flooded and the water depth ranged from 25 to 50 cm. Recently deposited, readily distinguishable plant detritus on the soil surface was gently removed before sampling. Five randomly located sediment samples were collected with a 5.5-cm diameter sharp edge PVC corer to a depth of




Fig. 1 Schematic of the experimental design and sampling strategy. Five soil cores (black dots) were randomly collected from control plots (white) and P-enriched plots dominated by

approximately 30 cm. Each core was sectioned into the upper (0–7 cm, mostly rhizosphere) and lower (7–30 cm, mostly bulk soil) based on color and texture. Respective layers of all the five cores were combined into a composite sample. Large plant debris, roots, and shells were removed; samples were homogenized and stored on ice until processing. Samples for acetylene reduction assay (ARA) were extracted the same day. Soil samples for the other assays were stored for a maximum of one month at 4°C, i.e., under the storage conditions recommended when samples could not be analyzed immediately (Alef & Nannipieri, 2005).

Interstitial water was collected into HCl-rinsed plastic bottles with a suction sampler according to McKee et al. (1988), filtered through a 0.45- μm filter and stored on ice until processing. $\text{NH}_4\text{-N}$ was determined colorimetrically by the modification of the indolphenol blue method (Liddicoat et al., 1975).

Sediment bulk density was measured for each core section (Singer & Munns, 1996). Dried soil samples were pretreated with 0.1 M HCl and analyzed on a Carlo-Erba series 5000 CHN-S analyzer to obtain the total soil organic carbon and nitrogen. Microbial biomass carbon was determined by fumigation extraction procedures of three 10-g subsamples coupled to a 0.5 M K_2SO_4 extraction (White & Reddy, 2001). The extracted organic C was determined on a Shimadzu total organic carbon analyzer (TOC-5050A). Microbial biomass C was calculated using the extraction efficiency factor $k_{\text{EC}} = 0.37$ as the difference between treated (fumigated) and untreated soils. Biomass nutrient values were expressed per gram soil dry weight. Total P was analyzed spectrophotometrically using ascorbic acid reduction of phosphomolybdate complex after combustion and consequent acid digestion (McNamara & Hill, 2000). The soil reactive P was analyzed using

Eleocharis (light gray) and *Typha* (dark gray). To avoid the discrepancy in plot size of controls; samples were collected in only a 5 \times 10 m area (indicated by dotted line)

ascorbic acid reduction of phosphomolybdenate complex in oxalate extracts (Owens et al., 1977).

Root stock biomass

Three additional cores were collected from each plot and sectioned as described above.

All the living and dead plant roots and rhizomes were removed from cores sections by sequentially washing them through a 1- and 0.5-mm sieve. The living and dead roots were separated from each sample using a two-phase decanting technique in Ludox TM-50, originally developed for mangrove roots (Robertson & Dixon, 1993). After separation, roots were oven dried at 80°C for 48 h and weighed for biomass determination.

Acetylene reduction assay (ARA)

We used the ARA method modified from Hardy et al. (1968). Fifty grams of freshly collected water saturated test soil (equal to about 15 g dry weight) were sealed in 100-ml glass bottles. All the assays were run in triplicates. One ml of 2% glucose solution (corresponding to the addition of 0.16 mg C per cm^3 of the sediment) was added to optimize the reaction (Dicker & Smith, 1980). Glucose is regarded as a good proxy for sucrose, which is likely to originate from wetland macrophyte root exudates (Nielsen et al., 2001). To establish the relationship between the glucose concentration and nitrogenase activity, we set up a separate experiment in which we incubated the samples with glucose concentrations ranging from 0 to 0.42 mg C cm^{-3} . One set of samples (low salinity marsh, upper sediment layers) was also assayed without the glucose enrichment. The bottles were equilibrated to the atmospheric pressure, 30 ml of the

headspace were removed and subsequently replaced with 20 ml of N_2 gas to lower the partial pressure of oxygen. Ten milliliters of acetylene, freshly prepared from CaC_2 , were added to each bottle, the bottles were vigorously shaken, and were incubated under dark conditions at 28–30°C. After 12 and 24 h the bottles were shaken again, and after 24 h, several milliliters of headspace were withdrawn with an airtight syringe and analyzed by gas chromatograph, using Shimadzu 14 GC with a flame ionization detector and a Porapak-T column at 80°C. Acetylene-free controls as well as blanks (empty bottles incubated with 10 ml of acetylene) showed no endogenous ethylene production. Gravimetric soil moisture was measured after the incubation. Rates of acetylene reduction were expressed as nmol of acetylene reduced per gram dry mass of sample per day of incubation.

Phospholipid fatty acid (PLFA) analysis

PLFA analyses were performed according to Frostegård et al. (1993). Phospholipids, in the form of fatty acid methyl esters, were quantified by gas chromatography using Hewlett Packard 6890 N with FID detector. Since the quantity of microbial PLFA is a function of microbial biomass (Hassett & Zak, 2005), we used the total amount of all bacterial PLFA identified in the samples as an indicator of live microbial biomass.

Data analysis

We used path analysis, according to Pedhazur (1997), to quantify the relationships among the variables of this study based on a path diagram that specifies the direct and indirect paths between sediment P and heterotrophic nitrogenase activity.

Results

All the results are presented for nitrogenase activity (NA) measured in samples enriched with glucose in the amount of $0.16 \text{ mg C cm}^{-3}$ soil. In the preliminary experiment, NA increased linearly with glucose additions, until about a concentration of 0.3 mg C cm^{-3} , and further additions had either a null or even negative effect (Fig. 2). NA without glucose was two to three orders of a magnitude lower.




Fig. 2 Mean nitrogenase activity (NA) of composite sediment samples amended with glucose, expressed as % of the highest value. Each NA assay was run in three replicates

Comparison of nitrogenase activity over all treatments, sediment layers, marsh salinity, and time

We found large differences in NA among control and P-enriched plots in samples from both low- and high-salinity marshes as well as between upper and lower sediment layers (Fig. 3). These patterns were very similar for both sampling periods (August 2005 and March 2007). There was a highly significant difference in NA among treatments and between the sediment layers (Table 1), while the effect of salinity was only marginal. NA in the *Typha* upper sediment layer was consistently lower, compared to the *Eleocharis* treatment, while the opposite was true for the lower layers (Fig. 3).

Relationship between nitrogenase activity and environmental variables

To explore how NA relates to changes in chemical and biological sediment variables, we calculated simple regressions (Fig. 4). NA was positively correlated with soil reactive P (Fig. 4A, $R^2 = 0.536$, $P < 0.001$), and, albeit less strongly, with total P ($R^2 = 0.44$, $P = 0.02$; data not shown). There was a strong negative relationship between NA and NH_4 -N in the interstitial water (Fig. 4B, $R^2 = 0.507$, $P = 0.009$). NA was also positively correlated with increasing content of total organic carbon, microbial carbon,




Fig. 3 Nitrogenase activity (NA) measured in August 2005 and March 2007 in the upper and lower layer sediments from controls (C) and P addition plots dominated by *Eleocharis cellulosa* (PE) and *Typha domingensis* (PT). The error bars indicate the standard error of the mean of three replicates,

which were taken from a composite sample from each plot. NA is expressed in nmols C₂H₄ per cm³ of sediment per day to account for different sediment densities among the upper and lower layers. The log scale was used to better visualize the differences between treatments

Table 1 Mean values (\pm SD) of nitrogenase activity (NA) and sediment characteristics for the upper and lower layer of control and P-enriched sites dominated by *Eleocharis* or *Typha*

	High salinity (7 mS)			Low salinity (0.2 mS)		
	Control	<i>Eleocharis</i>	<i>Typha</i>	Control	<i>Eleocharis</i>	<i>Typha</i>
<i>Upper layer</i>						
NA, nmol C ₂ H ₄ g ⁻¹ d ⁻¹	1.18 (0.54)	806 (52.5)	257 (186)	0.92 (0.25)	1189 (376)	985 (475)
Live root biomass, mg g ⁻¹	4.94 (0.54)	5.56 (0.64)	7.02 (1.37)	3.68 (1.2)	16.2 (1.16)	10.8 (0.95)
Bulk density, g cm ⁻³	0.22 (0.01)	0.23 (0.04)	0.26 (0.04)	0.37 (0.05)	0.21 (0.02)	0.18 (0.03)
Soil reactive P, μ g g ⁻¹	0.69 (0.07)	6.78 (0.49)	6.27 (0.02)	0.83 (0.09)	8.83 (0.31)	16.5 (0.25)
Total soil P, μ g g ⁻¹	219	1175	666	202	367	803
NH ₄ -N water, μ g l ⁻¹	2486 (47.9)	1108 (397)	951 (75.9)	1252 (436)	30.7 (13.4)	57.6 (20.4)
Total soil N, mg g ⁻¹	9.3 (0.10)	7.5 (0.40)	9.1 (3.60)	10.6 (0.30)	15.0 (0.80)	16.9 (0.80)
Microbial C, μ g g ⁻¹	122 (46.1)	154 (32.1)	148 (16.9)	239 (47.5)	409 (10.3)	752 (3.37)
Total soil org. C, mg g ⁻¹	98.5 (4.50)	165 (0.50)	146 (2.50)	115 (6.00)	187 (17.5)	205 (16.5)
Total PLFA, nmolFA g ⁻¹	248 (24.7)	283 (30.4)	293 (87.9)	145 (24.0)	208 (33.9)	282 (15.1)
<i>Lower layer</i>						
NA, nmol C ₂ H ₄ g ⁻¹ d ⁻¹	0.60 (0.40)	14.0 (2.00)	67.0 (36.0)	0.78 (0.24)	11.5 (3.50)	89.0 (24.0)
Live root biomass, mg g ⁻¹	0.76 (0.03)	1.31 (0.11)	1.17 (0.22)	0.9 (0.07)	1.54 (0.15)	3.65 (0.35)
Bulk density, g cm ⁻³	0.49 (0.03)	0.52 (0.07)	0.56 (0.01)	0.50 (0.04)	0.40 (0.01)	0.35 (0.02)
Soil reactive P, μ g g ⁻¹	0.38 (0.04)	2.00 (0.04)	0.16 (0.02)	2.65 (0.12)	2.59 (0.04)	3.72 (0.04)
Total soil P, μ g g ⁻¹	313	189	208	67	105	101
Total soil N, mg g ⁻¹	5.80 (0.40)	6.60 (1.30)	6.30 (3.10)	10.4 (3.60)	14.2 (0.20)	18.7 (3.30)
Microbial C, μ g g ⁻¹	6.39 (2.79)	13.5 (1.48)	12.7 (0.95)	45.5 (12.9)	131 (76.4)	77.2 (10.1)
Total soil org. C, mg g ⁻¹	97.5 (20.5)	112 (4.50)	123 (2.50)	60 (21.1)	155 (18.5)	158 (30.0)
Total PLFA, nmol FA g ⁻¹	46.7 (1.48)	46.4 (6.07)	46.2 (4.04)	54.5 (19.1)	34.7 (3.58)	141 (18.9)

The values are averages of 2005 and 2007 sampling dates (each date 3 replicated samples analyzed), except for root biomass that was sampled only in 2007 and total sediment P that was analyzed only in 2005




Fig. 4 Relationship between NA and: (A) soil reactive P (PO₄-P, µg g⁻¹); (B) NH₄-N in the interstitial water (µg l⁻¹); (C) total soil organic carbon (mg g⁻¹); (D) microbial carbon

(µg g⁻¹); (E) biomass of live roots (mg dry weight g⁻¹ of sediment); and (F) total PLFA (nmol FA g⁻¹). Each point represents the mean of three replicates

and total PLFA (Fig. 4C, D and F; $R^2 = 0.369$, $P = 0.002$; $R^2 = 0.469$, $P < 0.001$; and $R^2 = 0.353$, $P = 0.002$, respectively). We only had data on sediment live root biomass for 2007, but we found a highly significant correlation between sediment live root biomass and NA (Fig. 4E; $R^2 = 0.643$).

Path analysis

The effect of sediment P on NA is clearly very important and it can impact the N fixing bacteria directly by removing P deficiency, but also indirectly, through changes in other variables. While NA is




Fig. 5 Path diagram of the direct and indirect effects of sediment phosphorus concentrations on nitrogenase activity (NA). Path coefficients P_{21} , P_{31} , and P_{41} are standardized simple regression coefficients, path coefficients P_{Y1} , P_{Y2} , P_{Y3} , and P_{Y4} are standardized partial regression coefficients from the multiple regression. Products $P_{21}P_{Y2}$, $P_{31}P_{Y3}$, and $P_{41}P_{Y4}$ indicate the indirect effects of sediment P on NA through microbial carbon, live root biomass, and interstitial water $\text{NH}_4\text{-N}$, respectively. Path coefficients printed in bold are significant

at $P < 0.05$. If both path coefficients forming a product are significant at $P < 0.05$, the product is also printed in bold. The residual variable, U , includes all the unmeasured variables that affect the dependent variable and its path coefficient = $\sqrt{1 - R^2}$, where R is the proportion of variation in the dependent variable explained by the model. Besides individual path coefficients, the overall correlation between P sediment and NA are given in parentheses

positively correlated with total organic C, microbial C, total PLFA, and live root biomass, and negatively correlated with $\text{NH}_4\text{-N}$ (Fig. 4), all of these variables have also a significant relationship with sediment P (see standardized simple regression coefficients P_{21} , P_{31} , and P_{41} shown in the path analysis, Fig. 5; P values for all these regressions < 0.05). To elucidate the importance of the indirect effects of P on NA, we used a simple path analysis (Fig. 5). Because total organic C and total PLFA did not improve the significance of the multiple regression, only sediment P, microbial C, $\text{NH}_4\text{-N}$, and live root biomass were used in the model. The path analysis revealed that the indirect effect of P on NA through the root biomass is more important than the direct effect of P on NA (compare the product of path coefficients $P_{31}P_{Y3} = 0.624$ with $P_{Y1} = 0.453$ in Fig. 5).

Discussion

Response to phosphorus addition

The addition of P to our experimental plots significantly increased macrophyte primary production, which led to the rapid elimination of both floating and benthic cyanobacterial mats (Rejmánková et al.,

2008). A similar response has been described for other oligotrophic, P-limited wetlands such as the Florida Everglades (McCormick et al. 1996, 1998; Gaiser et al. 2005). While autotrophic cyanobacteria were eliminated, the biomass of heterotrophic sediment microorganisms increased due to higher C, input, as documented by increase in total organic C and, consequently, in microbial C, and total PLFA. No significant differences have been found in total N but the concentration of $\text{NH}_4\text{-N}$ in interstitial water, which is typically high in the unimpacted marshes, decreased several fold.

Heterotrophic N fixation significantly responded to these changes, particularly in the more active upper sediment layers where it was two orders of magnitude higher than in controls. Our first hypothesis stating that the elimination of cyanobacteria in P-enriched plots will shift the microbial activity from autotrophs to heterotrophs and increase the microbial activity in sediments, mainly in the rhizosphere was confirmed.

Heterotrophic N fixation is strongly dependent on available carbon, where carbon quality is more important for fixation than is quantity (Vitousek et al. 2002). We had two sediment carbon characteristics available: total organic C and microbial C, and NA was positively associated with both of them. Microbial biomass has also been shown to be closely

correlated with available C (Sparling et al. 1998; Ghani et al. 2003). However, the variable that was the most tightly correlated with NA was the biomass of live roots, which is in agreement with the fact that diazotrophs are most active and numerous in the environment that is rich in available C, e.g., in the rhizosphere. This “stimulation effect” of root exudates on NA is generally accepted in both terrestrial and wetland ecosystems (Van Berkum & Sloger, 1979; Welsh et al., 1996; Dakora & Drake, 2000; Nielsen et al., 2001). Even though our acetylene assays were conducted on sediments from which the roots were previously removed, the rhizosphere microflora was present, and sediment was apparently affected by previous flux of C from roots. As the root biomass increases in sediment, so also should the area that is enriched by available C. We found significant correlation between live root biomass and available C in both the upper and lower layers ($r^2 = 0.53$ and 0.83 , respectively).

The negative relationship between NA and the concentration of $\text{NH}_4\text{-N}$ in the interstitial water was in agreement with the widely accepted claim that the synthesis and/or NA is repressed by high levels of $\text{NH}_4\text{-N}/\text{NO}_3\text{-N}$ (Yoch & Whiting, 1986; Howarth et al., 1988b; Hansen et al., 2000).

Plant effects

We hypothesized that compared to *Eleocharis* dominated plots, the large quantity of litter and larger amount of C-rich root exudates in *Typha*-dominated plots would result in higher demand by sediment microorganisms for N and would lead to higher heterotrophic N fixation. Results from the upper sediment layer showed a completely opposite trend with consistently higher NA in *Eleocharis* plots rather than in *Typha* (Fig. 3). However, in the lower sediment layers, we found higher NA in *Typha* plots. We believe this discrepancy exists due to differences in stand age. Since *Typha* became fully established in plots only in 2005, its litter probably only impacts the upper sediment layers. Most importantly though, we found higher phenolic and lignin contents in *Typha* litter as opposed to *Eleocharis* litter (phenolics $5.68 \pm 1.5\%$ and $2.09 \pm 0.7\%$; lignin: $3.85 + 0.99\%$ and $1.52 + 0.38\%$ in dry leaf material, respectively; Rejmánková, unpublished data). Both phenolics and lignin have been found to limit N fixation (Pelegri & Twilley, 1998).

Thus it is conceivable, that the higher levels of these compounds constrained NA in *Typha* plots relative to *Eleocharis*.

Salinity effects

The effects of salinity on NA were not conclusive, and NA was not consistently lower in the high salinity plots. Finding lower NA in some plots from higher salinity marshes could be a result of a smaller, less active community of N fixers due to salinity stress. We can also speculate that N fixers in the high salinity plots would need to invest more energy into osmotic regulation than to the enzyme production. There are conflicting reports on the effects of salinity on NA. While some studies have found that concentrations of NaCl as low as 75 mM can have a negative impact on NA in symbiotic N fixers (Bolaños et al., 2006), others have reported no effects from highly saline environments (Zahran, 1997). The extent by which N fixers are constrained by salinity in our system will be explored in future research.

In conclusion, our data show that P addition controls heterotrophic N fixation mainly via root development, although litter input quality also appears to be important. The estimate of N balances indicates that microbial N fixation, under conditions of unlimited carbon supply could potentially cover and even exceed plant N demand.

Acknowledgments We would like to thank Ireneo Briceno and Russell King for their assistance in the field, Emily Carlson for laboratory assistance, Marcel Rejmánek for comments to the first draft of the article, and to the journal editor and the two anonymous reviewers for their helpful comments and suggestions. This research was supported by these grants: NSF grant # 0089211 to E. R.; ME 912, and MSM # 600 766 5801 to H.Š.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Alef, K. & P. Nannipieri, 2005. Methods in Applied Soil Microbiology and Biochemistry. Academic Press, London.
- Bagwell, C. E. & C. R. Lovell, 2000. Microdiversity of culturable diazotrophs from the rhizoplanes of the salt marsh

- grasses *Spartina alterniflora* and *Juncus roemerianus*. *Microbial Ecology* 39: 128–136.
- Bergholz, P. W., C. E. Bagwell & C. R. Lovell, 2001. Physiological diversity of rhizoplane diazotrophs of the saltmeadow cordgrass, *Spartina patens*: implications for host specific ecotypes. *Microbial Ecology* 42: 466–473.
- Bolaños, L., M. Martín, A. El-Hamdaoui, R. Rivilla & I. Bonilla, 2006. Nitrogenase inhibition in nodules from pea plants grown under salt stress occurs at the physiological level and can be alleviated by B and Ca. *Plant and Soil* 280: 135–142.
- Boyle, C. D. & D. G. Patriquin, 1981. Carbon metabolism of *Spartina-alterniflora* loisel in relation to that of associated nitrogen-fixing bacteria. *New Phytologist* 89: 275–288.
- Bristow, J. M., 1973. Nitrogen-fixation in rhizosphere of aquatic angiosperms. *Plant Physiology* 51: 34.
- Bürgmann, H., S. Meier, M. Bunge, F. Widmer & J. Zeyer, 2005. Effects of model root exudates on structure and activity of a soil diazotroph community. *Environmental Microbiology* 7: 1711–1724.
- Capone, D. G. & R. P. Kiene, 1988. Comparison of microbial dynamics in marine and fresh-water sediments – contrasts in anaerobic carbon catabolism. *Limnology and Oceanography* 33: 725–749.
- Casselman, M. E., W. H. Patrick & R. D. Delaune, 1981. Nitrogen-fixation in a gulf-coast salt-marsh. *Soil Science Society of America Journal* 45: 51–56.
- Coleman, D. C., 2008. From peds to paradoxes: linkages between soil biota and their influences on ecological processes, review. *Soil Biology and Biochemistry* 40: 271–289.
- Csonka, L. N., 1989. Physiological and genetic responses of bacteria to osmotic-stress. *Microbiological Reviews* 53: 121–147.
- Dakora, F. D. & B. G. Drake, 2000. Elevated CO₂ stimulates associative N₂ fixation in a C-3 plant of the Chesapeake Bay wetland. *Plant Cell and Environment* 23: 943–953.
- Dicker, H. J. & D. W. Smith, 1980. Acetylene reduction (nitrogen fixation) in a Delaware, USA salt marsh. *Marine Biology* 57: 241–250.
- Frostegård, A., A. Tunlid & E. Bååth, 1993. Phospholipid fatty-acid composition, biomass, and activity of microbial communities from 2 soil types experimentally exposed to different heavy-metals. *Applied and Environmental Microbiology* 59: 3605–3617.
- Galinski, E. A. & H. G. Trüper, 1994. Microbial behavior in salt-stressed ecosystems. *FEMS Microbiology Reviews* 15: 95–108.
- Gaiser, E. E., J. C. Trexler, J. H. Richards, D. L. Childers, D. Lee, A. L. Edwards, L. J. Scinto, K. Jayachandran, G. B. Noe & R. D. Jones, 2005. Cascading ecological effects of low-level phosphorus enrichment in the Florida Everglades. *Journal of Environmental Quality* 34: 717–723.
- Ghani, A., M. Dexter & K. W. Perrott, 2003. Hot-water extractable carbon in soils: a sensitive measurement for determining impacts of fertilisation, grazing and cultivation. *Soil Biology and Biochemistry* 35: 1231–1243.
- Hansen, J. W., J. W. Udy, C. J. Perry, W. C. Dennison & B. A. Lomstein, 2000. Effect of the seagrass *Zostera capricorni* on sediment microbial processes. *Marine Ecology-Progress Series* 199: 83–96.
- Hardy, R. W. F., R. D. Holsten, E. K. Jackson & R. C. Burns, 1968. Acetylene-ethylene assay for N₂ fixation – laboratory and field evaluation. *Plant Physiology* 43: 1185–1207.
- Hassett, J. E. & D. R. Zak, 2005. Aspen harvest intensity decreases microbial biomass, extracellular enzyme activity, and soil nitrogen cycling. *Soil Science Society of America Journal* 69: 227–235.
- Howarth, R. W., R. Marino, J. Lane & J. J. Cole, 1988a. Nitrogen-fixation in fresh-water, estuarine, and marine ecosystems. 1. Rates and importance. *Limnology and Oceanography* 33: 669–687.
- Howarth, R. W., R. Marino & J. J. Cole, 1988b. Nitrogen-fixation in fresh-water, estuarine, and marine ecosystems. 2. Biogeochemical controls. *Limnology and Oceanography* 33: 688–701.
- Howarth, R. W., F. Chan & R. Marino, 1999. Do top-down and bottom-up controls interact to exclude nitrogen-fixing cyanobacteria from the plankton of estuaries? An exploration with a simulation model. *Biogeochemistry* 46: 203–231.
- Inglett, P. W., K. R. Reddy & P. V. McCormick, 2004. Periphyton chemistry and nitrogenase activity in a northern Everglades ecosystem. *Biogeochemistry* 67: 213–233.
- Johnson, S. & E. Rejmánková, 2005. Impacts of land use on nutrient distribution and vegetation composition of freshwater wetlands in Northern Belize. *Wetlands* 25: 89–100.
- Liddicoat, M. I., S. Tibbitts & E. I. Butler, 1975. Determination of ammonia in seawater. *Limnology and Oceanography* 20: 131–132.
- McCormick, P. V., P. S. Rawlik, K. Lurding, E. P. Smith & F. H. Sklar, 1996. Periphyton-water quality relationships along a nutrient gradient in the northern Florida Everglades. *Journal of the North American Benthological Society* 15: 433–449.
- McCormick, P. V., R. B. E. Shuford, J. G. Backus & W. C. Kennedy, 1998. Spatial and seasonal patterns of periphyton biomass and productivity in the northern Everglades, Florida, USA. *Hydrobiologia* 362: 185–208.
- McKee, K. L., I. A. Mendelssohn & M. W. Hester, 1988. Reexamination of pore water sulfide concentrations and redox potentials near the aerial roots of *Rhizophora mangle* and *Avicennia germinans*. *American Journal of Botany* 75: 1352–1359.
- McNamara, A. E. & W. R. Hill, 2000. UV-B irradiance gradient affects photosynthesis and pigments but not food quality of periphyton. *Freshwater Biology* 43: 649–662.
- Moseman, S. M., 2007. Opposite diel patterns of nitrogen fixation associated with salt marsh plant species (*Spartina foliosa* and *Salicornia virginica*) in southern California. *Marine Ecology-an Evolutionary Perspective* 28: 276–287.
- Nielsen, L. B., K. Finster, D. T. Welsh, A. Donnelly, R. A. Herbert, R. de Wit & B. A. Lomstein, 2001. Sulphate reduction and nitrogen fixation rates associated with roots, rhizomes and sediments from *Zostera noltii* and *Spartina maritima* meadows. *Environmental Microbiology* 3: 63–71.

- Owens, L. B., D. W. Nelson & L. E. Sommers, 1977. Determination of inorganic phosphorus in oxalate extracts of soils. *Soil Science Society of America Journal* 41: 148–149.
- Paerl, H. W., B. M. Bebout, C. A. Currin, M. W. Fitzpatrick & J. L. Pickney, 1994. Nitrogen fixation in microbial mats. In Stal, L. J. & P. Caumette (eds), *Microbial mats. Structure, development and environmental significance*. NATO ASI Springer Verlag, Heidelberg: 325–337.
- Pedhazur, E. J., 1997. *Multiple regression in behavioral research: explanation and prediction*. Hartcourt Brace College Publishers, Fort Worth, Texas.
- Pelegri, S. P. & R. R. Twilley, 1998. Heterotrophic nitrogen fixation (acetylene reduction) during leaf-litter decomposition of two mangrove species from south Florida, USA. *Marine Biology* 131: 53–61.
- Rejmánková, E., K. O. Pope, M. D. Pohl & J. M. Rey Benayas, 1995. Fresh-water wetland plant-communities of Northern Belize – implications for paleoecological studies of Maya wetland agriculture. *Biotropica* 27: 28–36.
- Rejmánková, E. & J. Komárková, 2000. A function of cyanobacterial mats in phosphorus-limited tropical wetlands. *Hydrobiologia* 431: 135–153.
- Rejmánková, E., 2001. Effect of experimental phosphorus enrichment on oligotrophic tropical marshes in Belize, Central America. *Plant and Soil* 236: 33–53.
- Rejmánková, E., J. Komárek & J. Komárková, 2004. Cyanobacteria – a neglected component of biodiversity: patterns of species diversity in inland marshes of Northern Belize (Central America). *Diversity and Distributions* 10: 189–199.
- Rejmánková, E. & J. Komárková, 2005. Response of cyanobacterial mats to nutrient and salinity changes. *Aquatic Botany* 83: 87–107.
- Rejmánková, E. & K. Houdková, 2006. Wetland plant decomposition under different nutrient conditions: what is more important, litter quality or site quality? *Biogeochemistry* 80: 245–262.
- Rejmánková, E., P. Macek & K. Epps, 2008. Wetland ecosystem changes after three years of phosphorus addition. *Wetlands* 28: 914–927.
- Robertson, A. I. & P. Dixon, 1993. Separating live and dead fine roots using colloidal silica – an example from mangrove forests. *Plant and Soil* 157: 151–154.
- Roger, P. A., 1995. Biological N₂-fixation and its management in wetland rice cultivation. *Fertilizer Research* 42: 261–276.
- Roger, P. A. & J. K. Ladha, 1992. Biological N₂ fixation in wetland rice fields – estimation and contribution to nitrogen-balance. *Plant and Soil* 141: 41–55.
- Scott, J. T., R. D. Doyle & C. T. Filstrup, 2005. Periphyton nutrient limitation and nitrogen fixation potential along a wetland nutrient-depletion gradient. *Wetlands* 25: 439–448.
- Scott, J. T., R. D. Doyle, J. A. Back & S. I. Dworkin, 2007. The role of N₂ fixation in alleviating N limitation in wetland metaphyton: enzymatic, isotopic, and elemental evidence. *Biogeochemistry* 84: 207–218.
- Singer, M. J. & D. N. Munns, 1996. *Soils. An Introduction*, 3rd ed. Prentice-Hall, London: 77–78.
- Sparling, G., M. Vojvodic-Vukovic & L. A. Schipper, 1998. Hot-water-soluble C as a simple measure of labile soil organic matter: the relationship with microbial biomass C. *Soil Biology and Biochemistry* 30: 1469–1472.
- Steppe, T. F. & H. W. Paerl, 2005. Nitrogenase activity and Nifh expression in a marine intertidal microbial mat. *Microbial Ecology* 49: 315–324.
- Tjepkema, J. D. & H. J. Evans, 1976. Nitrogen-fixation associated with *Juncus-balticus* and other plants of Oregon wetlands. *Soil Biology and Biochemistry* 8: 505–509.
- Tyler, A. C., T. A. Mastronicola & K. J. McGlathery, 2003. Nitrogen fixation and nitrogen limitation of primary production along a natural marsh chronosequence. *Oecologia* 136: 431–438.
- Van Berkum, P. & C. Sloger, 1979. Immediate acetylene-reduction by excised grass roots not previously pre-incubated at low oxygen-tensions. *Plant Physiology* 64: 739–743.
- Vargas, R. & E. Novelo, 2007. Seasonal changes in periphyton nitrogen fixation in a protected tropical wetland. *Biology and Fertility of Soils* 43: 367–372.
- Vitousek, P. M., K. Cassman, C. Cleveland, T. Crews, C. B. Field, N. B. Grimm, R. W. Howarth, R. Marino, L. Martinelli, E. B. Rastetter & J. I. Sprent, 2002. Towards an ecological understanding of biological nitrogen fixation. *Biogeochemistry* 57: 1–45.
- Weidie, A. E., 1985. Geology of the Yucatan platform, part 1. In Ward, W. C., A. E. Weidie & W. Back (eds), *Geology and hydrogeology of the Yucatan and quaternary geology of northeastern Yucatan peninsula*. Geological Society, New Orleans, LA: 1–19.
- Welsh, D. T. & R. A. Herbert, 1999. Osmotically induced intracellular trehalose, but not glycine betaine accumulation promotes desiccation tolerance in *Escherichia coli*. *FEMS Microbiology Letters* 174: 57–63.
- Welsh, D. T., P. Wellsbury, S. Bourgues, R. de Wit & R. A. Herbert, 1996. Relationship between porewater organic carbon content, sulphate reduction and nitrogen fixation (acetylene reduction) in the rhizosphere of *Zostera noltii*. *Hydrobiologia* 329: 175–183.
- White, J. R. & K. R. Reddy, 2001. Influence of selected inorganic electron acceptors on organic nitrogen mineralization in Everglades soils. *Soil Science Society of America Journal* 65: 941–948.
- Whiting, G. J. & J. T. Morris, 1986. Nitrogen-fixation (C₂H₂ reduction) in a salt-marsh – its relationship to temperature and an evaluation of an in situ chamber technique. *Soil Biology and Biochemistry* 18: 515–521.
- Yoch, D. C. & G. J. Whiting, 1986. Evidence for NH₄⁺ switch-off regulation of nitrogenase activity by bacteria in salt-marsh sediments and roots of the grass *Spartina-alterniflora*. *Applied and Environmental Microbiology* 51: 143–149.
- Zahran, H. H., 1997. Diversity, adaptation and activity of the bacterial flora in saline environments. *Biology and Fertility of Soils* 25: 211–223.