

# UC Santa Cruz

## UC Santa Cruz Previously Published Works

**Title**

Closed Geodesics in Alexandrov Spaces of Curvature Bounded from Above

**Permalink**

<https://escholarship.org/uc/item/1226b2xb>

**Journal**

The Journal of Geometric Analysis, 21(2)

**ISSN**

1050-6926

**Author**

Lin, Longzhi

**Publication Date**

2011-04-01

**DOI**

10.1007/s12220-010-9153-0

Peer reviewed

# CLOSED GEODESICS IN ALEXANDROV SPACES OF CURVATURE BOUNDED FROM ABOVE

LONGZHI LIN

ABSTRACT. In this paper, we show a local energy convexity of  $W^{1,2}$  maps into  $CAT(K)$  spaces. This energy convexity allows us to extend Colding and Minicozzi's width-sweepout construction to produce closed geodesics in any closed Alexandrov space of curvature bounded from above, which also provides a generalized version of the Birkhoff-Lyusternik theorem on the existence of non-trivial closed geodesics in the Alexandrov setting.

## 0. INTRODUCTION

Closed geodesics have been investigated mainly in the case of closed (i.e., compact and without boundary) Riemannian manifolds, while various results were obtained for Finsler manifolds and in the more general case of metric spaces with certain special properties (known as Busemann  $G$ -spaces, see [Bu]). These studies were initiated by Hadamard [Had], Poincaré [Po] and Birkhoff [B1]. The method of finding non-trivial closed geodesics on simply-connected manifolds by curve shortening map and sweepouts goes back to Birkhoff in 1917: one pulls each curve in a sweepout (see Definition 3.2) on the manifold as tight as possible, in a continuous way and preserving the sweepout. See [B1],[B2], [Cr] and [CC] (page 533) for more about Birkhoff's ideas.

In this paper, we show the following local energy convexity of  $W^{1,2}$  maps into a  $CAT(K)$  space (also known as an  $\mathfrak{R}_K$  domain, see [BBI] or subsection 1.1 below).

**Theorem 0.1.** *Let  $\Sigma$  be a compact Riemannian domain and  $(X, d)$  be an Alexandrov space of curvature bounded from above by  $K$ . In an  $\mathfrak{R}_K$  domain of  $x \in X$ , there exists  $\rho = \rho(x, K) > 0$  such that for  $u, v \in W^{1,2}(\Sigma, X)$  with images staying in  $B_\rho(x) \subset \mathfrak{R}_K$ , the following holds:*

$$(0.2) \quad \frac{1}{4} \int_{\Sigma} |\nabla d(u, v)|^2 \leq E^u + E^v - 2E^w.$$

Here  $B_\rho(x)$  is the geodesic ball centered at  $x$  with radius  $\rho$ ,  $w = \frac{u+v}{2}$  is the mid-point map and  $E$  is the 2-energy of maps into metric spaces (see subsection 1.2).

We shall remark that Theorem 0.1 provides a stronger (quantitative) version of a result of Burago, Burago and Ivanov [BBI, Proposition 9.1.17]. Theorem 0.1 allows us to use Colding and Minicozzi's width-sweepout construction of closed geodesics to produce closed geodesics in another class of general metric spaces, namely, the (closed) Alexandrov spaces of curvature bounded from above. Our main result is the following useful property of good sweepouts in any closed Alexandrov space of curvature bounded from above: *each curve in the tightened sweepout whose length is close to the length of the longest curve in the sweepout must itself be close to a closed geodesic*; see Theorem 3.14 below and cf. [CM1]-[CM3], [LW], proposition 3.1 of [CD], proposition 3.1 of [Pi], and 12.5 of [Al]. Moreover, as an immediate

corollary of the existence of good sweepouts we present in section 3, we obtain a generalized Birkhoff-Lyusternik theorem on the existence of non-trivial closed geodesics, cf. [LyS], [Ly].

**Theorem 0.3.** (*Generalized Birkhoff-Lyusternik theorem*) *Let  $(X, d)$  be a closed Alexandrov space of curvature bounded from above by  $K$ . Suppose that the  $k_0$ -th homology group  $H_{k_0}(X)$  is nonzero for some  $k_0 \geq 1$ , then  $(X, d)$  admits at least one non-trivial closed geodesic.*

In [CM1], Colding and Minicozzi introduced the geometric invariant of closed Riemannian manifolds that they call the *width*. They succeeded in using the local linear replacement as Birkhoff's curve shortening map to construct good sweepouts that produce at least one closed geodesic, which realizes the width as its energy. In particular, there exist closed geodesics on any closed Riemannian manifold. The argument only produces non-trivial closed geodesics when the width is positive (see footnote 4). Their local linear replacement process is a discrete gradient flow (for the length functional of curves), and it depends solely on a local energy convexity for  $W^{1,2}$  maps into closed Riemannian manifolds (see Lemma 4.2 of [CM1]), which controls the distance of curves in the tightened sweepout from closed geodesics explicitly. (See [CM2] for 2-width of manifolds and sweepouts by 2-spheres instead of circles to produce minimal surfaces on manifolds.)

Recall that any compact smooth Riemannian manifold is an Alexandrov space of curvature bounded from above by some  $K$  (see Theorem 1.1). As mentioned, the local energy convexity is crucial in the width-sweepout construction of closed geodesics. Therefore it is reasonable that if we can find a similar energy convexity for maps into  $CAT(K)$  spaces, we can extend Colding and Minicozzi's width-sweepout construction to produce closed geodesics in any closed Alexandrov space of curvature bounded from above by  $K$ , which is locally a  $CAT(K)$  space. In theorem 2.2 of [KS] (see equation (2.2iv) of [KS]), Korevaar and Schoen provided an energy convexity of  $W^{1,2}$  maps from a compact Riemannian domain into an NPC space (when  $K = 0$ ) for which the model space<sup>1</sup> is  $\mathbf{R}^2$ . Since the model space for the case of  $K > 0$  is the standard Euclidean 2-hemisphere  $\mathbf{S}_K$ , which is locally  $\mathbf{R}^2$ , it is perhaps not surprising that a similar energy convexity should also hold locally. In fact, using the  $K$ -quadrilateral cosine  $\text{cos}q_K$  that Berg and Nikolaev defined in [BN] we will be able to show that, with a small image assumption, the same (up to a constant) energy convexity still holds for  $W^{1,2}$  maps into a  $CAT(K)$  space with  $K > 0$ . Namely, we have Theorem 0.1 above (see section 2 for the proof).

We remark that in [Se] (unpublished) Serbinowski provides a local energy convexity for maps into a  $CAT(K)$  space. His result is an analogue to (0.2) and leads to the uniqueness of the small-image solution to the Dirichlet problem from a Riemannian domain into an Alexandrov space of curvature  $\leq K$ , yet we cannot use his version of convexity to control the distance between curves directly and explicitly.

The paper is organized as follows. In section 1 we collect some basic definitions and properties of the Alexandrov spaces of curvature bounded from above and also the energy of maps into metric spaces introduced by Korevaar and Schoen [KS]. In section 2, we will prove the local energy convexity in  $CAT(K)$  spaces and we postpone the elementary proof of the main Lemma 2.2 in Appendix C. In section 3, we extend the delicate construction of good sweepouts used by Colding and Minicozzi [CM1] to our Alexandrov setting, while

---

<sup>1</sup>Also known as the  $K$ -plane, see footnote 2.

we show this sweepout construction satisfies the properties of the Birkhoff curve-shortening process in Appendixes A and B. Finally we give a proof of Theorem 0.3 in section 4.

**Acknowledgement.** The author would like to thank Professor William Minicozzi for his continued guidance. The author would also like to thank the referee for the suggestion on adding Theorem 0.3 to this paper.

## 1. PRELIMINARIES

1.1. **Alexandrov space of curvature  $\leq K$ .** In the 1950's, Alexandrov introduced spaces of curvature bounded from above in his papers [A1], [A2]. The terminology  $CAT(K)$  spaces was then coined by Gromov in 1987. The initials are in honor of Cartan, Alexandrov and Toponogov. For the self-containedness of this paper, we will recall some basic definitions here.

A metric  $d$  of the metric space  $(X, d)$  is called *intrinsic* if for every  $P, Q \in X$

$$d(P, Q) = \inf_{\mathcal{L}} \{ \text{Length}(\mathcal{L}) \},$$

where the inf is taken over all rectifiable curves  $\mathcal{L}$  joining the points  $P$  and  $Q$ , and  $\text{Length}(\mathcal{L})$  is the length of  $\mathcal{L}$  measured in the metric  $d$ .

A curve  $\mathcal{L}$  in a metric space  $(X, d)$  joining a pair of points  $A, B$  is called a *shortest arc* if its length is equal to  $d(A, B)$ .

A metric space is said to be *geodesically connected* or a *length space* if each pair of points in it can be joined by a shortest arc.

An  $\mathfrak{R}_K$  domain (also known as a  $CAT(K)$  space) of the metric space  $(X, d)$  is a metric space with the following properties:

- (i)  $\mathfrak{R}_K$  is a geodesically connected metric space.
- (ii) If  $K > 0$ , then the perimeter of each triangle in  $\mathfrak{R}_K$  is less than  $2\pi/\sqrt{K}$ .
- (iii)  $K$ -convexity: Each triangle  $\triangle ABC \subset \mathfrak{R}_K$  and its comparison triangle  $\triangle \overline{ABC}$  in the  $K$ -plane<sup>2</sup> have the  $CAT(K)$ -inequality:  $d(B, C) \leq d_{K\text{-plane}}(\overline{B}, \overline{C})$ , where  $\overline{D}$  is the point in the arc  $\overline{AC}$  such that  $d(A, D) = d_{K\text{-plane}}(\overline{A}, \overline{D})$ .

A metric space  $(X, d)$  is an Alexandrov space of curvature bounded from above by  $K$  if each point of  $X$  is contained in some neighborhood that is an  $\mathfrak{R}_K$  domain.

To see the candidates for such Alexandrov spaces of curvature bounded from above and relate the curvature in the sense of Alexandrov and the sectional curvature of a Riemannian manifold, we have the following theorem due to Alexandrov and Cartan.

**Theorem 1.1.** ([A1], [Ca]) *A smooth Riemannian manifold  $M$  is an Alexandrov space of curvature bounded from above by  $K$  if and only if the sectional curvature of  $M$  is  $\leq K$ .*

---

<sup>2</sup>The  $K$ -plane is the 2-dimensional model space of constant Gaussian curvature  $K$ , i.e.,  $\mathbf{R}^2$  if  $K = 0$ , the standard Euclidean 2-hemisphere  $\mathbf{S}_K$  of radius  $1/\sqrt{K}$  if  $K > 0$  and the hyperbolic plane of curvature  $K$  if  $K < 0$ . The comparison triangle means  $\triangle \overline{ABC}$  has the same length of corresponding side as  $\triangle ABC$ , measuring in respective metric. See [BBI].

**1.2. Energy of maps into metric spaces.** Let  $(\Sigma, g)$  be a  $n$ -dimensional compact Riemannian domain,  $d_\Sigma$  be the distance function on  $\Sigma$  induced by  $g$  and  $(X, d)$  be any complete metric space. A Borel measurable map  $f : \Sigma \rightarrow X$  is said to be in  $L^2(\Sigma, X)$  if

$$\int_{\Sigma} d^2(f(x), Q) d\mu < \infty$$

for some  $Q \in X$ . By the triangle inequality, this definition is independent of the choice of  $Q$ .

For  $\epsilon > 0$ , let  $\Sigma_\epsilon = \{\eta \in \Sigma : d_\Sigma(\eta, \partial\Sigma) > \epsilon\}$ ,  $S_\epsilon(\eta) = \{\xi \in \Sigma : d_\Sigma(\eta, \xi) = \epsilon\}$ ,  $d\sigma_{\eta, \epsilon}(\xi)$  be the  $(n-1)$ -dimensional surface measure on  $S_\epsilon(\eta)$  and  $w_n$  be the area form of the unit sphere. For  $u \in L^2(\Sigma, X)$ , construct an  $\epsilon$ -approximate energy function  $e_\epsilon : \Sigma \rightarrow \mathbf{R}$  by setting

$$(1.2) \quad e_\epsilon(\eta) = \begin{cases} \frac{1}{w_n} \int_{S_\epsilon(\eta)} \frac{d^2(u(\eta), u(\xi))}{\epsilon^2} \frac{d\sigma_{\eta, \epsilon}(\xi)}{\epsilon^{n-1}} & \text{for } \eta \in \Sigma_\epsilon, \\ 0 & \text{for } \eta \in \Sigma - \Sigma_\epsilon. \end{cases}$$

Define a linear functional  $E_\epsilon : C_c(\Sigma) \rightarrow \mathbf{R}$  on the set of continuous functions with compact support in  $\Sigma$  by setting

$$E_\epsilon(f) = \int_{\Sigma} f e_\epsilon d\mu.$$

**Definition 1.3.** ([KS], 1.3ii) The map  $u \in L^2(\Sigma, X)$  is said to have finite energy or equivalently  $u \in W^{1,2}(\Sigma, X)$  if

$$(1.4) \quad E^u = \sup_{0 \leq f \leq 1, f \in C_c(\Sigma)} \limsup_{\epsilon \rightarrow 0^+} E_\epsilon(f) < \infty.$$

The quantity  $E^u$  is defined to be the energy of the map  $u$ . It is shown in [KS] that if  $u$  has finite energy, then in fact there exists a function  $e(\eta) \in L^1(\Sigma)$  so that  $e_\epsilon(\eta) d\mu_g(\eta) \rightarrow e(\eta) d\mu_g(\eta)$  as measures. The function  $e(\eta)$  is called the energy density of  $u$  and we write it as  $|\nabla u|^2$  ( $|u'|^2$  when  $n = 1$ ) as an analogue of the Riemannian case. In particular

$$\text{Energy}(u) = E^u = \int_{\Sigma} |\nabla u|^2 d\mu \quad (= \int_{\Sigma} |u'|^2 d\mu \quad \text{when } n = 1).$$

**Remark 1.5.** By definition, if  $u, v \in W^{1,2}(\Sigma, X)$  then the pointwise distance function  $d(u, v) \in W^{1,2}(\Sigma, \mathbf{R})$  (see also theorem 1.12.2 of [KS]). For closed curves  $\alpha, \beta \in W^{1,2}(\mathbf{S}^1, X)$ , the fact that  $d(\alpha, \beta) \in W^{1,2}(\mathbf{S}^1, \mathbf{R})$  allows us to define the distance between  $\alpha$  and  $\beta$  in  $W^{1,2}(\mathbf{S}^1, X)$  by setting

$$\text{dist}(\alpha, \beta) = |d(\alpha, \beta)|_{W^{1,2}}.$$

Note that the Sobolev embedding  $C^0(\mathbf{S}^1, \mathbf{R}) \hookrightarrow W^{1,2}(\mathbf{S}^1, \mathbf{R})$  implies two  $W^{1,2}$  curves that are  $W^{1,2}$  close are also  $C^0$  close (cf.(3.13)).

## 2. LOCAL ENERGY CONVEXITY IN $CAT(K)$ SPACES

This section is devoted to prove Theorem 0.1. Equation (2.2iv) of [KS] already gave the case of  $K = 0$  (with any  $\rho > 0$ ). Our idea then follows from [KS] to prove the case of  $K > 0$ . We first provide a local distance convexity in the standard Euclidean 2-hemisphere  $\mathbf{S}_K$  and then apply Reshetnyak's majorization theorem to get the local energy convexity for  $W^{1,2}$  maps into a  $CAT(K)$  space with  $K \geq 0$ . In [BN], Berg and Nikolaev defined the

so-called  $K$ -quadrilateral cosine  $\text{cos}q_K$  in an Alexandrov space of curvature  $\leq K$  which has the property that  $|\text{cos}q_K| \leq 1$ . As we shall see in the following, this quantity is much related to the local distance convexity in  $\mathbf{S}_K$ .

**Lemma 2.1.** ([BN]) *Consider a quadruple  $\mathcal{Q} = \{A, B, C, D\}$  of order points (see Figure 1),  $A \neq B$  and  $C \neq D$ , in  $\mathbf{S}_K$  ( $\mathbf{R}^2$  if  $K = 0$ ). Let  $k = \sqrt{K}$  and  $d_{\mathbf{S}_K}$  be the distance function in  $\mathbf{S}_K$ . Let  $d_{\mathbf{S}_K}(A, B) = a$ ,  $d_{\mathbf{S}_K}(C, D) = b$ ,  $d_{\mathbf{S}_K}(A, D) = x$ ,  $d_{\mathbf{S}_K}(B, C) = y$ ,  $d_{\mathbf{S}_K}(A, C) = h$  and  $d_{\mathbf{S}_K}(B, D) = i$ . Then the limit of the  $K$ -quadrilateral cosine equals to the 0-quadrilateral cosine as  $K \rightarrow 0$ , i.e.,*

$$\begin{aligned} & \lim_{K \rightarrow 0} \text{cos}q_K(\overrightarrow{DA}, \overrightarrow{CB}) \\ &= \lim_{K \rightarrow 0} \frac{\cos ka + \cos ky \cos kx \cos kb + \cos ka \cos kb - \cos ky \cos kh - \cos kx \cos ki - \cos kh \cos ki}{(1 + \cos kb) \sin kx \sin ky} \\ &= \text{cos}q_0(\overrightarrow{DA}, \overrightarrow{CB}) = \frac{a^2 + b^2 - h^2 - i^2}{2xy}. \end{aligned}$$

Based on Lemma 2.1, the following lemma follows directly from an elementary computation (see Appendix C for the detailed computation).

**Lemma 2.2.** *For any  $x \in \mathbf{S}_K$ , there exists  $\tau = \tau(K) > 0$  such that if  $\{A, D, C, B\} \subset B_\tau(x) \subset \mathbf{S}_K$  is an ordered sequence and  $E, F$  are the mid-points of the shortest arcs  $AB$  and  $CD$  respectively, we have the following distance convexity:*

$$\frac{1}{4} (d_{\mathbf{S}_K}(A, B) - d_{\mathbf{S}_K}(C, D))^2 \leq d_{\mathbf{S}_K}^2(A, D) + d_{\mathbf{S}_K}^2(B, C) - 2d_{\mathbf{S}_K}^2(E, F).$$

We will next recall Reshetnyak's majorization theorem in 1968 for an Alexandrov space of curvature bounded from above, which is a far reaching generalization of the  $K$ -convexity that was established by Alexandrov.

**Theorem 2.3.** ([Re]) *Let  $(X, d)$  be an Alexandrov space of curvature  $\leq K$ . In an  $\mathfrak{R}_K$  domain of  $X$ , for every rectifiable closed curve  $\mathcal{L}$  with length less than  $2\pi/\sqrt{K}$  if  $K > 0$ , there is a convex domain  $\mathcal{V}$  in the  $K$ -plane and a map  $\varphi : \mathcal{V} \rightarrow \mathfrak{R}_K$  such that  $\varphi(\partial\mathcal{V}) = \mathcal{L}$ , the lengths of the corresponding arcs coincide, and  $d_{K\text{-plane}}(\eta, \xi) \geq d(\varphi(\eta), \varphi(\xi))$ , for  $\eta, \xi \in \mathcal{V}$ .*

**Remark 2.4.** In particular, for an ordered sequence of points  $\{A, D, C, B\}$  in an Alexandrov space of curvature bounded from above by  $K > 0$ , let  $0 \leq \lambda, \nu \leq 1$  be given. Define  $A_\lambda$  to be the point which is the fraction  $\lambda$  of the way from  $A$  to  $B$  (on the geodesic  $\gamma_{A,B}$ ). Let  $D_\nu$  be the point which is the fraction  $\nu$  of the way from  $D$  to  $C$  (along the opposite geodesic  $\gamma_{D,C}$ ). By Theorem 2.3, there exists an ordered sequence of points  $\{\bar{A}, \bar{D}, \bar{C}, \bar{B}\} \subset \mathbf{S}_K$  which are the consecutive vertices of a quadrilateral. We can construct the corresponding points in  $\mathbf{S}_K$  :

$$\bar{A}_\lambda = (1 - \lambda)\bar{A} + \lambda\bar{B}, \quad \bar{D}_\nu = (1 - \nu)\bar{D} + \nu\bar{C}.$$

Then by Theorem 2.3

$$\begin{aligned} d(A, B) &= d_{\mathbf{S}_K}(\bar{A}, \bar{B}), & d(C, D) &= d_{\mathbf{S}_K}(\bar{C}, \bar{D}), \\ d(A, D) &= d_{\mathbf{S}_K}(\bar{A}, \bar{D}), & d(B, C) &= d_{\mathbf{S}_K}(\bar{B}, \bar{C}), \\ d(A_\lambda, D_\nu) &\leq d_{\mathbf{S}_K}(\bar{A}_\lambda, \bar{D}_\nu). \end{aligned}$$

We call  $\{\overline{A}, \overline{D}, \overline{C}, \overline{B}\}$  the subembedding of  $\{A, D, C, B\}$ .

**Lemma 2.5.** *Let  $(X, d)$  be an Alexandrov space of curvature  $\leq K$ . In an  $\mathfrak{R}_K$  domain of  $x \in X$ , there exists  $\rho = \rho(x, K) > 0$  such that if  $\{A, D, C, B\} \subset B_\rho(x) \subset \mathfrak{R}_K$  is an ordered sequence and  $E, F$  are the mid-points of the shortest arcs  $AB$  and  $CD$  respectively, we have*

$$(2.6) \quad \frac{1}{4}(d(A, B) - d(C, D))^2 \leq d^2(A, D) + d^2(B, C) - 2d^2(E, F).$$

*Proof.* Equation (2.2iii) of [KS] gives the case of  $K = 0$  (with any  $\rho > 0$ ). For  $K > 0$ , let  $\varrho = \min\{d(x, y) \mid y \in \partial \mathfrak{R}_K\}$  and  $\rho = \min\{\tau/4, \varrho\}$  where  $\tau$  is from Lemma 2.2. Let  $\{\overline{A}, \overline{D}, \overline{C}, \overline{B}\} \subset \mathbf{S}_K$  be a subembedding of  $\{A, D, C, B\}$ . We see first of all that  $\overline{A}, \overline{D}, \overline{C}$  and  $\overline{B}$  have to be in a geodesic ball of radius at most  $4\rho \leq \tau$  in  $\mathbf{S}_K$  and thus satisfy the condition of Theorem 2.3. Then by Theorem 2.3 and Remark 2.4, letting  $\lambda = \nu = \frac{1}{2}$ , we obtain

$$(2.7) \quad \begin{aligned} \frac{1}{4}(d(A, B) - d(C, D))^2 &= \frac{1}{4}(d_{\mathbf{S}_K}(\overline{A}, \overline{B}) - d_{\mathbf{S}_K}(\overline{C}, \overline{D}))^2 \\ &\leq d_{\mathbf{S}_K}^2(\overline{A}, \overline{D}) + d_{\mathbf{S}_K}^2(\overline{B}, \overline{C}) - 2d_{\mathbf{S}_K}^2(\overline{A}_{\frac{1}{2}}, \overline{D}_{\frac{1}{2}}) \\ &\leq d^2(A, D) + d^2(B, C) - 2d^2(A_{\frac{1}{2}}, D_{\frac{1}{2}}), \end{aligned}$$

completing the proof.  $\square$

**Remark 2.8.** For general  $\lambda, \nu \in [0, 1]$ , a similar distance convexity still holds with coefficients in terms of  $\lambda$  and  $\nu$ .

*Proof.* (of **Theorem 0.1**) For  $u, v \in W^{1,2}(\Sigma, X)$  with images staying in  $B_\rho(x)$ , set  $\{A = u(\xi), B = v(\xi), C = v(\eta), D = u(\eta)\}$  as in Lemma 2.5, we have:

$$(2.9) \quad \frac{1}{4}(d(u(\eta), v(\eta)) - d(u(\xi), v(\xi)))^2 \leq d^2(u(\xi), u(\eta)) + d^2(v(\xi), v(\eta)) - 2d^2(w(\xi), w(\eta)),$$

where  $w(\xi) = \frac{u+v}{2}(\xi)$  is the mid-point of the geodesic connecting  $u(\xi)$  and  $v(\xi)$ .

Multiplying (2.9) by  $f(\eta)$  (where  $0 \leq f \leq 1$  and  $f \in C_c(\Sigma)$ ), averaging on the subset  $\{|\eta - \xi| < \varepsilon\}$  of  $\Sigma \times \Sigma$  and integrating over  $\Sigma$  (as in 1.3 of [KS] and see (1.2)), then first of all we conclude that  $w \in W^{1,2}(\Sigma, X)$ . By theorem 1.6.2 and theorem 1.12.2 of [KS] we obtain that for any  $f \in C_c(\Sigma), 0 \leq f \leq 1$ :

$$\frac{1}{4} \int_{\Sigma} f |\nabla d(u, v)|^2 \leq \int_{\Sigma} f |\nabla u|^2 + \int_{\Sigma} f |\nabla v|^2 - 2 \int_{\Sigma} f |\nabla w|^2.$$

Hence by definition (1.4), we have an analogue to (2.2iv) of [KS]:

$$(2.10) \quad \frac{1}{4} \int_{\Sigma} |\nabla d(u, v)|^2 \leq E^u + E^v - 2E^w.$$

$\square$

**Remark 2.11.** An immediate corollary of (2.10) is the uniqueness of the solution to the Dirichlet problem into a  $CAT(K)$  (with  $K > 0$ ) space with small image assumption, see [Se] and cf. theorem 2.2 of [KS].

**Corollary 2.12.** ([Se]) *Let  $(\Sigma, g)$  be a Lipschitz Riemannian domain and  $(X, d)$  be an Alexandrov space of curvature bounded from above by  $K > 0$ . Fix a point  $Q \in X$ , Let  $\phi \in W^{1,2}(\Sigma, X)$  with  $\phi(\Sigma) \subset B_\rho(Q)$  where  $\rho = \rho(Q, K)$  is given by Theorem 0.1. Define*

$$W_\phi^{1,2} = \{u \in W^{1,2}(\Sigma, X) \mid u(\Sigma) \subset B_\rho(Q) \text{ and } \text{tr}(u) = \text{tr}(\phi)\}.$$

*Then there exists a unique  $u \in W_\phi^{1,2}$  which satisfies*

$$E^u = \int_\Sigma |\nabla u|^2 d\mu = E_0 = \inf_{v \in W_\phi^{1,2}} E^v.$$

### 3. EXISTENCE OF GOOD SWEEPOUTS BY CURVES

Throughout the rest of this paper, we will let  $(X, d)$  be a closed Alexandrov space of curvature bounded from above by some  $K$ . Using the compactness of  $X$ , we let

$$(3.1) \quad \rho = \inf_{x \in X} \{\rho(x, K)\} > 0,$$

where  $\rho(x, K)$  is as in Theorem 0.1. Fix a large positive integer  $L$  and let  $\Lambda$  denote the space of piecewise linear maps (constant speed geodesics) from  $\mathbf{S}^1$  to  $X$  with exactly  $L^2$  breaks (possibly with unnecessary breaks) such that the length of each geodesic segment is at most  $\rho$  defined by (3.1), parametrized by a (constant) multiple of arclength and with Lipschitz bound<sup>3</sup>  $L$ . Let  $G \subset \Lambda$  denote the set of (possibly self-intersecting) closed geodesics in  $X$  of length at most  $\rho L^2$ . (The constant speed of a curve in  $\Lambda$  is equal to its length divided by  $2\pi$ ; and its energy is equal to its length squared divided by  $2\pi$ . In other words, energy and length are essentially equivalent, see (3.5) and (3.13)).

**3.1. The width.** In [CM1], Colding and Minicozzi introduced two crucial geometric concepts: sweepout and width. We will recall and extend these definitions to a closed Alexandrov space of curvature bounded from above.

**Definition 3.2.** A continuous map  $\sigma : \mathbf{S}^1 \times [-1, 1] \rightarrow X$  is called a *sweepout* in  $X$ , if for each  $s$  the map  $\sigma(\cdot, s)$  is in  $W^{1,2}(\mathbf{S}^1, X)$ , the map  $s \rightarrow \sigma(\cdot, s)$  is continuous (in the induced metric as in Remark 1.5) from  $[-1, 1]$  to  $W^{1,2}(\mathbf{S}^1, X)$ , and finally  $\sigma$  maps  $\mathbf{S}^1 \times \{-1\}$  and  $\mathbf{S}^1 \times \{1\}$  to points.

Let  $\Omega$  be the set of sweepouts in  $X$ . Given a map  $\hat{\sigma} \in \Omega$ , the homotopy class  $\Omega_{\hat{\sigma}}$  is defined to be the set of maps  $\sigma \in \Omega$  that are homotopic to  $\hat{\sigma}$  through maps in  $\Omega$ .

**Definition 3.3.** The *width*  $W = W(\hat{\sigma})$  associated to the homotopy class  $\Omega_{\hat{\sigma}}$  is defined by taking the infimum of the maximum of the energy of each slice. That is, set

$$(3.4) \quad W = \inf_{\sigma \in \Omega_{\hat{\sigma}}} \max_{s \in [-1, 1]} \text{Energy}(\sigma(\cdot, s)),$$

---

<sup>3</sup>Note that a  $W^{1,2}$  curve is also a  $C^{1/2}$  curve but not necessarily Lipschitz continuous, here the Lipschitz bound denotes the bound of the speed and is equivalent to the square of the  $C^{1/2}$  bound of the curve. See (3.13).



where the energy is the energy of maps into metric spaces given in subsection 1.2, namely, (3.5)

$$\text{Energy}(\sigma(\cdot, s)) = \sup_{\substack{f \in C_c(\mathbf{S}^1) \\ 0 \leq f \leq 1}} \limsup_{\epsilon \rightarrow 0^+} \int_{\mathbf{S}^1} f \left( \frac{d^2(\sigma(\eta - \epsilon, s), \sigma(\eta, s)) + d^2(\sigma(\eta, s), \sigma(\eta + \epsilon, s))}{2\epsilon^2} \right) d\eta.$$

We write  $\text{Energy}(\sigma(\cdot, s)) = E(\sigma(\cdot, s)) = \int_{\mathbf{S}^1} |\sigma'(x, s)|^2 dx$ . We shall see that a sweepout in  $X$  induces a map from sphere  $\mathbf{S}^2$  to  $X$  and the width is always non-negative and is positive if  $\tilde{\sigma}$  is in a non-trivial homotopy class<sup>4</sup>.

**Remark 3.6.** The  $\epsilon$ -approximate length function of  $\sigma$  converges to a  $L^1$  function, which coincides with the speed function of  $\sigma$ , as  $\epsilon \rightarrow 0^+$ , namely (see lemma 1.9.3 of [KS]),

$$\lim_{\epsilon \rightarrow 0^+} \frac{d(\sigma(\eta - \epsilon), \sigma(\eta)) + d(\sigma(\eta), \sigma(\eta + \epsilon))}{2\epsilon} = |\sigma'|(\eta) \quad \text{a.e. } \eta \in \mathbf{S}^1.$$

Throughout the rest of this paper we will use  $|\sigma'|$  to denote the speed function of a curve  $\sigma$  in  $X$ .

**3.2. Curve shortening  $\Psi$ .** The curve shortening is a map  $\Psi : \Lambda \rightarrow \Lambda$  so that (see also section 2 of [Cr])

- (1)  $\Psi(\gamma)$  is homotopic to  $\gamma$  and  $\text{Length}(\Psi(\gamma)) \leq \text{Length}(\gamma)$ .
- (2)  $\Psi(\gamma)$  depends continuously on  $\gamma$ .
- (3) There is a continuous function  $\phi : [0, \infty) \rightarrow [0, \infty)$  with  $\phi(0) = 0$  so that

$$(3.7) \quad \text{dist}^2(\gamma, \Psi(\gamma)) \leq \phi \left( \frac{\text{Length}^2(\gamma) - \text{Length}^2(\Psi(\gamma))}{\text{Length}^2(\Psi(\gamma))} \right).$$

- (4) Given  $\epsilon > 0$ , there exists  $\delta > 0$  so that if  $\gamma \in \Lambda$  with  $\text{dist}(\gamma, G) \geq \epsilon$ , then  $\text{Length}(\Psi(\gamma)) \leq \text{Length}(\gamma) - \delta$ .

We will use local linear replacement to define the curve shortening map  $\Psi$  which is identical to [CM1]: fix a partition of  $\mathbf{S}^1$  by choosing  $2L^2$  consecutive evenly spaced points<sup>5</sup>

$$(3.8) \quad x_0, x_1, x_2, \dots, x_{2L^2} = x_0 \in \mathbf{S}^1, \quad \text{so that } |x_j - x_{j+1}| = \frac{\pi}{L^2} \leq \frac{\rho}{2L}.$$

$\Psi(\gamma)$  is given in the following three steps:

*Step 1:* Replace  $\gamma$  on each *even* interval, i.e.,  $[x_{2j}, x_{2j+2}]$ , by the linear map with the same endpoints to get a piecewise linear curve  $\gamma_e : \mathbf{S}^1 \rightarrow X$ . Namely, for each  $j$ , we let  $\gamma_e|_{[x_{2j}, x_{2j+2}]}$  be the unique shortest (constant speed) geodesic from  $\gamma(x_{2j})$  to  $\gamma(x_{2j+2})$ .

*Step 2:* Replace  $\gamma_e$  on each *odd* interval by the linear map with the same endpoints to get the piecewise linear curve  $\gamma_o : \mathbf{S}^1 \rightarrow X$ .

*Step 3:* Reparametrize  $\gamma_o$  (fixing  $\gamma_o(x_0)$ ) to get the desired constant speed curve  $\Psi(\gamma) : \mathbf{S}^1 \rightarrow X$ .

It is easy to see that  $\Psi$  maps  $\Lambda$  to  $\Lambda$  and has property (1); cf. section 2 of [Cr]. The proof of properties (2), (3) and (4) for  $\Psi$  is virtually the same as [CM1]. We shall remark that there

<sup>4</sup>A particularly interesting example is when  $X$  is a topological 2-sphere with  $\pi_1(X) = \{0\}$  and the map induced by a sweepout from  $\mathbf{S}^2$  to  $X$  has degree one. In this case, the width is positive and realized by a non-trivial closed geodesic with index 1, see footnote 2 of [CM1].

<sup>5</sup>Note that this is not necessarily where the piecewise linear maps have breaks.

is a difficulty in the proofs of these properties: the second fundamental form (smoothness) of the manifold is used in the Riemannian case in [CM1], while we don't have the smoothness in an Alexandrov space of curvature bounded above. But note that the local energy convexity in Theorem 0.1 requires only that the two curves both stay in a small region, while the key lemma 4.2 in [CM1] requires that the two curves have the same endpoints. This fact allows us to get around this difficulty (see (A.7)). For the completeness of this paper, we include the proofs of properties (3) and (4) in Appendix A and the proof of property (2) in Appendix B. Throughout the rest of this section, we will assume these properties of  $\Psi$  and use them to prove the main theorem.

Combining properties (3) and (4) of  $\Psi$ , we have the following key lemma, which is crucial in producing the desired sequence of good sweepouts.

**Lemma 3.9.** *Given  $W \geq 0$  and  $\epsilon > 0$ , there exists  $\delta > 0$  so that if  $\gamma \in \Lambda$  and*

$$(3.10) \quad 2\pi(W - \delta) < \text{Length}^2(\Psi(\gamma)) \leq \text{Length}^2(\gamma) < 2\pi(W + \delta),$$

*then  $\text{dist}(\Psi(\gamma), G) < \epsilon$ .*

*Proof.* If  $W \leq \epsilon^2/6$ , then  $\delta = \epsilon^2/6$  gives  $\text{Length}(\Psi(\gamma)) \leq 2\epsilon$ . This tells us the bound on distance of  $\Psi(\gamma)$  to a point curve (e.g., its mid-point) which is a trivial closed geodesic in  $G$ .

Assume next that  $W > \epsilon^2/6$ . The triangle inequality gives

$$(3.11) \quad \text{dist}(\Psi(\gamma), G) \leq \text{dist}(\Psi(\gamma), \gamma) + \text{dist}(\gamma, G).$$

Since  $\Psi$  does not decrease the length of  $\gamma$  by much by the assumption, property (4) of  $\Psi$  bounds  $\text{dist}(\gamma, G)$  by  $\epsilon/2$  as long as  $\delta$  is sufficiently small. Similarly, property (3) of  $\Psi$  allows us to bound  $\text{dist}(\Psi(\gamma), \gamma)$  by  $\epsilon/2$  as long as  $\delta$  is sufficiently small.  $\square$

**3.3. Defining the good sweepouts.** Choose a sequence of maps  $\hat{\sigma}^j \in \Omega_\delta$  with

$$(3.12) \quad \max_{s \in [-1, 1]} \text{Energy}(\hat{\sigma}^j(\cdot, s)) < W + \frac{1}{j}.$$

Observe that (3.12) and the Cauchy-Schwarz inequality imply a uniform bound for the length and uniform  $C^{1/2}$  continuity for the slices, that are both independent of  $j$  and  $s$ . They follow immediately from the following: for any small  $\delta > 0$ ,  $[x, y] \subset [0, 2\pi]$  we pick  $f \in C_c([0, 2\pi])$ ,  $0 \leq f \leq 1$ , with  $f = 1$  on  $(x, y)$  and  $\text{supp}(f) \subset [x - \delta, y + \delta] \subset [0, 2\pi]$ , then

$$(3.13) \quad \begin{aligned} & d^2(\hat{\sigma}^j(x, s), \hat{\sigma}^j(y, s)) \leq \text{Length}^2(\hat{\sigma}^j(\cdot, s)|_{[x, y]}) \\ &= \lim_{\delta \rightarrow 0^+} \limsup_{\epsilon \rightarrow 0^+} \left( \int_{x-\delta}^{y+\delta} f \frac{d(\hat{\sigma}^j(\eta - \epsilon, s), \hat{\sigma}^j(\eta, s)) + d(\hat{\sigma}^j(\eta, s), \hat{\sigma}^j(\eta + \epsilon, s))}{2\epsilon} d\eta \right)^2 \\ &\leq |y - x| \lim_{\delta \rightarrow 0^+} \limsup_{\epsilon \rightarrow 0^+} \int_{x-\delta}^{y+\delta} f^2 \left( \frac{d^2(\hat{\sigma}^j(\eta - \epsilon, s), \hat{\sigma}^j(\eta, s)) + d^2(\hat{\sigma}^j(\eta, s), \hat{\sigma}^j(\eta + \epsilon, s))}{2\epsilon^2} \right) d\eta \\ &= |y - x| \text{Energy}(\hat{\sigma}^j(\cdot, s)|_{[x, y]}) \leq |y - x|(W + 1). \end{aligned}$$

In order to get started and be able to use the properties of  $\Psi$ , we would like all the initial curves to be in  $\Lambda$ . We will replace the  $\hat{\sigma}^j$ 's by sweepouts  $\sigma^j$  that, in addition to satisfying (3.12), also satisfy that the slices  $\sigma^j(\cdot, s)$  are in  $\Lambda$ . We will do this by using local linear

replacement similar to the construction of  $\Psi$ . Namely, the uniform  $C^{1/2}$  bound for the slices allows us to fix a partition of points  $y_0, \dots, y_N = y_0$  in  $\mathbf{S}^1$  so that each interval  $[y_i, y_{i+1}]$  is always mapped to a geodesic ball in  $X$  of radius at most  $\rho$ . Next, for each  $s$  and each  $j$ , we replace  $\hat{\sigma}^j(\cdot, s)|_{[y_i, y_{i+1}]}$  by the linear map (geodesic) with the same endpoints and call the resulting map  $\tilde{\sigma}^j(\cdot, s)$ . Reparametrize  $\tilde{\sigma}^j(\cdot, s)$  to have constant speed to get  $\sigma^j(\cdot, s)$ . It is easy to see that each  $\sigma^j(\cdot, s)$  satisfies (3.12). Furthermore, the length bound for  $\sigma^j(\cdot, s)$  also gives a uniform Lipschitz (speed) bound for the linear maps; let  $L$  be this bound and  $N \leq L^2$ .

We can see from the proof of property (2) for  $\Psi$  in Appendix B that  $\sigma^j$  is continuous in the transversal direction (i.e. with respect to  $s$ ) and homotopic to  $\hat{\sigma}$  in  $\Omega$ , cf. [B1], [B2], section 2 of [Cr] and appendix B of [CM1].

Finally, applying the replacement map  $\Psi$  to each  $\sigma^j(\cdot, s)$  gives a new sequence of sweepouts  $\gamma^j = \Psi(\sigma^j)$ . ( $\Psi$  depends continuously on  $s$  and preserves the homotopy class  $\Omega_{\hat{\sigma}}$ ; it is clear that  $\Psi$  fixes the constant maps at  $s = \pm 1$ .)

**3.4. Almost maximal implies almost critical.** We will show that the sequence  $\gamma^j = \Psi(\sigma^j)$  of sweepouts is tight in the sense of the Introduction. Namely, we have the following main theorem.

**Theorem 3.14.** *Given  $W \geq 0$  and  $\epsilon > 0$ , there exists  $\delta > 0$  so that if  $j > 1/\delta$  and for some  $s_0$*

$$(3.15) \quad 2\pi \text{Energy}(\gamma^j(\cdot, s_0)) = \text{Length}^2(\gamma^j(\cdot, s_0)) > 2\pi(W - \delta),$$

*then for this  $j$  we have  $\text{dist}(\gamma^j(\cdot, s_0), G) < \epsilon$ .*

*Proof.* Let  $\delta$  be given by Lemma 3.9. By (3.15), (3.12), and using that  $j > 1/\delta$ , we get

$$(3.16) \quad 2\pi(W - \delta) < \text{Length}^2(\gamma^j(\cdot, s_0)) \leq \text{Length}^2(\sigma^j(\cdot, s_0)) < 2\pi(W + \delta).$$

Thus, since  $\gamma^j(\cdot, s_0) = \Psi(\sigma^j(\cdot, s_0))$ , Lemma 3.9 gives  $\text{dist}(\gamma^j(\cdot, s_0), G) < \epsilon$ .  $\square$

#### 4. GENERALIZED BIRKHOFF-LYUSTERNIK THEOREM

**4.1. Parameter spaces.** Instead of using the interval  $[-1, 1]$ , as parameter space for the circles in the definition of sweepout (see Definition 3.2) and assuming that the curves start and end in point curves, one could have use any compact set  $\mathcal{P}$  and require that the curves are constant on  $\partial\mathcal{P}$  (or that  $\partial\mathcal{P} = \emptyset$ ). Then we let  $\Omega^{\mathcal{P}}$  be the set of continuous maps  $\sigma : \mathbf{S}^1 \times \mathcal{P} \rightarrow X$  so that for each  $s \in \mathcal{P}$  the curve  $\sigma(\cdot, s)$  is in  $W^{1,2}(\mathbf{S}^1, X)$ , the map  $s \rightarrow \sigma(\cdot, s)$  is continuous from  $\mathcal{P}$  to  $W^{1,2}(\mathbf{S}^1, X)$  and finally  $\sigma$  maps  $\partial\mathcal{P}$  to point curves. Given a map  $\hat{\sigma} \in \Omega^{\mathcal{P}}$ , the homotopy class  $\Omega_{\hat{\sigma}}^{\mathcal{P}} \subset \Omega^{\mathcal{P}}$  is defined to be the set of maps  $\sigma \in \Omega^{\mathcal{P}}$  that are homotopic to  $\hat{\sigma}$  through maps in  $\Omega^{\mathcal{P}}$ . Finally the *width*  $W = W(\hat{\sigma})$  is

$$(4.1) \quad W = \inf_{\sigma \in \Omega_{\hat{\sigma}}^{\mathcal{P}}} \max_{s \in \mathcal{P}} \text{Energy}(\sigma(\cdot, s)).$$

Theorem 3.14 holds for these general parameter spaces and the proof is virtually the same.

**4.2. Generalized Birkhoff-Lyusternik theorem.** The following is devoted to the proof of Theorem 0.3.

*Proof.* We will divide our proof into two cases. In the case of the fundamental group  $\pi_1(X) \neq 0$ , we can choose a non-contractible closed curve  $\sigma_0 : \mathbf{S}^1 \rightarrow X$ . Then by the definition of width (see Definition 3.3) we see that  $W > 0$ . It follows immediately from Theorem 3.14 that there exists at least one non-trivial closed geodesic in  $X$  if we apply the width-sweepout construction procedure as in section 3.

In the case of  $\pi_1(X) = 0$ , i.e.,  $X$  is simply-connected (or 1-connected), then it's well known that  $H_1(X) \cong \pi_1(X)/[\pi_1(X), \pi_1(X)]$  (see e.g. [Hat, Theorem 2A.1], page 166) and thus  $H_1(X) = 0$ . Then by the assumption of the theorem, there exists the first nonzero  $k_1$ -th homology group  $H_{k_1}(X) \neq 0$  for some integer  $k_1$  with  $2 \leq k_1 \leq k_0$ . Therefore by the Hurewicz theorem which states that the first nonzero homotopy and homology groups of a simply-connected space occur in the same dimension and are isomorphic (see e.g. [Hat, Theorem 4.32], page 366), we have

$$\pi_{k_1}(X) \cong H_{k_1}(X) \neq 0.$$

Thus there is a non-contractible map

$$\omega_0 : \mathbf{S}^{k_1} \rightarrow X$$

from the  $k_1$ -sphere  $\mathbf{S}^{k_1}$  to  $X$  for  $k_1 \geq 2$ . Note that  $\mathbf{S}^{k_1}$  is equivalent to  $\mathbf{S}^1 \times \bar{B}^{k_1-1} / \sim$ , where  $\sim$  is the equivalence relation  $(\theta_1, y) \sim (\theta_2, y)$  where  $\theta_1, \theta_2 \in \mathbf{S}^1$  and  $y \in \partial \bar{B}^{k_1-1}$ . Here  $\bar{B}^{k_1-1}$  is the closed unit ball in  $\mathbf{R}^{k_1-1}$ . We use this decomposition of  $\mathbf{S}^{k_1}$  to define the width of  $X$ .

Take  $\mathcal{P} = \bar{B}^{k_1-1}$  as the parameter space as in subsection 4.1 and define the width  $W$  as in (4.1). We see directly from the fact that  $\omega_0$  is non-contractible that  $W > 0$ . Again, it follows immediately from Theorem 3.14 that there exists at least one non-trivial closed geodesic in  $X$  if we apply the width-sweepout construction procedure as in section 3.  $\square$

## APPENDIX A. ESTABLISHING PROPERTIES (3) AND (4) OF $\Psi$

To prove property (3) of  $\Psi$ , we will use the following equivalent way to construct  $\Psi(\gamma)$  :

- (A<sub>1</sub>) Follow Step 1 to get  $\gamma_e$ .
- (B<sub>1</sub>) Reparametrize  $\gamma_e$  (fixing the image of  $x_0$ ) to get the constant speed curve  $\tilde{\gamma}_e$ . This reparametrization moves the points  $x_j$  to new points  $\tilde{x}_j$  (i.e.,  $\gamma_e(x_j) = \tilde{\gamma}_e(\tilde{x}_j)$ ).
- (A<sub>2</sub>) Do linear replacement on the odd  $\tilde{x}_j$  intervals to get  $\tilde{\gamma}_o$ .
- (B<sub>2</sub>) Reparametrize  $\tilde{\gamma}_o$  (fixing the image of  $x_0$ ) to get the constant speed curve  $\Psi(\gamma)$ .

One sees easily that this gives the same curve since  $\tilde{\gamma}_o$  is just a reparametrization of  $\gamma_o$ . We also see that each of the four steps is energy non-increasing<sup>6</sup>. Thus property (3) follows from the triangle inequality once we bound  $\text{dist}(\gamma, \gamma_e)$  and  $\text{dist}(\gamma_e, \tilde{\gamma}_e)$  in terms of the decrease in length (as well as the analogs for steps (A<sub>2</sub>) and (B<sub>2</sub>)).

The bound on  $\text{dist}(\gamma, \gamma_e)$  follows directly from the next corollary of Theorem 0.1.

<sup>6</sup>This is obvious for the linear replacements, since linear maps minimize energy. It follows from (3.13) for the reparametrizations, since for a curve  $\sigma : \mathbf{S}^1 \rightarrow X$  we have  $\text{Length}^2(\sigma) \leq 2\pi \text{Energy}(\sigma)$ , with equality if and only if its speed is a constant  $= \text{Length}(\sigma)/(2\pi)$  almost everywhere.

**Corollary A.1.** *There exists  $C$  so that if  $I$  is an interval of length at most  $\rho/L$ ,  $\sigma_1 : I \rightarrow X$  is a curve with Lipschitz bound  $L$ , and  $\sigma_2 : I \rightarrow X$  is the minimizing geodesic with the same endpoints, then*

$$\text{dist}^2(\sigma_1, \sigma_2) \leq C (E^{\sigma_1} - E^{\sigma_2}).$$

*Proof.* Let  $\Sigma = I \subset \mathbf{S}^1$  and note that  $w = \frac{\sigma_1 + \sigma_2}{2}$  has the same end points as  $\sigma_1$  and  $\sigma_2$ . Since  $d(\sigma_1, \sigma_2) \in W_0^{1,2}(I, \mathbf{R})$  (see theorem 1.12.2 of [KS]) and from Theorem 0.1, the Poincaré inequality and (2.10) imply

$$\text{dist}^2(\sigma_1, \sigma_2) \leq C(I) \int_I |\nabla d(\sigma_1, \sigma_2)|^2 d\mu \leq C (E^{\sigma_1} - E^{\sigma_2}),$$

where we used the minimality of  $\sigma_2$ . □

Applying Corollary A.1 on each of  $L^2$  intervals in step  $(A_1)$ , we get that

$$(A.2) \quad \text{dist}^2(\gamma, \gamma_e) \leq C (E^\gamma - E^{\gamma_e}) \leq \frac{C}{2\pi} (\text{Length}^2(\gamma) - \text{Length}^2(\Psi(\gamma))).$$

This gives the desired bound on  $\text{dist}(\gamma, \gamma_e)$  since  $\text{Length}(\Psi(\gamma)) \leq \rho L^2$ .

To bound  $\text{dist}(\gamma_e, \tilde{\gamma}_e)$ , we will use that  $\gamma_e$  is just the composition  $\tilde{\gamma}_e \circ P$ , where  $P : \mathbf{S}^1 \rightarrow \mathbf{S}^1$  is a monotone piecewise linear map<sup>7</sup> and let  $L$  be its Lipschitz bound as well. Using that the (piecewise constant) speed of  $\gamma_e$  is  $|\gamma'_e| = |(\tilde{\gamma}_e \circ P)'| = |\tilde{\gamma}'_e \circ P| \cdot |P'| \leq L^2$  (Note:  $|\tilde{\gamma}'_e \circ P|(x)$  denotes the speed of  $\tilde{\gamma}_e$  at point  $P(x)$ ) and the (constant) speed of  $\tilde{\gamma}_e = |\tilde{\gamma}'_e| = \text{Length}(\tilde{\gamma}_e)/(2\pi) \leq L$  (away from the breaks), and also that the integral of  $P'$  is  $2\pi$ , we have

$$(A.3) \quad \begin{aligned} \int_{\mathbf{S}^1} (P' - 1)^2 &= \int_{\mathbf{S}^1} (P')^2 - 2\pi = \int_{\mathbf{S}^1} \left( \frac{|\gamma'_e|}{|\tilde{\gamma}'_e \circ P|} \right)^2 - 2\pi = \frac{4\pi^2}{\text{Length}^2(\tilde{\gamma}_e)} \int_{\mathbf{S}^1} |\gamma'_e|^2 - 2\pi \\ &= 2\pi \frac{\text{Energy}(\gamma_e) - \text{Energy}(\tilde{\gamma}_e)}{\text{Energy}(\tilde{\gamma}_e)} \leq 2\pi \frac{\text{Energy}(\gamma) - \text{Energy}(\Psi(\gamma))}{\text{Energy}(\Psi(\gamma))}. \end{aligned}$$

Now divide  $\mathbf{S}^1$  into two sets,  $S_1$  and  $S_2$ , where  $S_1$  is the set of points within distance  $(\pi \int_{\mathbf{S}^1} |P' - 1|^2)^{1/2}$  of a break point for  $\tilde{\gamma}_e$ . Since  $P(x_0) = x_0$ , we have  $|P(x) - x| \leq (\pi \int_{\mathbf{S}^1} |P' - 1|^2)^{1/2}$ . Since  $\gamma_e$  and  $\tilde{\gamma}_e$  agree at  $x_0 = x_{2L^2}$ , the Wirtinger inequality<sup>8</sup> bounds  $\text{dist}^2(\gamma_e, \tilde{\gamma}_e)$  in terms of

$$(A.4) \quad \int_{\mathbf{S}^1} |\nabla d(\tilde{\gamma}_e \circ P, \tilde{\gamma}_e)|^2 \leq \int_{S_1} (|(\tilde{\gamma}_e \circ P)'| + |\tilde{\gamma}'_e|)^2 + \int_{S_2} |\nabla d(\tilde{\gamma}_e \circ P, \tilde{\gamma}_e)|^2,$$

where we used that the fact (3) in the proof of Lemma 2.2 (see last part of Appendix C) implies

$$(A.5) \quad \int_{S_1} |\nabla d(\tilde{\gamma}_e \circ P, \tilde{\gamma}_e)|^2 \leq \int_{S_1} (|(\tilde{\gamma}_e \circ P)'| + |\tilde{\gamma}'_e|)^2.$$

<sup>7</sup>The map  $P$  is Lipschitz, but the inverse map  $P^{-1}$  may not be if  $\gamma_e$  is constant on an interval.

<sup>8</sup>The Wirtinger inequality is just the usual Poincaré inequality which bounds the  $L^2$  norm in terms of the  $L^2$  norm of the derivative; i.e.,  $\int_0^{2\pi} f^2 dt \leq 4 \int_0^{2\pi} (f')^2 dt$  provided  $f(0) = f(2\pi) = 0$ .

We will bound both terms on the right hand side of (A.4) in terms of  $\int_{\mathbf{S}^1} |P' - 1|^2$  and then appeal to (A.3). To bound the first term, we have

$$(A.6) \quad \int_{S_1} (|(\tilde{\gamma}_e \circ P)'| + |\tilde{\gamma}'_e|)^2 \leq (L^2 + L)^2 \text{Length}(S_1) \leq 8L^6 \left( \pi \int_{\mathbf{S}^1} |P' - 1|^2 \right)^{1/2}.$$

We see that if  $(\pi \int_{\mathbf{S}^1} |P' - 1|^2)^{1/2} \geq \frac{\pi}{2L^2} = \frac{|x_j - x_{j+1}|}{2}$ , we are done since in this case  $S_2 = \emptyset$ .

On the other hand, suppose  $(\pi \int_{\mathbf{S}^1} |P' - 1|^2)^{1/2} < \frac{\pi}{2L^2}$ ; note that if  $x \in S_2$ , then  $\tilde{\gamma}_e(x)$  and  $\tilde{\gamma}_e \circ P(x)$  stay within the  $\rho$ -neighborhood between two break points (although  $\tilde{\gamma}_e$  and  $\tilde{\gamma}_e \circ P$  might not have the same endpoints) and  $|\tilde{\gamma}'_e \circ P| = |\tilde{\gamma}'_e| \leq L$  in this neighborhood. Thus, we can bound the second term by applying Theorem 0.1. Namely, by summing up the integral over each piece of  $S_2$ , we have

$$(A.7) \quad \begin{aligned} & \frac{1}{4} \int_{S_2} |\nabla d(\tilde{\gamma}_e \circ P, \tilde{\gamma}_e)|^2 \\ & \leq \text{Energy}((\tilde{\gamma}_e \circ P)|_{S_2}) + \text{Energy}(\tilde{\gamma}_e|_{S_2}) - 2\text{Energy}\left(\left(\frac{\tilde{\gamma}_e \circ P + \tilde{\gamma}_e}{2}\right)|_{S_2}\right) \\ & = \int_{S_2} |(\tilde{\gamma}'_e \circ P)P'|^2 + \int_{S_2} |\tilde{\gamma}'_e|^2 - 2 \int_{S_2} \left(\frac{|(\tilde{\gamma}'_e \circ P)P'| + |\tilde{\gamma}'_e|}{2}\right)^2 \\ & = \int_{S_2} \frac{(|\tilde{\gamma}'_e \circ P| \cdot |P'| - |\tilde{\gamma}'_e|)^2}{2} \leq \frac{L^2}{2} \int_{\mathbf{S}^1} |P' - 1|^2, \end{aligned}$$

completing the proof of property (3).

To prove property (4) of  $\Psi$ , suppose it is not true, namely, there exist  $\epsilon > 0$  and a sequence  $\gamma_j \in \Lambda$  with  $\text{Energy}(\Psi(\gamma_j)) \geq \text{Energy}(\gamma_j) - 1/j$  and  $\text{dist}(\gamma_j, G) \geq \epsilon > 0$ ; note that the second condition implies a positive lower bound for  $\text{Energy}(\gamma_j)$ . Observe next that the space  $\Lambda$  is compact<sup>9</sup> and, thus, a subsequence of the  $\gamma_j$ 's must converge to some  $\gamma \in \Lambda$ . Since property (3) implies that  $\text{dist}(\gamma_j, \Psi(\gamma_j)) \rightarrow 0$ , the  $\Psi(\gamma_j)$ 's also converge to  $\gamma$ . The continuity of  $\Psi$ , i.e., property (2) of  $\Psi$ , then implies that  $\Psi(\gamma) = \gamma$ . However, this implies that  $\gamma \in G$  since the only fixed points of  $\Psi$  are (possibly self-intersected) closed geodesics. This last fact follows immediately from Corollary A.1 and (A.3). However, this would contradict that the  $\gamma_j$ 's remain a fixed distance from any such closed geodesic, completing the proof of (4).

## APPENDIX B. THE CONTINUITY OF $\Psi$

**Lemma B.1.** *Let  $\gamma : \mathbf{S}^1 \rightarrow (X, d)$  be a  $W^{1,2}$  map with  $\text{Energy}(\gamma) \leq \rho L$ . If  $\gamma_e$  and  $\tilde{\gamma}_e$  are given by applying (A<sub>1</sub>) and (B<sub>1</sub>) to  $\gamma$ , then the map  $\gamma \rightarrow \tilde{\gamma}_e$  is continuous from  $W^{1,2}$  to  $\Lambda$  equipped with the  $W^{1,2}$  norm as in Remark 1.5.*

*Proof.* It follows from (3.13) and the energy bound that  $d(\gamma(x_{2j}), \gamma(x_{2j+2})) \leq \rho$  for each  $j$ , and thus we can apply step (A<sub>1</sub>). Now suppose that  $\gamma^1$  and  $\gamma^2$  are non-constant curves in  $\Lambda$  (continuity at the constant maps is obvious). For  $i = 1, 2$  and  $j = 0, 1, 2, \dots, L^2 - 1$ , let  $a_j^i = d(\gamma^i(x_{2j}), \gamma^i(x_{2j+2}))$ . Let  $S^i = \frac{1}{2\pi} \sum_{j=0}^{L^2-1} a_j^i$  be the speed of  $\tilde{\gamma}_e^i$ , so that  $|(\tilde{\gamma}_e^i)'| = S^i$  except at the  $L^2$  break points. Since, by Remark 1.5,  $W^{1,2}$  close curves are also  $C^0$  close, it

<sup>9</sup>Compactness of  $\Lambda$  follows since  $\sigma \in \Lambda$  depends continuously on the images of the  $L^2$  break points in the compact metric space  $X$ .

follows that the points  $\gamma_e(x_{2j}) = \gamma(x_{2j})$  (identity map) are continuous with respect to the  $W^{1,2}$  norm. Thus the  $a_j^i$ 's are continuous functions of  $\gamma^i$ , and so is each  $S^i$ . Moreover, the local energy convexity in Theorem 0.1 implies that the  $\gamma_e^i$ 's are indeed  $W^{1,2}$  close on each interval  $[x_{2j}, x_{2j+2}]$  if the  $\gamma^i$ 's are (since  $\gamma_e^i|_{[x_{2j}, x_{2j+2}]}$ 's also stay within a  $\rho$ -neighborhood and the right-hand side of the energy convexity for them is just a continuous function of  $S^i$ 's). Thus, we have shown  $\gamma \rightarrow \gamma_e$  is continuous.

To show  $\gamma_e \rightarrow \tilde{\gamma}_e$  is also continuous, it suffices to show that the  $\tilde{\gamma}_e^i$ 's are close when the  $\gamma_e^i$ 's are. Since the point  $x_0 = x_{2L^2}$  is fixed under the reparametrization, this will follow from applying Wirtinger's inequality to  $d(\tilde{\gamma}_e^1, \tilde{\gamma}_e^2) - d(\tilde{\gamma}_e^1(x_0), \tilde{\gamma}_e^2(x_0))$  once we show that  $\int_{\mathbf{S}^1} |\nabla d(\tilde{\gamma}_e^1, \tilde{\gamma}_e^2)|^2$  can be made small.

The piecewise linear curve  $\tilde{\gamma}_e^i$  is linear on the intervals

$$(B.2) \quad I_j^i = \left[ \frac{1}{S^i} \sum_{\ell < j} a_\ell^i, \frac{1}{S^i} \sum_{\ell \leq j} a_\ell^i \right].$$

Set  $I_j = I_j^1 \cap I_j^2$ . Observe first that since the intervals  $I_j^i$  in (B.2) depend continuously on  $\gamma_e^i$ , the measure of the complement  $\mathbf{S}^1 \setminus \left[ \bigcup_{j=0}^{L^2-1} I_j \right]$  can be made small, so that

$$(B.3) \quad \int_{\mathbf{S}^1 \setminus [\cup I_j]} |\nabla d(\tilde{\gamma}_e^1, \tilde{\gamma}_e^2)|^2 \leq \int_{\mathbf{S}^1 \setminus [\cup I_j]} (|(\tilde{\gamma}_e^1)'| + |(\tilde{\gamma}_e^2)'|)^2 \leq 4L^2 \text{Length}(\mathbf{S}^1 \setminus [\cup I_j])$$

can also be made small. We will divide the  $I_j$ 's into two groups, depending on the size of  $a_j^1$ . Fix some  $\epsilon > 0$  and suppose first that  $a_j^1 < \epsilon$ ; by continuity, we can assume that  $a_j^2 < 2\epsilon$ . For such  $j$ , we get

$$(B.4) \quad \int_{I_j} |\nabla d(\tilde{\gamma}_e^1, \tilde{\gamma}_e^2)|^2 \leq 2 \int_{I_j^1} |(\tilde{\gamma}_e^1)'|^2 + 2 \int_{I_j^2} |(\tilde{\gamma}_e^2)'|^2 \leq 2L(a_j^1 + a_j^2) \leq 6\epsilon L.$$

Since there are at most  $L^2$  breaks, summing over these intervals contributes at most  $6\epsilon L^3$ .


On the other hand, suppose now  $a_j^1 \geq \epsilon$ ; by continuity we can assume that  $a_j^2 \geq \epsilon/2$ . In this case,  $\tilde{\gamma}_e^i$  can be written on  $I_j$  as the composition  $\gamma_e^i \circ P_j^i$  where  $|(P_j^i)'| = 2\pi S^i / (L^2 a_j^i)$ . Furthermore,  $P_j^1$  and  $P_j^2$  both map  $I_j$  into  $[x_{2j}, x_{2j+2}]$  and arguing as (A.7) we have

$$\frac{1}{4} \int_{I_j} |\nabla d(\tilde{\gamma}_e^1, \tilde{\gamma}_e^2)|^2 = \frac{1}{4} \int_{I_j} |\nabla d(\gamma_e^1 \circ P_j^1, \gamma_e^2 \circ P_j^2)|^2 \leq \frac{1}{2} \int_{I_j} (|(\gamma_e^1)'| \cdot |(P_j^1)'| - |(\gamma_e^2)'| \cdot |(P_j^2)'|)^2.$$

This can be made small since the speed  $|(P_j^i)'|$  is continuous in  $\gamma^i$  and the (piecewise constant) speeds  $|(\gamma_e^i)'|$ 's are close when  $\gamma_e^i$ 's are. Therefore, the integral over these intervals can also be made small since there are at most  $L^2$  of them.  $\square$

## APPENDIX C. PROOF OF LEMMA 2.2

If  $(X, d)$  has curvature bounded from above by  $K > 0$  in the sense of Alexandrov, then  $(X, \frac{\sqrt{Kd}}{\sqrt{\epsilon}})$  has curvature bounded from above by  $\epsilon$ , so that the local distance convexity in Lemma 2.2 is homogenous w.r.t.  $K$ . Hence, it suffices to assume the metric space has curvature bounded from above by  $\epsilon$  which is sufficiently small.



AE=BE=1/2 a, CF=DF=1/2 b, AD=x, BC=y, EF=g, ED=c, AF=d, BF=e, EC=f

FIGURE 1.

Suppose now  $K > 0$  is sufficiently small. Let  $d_{\mathbf{S}_K}(A, B) = a$ ,  $d_{\mathbf{S}_K}(C, D) = b$ ,  $d_{\mathbf{S}_K}(A, D) = x$ ,  $d_{\mathbf{S}_K}(B, C) = y$ ,  $d_{\mathbf{S}_K}(E, F) = g$ ,  $d_{\mathbf{S}_K}(E, D) = c$ ,  $d_{\mathbf{S}_K}(A, F) = d$ ,  $d_{\mathbf{S}_K}(B, F) = e$  and  $d_{\mathbf{S}_K}(E, C) = f$  (see Figure 1). In the rest of this section we aim to prove the following that gives Lemma 2.2: for  $a, b, x, y$  small enough (i.e., under small region assumption), we have the following inequality:

$$(C.1) \quad \frac{1}{4}(a - b)^2 \leq x^2 + y^2 - 2g^2.$$

Based on Lemma 2.1, we first provide three key observations.

**Lemma C.2.** *For  $x, y$  sufficiently small and some  $\alpha, \beta \in \mathbf{R}$ , we have:*

$$W(x^2 + y^2 - 2g^2) = Sx^2 + Ty^2 + U + V - Rg^2 + O(x^2g^2) + O(y^2g^2) + O(g^4 + x^4 + y^4),$$

where

- (1)  $W = \frac{k^4}{6}(2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b)(c^2 + d^2 + e^2 + f^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg - 2\beta yg) + 2k^2(\cos \frac{k}{2}a + \cos \frac{k}{2}b) - \frac{k^2}{2}(\cos kc + \cos kd + \cos ke + \cos kf)$ ,
- (2)  $R = \frac{k^4}{2}(2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b)(c^2 + d^2 + e^2 + f^2 - \frac{1}{3}(a^2 + b^2) - 2\alpha xg - 2\beta yg) + 6k^2(\cos \frac{k}{2}a + \cos \frac{k}{2}b) - 2k^2(\cos kc + \cos kd + \cos ke + \cos kf)$ ,
- (3)  $S = \frac{k^4}{6}(2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b)(e^2 + f^2 - 2\beta yg) + k^2(\cos \frac{k}{2}a + \cos \frac{k}{2}b) + \frac{k^2}{2}(\cos kc + \cos kd - \cos ke - \cos kf)$ ,
- (4)  $T = \frac{k^4}{6}(2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b)(c^2 + d^2 - 2\alpha xg) + k^2(\cos \frac{k}{2}a + \cos \frac{k}{2}b) - \frac{k^2}{2}(\cos kc + \cos kd - \cos ke - \cos kf)$ ,
- (5)  $U = k^2(2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b)(c^2 + d^2 + e^2 + f^2 - \frac{1}{2}(a^2 + b^2) - 2\alpha xg - 2\beta yg)$ ,
- (6)  $V = 8(\cos \frac{k}{2}a + 1)(\cos \frac{k}{2}b + 1) - 4(\cos kc + 1)(\cos kd + 1) - 4(\cos ke + 1)(\cos kf + 1)$ .



*Proof.* Apply Lemma 2.1 to  $\{A, E, F, D\}$ , we can choose  $K$  sufficiently small to be determined later so that for some  $\alpha$  (with  $k = \sqrt{K}$ ,  $|\alpha|$  sufficiently small)

(C.3)

$$\alpha + \left(\frac{1}{4}a^2 + \frac{1}{4}b^2 - c^2 - d^2\right)/(2xg) = \cos q_K(\overrightarrow{AD}, \overrightarrow{EF}) = \cos q_K(\overrightarrow{DA}, \overrightarrow{FE})$$

(C.4)

$$= \frac{\cos \frac{k}{2}a + \cos kg \cos kx \cos \frac{k}{2}b + \cos \frac{k}{2}a \cos \frac{k}{2}b - \cos kg \cos kd - \cos kx \cos kc - \cos kc \cos kd}{(1 + \cos \frac{k}{2}b) \sin kx \sin kg}$$

(C.5)

$$= \frac{\cos \frac{k}{2}b + \cos kg \cos kx \cos \frac{k}{2}a + \cos \frac{k}{2}a \cos \frac{k}{2}b - \cos kx \cos kd - \cos kg \cos kc - \cos kc \cos kd}{(1 + \cos \frac{k}{2}a) \sin kx \sin kg}.$$

By Taylor series expansions for sine and cosine (in  $x$  and  $g$ ) and using (C.3)-(C.4), we have

$$\begin{aligned} & \left[ \frac{1}{4}(a^2 + b^2) - c^2 - d^2 + 2\alpha xg \right] (1 + \cos \frac{k}{2}b)(kx - \frac{1}{6}(kx)^3 + O(x^5))(kg - \frac{1}{6}(kg)^3 + O(g^5)) \\ = & 2xg \left[ (\cos \frac{k}{2}a)(1 + \cos \frac{k}{2}b) + (\cos \frac{k}{2}b)(1 - \frac{1}{2}(kx)^2 + \frac{1}{24}(kx)^4 + O(x^6))(1 - \frac{1}{2}(kg)^2 + \frac{1}{24}(kg)^4 \right. \\ & \left. + O(g^6)) - (\cos kd)(1 - \frac{1}{2}(kg)^2 + \frac{1}{24}(kg)^4 + O(g^6)) - (\cos kc)(1 - \frac{1}{2}(kx)^2 + \frac{1}{24}(kx)^4 \right. \\ & \left. + O(x^6)) - \cos kc \cos kd \right]. \end{aligned}$$

Combining the terms in  $x^2$  and  $g^2$  yields

$$\begin{aligned} & \left[ \frac{k^4}{6}(1 + \cos \frac{k}{2}b)(c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) - k^2 \cos kd + k^2 \cos \frac{k}{2}b \right] g^2 \\ = & - \left[ \frac{k^4}{6}(1 + \cos \frac{k}{2}b)(c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) - k^2 \cos kc + k^2 \cos \frac{k}{2}b \right] x^2 \\ & + k^2(1 + \cos \frac{k}{2}b)(c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) + 2(1 + \cos \frac{k}{2}a)(1 + \cos \frac{k}{2}b) \\ & - 2(1 + \cos kc)(1 + \cos kd) + O(x^2g^2) + O(x^4 + g^4). \end{aligned}$$

Similarly, using (C.3)-(C.5), we have

$$\begin{aligned} & \left[ \frac{k^4}{6}(1 + \cos \frac{k}{2}a)(c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) - k^2 \cos kc + k^2 \cos \frac{k}{2}a \right] g^2 \\ = & - \left[ \frac{k^4}{6}(1 + \cos \frac{k}{2}a)(c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) - k^2 \cos kd + k^2 \cos \frac{k}{2}a \right] x^2 \\ & + k^2(1 + \cos \frac{k}{2}a)(c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) + 2(1 + \cos \frac{k}{2}a)(1 + \cos \frac{k}{2}b) \\ & - 2(1 + \cos kc)(1 + \cos kd) + O(x^2g^2) + O(x^4 + g^4). \end{aligned}$$

Therefore,

$$\begin{aligned}
& \left[ \frac{k^4}{6} (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) + k^2 (\cos \frac{k}{2}a + \cos \frac{k}{2}b \right. \\
& \quad \left. - \cos kc - \cos kd) \right] g^2 \\
= & - \left[ \frac{k^4}{6} (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) + k^2 (\cos \frac{k}{2}a + \cos \frac{k}{2}b \right. \\
& \quad \left. - \cos kc - \cos kd) \right] x^2 + k^2 (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) \\
& + 4(1 + \cos \frac{k}{2}a)(1 + \cos \frac{k}{2}b) - 4(1 + \cos kc)(1 + \cos kd) + O(x^2g^2) + O(x^4 + g^4).
\end{aligned}$$

Similarly, in  $\{B, E, F, C\}$  we have for some  $\beta$  (with  $|\beta|$  sufficiently small)

$$\begin{aligned}
& \left[ \frac{k^4}{6} (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (e^2 + f^2 - \frac{1}{4}(a^2 + b^2) - 2\beta yg) + k^2 (\cos \frac{k}{2}a + \cos \frac{k}{2}b \right. \\
& \quad \left. - \cos ke - \cos kf) \right] g^2 \\
= & - \left[ \frac{k^4}{6} (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (e^2 + f^2 - \frac{1}{4}(a^2 + b^2) - 2\beta yg) + k^2 (\cos \frac{k}{2}a + \cos \frac{k}{2}b \right. \\
& \quad \left. - \cos ke - \cos kf) \right] y^2 + k^2 (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (e^2 + f^2 - \frac{1}{4}(a^2 + b^2) - 2\beta yg) \\
& + 4(1 + \cos \frac{k}{2}a)(1 + \cos \frac{k}{2}b) - 4(1 + \cos ke)(1 + \cos kf) + O(y^2g^2) + O(y^4 + g^4).
\end{aligned}$$

Adding up the above two equations then yields

$$\begin{aligned}
& \left[ \frac{k^4}{6} (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (c^2 + d^2 + e^2 + f^2 - \frac{1}{2}(a^2 + b^2) - 2\alpha xg - 2\beta yg) \right. \\
& \quad \left. + k^2 (2 \cos \frac{k}{2}a + 2 \cos \frac{k}{2}b - \cos kc - \cos kd - \cos ke - \cos kf) \right] g^2 \\
= & - \left[ \frac{k^4}{6} (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (c^2 + d^2 - \frac{1}{4}(a^2 + b^2) - 2\alpha xg) + k^2 (\cos \frac{k}{2}a + \cos \frac{k}{2}b \right. \\
& \quad \left. - \cos kc - \cos kd) \right] x^2 \\
& - \left[ \frac{k^4}{6} (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (e^2 + f^2 - \frac{1}{4}(a^2 + b^2) - 2\beta yg) + k^2 (\cos \frac{k}{2}a + \cos \frac{k}{2}b \right. \\
& \quad \left. - \cos ke - \cos kf) \right] y^2 \\
& + k^2 (2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b) (c^2 + d^2 + e^2 + f^2 - \frac{1}{2}(a^2 + b^2) - 2\alpha xg - 2\beta yg) \\
& + 8(1 + \cos \frac{k}{2}a)(1 + \cos \frac{k}{2}b) - 4(1 + \cos kc)(1 + \cos kd) - 4(1 + \cos ke)(1 + \cos kf) \\
& + O(x^2g^2) + O(y^2g^2) + O(g^4 + x^4 + y^4).
\end{aligned}$$

The lemma follows immediately by rearranging the terms above and using the definitions of  $W, R, S, T, U$ .  $\square$

**Remark C.6.** For fixed  $a$ , as  $x, y \rightarrow 0$  (and thus  $g \rightarrow 0, b \rightarrow a$ , and  $c, d, e, f \rightarrow \frac{1}{2}a$ ), we have  $\frac{S}{W} \rightarrow 1$  and  $\frac{T}{W} \rightarrow 1$ .

**Lemma C.7.** For  $a, b, x, y, k$  small enough, we have

$$U - Rg^2 \geq \frac{k^2}{2} g^2 - 4k^2(2|\alpha|xg + 2|\beta|yg).$$

*Proof.* Let  $|\cdot|$  denote the Euclidean distance in  $\mathbf{R}^3$  and  $\angle EF'D = \theta$  (see Figure 2). Then

$$|FF'| = \frac{1}{k}(1 - \cos \frac{k}{2}b), \quad |FF''| = \frac{1}{k}(1 - \cos kg) = \frac{2}{k} \sin^2 \left( \frac{k}{2}g \right), \quad |EF''| = \frac{1}{k} \sin kg,$$

and

$$|EF'|^2 = |F'F''|^2 + |EF''|^2 = \frac{1}{k^2} \left( 1 - \cos \frac{k}{2}b - 2 \sin^2 \left( \frac{k}{2}g \right) \right)^2 + \frac{1}{k^2} \sin^2(kg).$$

Thus,

$$\begin{aligned} & |CE|^2 + |DE|^2 - (|CF|^2 + |DF|^2) \\ &= (|CF'| + |EF'| \cos \theta)^2 + (|EF'| \sin \theta)^2 \\ &\quad + (|CF'| - |EF'| \cos \theta)^2 + (|EF'| \sin \theta)^2 - 2|CF'|^2 - 2|FF'|^2 \\ (C.8) \quad &= 2(|EF'|^2 - |FF'|^2) \\ &= \frac{2}{k^2} \left[ \left( 1 - \cos \frac{k}{2}b - 2 \sin^2 \left( \frac{k}{2}g \right) \right)^2 + \sin^2(kg) - \left( 1 - \cos \frac{k}{2}b \right)^2 \right] \\ &= \frac{2}{k^2} \left[ \sin^2(kg) - 4 \left( 1 - \cos \frac{k}{2}b \right) \sin^2 \left( \frac{k}{2}g \right) + 4 \sin^4 \left( \frac{k}{2}g \right) \right] \\ &= 2 \left( \cos \frac{k}{2}b \right) g^2 + O(g^4) > \left( \cos \frac{k}{2}b \right) g^2, \end{aligned}$$

for  $b, g$  small, which also implies  $|EF'| \geq |FF'|$ .

Similarly, for  $n \geq 2$ ,

$$\begin{aligned} & |CE|^{2n} + |DE|^{2n} - (|CF|^{2n} + |DF|^{2n}) \\ &= \left[ (|CF'| + |EF'| \cos \theta)^2 + (|EF'| \sin \theta)^2 \right]^n \\ (C.9) \quad &+ \left[ (|CF'| - |EF'| \cos \theta)^2 + (|EF'| \sin \theta)^2 \right]^n - 2 \left[ |CF'|^2 + |FF'|^2 \right]^n \\ &\geq 2 \left[ (|CF'|^2 + |EF'|^2)^n - (|CF'|^2 + |FF'|^2)^n \right] \geq 0. \end{aligned}$$

Now, note that

$$c = \frac{2}{k} \arcsin \left( \frac{k|DE|}{2} \right), \quad f = \frac{2}{k} \arcsin \left( \frac{k|CE|}{2} \right),$$

and

$$\frac{1}{2}b = \frac{2}{k} \arcsin \left( \frac{k|CF|}{2} \right) = \frac{2}{k} \arcsin \left( \frac{k|DF|}{2} \right).$$




FIGURE 2.

The Taylor series expansion

$$\left(2 \arcsin \left(\frac{x}{2}\right)\right)^2 = x^2 + \sum_{n=2} C_n x^{2n} \quad (C_n \geq 0)$$

and (C.8), (C.9) then imply: for  $x, y$  small enough (thus  $g$  is small enough),

$$\begin{aligned} & c^2 + f^2 - \frac{1}{2}b^2 \\ &= |CE|^2 + |DE|^2 - (|CF|^2 + |DF|^2) + \sum_{n=2} C_n k^{2n-2} (|CE|^{2n} + |DE|^{2n} - (|CF|^{2n} + |DF|^{2n})) \\ &> \left(\cos \frac{k}{2}b\right) g^2. \end{aligned}$$

Similarly,  $d^2 + e^2 - \frac{1}{2}a^2 > (\cos \frac{k}{2}a) g^2$ .

Therefore,

$$(C.10) \quad c^2 + d^2 + e^2 + f^2 - \frac{1}{2}(a^2 + b^2) > \left(\cos \frac{k}{2}a + \cos \frac{k}{2}b\right) g^2.$$

Recall the facts that as  $x, y \rightarrow 0$

- (1)  $c^2 + d^2 + e^2 + f^2 - \frac{1}{3}(a^2 + b^2) \rightarrow \frac{1}{6}(a^2 + b^2)$ ,
- (2)  $\cos kc + \cos kd + \cos ke + \cos kf \rightarrow 2(\cos \frac{k}{2}a + \cos \frac{k}{2}b)$ .

One observes that in a small geodesic ball  $B_\tau$  (thus  $a, b, x, y$  are small enough), we have:

- (1)  $\cos \frac{k}{2}a + \cos \frac{k}{2}b > \frac{15}{8}$ ,
- (2)  $c^2 + d^2 + e^2 + f^2 - \frac{1}{3}(a^2 + b^2) - 2\alpha xg - 2\beta yg < \frac{1}{2}$ ,
- (3)  $\cos kc + \cos kd + \cos ke + \cos kf > \frac{3}{2}(\cos \frac{k}{2}a + \cos \frac{k}{2}b)$ .

Therefore, using (C.10) and the definition of  $R$ , we obtain that for  $a, b, x, y$  small,  $k \leq 1$  :

$$\begin{aligned}
& U - Rg^2 \\
&= k^2 \left( 2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b \right) \left( c^2 + d^2 + e^2 + f^2 - \frac{1}{2}(a^2 + b^2) - 2\alpha xg - 2\beta yg \right) - Rg^2 \\
&\geq \frac{15k^2}{8} \left( 2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b \right) g^2 - \left( \frac{k^4}{4} \left( 2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b \right) + 3k^2 \left( \cos \frac{k}{2}a + \cos \frac{k}{2}b \right) \right) g^2 \\
&\quad - k^2 \left( 2 + \cos \frac{k}{2}a + \cos \frac{k}{2}b \right) (2\alpha xg + 2\beta yg) \\
&\geq k^2 \left( \frac{13}{4} - \frac{11}{8} \left( \cos \frac{k}{2}a + \cos \frac{k}{2}b \right) \right) g^2 - 4k^2(2|\alpha|xg + 2|\beta|yg) \\
&\geq \frac{k^2}{2} g^2 - 4k^2(2|\alpha|xg + 2|\beta|yg).
\end{aligned}$$

□

**Lemma C.11.**  $V \geq 0$  for  $a, b, x, y$  small enough.

*Proof.* By the triangle inequality, we know  $\frac{1}{2}(a+b) < c+d < \frac{1}{2}(a+b) + x+g$ ,  $\frac{1}{2}(a+b) < e+f < \frac{1}{2}(a+b) + y+g$ ,  $c+f > b$ , and  $d+e > a$ . If  $c \geq \frac{1}{2}b$ ,  $f \geq \frac{1}{2}b$  and  $d \geq \frac{1}{2}a$ ,  $e \geq \frac{1}{2}a$  then one easily sees  $V \geq 0$ . Now without loss of generality we suppose that there exists  $\varsigma > \sigma > 0$  such that

$$c = \frac{1}{2}b - \sigma, \quad d = \frac{1}{2}a + \varsigma, \quad e > \frac{1}{2}a - \varsigma, \quad f > \frac{1}{2}b + \sigma.$$

Assume now  $k = 1$ , then for  $a, b$  small:

$$\begin{aligned}
V &= 8(\cos \frac{1}{2}a + 1)(\cos \frac{1}{2}b + 1) - 4(\cos c + 1)(\cos d + 1) - 4(\cos e + 1)(\cos f + 1) \\
&= 8 \cos \frac{1}{2}b - 4(\cos c + \cos f) + 8 \cos \frac{1}{2}a - 4(\cos d + \cos e) + 8(\cos \frac{1}{2}a)(\cos \frac{1}{2}b) \\
&\quad - 4(\cos c)(\cos d) - 4(\cos e)(\cos f) \\
&\geq 8 \cos \frac{1}{2}b - 4(\cos(\frac{1}{2}b - \sigma) + \cos(\frac{1}{2}b + \sigma)) + 8 \cos \frac{1}{2}a - 4(\cos(\frac{1}{2}a + \varsigma) + \cos(\frac{1}{2}a - \varsigma)) \\
&\quad + 8(\cos \frac{1}{2}a)(\cos \frac{1}{2}b) - 4 \left[ \cos(\frac{1}{2}b - \sigma) \cos(\frac{1}{2}a + \varsigma) + \cos(\frac{1}{2}b + \sigma) \cos(\frac{1}{2}a - \varsigma) \right] \\
&= 8(1 - \cos \sigma) \cos \frac{1}{2}b + 8(1 - \cos \varsigma) \cos \frac{1}{2}a + 4(\cos \frac{a+b}{2} + \cos \frac{a-b}{2}) \\
&\quad - 2 \left[ \cos(\frac{a+b}{2} + \varsigma - \sigma) + \cos(\frac{b-a}{2} + \varsigma + \sigma) + \cos(\frac{a+b}{2} - \varsigma + \sigma) + \cos(\frac{b-a}{2} - \varsigma - \sigma) \right] \\
&\geq 4(1 - \cos(\varsigma - \sigma)) \cos \frac{a+b}{2} + 4(1 - \cos(\varsigma + \sigma)) \cos \frac{b-a}{2} \geq 0.
\end{aligned}$$

One sees that the above argument works for all  $k > 0$ , completing the proof. □

**Remark C.12.** From the proofs of these lemmas we also see that for  $a, b, x, y$  small enough,  $W, S, T, R > 0$ .

*Proof.* (of **Lemma 2.2**) Combining previous lemmas we get

$$W(x^2+y^2-2g^2) \geq Sx^2+Ty^2+\frac{k^2}{2}g^2-4k^2(2|\alpha|xg+2|\beta|yg)+O(x^2g^2)+O(y^2g^2)+O(g^4+x^4+y^4).$$

Now using nothing but the facts

- (1)  $0 < k^2 \leq W \leq 4k^2$  for  $a, b, x, y, k$  sufficiently small,
- (2)  $\frac{S}{W} \rightarrow 1$  and  $\frac{T}{W} \rightarrow 1$  uniformly as  $x, y \rightarrow 0$  (see Remark C.6),
- (3)  $|a - b| \leq x + y$ ,

we obtain for  $a, b, x, y$  sufficiently small:

$$(C.13) \quad x^2 + y^2 - 2g^2 \geq \frac{3}{4}(x^2 + y^2) + \frac{1}{16}g^2 - 4(2|\alpha|xg + 2|\beta|yg).$$

Now choose  $K$  sufficiently small so that  $\max\{|\alpha|, |\beta|\} \leq \frac{1}{128}$  and thus by Cauchy-Schwarz inequality

$$4(2|\alpha|xg + 2|\beta|yg) \leq \frac{1}{16}(x^2 + y^2 + g^2),$$

and therefore

$$x^2 + y^2 - 2g^2 \geq \frac{1}{2}(x^2 + y^2) \geq \frac{1}{4}(x + y)^2 \geq \frac{1}{4}(a - b)^2,$$

completing the proof of Lemma 2.2. □

## REFERENCES

- [A1] A.D. Alexandrov, A theorem on triangles in a metric space and some of its applications, *Trudy Mat.Inst.Steklova* 38 (1951), 5–23.
- [A2] A.D. Alexandrov, Über eine Verallgemeinerung der Riemannschen Geometrie, *Schr. Forschunginst. Math.* 1 (1957), 33–84.
- [Al] F.J. Almgren, The theory of varifolds, Mimeographed notes, Princeton, 1965.
- [BN] I.D. Berg and I.G. Nikolaev, On an extremal property of quadrilaterals in an Alexandrov space of curvature  $\leq K$ , *Contemporary mathematics*; 424. Providence, R.I: AMS, c2007.
- [B1] G.D. Birkhoff, Dynamical systems with two degrees of freedom. *TAMS* 18 (1917), no. 2, 199–300.
- [B2] G.D. Birkhoff, *Dynamical systems*, AMS Colloq. Publ. vol 9, Providence, RI, 1927.
- [BBI] D. Burago, Y. Burago and S. Ivanov, *A Course in Metric Geometry*, American Mathematical Society, 2001.
- [Bu] H. Busemann, *The geometry of geodesics*, Acad. Press, 1955.
- [Ca] E. Cartan, *Leçons sur la Géométrie des Espaces de Riemann.*, Gauthier-Villars, Paris, 1928, 2nd edition 1951.
- [CC] E. Calabi and J. Cao, Simple closed geodesics on convex surfaces, *J. Differential Geom.* 36 (1992), no. 3, 517–549.
- [CD] T.H. Colding and C. De Lellis, The min-max construction of minimal surfaces, *Surveys in differential geometry*, Vol. 8, Lectures on Geometry and Topology held in honor of Calabi, Lawson, Siu, and Uhlenbeck at Harvard University, May 3–5, 2002, Sponsored by JDG, (2003) 75–107.
- [CM1] T.H. Colding and W.P. Minicozzi II, Width and mean curvature flow, *Geometry and Topology*, 12 (2008), no. 5, 2517–2535.
- [CM2] T.H. Colding and W.P. Minicozzi II, Width and finite extinction time of Ricci flow, *Geometry and Topology*, 12 (2008), no. 5, 2537–2586.
- [CM3] T.H. Colding and W.P. Minicozzi II, Estimates for the extinction time for the Ricci flow on certain 3-manifolds and a question of Perelman, *JAMS*, 18 (2005), no. 3, 561–569.
- [CM4] T.H. Colding and W.P. Minicozzi II, *Minimal surfaces*, Courant Lecture Notes in Mathematics, 4. New York University, Courant Institute of Mathematical Sciences, New York, 1999.

- [Cr] C.B. Croke, Area and the length of the shortest closed geodesic, *J. Differential Geom.*, 27 (1988), no. 1, 1–21.
- [Had] J. Hadamard, Les surfaces à courbures opposées et leurs lignes géodésique, *J. Math. Pure Appl.*, 4 (1898) pp. 27-75.
- [Hat] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. xii+544 pp. ISBN: 0-521-79160-X; 0-521-79540-0.
- [KS] N. Korevaar and R. Schoen, Sobolev Spaces and Harmonic Maps for Metric Space Targets, *Communications in Analysis and Geometry*, 1 (1993), 561–659.
- [Ly] L.A. Lyusternik, Topology of functional spaces and calculus of variations in the large (Russian), *Trudy Inst. Steklov.* 19, Izdat. Akad. Nauk SSSR, Moscow, 1947; The topology of the calculus of variations in the large, translated from the Russian by J. M. Danskin. *Translations of Mathematical Monographs*, Vol. 16 American Mathematical Society, Providence, R.I. 1966 vii+96 pp.
- [LyS] L.A. Lyusternik and L. Šnirel'man, Topological methods in variational problems and their application to the differential geometry of surfaces, (Russian) *Uspehi Matem. Nauk (N.S.)* 2, (1947). no. 1(17), 166–217.
- [LW] L. Lin and L. Wang, Existence of good sweepouts on closed manifolds, *Proc. Amer. Math. Soc.*, to appear.
- [Pi] J.T. Pitts, Existence and regularity of minimal surfaces on Riemannian manifolds, Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo 1981.
- [Po] H. Poincaré, Sur les lignes géodésiques des surfaces convexes, *Trans. Amer. Math. Soc.* , 6 (1904) pp. 237-274.
- [Re] Y.G. Reshetnyak, Non-expanding mappings in a space of curvature not greater than  $K$ , *Sibirsk.Mat. Zh.*, s9, 918C927 (1968) (in Russian); English translation: *Sib.Math. J.* 9, 683–689 (1968).
- [Se] T. Serbinowski, Harmonic maps into metric spaces with curvature bounded above, Ph.D. thesis, University of Utah, 1995.

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, 3400 N. CHARLES ST., BALTIMORE, MD 21218

*E-mail address:* lzlin@math.jhu.edu