

UCSF

UC San Francisco Previously Published Works

Title

Pincer Morphology Is A Risk Factor For Developing Radiographic Hip Osteoarthritis; Data From The World Coach Consortium

Permalink

<https://escholarship.org/uc/item/12d7t13x>

Authors

Riedstra, NS
Van Buuren, MM
Boel, F
[et al.](#)

Publication Date

2023-03-01

DOI

10.1016/j.joca.2023.01.252

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

this effect. Future studies within the World COACH consortium will elucidate whether this is an overall effect or if specific high-risk subgroups, for instance the younger individuals, are responsible for the association found.

235

THE LONGITUDINAL ASSOCIATION OF HAND OSTEOARTHRITIS WITH PAID AND UNPAID WORK IMPAIRMENT AND RELATED COSTS: THE HAND OSTEOARTHRITIS IN SECONDARY CARE COHORT

S.E. Terpstra¹, L.A. van de Stadt¹, A.E. Boonen², F.R. Rosendaal¹, M. Kloppenburg¹. ¹Leiden Univ. Med. Ctr., Leiden, Netherlands; ²Maastricht Univ. Med. Ctr., Maastricht, Netherlands

Purpose: Hand osteoarthritis (OA) presence is associated with impairment in paid and unpaid work and contributes to societal burden and costs. However, the longitudinal development of hand OA-related paid and unpaid work impairment and related societal costs are unknown. Therefore, we aimed to investigate the association of hand OA with paid and unpaid work production loss, hinder and societal costs longitudinally. **Methods:** We used annual data of the Dutch Hand OSTeoArthritis in Secondary care (HOSTAS) cohort from baseline to four years of follow-up. Presence of primary hand OA was defined by the treating rheumatologist. Patient and OA characteristics were assessed with validated questionnaires on patient and hand OA characteristics and by physical examination. The Health and Labour Questionnaire (HLQ) - having a recall period of two weeks - was assessed each year to investigate hinder, hours of sick leave (absenteeism) and unproductive hours while at work as well as hinder and hours required to replace unpaid work tasks in relation to hand OA. The HLQ was added to the questionnaires after part of the cohort already had first or second follow-up assessment. Patients with HLQ data on >1 timepoint were included in this study. Societal costs of paid work loss (=all costs an employer makes in order to employ a worker, such as salary and social premiums) were estimated by multiplying the number of unproductive and sick leave hours due to hand OA by the average hourly costs of paid work in The Netherlands. Costs of unpaid work were estimated by multiplying the hours of unpaid work replaced by others by the Dutch gross average hourly salary of a household help (€12.50). Costs were adjusted to 2021 values and extrapolated to costs per year using conversion factors.

Results: Data on more than one timepoint was available for 470 patients. Baseline data were available for 381 patients, of whom 256 patients (67%) completed four years of follow-up, and 215 (56%) completed all five follow-up moments. Those who had baseline data but did not complete four years of follow-up (n=125), and those with data on all timepoints (n=215) did not differ substantially from those with at least data on baseline and four years of follow-up (n=256) in terms of baseline age, sex and AUSCAN pain score (data not shown). We focused on the group with at least data on baseline and four years, as patients in this group had a comparable and substantial follow-up duration. Baseline and four years follow-up data was present for 256 patients, of which 113 (52%) had paid work at baseline and 104 (41%) at four years (table 1 for characteristics of these patients). At baseline, 25/133 (19%) of working patients had unproductive hours due to hand OA (absenteeism or unproductive at work); 10/25 (40%) of them still worked at four years without unproductive hours, 7/25 (28%) still worked with unproductive hours, and 8/25 (32%) had retired. Median paid work hinder score (score range: 6-24) remained stable at 7 (interquartile range (IQR): 6;8) over the study period. Patients who transitioned from "no paid work hinder" at baseline to "paid work hinder" or "paid work unproductive hours" at four years (n=14) did not differ substantially from those who did not (n=90) in terms of baseline mean age, AUSCAN pain and AUSCAN function (data not shown). Unproductive hours at any timepoint were present for 84/470 patients (18%). Presence of these unproductive hours for individual patients fluctuates over time (figure 1). Regarding unpaid work, 105/255 (41%) required unpaid task replacement by others due to hand OA at baseline, of which 24 reported this no more at four years. Of those that had no replacement at baseline, 28 developed this at four years. Consequently, 108 (42%) had unpaid work replacement at four years. Unpaid work hinder was reported by 100/256 patients (39%) at baseline and 97/255 (36%) at four years. Societal paid and unpaid work-related costs were present at baseline for 136/256 patients, with a median of €62.50 ((IQR) 37.50;125) per two weeks (€1630 (978;3261 per year). At four years, these were present for 120/256 patients, with a median of €50 (37.50;100) per two weeks (€1304 (978;2609) per year).




Figure 1. Heatmap of adverse paid work outcomes over time per patient per year, for all patients having data for at least two timepoints and any production loss at any timepoint (baseline, 1, 2, 3, 4 year follow-up, n=84)

	Baseline, all patients (n = 256)	Baseline, paid work (133/255, 52%)	Four years, all patients (n = 256)	Four years, paid work (104/255, 41%)
Age, years	61.3 (8.0)	60.2 (7.5)	64.6 (8.4)	62.0 (8.1)
Sex, women, n (%)	204 (80%)	108 (82%)	204 (80%)	87 (84%)
BMI, kg/m ²	27.3 (4.7)	27.2 (4.7)	27.4 (5.0)	27.1 (4.7)
AUSCAN hand pain (0–20)	9 (4.3)	9.1 (4.6)	9.3 (4.3)	7.9 (4.5)
AUSCAN hand function (0–36)	15.1 (8.4)	14.6 (8.5)	15.2 (8.4)	14.5 (7.7)
HADS anxiety score (range 0–21) [^]	4 (2;6)	4 (2;6)	4 (2;6)	3 (2;6)
HADS depression score (range 0–21) [^]	2 (1;5)	2 (1;4)	2 (1;5)	1 (1;4)
Retired, n (%)	83 (32%)	–	95 (37%)	–

Numbers represent (SD) mean unless otherwise specified. median (interquartile range). Abbreviations: SD = standard deviation, BMI = Body Mass Index, HADS = Hospital Anxiety and Depression Scale, AUSCAN = Australian Canadian Osteoarthritis Hand Index.

Table 1

Osteoarthritis and Cartilage

Demographics and hand OA characteristics of patients with at least baseline and four years of follow-up data.

Conclusions: Patients with hand OA experience significant impairment in paid and unpaid work participation over the years, which translates into substantial societal costs. Impairment and costs fluctuate on individual level but remarkably seem to decrease over time on group level. Whether for example retirement, coping strategies, OA severity or survivorship bias might play a role in this needs further investigation.

236

PINCR MORPHOLOGY IS A RISK FACTOR FOR DEVELOPING RADIOGRAPHIC HIP OSTEOARTHRITIS; DATA FROM THE WORLD COACH CONSORTIUM

N.S. Riedstra¹, M.M. Van Buuren¹, F. Boel¹, H. Ahedi², V. Arbabi³, N. Arden⁴, S.M. Bierma-Zeinstra¹, C.G. Boer¹, F.M. Cicuttini⁵, T.F. Cootes⁶, D.T. Felson⁷, W.P. Gielis³, S. Kluzek⁸, N.E. Lane⁹, C. Lindner⁶, J. Lynch¹⁰, J. van Meurs¹, A.E. Nelson¹¹, M.C. Nevitt¹⁰, E.H. Oei¹, J. Runhaar¹, T.D. Spector¹², J. Tang¹, H.H. Weinans³, R. Agricola¹. ¹Erasmus Med. Ctr., Rotterdam, Netherlands; ²Univ. of Tasmania, Hobart, Australia; ³Univ. Med. Ctr. Utrecht, Utrecht, Netherlands; ⁴Oxford Univ., Oxford, United Kingdom; ⁵Monash Univ., Clayton, Australia; ⁶Univ. of Manchester, Manchester, United Kingdom; ⁷Boston Med. Ctr., Boston, MA; ⁸Univ. of Nottingham, Nottingham, United Kingdom; ⁹UC Davis, Davis, CA; ¹⁰Univ. of California, San Francisco, San Francisco, CA; ¹¹Univ. of North Carolina, Chapel Hill, NC; ¹²King's Coll. London, London, United Kingdom

Purpose: Osteoarthritis (OA) is the most prevalent joint disease with an estimated prevalence of 8% in the general population. Hip morphology has been marked as an important risk factor for the development of hip OA. One of the types thought to increase the risk of hip OA is pincer morphology, which is an overcoverage of the femoral head by the acetabulum. The overcoverage may result in the femoral neck impinging against the acetabulum during motion and possibly creating intra-articular damage over time. However, attempts to relate pincer morphology to the development of hip OA have yielded conflicting results. We studied the relationship between pincer morphology at baseline and the risk of developing radiographic hip OA within 8 years follow-up, using individual participant data from the Worldwide Collaboration on OsteoArthritis prediction for the Hip (World COACH).

Methods: The World COACH consortium was established to collect and harmonize all available individual participant data from prospective cohort studies (n=9) that have sequential pelvic or hip imaging available. For the current study, we included the 6 cohorts that had baseline pelvic radiographs and radiographic OA scores available within a maximum of 8 years. Standardized anteroposterior (AP) pelvic and/or hip radiographs were taken at baseline and at a follow-up visit between 4-8 years in each included cohort. Scores for radiographic OA were already available for each cohort, either by Kellgren & Lawrence grade (K&L), (modified) Croft grade, or an adaptation. For this analysis we harmonized these scores into "definitely no OA" (any score 0), "doubtful OA" (any score 1), or "definite OA" (any score ≥ 2 or total hip replacement (THR)). We only included hips without OA (any score 0) at baseline. An automatic Bonefinder[®] search model was used to annotate all baseline radiographs, outlining the bony shape per hip. The lateral center edge angle (LCEA) is a measurement for coverage of the femoral head by the acetabulum. The LCEA is formed by a vertical line perpendicular to the horizontal reference line and a line from the center of the femoral head to the most lateral point of the acetabulum (Figure 1). From the outline of the hip shape, the LCEA was calculated automatically. The threshold used for pincer morphology is a LCEA $\geq 45^\circ$. We excluded all hips with a LCEA $\leq 25^\circ$ in order to exclude hips with acetabular dysplasia as this is a hip shape that may also be related to hip OA. Development of radiographic hip OA was defined as a K&L grade ≥ 2 ,




Figure 1. AP pelvic radiograph with automatic Bonefinder search model of hip shape (blue), horizontal reference line (green) and automatic calculation of LCEA (black).

Table 1: Baseline characteristics of the study population

Characteristic	Study population (n=16655)
Women, n (%)	11729 (70.4)
Age, mean (sd)	62.7 (8.4)
BMI, mean (sd)	27.2 (4.4)
Cohort, n (%)	
CHECK	509 (3.1)
Chingford	691 (4.1)
Johnston County	427 (2.6)
OAI	5053 (30.3)
RS-I	1563 (9.4)
RS-II	1938 (11.6)
RS-III	2523 (15.1)
SOF	3951 (23.7)
LCEA $\geq 45^\circ$, n (%)	759 (4.6)

Croft score ≥ 2 , OA score =2 or THR, depending on available scores per cohort. The associations between baseline pincer morphology and development of radiographic hip OA were estimated using a logistic regression model with generalized mixed effects with 3 levels: hip side (left/right), individual and cohort. The results are expressed as odds ratios (OR) and were adjusted for baseline age, sex, and BMI.

Results: The six cohorts included yielded radiographic hip OA data on 51,363 hips (Table 1). We excluded 5,004 hips with definite OA and 18,250 hips with doubtful OA or missing OA scores at baseline. This left 28,109 hips without OA at baseline. After exclusion of hips with other missing data, insufficient quality radiographs or acetabular dysplasia, we included 16655 hips. At baseline 759 hips (5%) had pincer morphology. Within a maximum of 8 years (mean 6.1 \pm 1.7) follow-up, 683 hips (4.6%) developed radiographic hip OA. Pincer morphology was significantly associated with radiographic hip OA with an aOR of 1.59 (95% CI 1.156-2.195).

Conclusions: In hips without radiographic OA at baseline, the odds of developing hip OA within 8 years are 1.59 times higher in hips with pincer morphology than in hips without pincer morphology. The large and heterogeneous sample size allowed a robust estimate of this effect. Further studies within the World COACH consortium will elucidate whether this is an overall effect or if specific high-risk subgroups are responsible for the association found.

237

"SHORT QUESTIONNAIRE TO ASSESS HEALTH-ENHANCING PHYSICAL ACTIVITY" VERSUS ACTIHEART-MEASURED PHYSICAL ACTIVITY IN THE MIDDLE AGED GENERAL POPULATION: THE NETHERLANDS EPIDEMIOLOGY OF OBESITY STUDY

S.E. Terpstra, L.A. Hoogervorst, J.H. van der Velde, R. de Mutser, L.A. van de Stadt, F.R. Rosendaal, M. Kloppenburg. *Leiden Univ. Med. Ctr., Leiden, Netherlands*

Purpose: Adequate physical activity levels have been associated with wide-ranging beneficial health outcomes. It is therefore important to adequately measure physical activity. Physical activity is often measured by SQUASH (Short Questionnaire to Assess Health-enhancing physical activity), a patient-assessed questionnaire to measure physical activity. However, this is a subjective method. Data on validity of the SQUASH was not found a large, middle-aged population group. We therefore aimed to assess the construct validity of the SQUASH.

Methods: This is a cross-sectional analysis using data from the population-based Netherlands Epidemiology of Obesity (NEO) study, consisting of 6,671 participants aged 45-65 years. Participants reported the frequency and duration of their physical activity, which we expressed in hours per week of Metabolic Equivalent of Task (MET), using the SQUASH questionnaire based on a representative week in the last month. The SQUASH consists of eleven questions regarding four types of physical activity and consists of three main queries: 1) days per week; 2) average time per week; 3) frequency and duration. Patients with >0 queries missing in the SQUASH were excluded. The ActiHeart accelerometer was