UC San Diego UC San Diego Previously Published Works

Title

TLR-4 mediated group IVA phospholipase A2 activation is phosphatidic acid phosphohydrolase 1 and protein kinase C dependent

Permalink https://escholarship.org/uc/item/1452h15j

Journal Biochimica et Biophysica Acta, 1791(10)

ISSN 0006-3002

Authors

Grkovich, Andrej Armando, Aaron Quehenberger, Oswald <u>et al.</u>

Publication Date

2009-10-01

DOI

10.1016/j.bbalip.2009.02.002

Peer reviewed



NIH Public Access

Author Manuscript

Biochim Biophys Acta. Author manuscript; available in PMC 2010 October 1

Published in final edited form as:

Biochim Biophys Acta. 2009 October; 1791(10): 975–982. doi:10.1016/j.bbalip.2009.02.002.

TLR-4 mediated group IVA phospholipase A2 activation is phosphatidic acid phosphohydrolase 1 and protein kinase C dependent

Andrej Grkovich[†], Aaron Armando[★], Oswald Quehenberger^{*}, and Edward A. Dennis[†],[★],^{*} [†] Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, California 92093-0601

* Department of Pharmacology, University of California, San Diego, La Jolla, California 92093-0601

* Department of Medicine, University of California, San Diego, La Jolla, California 92093-0601

Abstract

Group IVA phospholipase A₂ (GIVA PLA₂) catalyzes the release of arachidonic acid (AA) from the *sn*-2 position of glycerophospholipids. AA is then further metabolized into terminal signaling molecules including numerous prostaglandins. We have now demonstrated the involvement of phosphatidic acid phosphohydrolase 1 (PAP-1) and protein kinase C (PKC) in the Toll-like receptor-4 (TLR-4) activation of GIVA PLA₂. We also studied the effect of PAP-1 and PKC on Ca⁺² induced and synergy enhanced GIVA PLA₂ activation. We observed that the AA release induced by exposure of RAW 264.7 macrophages to the TLR-4 specific agonist Kdo₂-Lipid A is blocked by the PAP-1 inhibitors bromoenol lactone (BEL) and propranolol as well as the PKC inhibitor Ro 31-8220; however these inhibitors did not reduce AA release stimulated by Ca⁺² influx induced by the P2X7 purinergic receptor agonist ATP. Additionally, stimulation of cells with diacylglycerol (DAG), the product of PAP-1 mediated hydrolysis, initiated AA release from unstimulated cells as well as restored normal AA release from cells treated with PAP-1 inhibitors. Finally, neither PAP-1 nor PKC inhibition reduced GIVA PLA₂ synergistic activation by stimulation with Kdo₂–Lipid A and ATP.

INTRODUCTION

Bacterial sepsis and septic shock result from the exacerbated production of inflammatory mediators by the immune system in response to bacterial endotoxins such as lipopolysaccharide (LPS) (1). Exposure of macrophages to LPS initiates the Toll-like receptor-4 (TLR-4) signaling pathway, resulting in the release of a host of cytokines and eicosanoids, including various prostaglandins (PGs). PGs are known to contribute to a number of normal and pathological physiological processes, including inflammation, pain, and vascular permeability and are also implicated as major contributors to endotoxic shock (2).

Group IVA phospholipase A_2 (GIVA PLA₂), also known as the cytosolic PLA₂ (cPLA₂), is the proinflammatory phospholipase responsible for the release of arachidonic acid (AA) which

[#]This work was supported by NIH grant GM64611 and utilized HPLC-MS equipment supported by LIPID MAPS NIH grant GM069335 *Corresponding author. Phone, 858-534-3055; FAX, 858-534-7390; E-mail, edennis@ucsd.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

is further metabolized into eicosanoids (3,4). GIVA PLA₂ catalyzes the hydrolysis of AA from the *sn*-2 position of membrane phospholipids, resulting in the release of free AA from the membrane and formation of lysophospholipids (5,6). GIVA PLA₂ has been implicated in a variety of inflammatory disease models, including collagen induced arthritis, autoimmune encephalomyelitis and acute lung injury; therefore, obtaining a complete understanding of the molecular mechanism of its activation is essential. (7–9).

Phosphatidic acid phosphohydrolase 1 (PAP-1) is a Mg⁺²-dependent, cytosolic enzyme that associates with phosphatidic acid (PA) containing membrane surfaces, where it converts PA into inorganic phosphate and diacylglycerol (DAG) (10-17). With the discovery by Carman and coworkers (17) in 2006 that yeast PAP-1 is the yeast ortholog of Lipin-1 discovered by Reue and coworkers (18); the interest in PAP-1 has escalated. For a recent review of the lipins, see (19). Our laboratory has utilized two different immortalized mammalian cell lines to characterize distinctive signaling pathways in which PAP-1 inhibition leads to the loss of cyclooxygenase 2 (COX-2) expression and PGE₂ release. The first study was conducted in the human amnionic WISH cell line utilizing the protein kinase C (PKC) activator phorbol myristate acetate (PMA) (18). The second study was conducted in the human U937 macrophage-like cell line stimulated with the Toll-like receptor (TLR)-4 agonist lipopolysaccharide (LPS) (19). In the macrophage cells, we (21) found in a Western blot that antibody to Lipin-1, and not antibodies to Lipin-2 or Lipin-3, crossreacted with protein at a molecular mass of 140 kDa on SDS-PAGE electrophoresis, the correct molecular mass for Lipin-1 (22), suggesting its presence and that the macrophage PAP-1 is Lipin-1, in analogy to the Carman finding in yeast (16,17). Drawing on these studies and other published reports, we hypothesized that DAG evolved from PAP-1 is likely to activate PKC, which has been shown to cause the phosphorylation of GIVA PLA2 in cellular studies conducted with phorbol esters (20-23).

In this manuscript, we provide further evidence that inhibition of PAP-1 or PKC results in reduced GIVA PLA₂ activation in macrophages stimulated with the TLR-4 specific agonist, Kdo₂-Lipid A (24). Furthermore, we observed that exposure of cells to the inhibitors did not reduce GIVA PLA₂ -mediated AA release initiated with Ca⁺² influx via ATP. Finally, our data suggests that inhibition of PAP-1 or PKC does not affect synergistic activation of GIVA PLA₂, despite reducing the absolute amount of AA released from cells that have been stimulated with both Kdo₂-Lipid A and ATP stimuli.

EXPERIMENTAL PROCEDURES

Materials

Murine RAW 264.7 macrophages were purchased from American Type Culture Collection (Manassas, VA). DMEM cell culture medium was obtained from Gibco (Grand Island, NY). Fetal bovine serum was from VWR International (Bristol, CT). DTT and lipopolysaccharide (LPS), from *E. coli* 0111:B4, were obtained from Sigma Chemical Company (St. Louis, MO). Kdo₂-Lipid A, 1-palmitoyl-2-arachidonoyl phosphatidylcholine was from Avanti Polar Lipids (Alabaster, AL). Propranolol, Dioctoyl-DAG and Ro 31-8220 were from Biomol (Plymouth Meeting, PA). Bromoenol lactone (BEL) was from Cayman Chemical (Ann Arbor, MI). 1-palmitoyl-2-(1-¹⁴C)-palmitoyl phosphatidylcholine, 1-palmitoyl-2-(1-¹⁴C)-arachidonoyl phosphatidylcholine, [5,6,8,9,11,12,14,15-³H] arachidonic acid (specific activity = 100 Ci/mmol) and [9,10-³H] oleic acid (specific activity = 23 Ci/mmol) were obtained from NEN Life Science Products (Boston, MA). Phosphatidylinositol 4,5-bisphosphate was from Roche (Basel, Switzerland). The specific cPLA₂ inhibitor, pyrrophenone, was kindly provided by Dr. Kohji Hanasaki (Shionogi Research Laboratories of Shionogi & Co., Ltd). 20 cm × 20 cm × 250 µm K6 Silica gel thin layer chromatography plates were from Whatman (Clifton, New Jersey).

Cell Culture and Stimulation Protocol

RAW 264.7 macrophages were maintained at 37 °C in a humidified atmosphere at 90% air and 5% CO2 DMEM supplemented with 10% fetal bovine serum, 2 mM glutamine, 100 units/ mL penicillin, 100 µg/ml streptomycin, and non-essential amino acids. Cells were plated at a confluency of 2×10^6 cells/6 well tissue culture plate and $1 \times 10^6/12$ well tissue culture plate at the time of experimentation. Following plating, they were allowed to adhere overnight, and then used for experiments the following day, with the exception of cells that were radiolabeled, described below. When DAG was added to the cells, it was done immediately after the addition of Kdo₂-Lipid A. Whenever cellular experiments required the usage of chemical inhibitors; the inhibitors were initially diluted into media at a stock concentration that was 100 times more concentrated than that of the final cellular concentration. From this stock concentration, the inhibitors were then applied to the cells to achieve the desired final cellular concentration. The final concentration of DMSO in the supernatants never exceeded .05% (volume/volume). When synergy experiments were performed, ATP was added to cells treated with Kdo2-Lipid A 50 minutes after Kdo₂-Lipid A stimulation, and supernatants were collected at 1 hour following stimulation. Cell viability was assessed visually by the Trypan Blue Dye exclusion assay (Gibco, Grand Island, NY) and through the usage of the CytoTox 96 Non-Radioactive Cytotoxicity Assay (Promega, Madison, WI).

Sample Preparation

Media was analyzed for extracellular eicosanoid release, as described previously (25). After stimulation, 1.8 ml of media was removed and supplemented with 100 μ l of internal standards (100 pg/ μ l, EtOH) and 100 μ l EtOH to bring the total EtOH to 10% by volume. Samples were centrifuged for 5 minutes at 3000 rpm to remove cellular debris, and the supernatants were decanted into solid phase extraction columns. All eicosanoid extractions were conducted using Strata-X SPE columns (Phenomenex). The columns were washed with 2 ml MeOH and then 2 ml H₂O prior to the samples being applied. After applying the sample, the columns were washed with 10% MeOH to remove non-adherent debris, and eicosanoids were then eluted off the column with 1 ml MeOH. The eluant was dried under vacuum and redissolved in 100 μ l of solvent A [water-acetonitrile-formic acid (63:37:0.02; v/v/v)] for LC/MS analysis.

Cell Quantitation

Eicosanoid levels were normalized to cell number using DNA quantitation. After the extracellular media was removed, the cells were scraped in 500 μ l PBS and stored at 4°C for DNA quantitation using the Broad Range DNA Quant-Kit (Invitrogen).

HPLC and Mass Spectrometry

The analysis of eicosanoids was performed by LC/MS/MS (26). Eicosanoids were separated by reverse-phase HPLC on a C18 column (2.1 mm \times 150 mm, Grace-Vydac) at a flow rate of 300 µl/min at 25°C. The column was equilibrated in Solvent A [water-acetonitrile-formic acid (63:37:0.02; v/v/v)], and samples were injected using a 50 µl injection loop and eluted with a linear gradient from 0%–20% solvent B [acetonitrile-isopropyl alcohol (50:50; v/v)] between 0 to 6 min; solvent B was increased to 55% from 6 to 6.5 min and held until 10 min. Solvent B was increased to 100% from 10 to 12 min and held until 13 min; solvent B was dropped to 0% by 13.5 min and held until 16 min.

Eicosanoids were analyzed using a tandem quadrupole mass spectrometer (ABI 4000 Q Trap®, Applied Biosystems) via multiple-reaction monitoring (MRM) in negative-ion mode. The electrospray voltage was -4.5 kV, the turbo ion spray source temperature was 525°C. Collisional activation of eicosanoid precursor ions used nitrogen as a collision gas. While arachidonic acid and its major metabolites (referred to as eicosanoids) were detected, by far

the major species at one hour was arachidonic acid, which was the focus of this study; for quantitation of the major eicosanoids produced, see (30).

Quantitative eicosanoid determination was performed by stable isotope dilution, as previously described (26). Results are determined as pmol of AA per million cells (mean \pm standard deviation).

RESULTS

PAP-1 inhibition reduces Kdo₂-Lipid A stimulated AA release

We initiated this study by characterizing GIVA PLA₂-mediated AA release from RAW 264.7 macrophages that were stimulated with Kdo₂-Lipid A. Kdo₂-Lipid A is a truncated LPS structure which is a specific binding-agonist of TLR-4 (24). The AA release time-course experiments were conducted over a period of 24 hours. As described in Figure 1, the maximal release of AA was observed to be 300–400 pmol AA/10⁶ cells at 1 hour after stimulation, after which AA release gradually returned to baseline levels. Subsequent AA release experiments were conducted at one hour following stimulation as this was determined to be the most optimal time point.

To ensure that GIVA PLA₂ is the enzyme that is responsible for mediating AA release under the experimental conditions employed, cells were pretreated with the GIVA PLA₂ specific inhibitor pyrrophenone (31) prior to stimulation. Pretreatment of the cells with pyrrophenone reduced Kdo₂-Lipid A stimulated AA release to background levels, as shown in Figure 2A; this result indicates that the Kdo₂-Lipid A stimulated AA release is mediated through the activity of GIVA PLA₂.

To evaluate whether PAP-1 may participate in the activation of GIVA PLA₂, we preincubated cells with the PAP-1 inhibitors BEL and propranolol prior to stimulation of the cells. Preincubation of the cells with bromoenol lactone (BEL) resulted in a ~70% decrease of Kdo₂-Lipid A stimulated AA release into the supernatant, as seen in Figure 2B. This result was corroborated by repeating the experiment with another inhibitor of PAP-1, propranolol, as shown in Figure 2C.

To address the possibility that PLC may play a role in this pathway, cells were treated with the classic PLC inhibitors D609 and U-71322 prior to stimulation with Kdo₂-Lipid A; neither inhibitor reduced AA release, suggesting that PLC is not involved in this signaling cascade (data not shown).

BEL and propranolol do not inhibit ATP stimulated AA release

The inhibitor data summarized in Figures 2B and 2C suggest that PAP-1 participates in the TLR-4 mediated activation of GIVA PLA₂. Cellular GIVA PLA₂ activity can also be induced by the treatment of cells with Ca^{+2} mobilizing agonists, such as the purinergic P2X7 receptor agonist ATP. Inhibited and uninhibited cells were treated with ATP prior to AA release quantitation. Stimulation of macrophages with ATP increased AA release into the supernatant (Figure 3). To ensure that the Ca^{+2} agonists had induced AA release via participation of GIVA PLA₂, cells were also treated with pyrrophenone prior to stimulation; exposure to pyrrophenone resulted in reduction of AA release to background levels. Neither BEL nor propranolol reduced AA release induced by this stimulus, indicating that the inhibitors do not affect GIVA PLA₂.

As an additional control, to ensure that an extracellular Ca^{+2} influx into the cell is not required in the TLR-4 mediated activation of the GIVA PLA₂, cells were cultured in the presence of EDTA and EGTA prior to stimulation with Kdo₂-Lipid A. Neither EDTA nor EGTA reduced

the AA release, indicating that Ca^{+2} influx is not required for TLR-4 mediated activation of the GIVA PLA₂ (Figure 4).

PAP-1 chemical inhibition does not reduce the synergistic activation of GIVA PLA₂

Recently, we reported that cells that were stimulated with both Kdo₂-Lipid A and ATP released more AA than the sum of AA released from cells stimulated separately, a phenomena referred to as synergy (27). The extent to which AA release is enhanced by joint Kdo₂-Lipid A and ATP stimuli can be calculated from:

Synergistic Activation - [Kdo ₂ - Lipid A+ATP]	
Synergistic Activation $\frac{1}{[Kdo_2 - Lipid A] + [ATP]}$	Equation 1

When synergy is present, the synergistic activation value is greater than 1.0, as joint Kdo₂-Lipid A and ATP stimulation result in AA release that exceeds the sum of the individual stimulations. If no synergy is present, and therefore joint stimulation equals the sum total of separate stimulations, the value will equal 1.0. Synergy-enhanced GIVA PLA₂ activation results in enhanced AA release that exceeds the sum total of TLR-4 and Ca⁺² mediated AA releases.

In the current study, sequential treatment of cells with Kdo₂-Lipid A and ATP resulted in a synergistic activation value of 2.5 ± 0.4 , as shown in Figure 5A. In this particular example, cells that were treated with Kdo₂-Lipid A and ATP released 1,600 pmol AA/10⁶ cells, which is four times greater than the amount of AA released by either Kdo₂-Lipid A or ATP alone and approximately 2.5 times greater than the sum total of separate Kdo₂-Lipid A and ATP stimulations. It is important to note that the absolute level of AA released via the [Kdo₂-Lipid A + ATP] stimulation shown in Figure 5A is not an indicator of synergy. To determine the extent to which synergy activation is present, one must compute the synergistic activation value through equation 1 and then evaluate it relative to a value of "1".

Figures 2 and 3 show that preincubation of macrophages with either BEL or propranolol prior to Kdo_2 -Lipid A treatment reduces AA release, whereas neither inhibitor affects ATP stimulated AA release. Similarly, preincubation of the cells with BEL or propranolol reduces the AA released from cells treated with [Kdo₂-Lipid A + ATP], as shown in Figure 5A. However, despite the reduction in [Kdo₂-Lipid A + ATP] stimulated AA release, the synergistic activation value was not reduced by the presence of the inhibitors (Figure 5B). This suggests that PAP-1 inhibition reduces AA release from cells treated with [Kdo₂-Lipid A + ATP] by reducing the Kdo₂-Lipid A portion of the stimulation, while not affecting the contribution of the synergy stimulus.

Exogenous DAG stimulates GIVA PLA2-mediated AA release

Since PAP-1 converts cellular PA into DAG, macrophages were treated with exogenous DAG to assess whether this would elicit AA release. The addition of DAG to unstimulated cells slightly increased AA release into the supernatant. Additionally, the supplementation of DAG to Kdo₂-Lipid A treated cells resulted in a further increased AA release from the cells relative to that of cells that were treated with only Kdo₂-Lipid A (Figure 6). To ensure that the enhanced AA release was due to additional activation of GIVA PLA₂, we preincubated cells with pyrrophenone prior to stimulation. Pyrrophenone inhibited AA release to background levels in both cases, indicating that the enhanced AA release elicited by DAG is mediated through the action of GIVA PLA₂.

Exogenous DAG restores AA release from macrophages inhibited by propranolol or BEL

We investigated whether the addition of exogenous DAG in an "add-back" fashion would restore AA release from cells that had been pretreated with BEL or propranolol. The addition of DAG to cells that had been pretreated with the PAP-1 inhibitors prior to stimulation with Kdo₂-Lipid A resulted in full restoration of AA release to levels seen in uninhibited cells (Figure 7A and B). This result suggests that the effect of PAP-1 inhibition is reversible by the addition of DAG, which is the product of PAP-1 mediated hydrolysis.

PKC chemical inhibition inhibits Kdo₂-Lipid A stimulated AA release, but not synergistic activation

Since our data suggests that inhibition of PAP-1 results in decreased Kdo₂-Lipid A stimulated AA release but does not affect the synergistic activation of GIVA PLA₂, we next investigated whether inhibition of PKC would corroborate the PAP-1 inhibitory data. Cells were treated with the general PKC inhibitor Ro 31-8220 prior to stimulation with Kdo₂-Lipid A, ATP and [Kdo₂-Lipid A + ATP] and the synergistic activation value was computed. Similar to that of the PAP-1 inhibitor results, inhibition of PKC resulted in baseline reduction of Kdo₂-Lipid A stimulated AA release and also significantly reduced [Kdo₂-Lipid A + ATP] stimulated AA release, while not having a significant affect on ATP stimulation (Figure 8A). Also, consistent with the PAP-1 inhibitory data, the synergistic activation value was not reduced relative to that of the uninhibited cells (Figure 8B).

DISCUSSION

The focus of this research was to investigate the relationship between PAP-1, PKC and the activation of GIVA PLA₂ in the TLR-4 signaling cascade. To evaluate cellular GIVA PLA₂ activity, macrophages were stimulated with the TLR-4 specific agonist Kdo₂-Lipid A and then AA release into the supernatant was quantitated as a metabolic indicator of GIVA PLA₂ function. In previous studies in which AA release was quantitated from macrophages, our laboratory and others made use of scintillation counting of radioactive AA as well as EIA quantitation of PGE₂ secreted to the supernatant as an indicator of cellular GIVA PLA₂ activity (19,28–31). Both techniques have important limitations that must be accounted for. Scintillation counting does not discriminate between AA release and AA-derived metabolites, such as eicosanoids. Utilization of commercial EIA kits to measure PGE₂ release as an indicator of AA release. In the current study, we make use of HPLC-MS methodology to quantitate AA release specifically from unlabelled cells.

We began this study by conducting time courses of AA release from macrophages stimulated with Kdo₂-Lipid A to evaluate the optimal time to measure AA release following cellular activation. Maximal AA release into the supernatant was observed at one hour following stimulation, and therefore this was chosen to be the time point at which subsequent experiments were conducted. Since AA release was utilized as an indicator of cellular GIVA PLA₂ activity, it was imperative to ensure that Kdo₂-Lipid A stimulated AA release was mediated via GIVA PLA₂. To achieve this, macrophages were pretreated with the GIVA PLA₂ specific inhibitor, pyrrophenone, prior to stimulation with Kdo₂-Lipid A. Pyrrophenone reduced AA release to background levels, indicating that GIVA PLA₂ is responsible for Kdo₂-Lipid A stimulated AA release.

To investigate whether PAP-1 participates in the Kdo₂-Lipid A stimulated GIVA PLA₂ activation, cells were treated with the PAP-1 inhibitors BEL and propranolol prior to stimulation. Preincubation of cells with BEL or propranolol resulted in a marked reduction of AA release into the supernatant. Since this result could be explained by direct inhibition of

GIVA PLA₂, we studied the effect of the inhibitors on AA release from cells stimulated with the Ca⁺² mobilizing agent, ATP. ATP activates GIVA PLA₂ by raising the cellular Ca⁺² levels, rather than through a TLR-4-mediated mechanism (30). Neither BEL nor propranolol reduced AA release from cells stimulated with ATP, indicating that the reduced Kdo₂-Lipid A stimulated AA release is not due to direct inhibition of cellular GIVA PLA₂ activity. It is worth noting that we have shown in a previous publication that activation of TLR-4 in macrophages does not increase intracellular Ca⁺² levels (32). Additionally, we confirmed in this manuscript that AA release from cells stimulated with ATP is mediated through GIVA PLA₂ by pyrrophenone inhibition. For the sake of completion, we investigated whether DAG generated by PLC may play a role in GIVA PLA₂ activation by dosing cells with the PLC inhibitors D609 and U-71322 prior to stimulation; neither inhibitor had an affect, suggesting that PLC is not involved.

Since our data suggests that PAP-1 inhibition reduced GIVA PLA2 activation in Kdo2-Lipid A treated cells, it was logical to assess what the effect of introducing DAG would be on GIVA PLA₂ activation. To evaluate whether DAG would directly activate cellular GIVA PLA₂, cells were stimulated with exogenous DAG prior to the measurement of AA release into the supernatant. The addition of DAG alone significantly increased AA release, relative to that of control cells. Additionally, supplementation of DAG to macrophages that were also pretreated with Kdo2-Lipid A resulted in enhanced AA release into the supernatant. The AA release was confirmed to be mediated through GIVA PLA₂ by pyrrophenone inhibition. Taken in conjunction with the PAP-1 inhibitory data, the fact that DAG evoked AA release suggests that DAG formation is an essential step in the stimulation of GIVA PLA₂ catalysis (33–37). Furthermore, considering the fact that the PAP-1 and PKC inhibitory data are similar would suggest that PKC is a likely molecular target of PAP-1 that also participates in the activation mechanism (38). Additionally, since PAP-1 inhibition reduced AA release, exogenous DAG was added to PAP-1 inhibited cells to discern whether this would restore AA release. It did restore AA release to levels released from uninhibited cells, suggesting that introduction of the DAG had compensated for the blocked PAP-1.

In a recent publication, our laboratory described the synergistic activation of GIVA PLA₂ in macrophages that had been treated with Kdo₂-Lipid A and ATP, resulting in AA release that exceeded the sum total of separate Kdo₂-Lipid A and ATP stimulations. This phenomenon has been referred to as synergy (27), whereby in combination Kdo₂-Lipid A activates GIVA PLA₂ by eliciting TLR-4 signaling and ATP does it through increased cellular Ca⁺²; but somehow leading to more product than the sum of each acting alone. Our current data suggest that these two activation mechanisms are quite distinct. First, it is shown that PAP-1 and PKC inhibition do not affect ATP stimulation, whereas they have a pronounced affect on Kdo₂-Lipid A mediated activation. Second, by adding EGTA and EDTA to chelate extracellular Ca⁺², we have shown that Kdo₂-Lipid A does not require an influx of Ca⁺², unlike ATP.

In the current manuscript, we have expanded upon our previous knowledge of GIVA PLA₂ synergistic activation by studying the chemical inhibition of the TLR-4 signaling while observing the effect upon synergy. Initially, PAP-1 chemical inhibitors were employed with the observation that PAP-1 inhibition resulted in a reduction of [Kdo₂-Lipid A and ATP] stimulated AA release, while the synergistic activation was unaffected. A likely explanation for this observation is that PAP-1 participates in the TLR-4 mediated signaling mechanism leading to GIVA PLA₂ activation, but does not participate in the synergistic activation. Possibilities for other independent effects of the TLR-4 activation are discussed elsewhere (30). Furthermore, this data suggests that blockage of the TLR-4 signaling mechanism does not inhibit activation of GIVA PLA₂ through synergy. To confirm this data, the general PKC inhibitor Ro 31-8220 was employed to block PKC, which lies downstream of PAP-1 in this pathway. Interestingly, inhibition of PKC resulted in total blockage of AA release elicited from

cells stimulated with Kdo₂-Lipid A while not reducing the synergistic activation value. This demonstrates that full inhibition of the TLR-4 signaling cascade leading to GIVA PLA₂ does not result in a reduction in synergistic activation, further suggesting that GIVA PLA₂ synergistic activation is the result of a unique signaling mechanism.

It is now known that the mammalian lipins 1, 2 and 3 exhibit PAP activity similar to the PAH1 gene product of yeast (43). Using qRT-PCR we have found in RAW cells that lipin 1 is expressed at significant levels. We also found that lipin 2 is expressed at even higher levels, but did not detect lipin 3 to any significant extent (data not shown). Therefore, we did not differentiate the extent to which each of the lipins contributes to the observed PAP-associated effects.

In summary, we present data that suggests that the activation of GIVA PLA₂ by Kdo₂-Lipid A requires the participation of PAP-1; it is likely that the function of PAP-1 in this model is to activate PKC. Four lines of evidence support such a theory. First, it is known that GIVA PLA₂ undergoes phosphorylation and activation involving PKC through a number of agonist-induced signaling pathways, including TLR-4 via LPS (42,44–46). Gene deletion or chemical inhibition of PKC in these studies resulted in loss of agonist-induced GIVA PLA₂ phosphorylation with reduced arachidonic acid release. Direct activation of PKC by the agonist PMA has been shown to result in the release of arachidonic acid and GIVA PLA₂ activation in a number of macrophage cell lines, including RAW 264.7 and U937 (20,42,43).

Second, all three proteins have been observed to co-localize on the nuclear membrane surface. PAP-1 contains a nuclear localization sequence and the protein expression is concentrated to the nuclear membrane region of HEK293 and 3T3-L1 cells (49,50). PKC has been shown to translocate to the nuclear membrane with rising membrane levels of DAG (51,52). Likewise, GIVA PLA₂ associates with the nuclear membrane after phosphorylation (53).

Third, PAP-1 was found to be associated with PKC ε and epidermal growth factor (EGF) receptor in EGF receptor immunoprecipitates. Following treatment with EGF and a second immunoprecipitation with an anti-PKC ε antibody, PAP-1 was found to be associated with PKC ε (54). GIVA PLA₂ is known to undergo phosphorylation and activation with subsequent AA release by way of an oxidative stress EGF receptor mediated pathway (55).

Finally, we have previously described that PAP-1 participates in TLR-4 mediated COX-2 expression, and our current results suggest that PAP-1 also participates in the TLR-4 activation of the GIVA PLA₂ (18,19). Collectively, these results suggest that PAP-1 plays a significant role in inflammatory signaling within the TLR-4 activated macrophage. Of course, these conclusions about the role of PAP-1 are based heavily on the inhibitor studies reported herein and, of course, "a specific inhibitor is one that has not yet been shown to lack specificity". While BEL and propranolol undoubtedly have other activities in cells, the overall argument for a PAP-1 role depends on their being specific for the specified steps in the TLR-4 activation of GIVA PLA₂. The observation that these inhibitors do not affect the activation with ATP bolsters the argument, but does not prove specificity, so additional approaches to test this hypothesis are certainly warranted.

References

- 1. Van Amersfoort ES, Van Berkel TJ, Kuiper J. Receptors, mediators, and mechanisms involved in bacterial sepsis and septic shock. Clin Microbiol Rev 2003;16:379–414. [PubMed: 12857774]
- Bos CL, Richel DJ, Ritsema T, Peppelenbosch MP, Versteeg HH. Prostanoids and prostanoid receptors in signal transduction. Int J Biochem Cell Biol 2004;36:1187–1205. [PubMed: 15109566]

- Jiang YJ, Lu B, Choy PC, Hatch GM. Regulation of cytosolic phospholipase A2, cyclooxygenase-1 and -2 expression by PMA, TNFalpha, LPS and M-CSF in human monocytes and macrophages. Mol Cell Biochem 2003;246:31–38. [PubMed: 12841340]
- 4. Dieter P, Kolada A, Kamionka S, Schadow A, Kaszkin M. Lipopolysaccharide-induced release of arachidonic acid and prostaglandins in liver macrophages: regulation by Group IV cytosolic phospholipase A2, but not by Group V and Group IIA secretory phospholipase A2. Cell Signal Mar; 2002 14(3):199–204. [PubMed: 11812647]
- Alonso F, Henson PM, Leslie CC. A cytosolic phospholipase in human neutrophils that hydrolyzes arachidonoyl-containing phosphatidylcholine. Biochim Biophys Acta 1986;878:273–280. [PubMed: 3092867]
- Kramer RM, Checani GC, Deykin A, Pritzker CR, Deykin D. Solubilization and properties of Ca2+dependent human platelet phospholipase A2. Biochim Biophys Acta 1986;878:394–403. [PubMed: 3756201]
- Nagase T, Uozumi N, Ishii S, Kume K, Izumi T, Ouchi Y, Shimizu T. Acute lung injury by sepsis and acid aspiration: a key role for cytosolic phospholipase A2. Nat Immunol 2000 Jul;1(1):42–6. [PubMed: 10881173]1:42–46 2000
- Marusic S, Leach MW, Pelker JW, Azoitei ML, Uozumi N, Cui J, Shen MW, DeClercq CM, Miyashiro JS, Carito BA, Thakker P, Simmons DL, Leonard JP, Shimizu T, Clark JD. Cytosolic phospholipase A2 alpha-deficient mice are resistant to experimental autoimmune encephalomyelitis. J Exp Med 2005;202:841–851. [PubMed: 16172261]
- Hegen M, Sun L, Uozumi N, Kume K, Goad ME, Nickerson-Nutter CL, Shimizu T, Clark JD. Cytosolic phospholipase A2alpha-deficient mice are resistant to collagen-induced arthritis. J Exp Med 2003;197:1297–1302. [PubMed: 12743172]
- Mitchell MP, Brindley DN, Hubscher G. Properties of phosphatidate phosphohydrolase. Eur J Biochem 1971;18:214–220. [PubMed: 5541505]
- Smith ME, Sedgwick B, Brindley DN, Hubscer G. The role of phosphatidate phosphohydrolase in glyceride biosynthesis. Eur J Biochem 1967;3:70–77. [PubMed: 4295052]
- Walton PA, Possmayer F. Translocation of Mg2+-dependent phosphatidate phosphohydrolase between cytosol and endoplasmic reticulum in a permanent cell line from human lung. Biochem Cell Biol 1986;64:1135–1140. [PubMed: 3030367]
- Walton PA, Possmayer F. The role of Mg2+-dependent phosphatidate phosphohydrolase in pulmonary glycerolipid biosynthesis. Biochim Biophys Acta 1984;796:364–372. [PubMed: 6095917]
- 14. Sturton RG, Brindley DN. Factors controlling the metabolism of phosphatidate by phosphohydrolase and phospholipase A-type activities. Effects of magnesium, calcium and amphiphilic cationic drugs. Biochim Biophys Acta 1980;619:494–505. [PubMed: 6257299]
- 15. Walton PA, Possmayer F. Mg2-dependent phosphatidate phosphohydrolase of rat lung: development of an assay employing a defined chemical substrate which reflects the phosphohydrolase activity measured using membrane-bound substrate. Anal Biochem 1985;151:479–486. [PubMed: 3006539]
- Carman GM, Han GS. Roles of phosphatidate phosphatase enzymes in lipid metabolism. Trends Biochem Sci 2006;31:694–699. [PubMed: 17079146]
- Han GS, Wu WI, Carman GM. The Saccharomyces cerevisiae Lipin homolog is a Mg2+-dependent phosphatidate phosphatase enzyme. J Biol Chem 2006;281:9210–8. [PubMed: 16467296]
- 18. Péterfy M, Phan J, Xu P, Reue K. Lipodystrophy in the *fld* mouse results from mutation of a new gene encoding a nuclear protein, lipin. Nat Genet 2001;27:121–124. [PubMed: 11138012]
- 19. Reue K, Dwyer JR. Lipin proteins and metabolic homeostasis. J Lipid Res. 200810.1194/jlr.R800052-JLR200Epub ahead of print
- Johnson CA, Balboa MA, Balsinde J, Dennis EA. Regulation of cyclooxygenase-2 expression by phosphatidate phosphohydrolase in human amnionic WISH cells. J Biol Chem 1999;274:27689– 27693. [PubMed: 10488110]
- Grkovich A, Johnson CA, Buczynski MW, Dennis EA. Lipopolysaccharide-induced cyclooxygenase-2 expression in human U937 macrophages is phosphatidic acid phosphohydrolase-1-dependent. J Biol Chem 2006;281:32978–32987. [PubMed: 16950767]

- Huffman TA, Mothe-Satney I, Lawrence JC Jr. Insulin-stimulated phosphorylation of lipin mediated by the mammalian target of rapamycin. Proc Natl Acad Sci U S A 2002;99:1047–1052. [PubMed: 11792863]
- 23. Jiang Y, Lu Z, Zang Q, Foster DA. Regulation of phosphatidic acid phosphohydrolase by epidermal growth factor. Reduced association with the EGF receptor followed by increased association with protein kinase Cepsilon. J Biol Chem 1996;271:29529–29532. [PubMed: 8939878]
- 24. Exton JH. Signaling through phosphatidylcholine breakdown. J Biol Chem 1990;265:1–4. [PubMed: 2104616]
- 25. Exton JH. Phosphatidylcholine breakdown and signal transduction. Biochim Biophys Acta 1994;1212:26–42. [PubMed: 8155724]
- Testerink C, Munnik T. Phosphatidic acid: a multifunctional stress signaling lipid in plants. Trends Plant Sci 2005;10:368–375. [PubMed: 16023886]
- 27. Raetz CR, Garrett TA, Reynolds CM, Shaw WA, Moore JD, Smith DC Jr, Ribeiro AA, Murphy RC, Ulevitch RJ, Fearns C, Reichart D, Glass CK, Benner C, Subramaniam S, Harkewicz R, Bowers-Gentry RC, Buczynski MW, Cooper JA, Deems RA, Dennis EA. Kdo2-Lipid A of Escherichia coli, a defined endotoxin that activates macrophages via TLR-4. J Lipid Res 2006;47:1097–1111. [PubMed: 16479018]
- Deems R, Buczynski MW, Bowers-Gentry R, Harkewicz R, Dennis EA. Detection and quantitation of eicosanoids via high performance liquid chromatography-electrospray ionization-mass spectrometry. Methods Enzymol 2007;432:59–82. [PubMed: 17954213]
- Hall LM, Murphy RC. Activation of human polymorphonuclear leukocytes by products derived from the peroxidation of human red blood cell membranes. Chem Res Toxicol 1998;11:1024–1031. [PubMed: 9760276]
- Buczynski MW, Stephens DL, Bowers-Gentry RC, Grkovich A, Deems RA, Dennis EA. TLR-4 and sustained calcium agonists synergistically produce eicosanoids independent of protein synthesis in RAW264.7 cells. J Biol Chem 2007;282:22834–47. [PubMed: 17535806]
- 31. Burke JE, Dennis EA. Phospholipase A2 Biochemistry. Cardiovascular Drugs and Therapy. 200810.1007/s10557-008-6132-9Epub ahead of print
- Balboa MA, Balsinde J, Dennis EA. Involvement of phosphatidate phosphohydrolase in arachidonic acid mobilization in human amnionic WISH cells. J Biol Chem 1998;273:7684–7690. [PubMed: 9516474]
- 33. Balboa MA, Perez R, Balsinde J. Amplification mechanisms of inflammation: paracrine stimulation of arachidonic acid mobilization by secreted phospholipase A2 is regulated by cytosolic phospholipase A2-derived hydroperoxyeicosatetraenoic acid. J Immunol 2003;171:989–994. [PubMed: 12847271]
- Balsinde J, Balboa MA, Dennis EA. Inflammatory activation of arachidonic acid signaling in murine P388D1 macrophages via sphingomyelin synthesis. J Biol Chem 1997;272:20373–20377. [PubMed: 9252342]
- Balsinde J. Roles of various phospholipases A2 in providing lysophospholipid acceptors for fatty acid phospholipid incorporation and remodeling. Biochem J 2002;364:695–702. [PubMed: 12049633]
- 36. Asmis R, Randriamampita C, Tsien RY, Dennis EA. Intracellular Ca2+, inositol 1,4,5-trisphosphate and additional signalling in the stimulation by platelet-activating factor of prostaglandin E2 formation in P388D1 macrophage-like cells. Biochem J 1994;298(Pt 3):543–551. [PubMed: 8141766]
- Balsinde J, Balboa MA, Li WH, Llopis J, Dennis EA. Cellular regulation of cytosolic group IV phospholipase A2 by phosphatidylinositol bisphosphate levels. J Immunol 2000;164:5398–5402. [PubMed: 10799904]
- Mosior M, Six DA, Dennis EA. Group IV cytosolic phospholipase A2 binds with high affinity and specificity to phosphatidylinositol 4,5-bisphosphate resulting in dramatic increases in activity. J Biol Chem 1998;273:2184–2191. [PubMed: 9442060]
- Casas J, Gijon MA, Vigo AG, Crespo MS, Balsinde J, Balboa MA. Phosphatidylinositol 4,5bisphosphate anchors cytosolic group IVA phospholipase A2 to perinuclear membranes and decreases its calcium requirement for translocation in live cells. Mol Biol Cell 2006;17:155–162. [PubMed: 16221889]

- 40. Pettus BJ, Bielawska A, Subramanian P, Wijesinghe DS, Maceyka M, Leslie CC, Evans JH, Freiberg J, Roddy P, Hannun YA, Chalfant CE. Ceramide 1-phosphate is a direct activator of cytosolic phospholipase A2. J Biol Chem 2004;279:11320–11326. [PubMed: 14676210]
- Pettus BJ, Bielawska A, Spiegel S, Roddy P, Hannun YA, Chalfant CE. Ceramide kinase mediates cytokine- and calcium ionophore-induced arachidonic acid release. J Biol Chem 2003;278:38206– 38213. [PubMed: 12855693]
- 42. Luo SF, Lin WN, Yang CM, Lee CW, Liao CH, Leu YL, Hsiao LD. Induction of cytosolic phospholipase A2 by lipopolysaccharide in canine tracheal smooth muscle cells: involvement of MAPKs and NF-kappaB pathways. Cell Signal 2006;18:1201–1211. [PubMed: 16278065]
- 43. Carman GM, Han GS. Phosphatidic acid phosphatase, a key enzyme in the regulation of lipid synthesis. J Biol Chem 2008;284:2593–2597. [PubMed: 18812320]
- 44. Xu J, Weng YI, Simonyi A, Krugh BW, Liao Z, Weisman GA, Sun GY. Role of PKC and MAPK in cytosolic PLA2 phosphorylation and arachadonic acid release in primary murine astrocytes. J Neurochem 2002;83:259–270. [PubMed: 12423237]
- 45. Xu J, Weng YI, Simonyi A, Krugh BW, Liao Z, Weisman GA, Sun GY. Role of PKC and MAPK in cytosolic PLA2 phosphorylation and arachadonic acid release in primary murine astrocytes. J Neurochem 2002;83:259–270. [PubMed: 12423237]
- Li Q, Subbulakshmi V, Oldfield CM, Aamir R, Weyman CM, Wolfman A, Cathcart MK. PKCalpha regulates phosphorylation and enzymatic activity of cPLA2 in vitro and in activated human monocytes. Cell Signal 2007;19:359–366. [PubMed: 16963226]
- 47. Lin WW, Chen BC. Distinct PKC isoforms mediate the activation of cPLA2 and adenylyl cyclase by phorbol ester in RAW264.7 macrophages. Br J Pharmacol 1998;125:1601–1609. [PubMed: 9884090]
- 48. Balsinde J, Balboa MA, Insel PA, Dennis EA. Differential regulation of phospholipase D and phospholipase A2 by protein kinase C in P388D1 macrophages. Biochem J 1997;321(Pt 3):805–809. [PubMed: 9032469]
- Peterfy M, Phan J, Reue K. Alternatively spliced lipin isoforms exhibit distinct expression pattern, subcellular localization, and role in Adipogenesis. J Biol Chem 2005;280:32883–32889. [PubMed: 16049017]
- 50. Peterfy M, Phan J, Xu P, Reue K. Lipodystrophy in the fld mouse results from mutation of a new gene encoding a nuclear protein, lipin. Nat Genet 2001;27:121–124. [PubMed: 11138012]
- 51. Deacon EM, Pettitt TR, Webb P, Cross T, Chahal H, Wakelam MJ, Lord JM. Generation of diacylglycerol molecular species through the cell cycle: a role for 1-stearoyl, 2-arachidonyl glycerol in the activation of nuclear protein kinase C-betaII at G2/M. J Cell Sci 2002;115:983–989. [PubMed: 11870217]
- 52. Divecha N, Banfic H, Irvine RF. The polyphosphoinositide cycle exists in the nuclei of Swiss 3T3 cells under the control of a receptor (for IGF-I) in the plasma membrane, and stimulation of the cycle increases nuclear diacylglycerol and apparently induces translocation of protein kinase C to the nucleus. EMBO J 1991;10:3207–3214. [PubMed: 1655412]
- Marshall J, Krump E, Lindsay T, Downey G, Ford DA, Zhu P, Walker P, Rubin B. Involvement of cytosolic phospholipase A2 and secretory phospholipase A2 in arachidonic acid release from human neutrophils. J Immunol 2000;164:2084–2091. [PubMed: 10657662]
- 54. Jiang Y, Lu Z, Zang Q, Foster DA. Regulation of phosphatidic acid phosphohydrolase by epidermal growth factor. Reduced association with the EGF receptor followed by increased association with protein kinase Cepsilon. J Biol Chem 1996;271:29529–29532. [PubMed: 8939878]
- 55. Pawliczak R, Huang XL, Nanavaty UB, Lawrence M, Madara P, Shelhamer JH. Oxidative stress induces arachidonate release from human lung cells through the epithelial growth factor receptor pathway. Am J Respir Cell Mol Biol 2002;27:722–731. [PubMed: 12444032]



Figure 1. Kdo₂-Lipid A stimulated AA release time-course

RAW 264.7 macrophages were cultured in the presence (•) or absence (\circ) of 100 ng/mL Kdo₂-Lipid A, the media was collected at the indicated time points and then subsequently analyzed for AA release by HPLC-MS. The insert is a separate experiment showing Kdo₂–Lipid A stimulated AA release over a 60 minute time-course. A representative experiment of three individual experiments is shown. Data are expressed as mean values ± S.D. of three individual replicates.



Figure 2. Inhibition of PAP-1 blocks the Kdo2-Lipid A stimulated release of AA

AA release was measured from RAW 264.7 macrophages preincubated with and without (A) 1 μ M pyrrophenone, (B) 25 μ M BEL, or (C) 50 μ M propranolol prior to stimulation with (\blacksquare) and without (\Box) 100 ng/mL Kdo₂-Lipid A. The media was collected at 1 hour following Kdo₂-Lipid A stimulation and analyzed for AA release by HPLC-MS. A representative experiment of three individual experiments is shown. Data are expressed as mean values \pm S.D. of three individual replicates.



Figure 3. Inhibition of PAP-1 does not block the ATP stimulated release of AA

AA release was measured from RAW 264.7 macrophages preincubated with and without 1 μ M pyrrophenone, 25 μ M BEL or 50 μ M propranolol prior to stimulation with (**n**) and without (**n**) 2 mM ATP. The media was collected at 10 minutes following stimulation and analyzed for AA release. A representative experiment of three individual experiments is shown. Data are expressed as mean values \pm S.D. of three individual replicates.



Figure 4. The Ca⁺² chelators EDTA and EGTA do not reduce Kdo₂-Lipid A stimulated AA release *A*, AA release was measured from RAW 264.7 macrophages preincubated with and without 2 mM EDTA or EGTA prior to stimulation with (\blacksquare) and without (\Box) 100 ng/mL Kdo₂-Lipid A. The media was collected at 1 hour following stimulation and analyzed for AA release. A representative experiment of three individual experiments is shown. Data are expressed as mean values \pm S.D. of three individual replicates.

Grkovich et al.



Figure 5. Inhibition of PAP-1 blocks synergy-enhanced AA release, but does not reduce the synergistic activation value

A, AA release was measured from RAW 264.7 macrophages preincubated with and without $25 \,\mu\text{M}$ BEL or $50 \,\mu\text{M}$ propranolol prior to stimulation with and without $100 \,\text{ng/mL}$ Kdo₂-Lipid A for 50 minutes and with and without 2 mM ATP for 10 minutes (as indicated in the legend). The media was collected at 1 hour following the addition of the 100 ng/mL Kdo₂-Lipid A agonist and analyzed for AA release. *B*, Synergistic activation values were calculated for uninhibited cells, preincubated with 25 μ M BEL or 50 μ M propranolol. A representative experiment of three individual experiments is shown. Data are expressed as mean values \pm S.D. of three individual replicates.



Figure 6. Exogenous DAG bolsters the Kdo2-Lipid A stimulated release of AA

AA release into the media was measured from RAW 264.7 macrophages that were stimulated with and without 100 ng/mL Kdo₂-Lipid A (as indicated) and in the absence (white bars) and presence (black bars) of 50 μ M DAG for 1 hour. Additionally, grey bars indicate AA release from cells preincubated with 1 μ M pyrrophenone for 30 minutes prior to the addition of DAG. A representative experiment of three individual experiments is shown. Data are expressed as mean values \pm S.D. of three individual replicates.

Grkovich et al.





AA release was measured from cells preincubated with and without (A) 25 μ M BEL or (B) 50 μ M propranolol (as indicated) prior to stimulation with 100 ng/mL Kdo₂-Lipid A (black bars) and 50 μ M DAG (grey bars). The media was collected at 1 hour following stimulation and analyzed for AA release. A representative experiment of three individual experiments is shown. Data are expressed as mean values \pm S.D. of three individual replicates.

Grkovich et al.



Figure 8. Inhibition of PKC blocks synergy-enhanced AA release, but does not reduce the synergistic activation value

A, AA release was measured from RAW 264.7 macrophages preincubated with and without 5 μ M Ro 31-8220 prior to stimulation with and without 100 ng/mL Kdo₂-Lipid A for 50 minutes and with and without 2 mM ATP for 10 minutes (as indicated in the legend). The media was collected at 1 hour following the addition of the 100 ng/mL Kdo₂-Lipid A agonist and analyzed for AA release. *B*, Synergistic activation values were calculated for uninhibited cells and those preincubated with 5 μ M Ro 31-8220. A representative experiment of three individual experiments is shown. Data are expressed as mean values ± S.D. of three individual replicates.