

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

A STUDY OF Ke4 DECAYS

### Permalink

<https://escholarship.org/uc/item/16d8c94d>

### Authors

Ely, Robert P.  
Gidal, George  
Hagopian, Vasken  
et al.

### Publication Date

1968-10-01

LIBRARY AND  
DOCUMENTS SECTION

UCRL-18626

*g-2*

A STUDY OF  $K_{e4}$  DECAYS

Robert P. Ely, Jr., George Gidal,  
Vasken Hagopian, George E. Kalmus,  
Kelvin Billing, Frederick W. Bullock, Michael J. Esten,  
M. Govan, Cyril Henderson, William L. Knight,  
F. Russell Stannard, Ortwin Treutler, Ugo Camerini,  
David Cline, William F. Fry, Herman Haggerty,  
Robert H. March, and William J. Singleton

October 1968

TWO-WEEK LOAN COPY

This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545

**LAWRENCE RADIATION LABORATORY**  
**UNIVERSITY of CALIFORNIA BERKELEY**

UCRL-18626

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Physical Review

UCRL-18626  
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

A STUDY OF  $K_{e4}$  DECAYS

Robert P. Ely, Jr., George Gidal,  
Vasken Hagopian, George E. Kalmus,  
Kelvin Billing, Frederick W. Bullock, Michael J. Esten,  
M. Govan, Cyril Henderson, William L. Knight,  
F. Russell Stannard, Ortwin Treutler, Ugo Camerini,  
David Cline, William F. Fry, Herman Haggerty,  
Robert H. March, and William J. Singleton

October 1968

A STUDY OF  $K_{e4}$  DECAYS

Robert P. Ely, Jr., George Gidal,  
Vasken Hagopian,\* and George E. Kalmus

Lawrence Radiation Laboratory  
University of California  
Berkeley, California

Kelvin Billing, Frederick W. Bullock, Michael J. Esten,  
M. Govan, Cyril Henderson,† William L. Knight,‡  
F. Russell Stannard, and Ortwin Treutler

Department of Physics  
University College London  
London

Ugo Camerini, David Cline, William F. Fry,  
Herman Haggerty, Robert H. March, and William J. Singleton

Department of Physics  
University of Wisconsin  
Madison, Wisconsin

October 1968

ABSTRACT

A study of  $13.3 \times 10^6$  stopped  $K^+$  in a heavy-liquid bubble chamber yielded 269  $K_{e4}$  decays of the type  $e^+ \pi^+ \pi^- \nu$ , a total greater by a factor of 4 than the number found in a previous experiment. No examples of  $e^- \pi^+ \pi^- \bar{\nu}$  were found. With 95% confidence the upper limit for the decay rate of  $K_{e4}(e^-)$  was found to be  $56 \text{ sec}^{-1}$ . The rate for  $K_{e4}(e^+)$  was found to be  $(2.60 \pm 0.30) \times 10^3 \text{ sec}^{-1}$ .

The angular distributions and the dipion invariant-mass plot have been fitted by varying the form factors  $f_s$ ,  $f_p$ ,  $g$ , and  $h$ , and the difference between s- and p-wave  $\pi$ - $\pi$  phase-shift. Two acceptable solutions have been found. Both agree that the vector form factor,  $h$ , is significantly nonzero and that its sign is negative with respect to that of  $f_s$ .

Also it has been found necessary to include  $f_p$  in order to obtain a good fit. Although both solutions give the magnitude of the phase difference to be  $25 \pm 9$  deg, the two estimates have opposite sign.

No evidence of a  $\sigma$  resonance is seen. The angular distributions are found to be consistent with time-reversal invariance, and with the locality of lepton production.

## I. INTRODUCTION

The rarity of the  $K_{e4}$  decay makes it a difficult process to study. Thus, in spite of its considerable theoretical interest, there has to date been only one experimental investigation,<sup>1</sup> based on 69 events.

The experiment presented here represents an extension of this work with statistics increased by a factor 4. This has been made possible through the use of a larger heavy-liquid bubble chamber, permitting a greater number of decays per picture. Also the number of pictures taken this time was greater by a factor of more than 2.

Progress reports on this work have already been presented at various conferences,<sup>2</sup> and a preliminary analysis of the data has been given by Berends, Donnachie, and Oades.<sup>3</sup>

The decay modes of interest are

$$K_{e4}(e^+): \quad K^+ \rightarrow \pi^+ \pi^- e^+ \nu; \quad (1)$$

$$K_{e4}(e^-): \quad K^+ \rightarrow \pi^+ \pi^+ e^- \bar{\nu}. \quad (2)$$

The general form of the matrix element has been discussed by several authors,<sup>4</sup> assuming a V - A theory.<sup>5</sup> Reaction (2) has been shown to proceed almost entirely through the axial-vector current, whereas reaction (1) is a mixture of vector and axial-vector.

Several interesting topics are investigated in this paper. Firstly, the  $\Delta Q = \Delta S$  selection rule forbids  $K_{e4}(e^-)$ , so a search for this reaction permits a test of the rule for axial-vector currents in weak interactions.<sup>4, 6</sup> The rate for  $K_{e4}(e^+)$  is compared with several predictions,<sup>4, 6-11</sup> some of which include the effects of final-state interactions.

The angular correlations between the decay products lead to a determination of the form factors, and these are checked against theoretical prediction.<sup>11</sup> They also permit a test of time-reversal invariance.

Lastly,  $K_{e4}$  decays are notable for their unique property of allowing one to investigate the s-wave  $\pi$ - $\pi$  interaction in the absence of any additional strongly interacting particles.<sup>12-16</sup> This arises because in the final state of the decay, the two pions are accompanied only by two leptons.

## II. THEORY

In this section we are following the treatment of the theory given in Pais and Treiman's paper.<sup>16</sup>

The matrix element in first-order perturbation theory (aside from the usual kinematic factors) is

$$M[K_{e4}(e^+)] = |G/\sqrt{2}| [\bar{\nu}\gamma^\lambda(1+\gamma_5)e] \langle \pi^+\pi^- | J_\lambda^V + J_\lambda^A | K^+ \rangle,$$

where  $J^V$  and  $J^A$  are the strangeness-changing vector and axial-vector currents of the hadrons.

From invariance considerations

$$\langle \pi^+\pi^- | J_\lambda^V | K^+ \rangle = \frac{h}{M^3} \epsilon_{\lambda\mu\nu\sigma} K_\mu P_\nu Q_\sigma$$

and

$$\langle \pi^+\pi^- | J_\lambda^A | K^+ \rangle = \frac{1}{M} \{fP_\lambda + gQ_\lambda + r(K-P)_\lambda\},$$

where  $M$  is the mass of the  $K$  meson, and  $P = (p_+ + p_-)$ ,  $Q = (p_+ - p_-)$ , where  $p_+$  and  $p_-$  are the  $\pi^+$  and  $\pi^-$  four-momenta and  $K$  is the kaon four-momentum. The dimensionless form factors,  $f$ ,  $g$ ,  $r$ , and  $h$ , are functions of the invariant quantities  $P^2$ ,  $(K-P)^2$ , and  $(K \cdot Q)$ , or equivalently,  $S_\pi$ ,  $S_f$ , and  $\theta_\pi$ , where  $S_\pi$  is the square of the  $\pi^+\pi^-$  invariant mass



(i. e.,  $S_\pi = -P^2$ ) and  $S_\ell$  is the square of the  $e^+ \nu$  invariant mass (i. e.,  $S_\ell = -L^2$ , where  $L = p_e + p_\nu$ ;  $p_e$  and  $p_\nu$  are the positron and neutrino four-momenta). In Fig. 1 we have the definition of  $\theta_\pi$ , which is the angle of the  $\pi^+$  in the dipion center-of-mass system with respect to the dipion line of flight.

Apart from spin,  $K_{e4}$  decay is kinematically parameterized by five independent variables. We have chosen these five to be  $S_\pi$ ,  $S_\ell$ ,  $\theta_\pi$ ,  $\theta_\ell$ , and  $\phi$ . The latter two angles are also defined in Fig. 1.

The probability distribution summed over lepton spins is of the form

$$d^5W = \frac{\pi^2 G^2 \sin^2 \theta_c}{(2\pi)^8 16M^5} X \left( \frac{Q^2}{S_\pi} \right)^{1/2} \left( 1 - \frac{m^2}{S_\ell} \right)^2 I(S_\pi, S_\ell, \theta_\pi, \theta_\ell, \phi) \\ \times dS_\pi dS_\ell d\cos\theta_\pi d\cos\theta_\ell d\phi, \quad (3)$$

where  $\theta_c$  is the Cabbibo angle,  $m$  is the electron mass, and  $X$  is defined as  $X = [(P \cdot L)^2 - S_\pi \cdot S_\ell]^{1/2}$ .

The distribution function  $I$  has an explicit structure in the variables  $\theta_\ell$  and  $\phi$ , which do not enter into the form factors. By grouping terms according to their behavior with respect to these variables we can examine the spectra more readily.

For  $I$  we have

$$I = I_1 + I_2 \cos 2\theta_\ell + I_3 \sin^2 \theta_\ell \cos 2\phi \\ + I_4 \sin 2\theta_\ell \cos \phi + I_5 \sin \theta_\ell \cos \phi + I_6 \cos \theta_\ell + I_7 \sin \theta_\ell \sin \phi \\ + I_8 \sin 2\theta_\ell \sin \phi + I_9 \sin^2 \theta_\ell \sin 2\phi. \quad (4)$$

Neglecting terms involving  $m^2/S_l$ , the functions  $I_1 \cdots I_9$  depend on  $S_\pi$ ,  $S_l$ , and  $\theta_\pi$ , in the following manner (for the complete expressions see Ref. 16):

$$\begin{aligned}
 I_1 &= \frac{1}{4} [ |F_1|^2 + \frac{3}{2} \sin^2 \theta_\pi ( |F_2|^2 + |F_3|^2 ) ], \\
 I_2 &= -\frac{1}{4} [ |F_1|^2 - \frac{1}{2} \sin^2 \theta_\pi ( |F_2|^2 + |F_3|^2 ) ], \\
 I_3 &= -\frac{1}{4} [ |F_2|^2 - |F_3|^2 ] \sin^2 \theta_\pi, \\
 I_4 &= \frac{1}{2} \operatorname{Re} F_1^* F_2 \sin \theta_\pi, \\
 I_5 &= -\operatorname{Re} F_1^* F_3 \sin \theta_\pi, \\
 I_6 &= -\operatorname{Re} F_2^* F_3 \sin^2 \theta_\pi, \\
 I_7 &= -\operatorname{Im} F_1^* F_2 \sin \theta_\pi, \\
 I_8 &= \frac{1}{2} \operatorname{Im} F_1^* F_3 \sin \theta_\pi, \\
 I_9 &= -\frac{1}{2} \operatorname{Im} F_2^* F_3 \sin^2 \theta_\pi.
 \end{aligned} \tag{5}$$

$F_1$ ,  $F_2$ , and  $F_3$  are the following combinations of kinematic factors and form factors:

$$\begin{aligned}
 F_1 &= Xf - (PL)(Q^2/S_\pi)^{1/2} g \cos \theta_\pi, \\
 F_2 &= (Q^2 S_l)^{1/2} g, \\
 F_3 &= (Q^2 S_l)^{1/2} X(h/M^2).
 \end{aligned} \tag{6}$$

Note that the r form factor is unimportant in  $K_{e4}$  decay, as it is always multiplied by  $m^2/S_l$ , and so does not appear in (6).

The form factors  $f$ ,  $g$ , and  $h$  carry direct strong-interaction information, assuming time-reversal invariance holds. In a partial-wave expansion of the form factors with respect to the angular momentum of the dipion system, a partial-wave amplitude of definite angular momentum and isospin must have the phase of the corresponding pion-pion scattering amplitude. The odd partial waves have  $I = 1$  for the dipion system, whereas even partial waves contribute to both  $I = 0$  and  $I = 2$  states. Invoking the  $\Delta I = 1/2$  rule for semileptonic decays, we are left with  $\ell = 0$ ,  $I = 0$ , and  $\ell = 1$ ,  $I = 1$  states, assuming that states with  $\ell \geq 2$  are not important at these low energies.

The terms in the partial-wave expansion up to and including  $\ell = 1$  can now be written

$$\begin{aligned} f &= \tilde{f}_s e^{i\delta_s} + \tilde{f}_p e^{i\delta_p} \cos\theta_\pi, \\ g &= \tilde{g} e^{i\delta_p}, \\ h &= \tilde{h} e^{i\delta_p}, \end{aligned} \quad (7)$$

where  $\tilde{f}_s$ ,  $\tilde{f}_p$ ,  $\tilde{g}$ , and  $\tilde{h}$  are real functions of the variables  $S_\pi$  and  $S_\ell$ , and the phases  $\delta_s$  and  $\delta_p$  are the pion-pion phase shifts, and are functions of  $S_\pi$ .

By substituting (7) into (6) we obtain

$$\begin{aligned} F_1 &= X \tilde{f}_s e^{i\delta_s} + X \tilde{f}_p e^{i\delta_p} \cos\theta_\pi - (P \cdot L)(Q^2/S_\pi)^{1/2} \tilde{g} e^{i\delta_p} \cos\theta_\pi, \\ F_2 &= (Q^2 S_\ell)^{1/2} \tilde{g} e^{i\delta_p}, \\ F_3 &= (Q^2 S_\ell)^{1/2} X (\tilde{h} e^{i\delta_p}/M^2). \end{aligned} \quad (8)$$

Although this is a perfectly valid substitution as long as  $\tilde{f}_s$ ,  $\tilde{f}_p$ ,  $\tilde{g}$ , and  $\tilde{h}$  are functions of  $S_\pi$  and  $S_\ell$ . In the case of the p-wave form factors  $\tilde{f}_p$ ,  $\tilde{g}$ , and  $\tilde{h}$ , however, there is a known energy dependence due to the angular momentum barrier, which may be explicitly taken out of the form factors. In fact this has been done for  $\tilde{g}$  and  $\tilde{h}$ , but not yet for  $\tilde{f}_p$ . To do this for  $\tilde{f}_p$  we somewhat arbitrarily substitute  $\beta X \tilde{f}'_p / M^2$ , where  $\beta \equiv (Q^2 / S_\pi)^{1/2}$ . The exact form the expressions should take is unknown, but our choice has the essential feature of forcing  $\tilde{f}_p$  to zero when  $S_\pi$  equals  $4\mu^2$ , where  $\mu$  is the mass of the pion.

$$\therefore F_1 = X \tilde{f}_s e^{i\delta s} + \frac{\beta X^2}{M^2} \tilde{f}'_p e^{i\delta p} \cos \theta_\pi - (\mathbf{P} \cdot \mathbf{L}) \beta \tilde{g} e^{i\delta p} \cos \theta_\pi. \quad (8')$$

By substituting Eqs. (8) and (8') into (5) we obtain the distribution function  $I$  in terms of kinematic factors, the form factors  $\tilde{f}_s$ ,  $\tilde{f}'_p$ ,  $\tilde{g}$ ,  $\tilde{h}$ , and the s- and  $\tilde{p}$ -wave phase shifts  $\delta s$  and  $\delta p$ . In fact, since the absolute phase is arbitrary, the difference of the phases ( $\delta s - \delta p$ ) is the measured quantity.

In this treatment we have assumed that the leptons are produced at a point. This implies that the  $\theta_\ell$  distribution can be fitted by an expression of the form

$$dW/d\cos\theta_\ell = A(1 + B \cos\theta_\ell + C \cos 2\theta_\ell), \quad (9)$$

and the  $\phi$  distribution is fitted by

$$dW/d\phi = \alpha(1 + \beta \cos\phi + \gamma \sin\phi + \delta \sin^2\phi + \epsilon \sin 2\phi) \quad (10)$$

[these distributions are obtained by integrating (3) over all variables except  $\theta_\ell$  and  $\phi$ ].

Should extra terms be needed in either of these distributions, this would be a violation of the assumption of locality for the lepton production.

Four approaches were used in the analysis of the data from this experiment.

### 1st Method

Expression (3) was used to generate Monte Carlo  $K_{e4}$  events for particular values of  $\nu (\equiv \tilde{f}'_p / \tilde{f}_s)$ ,  $\eta (\equiv \tilde{g} / \tilde{f}_s)$ , and  $\gamma (\equiv \tilde{h} / \tilde{f}_s)$ , and of  $a_0$ , where  $a_0$  is related to the s-wave phase shift by the Chew-Mandelstam effective range formula,

$$\cot \delta s = \frac{1}{a_0 \beta} + \frac{2}{\pi} \ln \left\{ \frac{(S_\pi)^{1/2} (1+\beta)}{2\mu} \right\}.$$

We have assumed that  $\delta p$  is due to the tail of the  $\rho$  meson and is therefore very small in our mass range and can be neglected. We have generally taken the form factors to be constant, but have also investigated the effect of allowing  $\tilde{f}_s$  to be enhanced. When it has been enhanced we have used the relativistic Watson enhancement factor,

$$\tilde{f}_s = \frac{\tilde{f}_s^0 \sin \delta s}{a_0 \beta},$$

where  $a_0$  is in pion Compton wavelengths.

The Monte Carlo-generated events were subjected to the same cuts as the experimental data (the cuts arise from event-selection criteria to be discussed later). These events were then used to obtain the appropriately modified theoretical distributions of the measured variables, which were then compared with the experimental plots by a  $\chi^2$  technique. By changing the values of the input parameters, the gross features of

the  $\chi^2$  map were studied. However, this procedure, when dealing with four variables, is costly in computer time, and therefore the optimum set of parameters necessary to minimize  $\chi^2$  was not obtained by this technique.

### 2nd Method

The optimization was performed by a program which simultaneously fitted the  $\cos\theta_\pi$ ,  $\cos\theta_l$ ,  $\phi$ , and  $(S_\pi)^{1/2}$  distributions. The sum of the  $\chi^2$  for the four distributions was minimized by varying the form factors  $\nu$ ,  $\eta$ , and  $\gamma$ , and the average phase shift  $\langle \delta s - \delta p \rangle$ . The latter is the average value of  $(\delta s - \delta p)$  taken over our mass spectrum. The fit was made to the theoretical one-dimensional plots suitably modified by the Monte Carlo program for the effects of cuts in the data.

For the conditions prevailing in this particular experiment, these modified distributions have the forms given in the Appendix.

### 3rd Method

Expression (3) was evaluated for each event and the results were used to construct a likelihood function. A search program was employed to obtain the maximum value of the likelihood as a function of the free parameters, i. e., the form factors and the scattering length. Because of biases introduced in certain variables by the selection criteria, a restricted subset of the data was utilized and the range of integration suitably modified.

### 4th Method

Pais and Treiman<sup>16</sup> point out that information on  $\langle \delta s - \delta p \rangle$  can be obtained by looking at the average values of certain I coefficients (Eq. 5).

This method, unlike the first three approaches, yields information independent of the values of the form factors, and of assumptions regarding their energy dependence. It can be seen that the values of the  $I$ 's are governed only by the number of events in various regions of the two-dimensional plot of  $\phi$  vs  $\cos\theta_\ell$ .

Specifically, we have

$$\tan \langle \delta s - \delta p \rangle = \frac{1}{2} \langle I_7 \rangle / \langle I_4 \rangle, \quad (11a)$$

also

$$\tan \langle \delta s - \delta p \rangle = 2 \langle I_8 \rangle / \langle I_5 \rangle, \quad (11b)$$

where  $\langle I_n \rangle$  are defined in Eq. (5).

Should these two expressions for calculating  $\langle \delta s - \delta p \rangle$  not yield the same answer, this would mean that (i) time-reversal invariance is violated, or (ii) there is an  $I = 2$  admixture in the  $s$  wave, or (iii) there is a significant  $d$ -wave  $\pi$ - $\pi$  contribution.

Furthermore, Pais and Treiman point out that  $\langle I_9 \rangle$  should vanish under the assumptions of time-reversal invariance and the  $\Delta I = 1/2$  rule, and the absence of waves with  $\ell \geq 3$ . One may also test the assumption regarding the absence of waves with  $\ell \geq 2$  by examining the  $\cos\theta_\pi$  spectrum, which should be fitted by an expression of the form

$$dW/d\cos\theta_\pi = a(1 + b\cos\theta_\pi + c\cos^2\theta_\pi). \quad (12)$$

The need for any additional terms would imply the presence of higher waves.

### III. EXPERIMENTAL PROCEDURE

#### A. Details of Exposure

The beam used was a conventional two-stage separated beam of overall length approximately 25 meters, produced at 15 deg from an internal target in the CERN proton synchrotron. The beam was transported at 800 MeV/c, with a momentum bite of  $\pm 1.3\%$ , and degraded at the bubble chamber entrance window so that the  $K^+$  stopping points were well spread out in the center of the chamber.

The CERN enlarged 1.1-meter-diameter heavy-liquid chamber<sup>17</sup> was used. It was filled with  $C_2F_5Cl$ , which has a density of  $1.2 \text{ g/cm}^3$  and a radiation length of 25 cm under operating conditions. In this exposure care was taken to keep the bubble size small in order to be able to see the decay origins clearly. This was also important in seeing the  $\mu^+$  from decay of stopping  $\pi^+$  (range of  $\mu^+ = 0.15 \text{ cm}$ ).

We took 551 000 pictures, yielding a total of  $13.3 \times 10^6$  stopping  $K^+$  in the fiducial region (24 stopping  $K^+$ /picture). The film was divided equally between the three institutions (LRL, UW, and UCL) for scanning and preliminary selection of events.

#### B. Scanning

The film was scanned for  $K_{e4}$  candidates, and approximately two-thirds was rescanned in order to obtain the scanning efficiency. In order to estimate the total number of stopped  $K^+$  every 50th picture was scanned for  $\tau$  decays.

To pass the preliminary criteria used at the scan table, a  $K_{e4}$  candidate had to satisfy four conditions:



(i) The decay point had to lie within the fiducial region. This was a region specified to avoid the immediate vicinity of the beam entry window, which was difficult to scan because of a particularly high density of tracks.

(ii) The ionization of the incoming track near the decay point had to be consistent with that of a stopping K meson.

(iii) There had to be three tracks from the origin.

(iv) One track had to appear to be an electron, identified by spiralization, and the other two had to be consistent with being pions.

These requirements were not very stringent, and 2000 of the events passed by the scanners were measured. The momentum of the electrons was estimated by the Behr-Mittner procedure.<sup>18</sup> Photographic prints of all these events, together with the results of measurement, were then examined by physicists, who applied the more demanding criteria that follow.

### C. Selection Criteria for $K_{e4}$

#### 1. Exclusion of Tau Decays at Rest

Most events recorded as candidates by scanners are merely  $\tau$  decays, for which one of the  $\pi^+$  is too short to be seen. If this  $\pi$  decays to a  $\mu^+$  which then emits an  $e^+$  in approximately the same direction as the  $\mu^+$  track, the latter can be mistaken for the first part of the electron track. The electron thus appears to originate from the  $K^+$  decay point. Even though such events outnumber genuine  $K_{e4}$  by approximately 5 to 1, they present no problem of identification. This comes about because they are characterized by a  $\pi^-$  and a  $\pi^+$  going off nearly opposite to each other (at 155 to 180 deg), and with nearly equal momenta. This configuration,

though kinematically allowed for  $K_{e4}$  decays, is not particularly favored. Criteria can be set up, then, that exclude these  $\tau$  decays, while at the same time they reject only a few genuine  $K_{e4}$ . These criteria involve measurements on the two long pion tracks. For a  $\tau$  decay the missing mass,  $M_N$ , should be that of a  $\pi^+$ , viz 139.6 MeV. This estimate will be subject to some measurement error, so in practice one must reject as  $\tau$  those events with  $M_N$  lying in a certain finite range. The extent of this range was determined by plotting  $M_N$  for a random sample of  $K_{e4}$  candidates (Fig. 2). These events were selected so that the missing momentum was less than 50 MeV/c, since a  $\pi^+$  with a momentum exceeding this value would have a range greater than 4.5 mm and its track would be clearly visible even if steeply dipping in the chamber. It is seen that most of the events are centered closely about 139 MeV. On the basis of this figure it was decided to exclude all events for which  $130 \text{ MeV} < M_N < 150 \text{ MeV}$  and at the same time the missing momentum was  $< 50 \text{ MeV/c}$ . This rejected about 1450 events.

## 2. Exclusion of Tau Decays in Flight

Although the candidates had to have an incoming track whose ionization was consistent with the  $K^+$  meson's having come to rest, this requirement does not exclude slow primaries with momenta  $< 200 \text{ MeV/c}$ . A  $\tau$  decay in flight with a short  $\pi^+$  track and a  $\mu$  mistaken as part of the electron track would no longer necessarily have its two visible pions approximately collinear. Thus all candidates were tested to see whether the two pion tracks were consistent with the pions' having come from a  $\tau$  decay in flight with a  $K^+$  momentum  $< 200 \text{ MeV/c}$ . If so, and if it were

further found that the third pion had a momentum  $< 50 \text{ MeV}/c$ , the event was rejected. This criterion affected about 100 events.

### 3. Kinematical Constraint

For an event to be accepted it did, of course, have to fit satisfactorily the  $K_{e4}$  hypothesis. A further 150 events failed to meet this requirement and were rejected. On examination of the prints by a physicist, alternative interpretations were found in all cases.

### 4. Exclusion of Negative Secondaries with Steep Tracks

For some events, although the negatively charged secondary was consistent with being a pion, there was still some doubt as to its identification. This was the case when the track was either particularly steep or short. Such a track could be an electron and so when combined with the  $e^+$  would constitute a Dalitz pair. The event would therefore not be a  $K_{e4}$  but a  $\tau^+$ ,  $K_{\mu 3}$ , or  $K_{\pi 2}$ .

It was decided in consequence to reject the 21 events for which the negative track had a dip angle  $> 70 \text{ deg}$ .<sup>19</sup>

### 5. Exclusion of Positrons with Steep Tracks

It was sometimes difficult to decide whether a steep track was an  $e^+$  from a  $K_{e4}$  decay or a  $\pi^+$  from a  $\tau$  decay. Thus all events with an  $e^+$  dipping at an angle  $> 70 \text{ deg}$  were rejected. This accounted for a further two events.

### 6. Exclusion of Pion Secondaries with Short Tracks

As indicated above under 4, a very short  $\pi^-$  (if it did not produce a visible star at the end of its range) could be confused with a  $e^-$ . Not so much difficulty was encountered in the identification of a short  $\pi^+$ ,

because in general even if the track of the  $\pi^+$  itself were not visible one could see its decay  $\mu^+$ . Nevertheless if either pion had a short track, whether one could unambiguously identify it or not, it was difficult to measure the direction of the track accurately. In the circumstances it was decided to reject the 22 events for which the  $\pi^-$  or the  $\pi^+$  had a momentum  $< 48$  MeV/c (corresponding to a range of approximately 4 mm).

#### IV. RESULTS

##### A. Branching Ratio for $K_{e4}(e^+)$

After application of the various criteria outlined in the previous section, 269 events remain as  $K_{e4}(e^+)$  decays. These must be corrected for various losses before the branching ratio can be obtained.

##### 1. Correction for $\tau$ -Decay-at-Rest Criterion

Figure 3 shows a scatter plot of the unfitted values of momentum vs invariant mass of the dilepton system for all accepted  $K_{e4}$ . No events are to be seen in the rectangular region, owing to the criterion for eliminating  $\tau$  decays with approximately collinear configuration. The figure demonstrates that the majority of  $K_{e4}$  are far removed from this region, and that a correction of only two events should be applied.

The magnitude of this correction has been confirmed by use of the Monte Carlo program previously described.

##### 2. Correction for $\tau$ -Decay-in-Flight Criterion

Some  $K_{e4}$  at rest have a configuration such that for certain directions for the incoming K meson the event would be thrown out by the  $\tau$ -decay-in-flight criterion. The  $K_{e4}$  found in this experiment were examined with a view to determining the chance that the event would have

been rejected had the incoming K meson assumed some other direction. This study led to the conclusion that between 0.5% and 1% of genuine  $K_{e4}$  decays had been lost.

### 3. Correction for $\pi^-$ Having Steep Tracks

This is a purely geometrical correction based on the solid angle available. Its value is 6.0%.

### 4. Correction for Positrons Having Steep Tracks

This is also a geometrical correction, but allowance has been made for events having both a steep  $\pi^-$  track and a steep positron track. The correction is 5.6%.

### 5. Correction for $\pi^\pm$ Having Short Tracks

The same Monte Carlo program yields a correction of 15% for short  $\pi^\pm$  tracks.

### 6. Correction for Scanning Efficiency

Based on a rescan of two-thirds of the film, the average efficiency of the first scan was 66% and that of the second scan 73%. This meant that  $83 \pm 5\%$  of events passing the criteria were found. The events were divided up in various ways according to their configuration, but no preferential bias against any particular type could be discerned.<sup>20</sup>

### 7. Other Corrections

No other important means of losing  $K_{e4}$  have been found.<sup>21</sup> There are in addition, however, some corrections to be considered for effects leading to the acceptance of spurious events as genuine  $K_{e4}$ .

For example, a  $\tau$  decay at rest that would normally be rejected by Step 1 could be accepted as a  $K_{e4}$  if one of the pions scattered very early

so that its scattered direction were mistaken for its original direction at the K decay. Consequently a study was made of the visible scatterings to be found on the first 1.5 cm of the tracks associated with events rejected by Step 1. This permitted an estimate of the chance that such scatterings could occur in the first 2 mm of the track (and thus escape observation), and of the likelihood that the scatterings would be of a size and in a direction such as to lead to acceptance of the events as  $K_{e4}$ . The resulting correction was very small, viz., about 1 event.

Events normally rejected under (1) have a second way of evading the criterion. This arises when the  $\pi^+$  decays in flight to give a combined  $\pi^+, \mu^+$  length significantly greater than the range that the  $\pi^+$  would have had if it had come to rest. If the  $\pi^+, \mu^+$  track is mistaken as being entirely due to the  $\pi^+$  (this is quite likely, as  $\mu^+$  from  $\pi^+$  decay at rest do not always have readily visible tracks in heavy-liquid chambers) the event may be accepted as a  $K_{e4}$ . From the moderation times of the pions and the known lifetime it is calculated that 1 spurious event has been accepted in this manner.

The last background we mention also concerns  $\tau$  decay. We consider the  $\pi^+$  to decay while still lightly ionizing, and to give a very short track (i. e., the  $\mu^+$  goes backwards in the pion center-of-mass system). If the electron from the  $\mu^+$  decay is emitted approximately in the same direction as that of the original  $\pi^+$ , the whole  $\pi^+, \mu^+, e^+$  combined track can be mistaken for an  $e^+$  emitted from the K decay. From a consideration of the average moderation time of a pion from a  $\tau$  decay, the lifetime of the  $\pi^+$ , the solid angle available to the  $\mu^+$  in the pion c. m. system, and the solid angle available to the  $e^+$ , a correction of about 1 event is indicated.

Several other sources of background were investigated and were found to be small. The overall result is that our final sample of 269 accepted events contains about 3 events that are not genuine  $K_{e4}$ .

#### 8. Estimation of the Total Number of $K^+$ Based on the Scan for $\tau$ Decay

The average number of  $\tau$  decays found in the sample scan of every 50th picture throughout the run was  $1.34 \pm 0.02$   $\tau$ /picture. The error is dominated by the statistical uncertainty. The scanning efficiency was determined to be 99.5%. The total number of  $\tau$  on the 551 000 pictures is

$$551\,000 \times 1.34 \times \frac{100}{99.5} = 743\,000.$$

Using the branching ratio<sup>22</sup>

$$K^+ \rightarrow \tau^+ / \text{all } K^+ \text{ decays} = (5.57 \pm 0.04)\%,$$

one finds the total number of  $K^+$

$$= (13.3 \pm 0.3) \times 10^6.$$

#### 9. Estimation of the Branching Ratio

After application of the various corrections, the estimated total number of  $K_{e4}(e^+)$  is 431 (see Table I). The branching ratio is therefore

$$K_{e4}(e^+) / \text{all } K^+ \text{ decays} = (3.25 \pm 0.35) \times 10^{-5}.$$

This compares well with the value  $(3.6 \pm 0.8) \times 10^{-5}$  found in the previous experiment.<sup>1</sup> The corresponding decay rate is  $(2.60 \pm 0.30) \times 10^3 \text{ sec}^{-1}$ .

### B. Upper Limit to the Branching Ratio for $K_{e4}(e^-)$ Decay

No candidates for the  $K_{e4}(e^-)$  mode were found, either in this or in the previous experiment.<sup>1</sup>

Although the scanning criteria for this mode were similar to those for  $K_{e4}(e^+)$ , the detection efficiencies for the two modes are not equal. The sources of background are different, so one cannot make a straight comparison.

For example, a  $\pi^- \rightarrow \mu^- \rightarrow e^-$  decay is less probable, so corrections IV A 1 and IV A 2 can be relaxed.

Instead of rejecting one of the pions--viz. the negative one--when its track is steep, we now under IV A 4 must reject either  $\pi^+$  if its track is steep. This is because the  $\pi^+$  may be ambiguous with an  $e^+$  and the K decay could then be a  $\tau^+$  with a Dalitz electron pair.

Events must be rejected if the two  $\pi^+$  are consistent with having come from a  $\tau$  decay and the  $e^-$  lies in approximately the direction of the expected third pion. This is necessary to avoid a  $\tau$  decay in which the  $\pi^-$  charge-exchanges, and one of the gammas from the  $\pi^0$  decay gives a Compton electron almost immediately, the combined  $\pi^-, e^-$  track appearing to be an  $e^-$  from the K decay.

Considerations of this nature lead to an overall detection efficiency for  $K_{e4}(e^-)$  which is 70% of that appropriate for  $K_{e4}(e^+)$ .

Thus this experiment yields an upper limit

$$K_{e4}(e^-)/\text{all } K^+ \text{ decays} < 7 \times 10^{-7}$$

at the 95% confidence level.



### C. Mass and Angular Distributions

Figure 4a shows the distribution of  $\phi$ , the angle between the dipion and dilepton planes. Figure 4b gives the same distribution with the events from the previous experiment included. Both are adequately fitted by a function of the form  $\alpha(1 + \beta \cos \phi + \gamma \sin \phi + \delta \sin^2 \phi)$ , where for Fig. 4b  $\beta = 0.20 \pm 0.08$ ,  $\gamma = 0.26 \pm 0.08$ , and  $\delta = -0.03 \pm 0.11$ . The value of  $\chi^2$  is 3.5 for eight degrees of freedom. (The curves shown refer not to this solution but to solutions involving simultaneous fits to several histograms. These are described later.)

The angle that the  $\pi^+$  makes with the dipion line of flight in the dipion rest system is plotted in Fig. 5. Unlike  $\phi$ ,  $\cos \theta_\pi$  is strongly affected by the cut on short-track pions. An impression of the size of the corrections needed to each bin is given by the cross-hatched events. These were estimated by the Monte Carlo program (mentioned in Section II, Theory) which generated  $K_{e4}$  events by using our most probable set of values of the form factors and the  $\pi$ - $\pi$  phase shift (to be obtained in the next section).

The  $\cos \theta_\pi$  distribution should be of the form  $a(1 + b \cos \theta_\pi + c \cos^2 \theta_\pi)$ . From Fig. 5b, we estimate  $b$  to be  $(0.61 \pm 0.12)$ . Evaluation of the coefficient  $c$  depends sensitively upon the numbers of events with large values of  $|\cos \theta_\pi|$ . These are the values for which the correction for short-track pions is greatest, and so it is difficult to estimate  $c$  with any degree of reliability. We have considered it desirable to base conclusions to be drawn later from this distribution solely on that part which is largely unaffected by the cut, i. e., the region lying between the dotted lines.

It can be seen from Fig. 6 that the slope of the  $\cos\theta_\pi$  plot changes markedly as a function of  $S_\pi$ . This behavior can be understood if the  $f_p$  form factor goes to zero as  $\beta$  goes to zero at small values of  $S_\pi$ . The quantity  $[(\beta X/M^2) \tilde{f}'_p]$ , which replaces  $\tilde{f}_p$  in the analysis to allow for the effects of the angular momentum barrier, has just this kind of behavior.

Figure 7 shows the invariant-mass distribution of the dipion system. The cross-hatched events in the first three bins have been added to compensate for the loss of short pions. Similarly the estimated two events excluded by the  $\tau$ -decay-at-rest criterion (Section IV A 1) are kinematically constrained to fall within the fourth bins.

Figure 8 gives the histogram of  $\cos\theta_\ell$ , the angle between the  $e^+$  and the dilepton line of flight in the dilepton rest frame.  $\cos\theta_\ell$  is largely unaffected by the short-track pion correction, but for this angle a new effect becomes significant. It concerns events for which the laboratory-system velocity,  $\beta_{ev}$ , of the dilepton system is close to unity. For this class of events the errors on  $\cos\theta_\ell$  arising from measurement errors on angles and momenta can become highly asymmetric. This introduces a systematic bias as events are moved more readily towards negative values of  $\cos\theta_\ell$  than towards positive ones. This effect is readily reduced to negligible proportions by removing the 16% of events for which  $\beta_{ev} > 0.95$ . This cut explains why the numbers of events in Fig. 8 are less than those for the other plots.

Figure 9 is a two-dimensional scatter diagram showing the correlation between  $\cos\theta_\ell$  and  $\phi$ , for events with  $\beta_{ev} < 0.95$ . As was indicated in the Section II, on Theory, correlations between these two angles have a particular significance in the Pais-Treiman method of analysis. The average values of the relevant correlations are shown in Table II.

It is to be noted that the quantity  $\langle \sin^2 \theta_l \sin 2\phi \rangle$ , which determines  $\langle I_9 \rangle$ , is consistent with zero, as is required by time-reversal invariance.

D. Determination of the  $\pi$ - $\pi$  Phase-Shift Difference  
and the Form Factors

In principle the most satisfactory way of determining the s-wave  $I = 0$   $\pi$ - $\pi$  phase shift is to use the ratios of the I coefficients given in Eq. (11). The theory upon which such estimates are based rests on very few assumptions, viz. the absence of d waves and of an  $I = 2$  admixture in the s wave, and time-reversal invariance.

Unfortunately the correlations concerned are not very strong, and the statistical accuracy obtainable with 300 events is exceedingly poor. We are not even able unambiguously to determine the sign, as both expressions (11a) and (11b) involve quantities (viz.,  $I_4$  and  $I_8$ ) which are not significantly different from zero. Taking the data at face value, the estimates of the magnitude of  $\langle \delta s - \delta p \rangle$

$$\begin{aligned} |\langle \delta s - \delta p \rangle| &= \tan^{-1} \left( \frac{1}{2} \langle I_7 \rangle / \langle I_4 \rangle \right) = 90 \pm 40 \text{ deg} \\ \text{and} \quad |\langle \delta s - \delta p \rangle| &= \tan^{-1} \left( 2 \langle I_8 \rangle / \langle I_5 \rangle \right) = 0 \pm 40 \text{ deg.} \end{aligned} \tag{12}$$

One observation we can make, however, is that  $\langle I_7 \rangle$ , and consequently the numerator of expression (11a), is almost 3 standard deviations from zero. [Note the errors as quoted in (12) are not Gaussian.] Thus although the large fractional error on the denominator makes it difficult to establish a magnitude for  $\tan \langle \delta s - \delta p \rangle$ , our result is not easily reconcilable with a value close to zero.

Until it becomes possible to perform experiments yielding larger numbers of events, the above method for extracting the phase shift is

not very promising, so in order to proceed further we need to introduce values for the form factors, and assumptions regarding their energy dependence.

As mentioned in Section II (Theory), three methods of obtaining acceptable sets of values for the form factors and phase-shift difference have been used. They each have certain advantages. The maximum-likelihood technique is able to extract information contained in correlations between the different variables. This information is lost in performing a simultaneous least-squares fit to the histograms of  $\phi$ ,  $\cos\theta_\pi$ ,  $\cos\theta_\ell$ , and  $(S_\pi)^{1/2}$ . However, this second method is able to make use of events with  $|\cos\theta_\pi| > 0.6$  and  $\beta_{e\nu} > 0.95$  in those plots where they introduce no bias. The Monte Carlo method, on the other hand, is the most effective way of checking the influence of cuts and biases in the data.

Assuming all form factors to be independent of  $S_\pi$  and  $S_\ell$ , the least-squares-fit program yields four solutions with acceptable values of  $\chi^2$ . These are listed in Table III.

Nominally there are 29 degrees of freedom. However, the fact that the same events are used in all histograms imposes restrictions that tend to reduce the effective number of degrees of freedom. By fitting 50 batches of Monte Carlo events generated with solution A parameters, and examining the resulting  $\chi^2$  values, we estimate that the effective number of degrees of freedom to which the  $\chi^2$  values of Table III refer lies between 28 and 29. This implies that these histograms are almost independent projections. In considering the results of Table III, it should be noted that the Pais-Treiman form factor  $f_p$  may be obtained

from the value of  $v$  by multiplying it by  $\approx 0.11$ .

With the maximum-likelihood program we can in fact exclude solutions C and D. This is rendered possible by a study of  $\theta_1$  vs  $\phi$  correlations, which are not open to examination by the program that performs a fit to the one-dimensional plots. Specifically, it is found that the large values of  $\gamma$  and of  $\sin(\delta_s - \delta_p)$  lead to values of  $I_g$  which are 3 to 4 standard deviations removed from the experimental number.

Of the two remaining solutions, A is somewhat favored over B, but both have acceptable  $\chi^2$  values. Unfortunately, with respect to determining the phase shift, although the two solutions have the same magnitude for  $(\delta_s - \delta_p)$ , the signs are opposite. We postpone discussion of this point until later in the paper.

We have also investigated the effect of an  $S_\pi$  energy dependence in  $\tilde{f}_s$  proportional to the Watson enhancement factor, and solutions corresponding to solutions A and B in Table III are given as solutions  $A_e$  and  $B_e$  in Table IV. The differences are not large.

The maximum-likelihood method yielded essentially the same solutions, within errors, and the Monte Carlo approach also located the two minima in the  $\chi^2$  map in regions corresponding closely to solutions A and B.

## V. DISCUSSION

The rate for  $K_{e4}(e^+)$  decay has been found to be  $(2.60 \pm 0.30) \times 10^3 \text{ sec}^{-1}$ . Theoretical calculations that do not include final-state interactions all contain an adjustable parameter, and can be brought into agreement with our rate with a reasonable value of the parameter. Brown

and Faier<sup>9</sup> allow for the final-state interactions by assuming that the decay proceeds through a  $\sigma$  meson. Although their rate is in agreement with our value, the  $(S_\pi)^{1/2}$  spectrum does not furnish any evidence of the presence of a  $\sigma$  meson. Weinberg,<sup>11</sup> using current algebra and soft-pion techniques, is able to relate the  $K_{e4}$  form factors to those of  $K_{e3}$ , and hence predicts a rate of  $(1.88 \pm 0.23) \times 10^3 \text{ sec}^{-1}$ . In regard to this, however, we point out that for solution A, where  $\nu$  is large,  $f_p$  contributes  $\approx 30\%$  to the rate.

The upper limit, at the 95% confidence level of  $7 \times 10^{-7}$  for the  $K_{e4}(e^-)$  branching ratio, adds weight to the  $\Delta Q = \Delta S$  rule.  $K_{e4}(e^-)$  decay, if it occurs, proceeds through the axial-vector current.  $K_{e4}(e^+)$  decay involves a mixture of both axial-vector and vector, but although the latter gives rise to strong interference terms, its contribution to the rate is small. Thus our conclusions refer specifically to the rule as it applies to the axial-vector currents in weak interactions. In interpreting our result one must be careful to make allowance for the differing dipion interactions in  $K_{e4}(e^-)$  decay and in  $K_{e4}(e^+)$  decay. For  $K_{e4}(e^-)$  the dipion state is pure  $I = 2$ , whereas for  $K_{e4}(e^+)$  we have  $I = 0$  or  $1$ . Because of the relatively low energy of the interaction, one further assumes only s and p waves to be important in  $K_{e4}(e^+)$ , whereas only s waves would be present in  $K_{e4}(e^-)$ . The violation parameter is defined as the ratio of the amplitudes of the currents,

$$x = \frac{A(\Delta Q = -\Delta S)}{A(\Delta Q = +\Delta S)}.$$

In order to evaluate  $x$  one needs to know the enhancement factor due to the final-state interaction. For no s-wave interaction our branching

ratios lead to  $x < 0.15$  at the 95% confidence level, whereas for enhancement factors variously estimated as extending up to a value of 4 (Refs. 8 and 10) we obtain

$$x < 0.3 \quad (95\% \text{ confidence level}).$$

The three methods that yielded values for both the form factors and the phase-shift difference agreed that there are two acceptable solutions.

We should mention that in the paper based on the previous experiment<sup>1</sup> a single unambiguous solution was reported. This was because the importance of the  $\cos\theta_\ell$  distribution and the form factor  $f_p$  had not at that time been recognized. The differing conclusions of that paper and this, it should be stressed, are not due to any contradiction between the two sets of data, but rather to the method of analysis now including the  $f_p$  form factor. For the same reason, the analyses given in the first three preliminary reports<sup>2</sup> of this experiment are to be regarded as inadequate.

That we should have obtained two solutions that fit the data is not entirely unexpected. Berends, Donnachie, and Oades<sup>23</sup> have pointed out that if the original type of analysis proposed by Cabibbo and Maksymowicz<sup>13</sup> were applied to the  $(S_\pi)^{1/2}$ ,  $\cos\theta_\pi$ , and  $\phi$  data with the modification that  $f_p$  were no longer put to zero, then one must obtain two solutions that are equally satisfactory. These are related to each other in the following way:

$$\begin{aligned} {}_1\tilde{f}_s &= {}_2\tilde{f}_s, \\ {}_1\tilde{f}_p &= {}_2\tilde{f}_p + \xi_2\tilde{g}, \\ {}_1\tilde{g} &= -{}_2\tilde{g}, \\ {}_1\tilde{h} &= {}_2\tilde{h}, \\ {}_1\langle \delta s - \delta p \rangle &= -{}_2\langle \delta s - \delta p \rangle, \end{aligned}$$

where  $\xi$  is a kinematic factor, having an approximate value of 9.5, and is appropriate to our particular method of introducing an angular momentum barrier effect for  $f_p$ . They went on to point out that this ambiguity could be resolved by studying the variable  $\theta_\ell$ . Specifically the two solutions lead to coefficients for the  $\cos\theta_\ell$  term that are equal in magnitude but opposite in sign.

An essential conclusion of this experiment is that the information contained in our  $\theta_\ell$  plot is insufficient to distinguish which solution is correct. The two possible values of the coefficient of  $\cos\theta_\ell$  are  $-0.09$  and  $+0.03$ . The experimental value is  $-0.14 \pm 0.09$ .

Thus although our experiment favors the former value, we would be unwise to consider this solution as established. The need for such caution is particularly emphasized by the fact that by allowing  $\tilde{f}_s$  to be enhanced we can materially alter the relative probabilities of the two solutions. Nevertheless, certain definite statements can be extracted from the data.

The most statistically significant angular correlation is the slope of the  $\cos\theta_\pi$  plot, which was seen in Section IVC to have a value  $0.61 \pm 0.12$ . This slope is governed by the values of  $\nu$  and  $\eta$ , and our measurement leads to the approximate relation

$$2\nu + 9\eta = 7,$$

which holds when the magnitudes of the form factors and phase shift are not greatly in excess of those of solutions A and B.

It is clear from this relation that if  $\nu$  could be determined, the value of  $\eta$  would be established and consequently  $\tilde{g}$ . The ambiguity of the two



solutions would then be resolved. However,  $\nu$  has little significance for the other angular distributions and has essentially to be determined from the  $\cos\theta_\pi$  distribution.

Next we note that both solutions require the vector form factor  $\tilde{h}$  ( $\gamma = \tilde{h}/\tilde{f}_s$ ) to be significantly different from zero, and its sign is negative. It is true that in solution B it is not so far removed from zero. However, the poorer  $\chi^2$  for this solution is largely due to an internal conflict in the data in which the large value of  $\tilde{h}$  required by the coefficient of  $\cos\phi$  is set against other requirements of the fit. Thus  $\gamma$  for this solution has already been reduced to a level that makes the overall probability less than that of solution A. That  $\tilde{h}$  is significant is a conclusion that was not justified on the statistics of the previous experiment.

Finally, we point out that the quantity  $\eta \sin\langle\delta s - \delta p\rangle$  is a constant for both types of solution given by Berends, Donnachie, and Oades, a change of sign of  $\eta$  being compensated by a similar change in  $\sin\langle\delta s - \delta p\rangle$ . This constant relationship is seen to hold true for the two solutions A and B, and also the solutions in which  $\tilde{f}_s$  is enhanced,  $A_e$  and  $B_e$ . The values of the product are respectively  $(0.68 \pm 0.22)$ ,  $(0.57 \pm 0.20)$ ,  $(0.46 \pm 0.20)$ , and  $(0.56 \pm 0.13)$ . The quantity is almost entirely governed by the coefficient of  $\sin\phi$ .

Weinberg<sup>11</sup> estimates  $\sin\delta s = 0.1$  (giving  $a_0 = 0.2\lambda_\pi$ ) and  $\eta = 1$ , using current algebra. His value for the product is thus 0.1. His requirement of a positive value for  $\eta$  leads to a choice of solutions B or  $B_e$ . Our evaluation of the product  $\eta \sin\langle\delta s - \delta p\rangle$  is thus just over 2 standard deviations from his estimate.

In conclusion we consider the possibility of resolving the two-solution ambiguity in later experiments. In principle the ideal way to extract the phase shift is to use either equation (11a) or (11b). This, as Pais and Treiman have explained, could yield estimates independent of assumptions regarding the form factors. That we were not able even to obtain the sign of the phase shift in this experiment was because our statistics were inadequate to obtain values of  $I_8$  or  $I_4$  that were significantly removed from zero.

Likewise the sign of the phase shift was unresolved in the alternative approach because of the smallness of the coefficient of  $\cos\theta_l$ , which is related to  $I_6$ . Indeed, in order to make an unambiguous measurement of  $\langle \delta s - \delta p \rangle$  it is necessary to determine that one of the odd moments of  $\cos\theta_l$ ,  $I_4$ ,  $I_6$ , or  $I_8$  is significantly different from zero. It should be noted that, this being the case, the problem is not merely one of accumulating larger statistics. The  $\theta_l$  plot, as we have pointed out, is liable to suffer from biases arising from measurement inaccuracy. Increased measuring errors in any subsequent experiment would of course render it necessary to make even more drastic cuts in  $\beta_{ev}$  than the one employed in this investigation.

### ACKNOWLEDGMENTS

We express our deep gratitude to the CERN personnel, and especially to Dr. C. A. Ramm and his colleagues in the N. P. A. Division, for making this exposure possible.

We should like to acknowledge the help of Dr. R. W. Birge, Dr. W. M. Powell and Dr. D. J. Miller during the early part of this work. We also acknowledge many helpful discussions on the theoretical aspects of the work, especially those with Dr. A. Pais, Dr. R. Cence, Dr. S. Treiman, Dr. A. Donnachie, Dr. G. Oades, Dr. S. M. Berman, and Dr. J. Yellin.

The members of the UCL group thank the Science Research Council for its financial support, and one of them (W. L. K.) acknowledges the receipt of a Turner & Newall fellowship.

The work of the LRL group was done under auspices of the U. S. Atomic Energy Commission.

## APPENDIX

These are the equations used in the simultaneous fit to the  $\phi$ ,  $\cos\theta_\pi$ ,  $\cos\theta_\ell$ , and  $(S_\pi)^{1/2}$  histograms. The expressions include corrections appropriate to the selection criteria adopted in this investigation.

 $\phi$  Distribution

$$\frac{\partial\Gamma}{\partial\phi} = \alpha(1 + \beta\cos\phi + \lambda\sin\phi + \kappa\cos^2\phi),$$

where  $\alpha = 56.3/(2+\kappa),$

$$\beta = -12.38 \gamma \cos\langle\delta s - \delta p\rangle M,$$

$$\lambda = 60.7 \eta \sin\langle\delta s - \delta p\rangle M,$$

$$\kappa = (0.29 \gamma^2 - 8.9 \eta^2) M,$$

and  $M = [127 + 0.37 \nu^2 + 23.3 \eta^2 + 0.14 \gamma^2 + 3.68 \eta\nu]^{-1}.$

 $\cos\theta_\pi$  Distribution

$$\frac{\partial\Gamma}{\partial\cos\theta_\pi} = a(1 + b\cos\theta_\pi + c\cos^2\theta_\pi),$$

where  $a = 246/(6 + 0.72 c),$

$$b = (33 \nu + 162.5 \eta) \cos\langle\delta s - \delta p\rangle L,$$

$$c = (1.81 \nu^2 + 33.5 \eta^2 - 0.46 \gamma^2 + 18.1 \eta\nu)L,$$

and  $L = [154 + 14.4 \eta^2 + 0.46 \gamma^2]^{-1}.$

 $\cos\theta_\ell$  Distribution

$$\frac{\partial\Gamma}{\partial\cos\theta_\ell} = A(1 + B\cos\theta_\ell + C\cos^2\theta_\ell),$$

where  $A = 28.3/(1 + 0.333 C),$

$$B = -4.9 \eta\gamma N,$$

$$C = (-190 - 0.552 \nu^2 - 7.9 \eta^2 + 0.21 \gamma^2 - 5.52 \eta\nu)N,$$

and 
$$N = [190 + 0.552 \nu^2 + 21.3 \eta^2 + 0.21 \gamma^2 + 5.52 \eta\nu]^{-1}.$$

$(S_{\pi})^{1/2}$  Distribution

280 to 300 MeV:  $(122 + 0.23 \nu^2 + 6.3 \eta^2 + 0.10 \gamma^2 + 1.92 \eta\nu)F,$

300 to 320 MeV:  $(135 + 0.53 \nu^2 + 18.7 \eta^2 + 0.26 \gamma^2 + 4.90 \eta\nu)F,$

320 to 340 MeV:  $(108 + 0.54 \nu^2 + 23.3 \eta^2 + 0.28 \gamma^2 + 5.50 \eta\nu)F,$

340 to 360 MeV:  $(73 + 0.40 \nu^2 + 21.0 \eta^2 + 0.21 \gamma^2 + 4.42 \eta\nu)F,$

360 to 380 MeV:  $(43 + 0.21 \nu^2 + 15.2 \eta^2 + 0.12 \gamma^2 + 2.69 \eta\nu)F,$

380 to 400 MeV:  $(21 + 0.07 \nu^2 + 8.9 \eta^2 + 0.06 \gamma^2 + 1.19 \eta\nu)F,$

400 to 420 MeV:  $(9 + 0.014 \nu^2 + 4.1 \eta^2 + 0.01 \gamma^2 + 0.35 \eta\nu)F,$

420 to 440 MeV:  $(3 + 0.007 \nu^2 + 1.4 \eta^2 + 0.004 \gamma^2 + 0.15 \eta\nu)F,$

where 
$$F = [1.25 + 0.0051 \nu^2 + 0.264 \eta^2 + 0.003 \gamma^2 + 0.055 \eta\nu]^{-1}.$$

REFERENCES

\*Now at University of Pennsylvania.

†Now at University of Aberdeen.

‡Now at CERN.

1. R. W. Birge, R. P. Ely, Jr., G. Gidal, G. E. Kalmus, A. Kernan, W. M. Powell, U. Camerini, D. Cline, W. F. Fry, J. G. Gaidos, D. Murphree, and C. T. Murphy, *Phys. Rev.* 139, B1600 (1965).
2. Proceedings of the XIIIth International Conference on High Energy Physics, Berkeley, 1966 (University of California Press, Berkeley and Los Angeles, 1967);  
Meeting of the Physical Society and Institute of Physics, held at University College, London, May 1967 (unpublished);  
Princeton Conference on K Mesons, November 1967; Proceedings of the XIVth International Conference on High Energy Physics, Vienna, 1968 (to be published).
3. F. A. Berends, A. Donnachie, and G. C. Oades, *Nucl. Phys.* B3, 569 (1967).
4. V. S. Mathur, *Nuovo Cimento* 14, 1322 (1959); L. B. Okun and E. P. Shabalin, *Zh. Eksperim. i Teor. Fiz.* 37, 1775 (1959) [English translation: *Soviet Phys. - JETP* 10, 1252 (1960)];  
A. Sirlin, *Phys. Rev.* 129, 1377 (1963).
5. R. P. Feynman and M. Gell-Mann, *Phys. Rev.* 109, 193 (1958).
6. K. Chadan and S. Oneda, *Phys. Rev. Letters* 3, 292 (1959);  
E. P. Shabalin, *Zh. Eksperim i Teor. Fiz.* 39, 345 (1960) [English transl.: *Soviet Phys. - JETP* 12, 245 (1961)]; R. E. Berends and A. Sirlin, *Phys. Rev. Letters* 8, 221 (1962);

- B. Sakita, M. Kato, and E. McClement, University of Wisconsin (unpublished report); B. A. Arbazov, Nguyen Van Hieu, and R. N. Faustov, Zh. Eksperim. i. Teor. Fiz. 44, 329 (1963) [English transl.: Soviet Phys. - JETP 17, 225 (1963)].
7. J. Iliopoulos, Orsay Report No. TH/84 and Erratum (unpublished).
  8. G. Ciocchetti, Nuovo Cimento 25, 385 (1962).
  9. L. M. Brown and H. Faier, Phys. Rev. Letters 12, 514 (1964).
  10. C. Kacser, P. Singer, and T.N. Truong, Phys. Rev. 137, B1605 (1965); 139, AB5(E) (1965).
  11. S. Weinberg, Phys. Rev. Letters 17, 336 (1966) [erratum: 18, 1178 (1967)].
  12. E. P. Shabalin, Zh. Eksperim. i Teor. Fiz. 44, 765 (1963) [English translation: Soviet Phys. - JETP 17, 517 (1963)].
  13. N. Cabibbo and A. Maksymowicz, Phys. Rev. 137, B438 (1965).
  14. R. H. Dalitz, in Proceedings of the International School of Physics "Enrico Fermi," Varenna Lectures, 1964.
  15. S. Weinberg, Phys. Rev. Letters 17, 616 (1966).
  16. A. Pais and S. B. Treiman, Phys. Rev. 168, 1858 (1968).
  17. C. A. Ramm and L. Resegotti, in Proceedings of an International Conference on Instrumentation for High Energy Physics, Lawrence Radiation Laboratory, September 12-14, 1960 (Lawrence Radiation Laboratory, Berkeley, 1961), p. 127.
  18. L. Behr and P. Mittner, Nucl. Instr. Methods 20, 446 (1963).
  19. This angle is larger than the previously adopted value (Ref. 1) of 60 deg, due to the more favorable stereo angle of the cameras in the chamber used in this work.

20. We used the usual method for determining the scanning efficiency of the two scans, i. e.,  $\epsilon_1 = N_{12}/N_2$ ,  $\epsilon_2 = N_{12}/N_1$ , and  $\epsilon_{12} = 1 - (1-\epsilon_1)(1-\epsilon_2)$ , where  $N_1$  and  $N_2$  are number of events found on the first and second scan respectively, and  $N_{12}$  is the number of events found in common. It has recently been pointed out by S. E. Derenzo and R. H. Hildebrand (in a paper to be submitted to Nucl. Instr. Methods) that this method, which assumes that each event has the same "visibility," is open to considerable doubt. However, in order to do their analysis three scans of the film are needed. Should the conclusion reached by Derenzo and Hildebrand be applicable to our experiment, the effect would be to lower our value of  $\epsilon_{12}$ , i. e., to increase the rate.
21. In the previous experiment corrections were required for  $K_{e4}$  that were unidentified because one of the secondaries left the chamber. Because of the much larger chamber used this time, these corrections are now negligible.
22. See summary by A. H. Rosenfeld et al., Rev. Mod. Phys. 40, 1 (1968) [Lawrence Radiation Laboratory Report UCRL-8030 (Rev.), Jan. 1968].
23. F. A. Berends, A. Donnachie, and G. C. Oades, A Study of  $K_{l4}$  Decay, University of Glasgow Preprint (to be published in Phys. Rev.).



Table I. Summary of corrections  
due to scanning and selection criteria.

<u>Section number</u>	<u>Fractional acceptance</u>
IV A 1	0.993
IV A 2	0.993
IV A 3	0.94
IV A 4	0.944
IV A 5	0.85
IV A 6	0.83
IV A 7	1.011
Product	0.624

Table II. Mean values of the  $\phi$  vs  $\theta_\ell$  correlations relevant to the determination  $I_1 \cdots I_9$ .

$\langle \cos 2\theta_\ell \rangle$	$-0.502 \pm 0.035$
$\langle \sin^2 \theta_\ell \cos 2\phi \rangle$	$-0.011 \pm 0.034$
$\langle \sin 2\theta_\ell \cos \phi \rangle$	$-0.001 \pm 0.028$
$\langle \sin \theta_\ell \cos \phi \rangle$	$+0.093 \pm 0.036$
$\langle \cos \theta_\ell \rangle$	$-0.048 \pm 0.029$
$\langle \sin \theta_\ell \sin \phi \rangle$	$+0.102 \pm 0.036$
$\langle \sin 2\theta_\ell \sin \phi \rangle$	$-0.004 \pm 0.028$
$\langle \sin^2 \theta_\ell \sin 2\phi \rangle$	$-0.041 \pm 0.033$

Table III. Values of the form factors and phase shifts for the four solutions with acceptable  $\chi^2$  ( $\tilde{f}_s$  unenhanced).

	Solutions			
	A	B	C	D
$\tilde{f}_s$ (Note a)	$4.32 \pm 0.26$	$4.52 \pm 0.29$	$4.25 \pm 0.33$	$4.14 \pm 0.32$
$\tilde{g}$	$-6.96 \pm 0.77$	$6.01 \pm 0.86$	$1.06 \pm 0.60$	$-2.36 \pm 0.65$
$\tilde{h}$	$-10.4 \pm 3.8$	$-4.92 \pm 3.67$	$-42.5 \pm 8.3$	$-42.2 \pm 8.5$
$\tilde{f}'_p$	$54.0 \pm 6.1$	$-14.0 \pm 5.1$	$21.7 \pm 9.5$	$41.4 \pm 9.3$
$\eta (\equiv \tilde{g}/\tilde{f}_s)$	$-1.61 \pm 0.15$	$1.33 \pm 0.17$	$0.25 \pm 0.14$	$-0.57 \pm 0.15$
$\gamma (\equiv \tilde{h}/\tilde{f}_s)$	$-2.41 \pm 0.86$	$-1.09 \pm 0.81$	$-10.0 \pm 1.8$	$-10.2 \pm 1.9$
$\nu (\equiv \tilde{f}'_p/\tilde{f}_s)$	$12.5 \pm 1.2$	$-3.1 \pm 1.1$	$5.1 \pm 2.2$	$10.0 \pm 2.1$
$\langle \delta_s - \delta_p \rangle$ (rads)	$-0.44 \pm 0.14$	$0.44 \pm 0.15$	$1.21 \pm 0.10$	$-1.17 \pm 0.10$
$a_0(\lambda_\pi)$	$-0.89 \pm 0.28$	$1.26^{+0.68}_{-0.52}$	Very large	$-2.83^{+0.40}_{-0.50}$
$\chi^2$	26.6	38.6	36.3	26.0

a. We have used the value of 0.26 for  $\sin\theta_c$  [N. Brene, M. Ross, and A. Sirlin, Nucl. Phys. B6, 256 (1968)] in order to obtain these values.


Table IV. Values of the form factors and phase shifts corresponding to solutions A and B of Table III when  $\tilde{f}_s$  is enhanced.

	Solutions	
	$A_e$	$B_e$
$\tilde{f}_s$ (Note a)	$4.08 \pm 0.23$	$6.1 \pm 0.66$
$\tilde{g}$	$-6.77 \pm 0.74$	$6.8 \pm 1.09$
$\tilde{h}$	$-9.35 \pm 3.42$	$-5.02 \pm 3.89$
$\tilde{f}'_p$	$52.2 \pm 5.8$	$-14.7 \pm 5.7$
$\eta (\equiv \tilde{g}/\tilde{f}_s)$	$-1.66 \pm 0.15$	$1.12 \pm 0.13$
$\gamma (\equiv \tilde{h}/\tilde{f}_s)$	$-2.29 \pm 0.83$	$-0.82 \pm 0.63$
$\nu (\equiv \tilde{f}'_p/\tilde{f}_s)$	$12.8 \pm 1.2$	$-2.41 \pm 0.90$
$\langle \delta_s - \delta_p \rangle$ (rads)	$-0.28 \pm 0.12$	$0.50 \pm 0.10$
$a_0(\lambda_\pi)$	$-0.58 \pm 0.24$	$1.50 \pm 0.48$
$\chi^2$	31.8	33.0

a. We have used the value of 0.26 for  $\sin \theta_c$  [N. Brene, M. Ross, and A. Sirlin, Nucl. Phys. B6, 256 (1968)] in order to obtain these values.


## FIGURE CAPTIONS

- Fig. 1. A diagram illustrating the various angles referred to in the text.
- Fig. 2. The missing mass,  $M_N$ , calculated on the basis of measurements on the two pions, for a random sample of  $K_{e4}$  candidates. For these events the missing momentum is less than 50 MeV/c.
- Fig. 3. Scatter plot of the unfitted momentum vs invariant mass of the dilepton system for all accepted  $K_{e4}$ . Events in the rectangular region have been excluded by the  $\tau$ -decay-at-rest criterion.
- Fig. 4. Distribution of the angle  $\phi$ , (a) for this experiment, and (b) for both experiments combined.
- Fig. 5. Distribution of the angle  $\theta_\pi$ , (a) for this experiment, and (b) for both experiments combined. The cross-hatched events show the extent of the correction needed for the short-track-pion criterion. Only events lying within the range  $-0.6 < \cos\theta_\pi \leq +0.6$  are used in the overall fit.
- Fig. 6. Distribution of  $\theta_\pi$ , uncorrected for the loss of short-track pions (a) for events with  $(S_\pi)^{1/2} < 320$  MeV, and (b) for events with  $(S_\pi)^{1/2} > 320$  MeV.
- Fig. 7. Distribution of the invariant mass of the dipion system, (a) for this experiment, and (b) for both experiments combined. The cross-hatched events show the extent of the correction needed mainly for the criterion excluding short pions.
- Fig. 8. Distribution of the angle  $\theta_\ell$ , (a) for this experiment, and (b) for both experiments combined. Events with  $\beta_{e\nu} > 0.95$  have been excluded from these histograms.
- Fig. 9. Scatter plot of  $\cos\theta_\ell$  vs  $\phi$  for both experiments combined. Events with  $\beta_{e\nu} > 0.95$  have been excluded.




XBL68II-7287

Fig. 1



XBL6811-7288


Fig. 2



XBL6811-7289


Fig. 3






XBL6811-7290

Fig. 4




XBL6811-7291

Fig. 5




XBL 6811-7292

Fig. 6




XBL6811-7293

Fig. 7



XBL6811-7294

Fig. 8



XBL6811-7295

Fig. 9

LEGAL NOTICE

*This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:*

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

*As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE RADIATION LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720