

UC Irvine

ICTS Publications

Title

A phase II evaluation of the potent, highly selective PARP inhibitor veliparib in the treatment of persistent or recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who carry a germline BRCA1 or BRCA2 mutation - An...

Permalink

<https://escholarship.org/uc/item/18c9j72s>

Journal

Gynecologic Oncology, 137(3)

ISSN

00908258

Authors

Coleman, Robert L
Sill, Michael W
Bell-McGuinn, Katherine
et al.

Publication Date

2015-06-01

DOI

10.1016/j.ygyno.2015.03.042

Peer reviewed

changes, similar to what others have done with scars and pigmented lesions[22, 25], in order to examine the effectiveness of patient monitoring and compare it to results obtained using SFDI.

There was significant variability in the color changes seen from animal to animal for a given occlusion level. While most animals had noticeable flap color changes during complete 100% occlusions (no measurable blood flow) in the arterial and venous groups, there were some animals in both groups that did not. During the partial occlusion experiments most of the animals did not have noticeable changes in flap color. However, at the 75% occlusion level in the venous group, some changes in color were slightly above the JND line. Other groups using similar techniques in pigs have seen larger changes in color during occlusion[7], but this is probably attributed to the fact that they occluded the flaps for a much longer period of time. While it is likely that we could have created larger changes in color by extending occlusion times, ultimately we were interested in quantifying the ability of human observation to detect early signs of flap failure. We have previously shown the ability of SFDI to detect multiple physiological parameters related to the perfusion status of a given flap[21]. The use of those parameters, specifically ctTHb for monitoring venous occlusions and stO₂ for monitoring arterial occlusions, were significantly better at detecting changes in the flap. In addition to being more reliable at detecting changes at the 100% occlusion level, SFDI was also effective at detecting changes during the partial occlusions. When the data for this experiment was collected, it was analyzed at a later date, but the current version of the system can output results immediately after acquisition. The current standard of care is post-operative visual examination on an hourly basis. SFDI has the potential to be useful as a quantitative tool during this inspection.

It is important to note that the digital color camera used here was more effective at detecting flap color changes than unaided human perception. The camera was able to record changes below the JND threshold. This suggests the potential for cameras to not only be used for remote monitoring purposes, but also for actual analysis of flap health. One drawback to using digital color photography is that it is not effective at analyzing a flap at a single point in time. It must continuously monitor a flap with essentially calibrated, controlled lighting conditions and assumes that the baseline condition of the flap is healthy in order to be effective. Ultimately, the ability to monitor physiological parameters like oxygen saturation and total hemoglobin levels that are directly related to flap health are better indicators of early flap failure.

5. Conclusion

The clinical appearance of a free flap in the early postoperative period is not always representative of the underlying microvascular condition. As demonstrated by the color imaging data in our study, clinically apparent changes in flap color that are within the range of human detection are often not visible until significant vascular occlusion has occurred. Digital analysis of flap photos over time appears to be more sensitive in detecting changes compared human perception alone in a swine model. Further study should be done to determine the efficacy of this technique in a clinical setting. By comparison, SFDI is capable of detecting changes in tissue oxygen saturation as a result of partial blood flow occlusions

to pedicle flaps before they are visible to the human eye. The early detection of flaps with vascular compromise has the potential to improve response times and salvage rates, making photographic flap analysis and SFDI useful tools for free flap management.

Acknowledgments

The authors thank Hak-Su Kim, Earl Steward and Roger Geertsema for their assistance and guidance with respect to the animals in the study. We also thank Pierre Khoury for his help collecting data.

We thankfully recognize support from the Beckman Foundation and the NIH, including P41EB015890 (A Biomedical Technology Resource) from NIBIB and R42GM077713 from NIGMS. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIBIB or NIH. Dr. Mazhar, Dr. Cuccia, and Dr. Durkin have a financial interest in Modulated Imaging, Inc., which developed the Tissue OxImager and provided the device used in this study.

References

1. Chubb D, Rozen WM, Whitaker IS, Acosta R, Grinsell D, Ashton MW. The efficacy of clinical assessment in the postoperative monitoring of free flaps: a review of 1140 consecutive cases. *Plast Reconstr Surg*. 2010; 125(4):1157–1166. [PubMed: 20335866]
2. Pohlentz P, Blessmann M, Blake F, Li L, Schmelzle R, Heiland M. Outcome and complications of 540 microvascular free flaps: the Hamburg experience. *Clin Oral Inves*. 2007; 11(1):89–92.
3. Chen K-T, Mardini S, Chuang DC-C, Lin C-H, Cheng M-H, Lin Y-T, Huang W-C, Tsao C-K, Wei F-C. Timing of presentation of the first signs of vascular compromise dictates the salvage outcome of free flap transfers. *Plast Reconstr Surg*. 2007; 120(1):187–195. [PubMed: 17572562]
4. Bui DT, Cordeiro PG, Hu Q-Y, Disa JJ, Pusic A, Mehrara BJ. Free flap reexploration: indications, treatment, and outcomes in 1193 free flaps. *Plast Reconstr Surg*. 2007; 119(7):2092–2100. [PubMed: 17519706]
5. Kroll SS, Schusterman MA, Reece GP, Miller MJ, Evans GR, Robb GL, Baldwin BJ. Timing of pedicle thrombosis and flap loss after free-tissue transfer. *Plast Reconstr Surg*. 1996; 98(7):1230–1233. [PubMed: 8942909]
6. Steele MH. Three-Year Experience Using Near Infrared Spectroscopy Tissue Oximetry Monitoring of Free Tissue Transfers. *Ann Plast Surg*. 2011; 66(5):540–545. [PubMed: 21301288]
7. Russell JA, Conforti ML, Connor NP, Hartig GK. Cutaneous Tissue Flap Viability following Partial Venous Obstruction. *Plast Reconstr Surg*. 2006; 117(7):2259–2266. [PubMed: 16772926]
8. Chan RK, Liu A, Bojovic B, Chun Y, Guo L, Caterson SA, Orgill DP, Pribaz JJ. Venous congestion in abdominal flap breast reconstructions--a simple treatment for a temporary problem. *J Plast Reconstr Aesthet Surg*. 2011; 64(5):135–136.
9. Chubb D, Whitaker IS, Rozen WM, Ashton MW. Continued Observations in the Postoperative Monitoring of Free Flaps. *Plast Reconstr Surg*. 2012; 129(1):222–223.
10. Gimbel ML, Rollins MD, Fukaya E, Hopf HW. Monitoring Partial and Full Venous Outflow Compromise in a Rabbit Skin Flap Model. *Plast Reconstr Surg*. 2009; 124(3):796–803. [PubMed: 19730298]
11. Lorenzetti F, Ahovuo J, Suominen S, Salmi A, Asko-Seljavaara S. Colour Doppler ultrasound evaluation of haemodynamic changes in free tram flaps and their donor sites. *Scand J Plast Reconstr Surg Hand Surg*. 2002; 36(4):202–206. [PubMed: 12426993]
12. Vakharia KT, Henstrom D, Lindsay R, Cunnane MB, Cheney M, Hadlock T. Color Doppler ultrasound: effective monitoring of the buried free flap in facial reanimation. *Otolaryngol Head Neck Surg*. 2012; 146(3):372–376. [PubMed: 22261491]
13. Thorniley MS, Sinclair JS, Barnett NJ, Shurey CB, Green CJ. The use of near-infrared spectroscopy for assessing flap viability during reconstructive surgery. *Br J Plast Surg*. 1998; 51(3):218–226. [PubMed: 9664881]

14. Pharaon MR, Scholz T, Bogdanoff S, Cuccia D, Durkin AJ, Hoyt DB, Evans GRD. Early Detection of Complete Vascular Occlusion in a Pedicle Flap Model Using Quantitation Spectral Imaging. *Plast Reconstr Surg*. 2010; 126(6):1924–1935. [PubMed: 21124132]
15. Yafi A, Vetter TS, Scholz T, Patel S, Saager RB, Cuccia DJ, Evans GR, Durkin AJ. Postoperative quantitative assessment of reconstructive tissue status in a cutaneous flap model using spatial frequency domain imaging. *Plast Reconstr Surg*. 2011; 127(1):117–130. [PubMed: 21200206]
16. Whitaker I, Pratt G, Rozen W, Cairns S, Barrett M, Hiew L, Cooper M, Leaper D. Near Infrared Spectroscopy for Monitoring Flap Viability Following Breast Reconstruction. *J Reconstr Microsurg*. 2011
17. Matsui A, Lee BT, Winer JH, Laurence RG, Frangioni JV. Quantitative Assessment of Perfusion and Vascular Compromise in Perforator Flaps Using a Near-Infrared Fluorescence-Guided Imaging System. *Plast Reconstr Surg*. 2009; 124(2):451–460. [PubMed: 19644259]
18. Nguyen JT, Lin SJ, Tobias AM, Gioux S, Mazhar A, Cuccia DJ, Ashitate Y, Stockdale A, Oketokoun R, Durr NJ, Moffitt LA, Durkin AJ, Tromberg BJ, Frangioni JV, Lee BT. A novel pilot study using spatial frequency domain imaging to assess oxygenation of perforator flaps during reconstructive breast surgery. *Ann Plast Surg*. 2013; 71(3):308–315. [PubMed: 23945533]
19. Nguyen TT, Ramella-Roman JC, Moffatt LT, Ortiz RT, Jordan MH, Shupp JW. Novel application of a spatial frequency domain imaging system to determine signature spectral differences between infected and noninfected burn wounds. *J Burn Care Res*. 2013; 34(1):44–50. [PubMed: 23292572]
20. Gioux S, Mazhar A, Lee BT, Lin SJ, Tobias AM, Cuccia DJ, Stockdale A, Oketokoun R, Ashitate Y, Kelly E, Weinmann M, Durr NJ, Moffitt LA, Durkin AJ, Tromberg BJ, Frangioni JV. First-in-human pilot study of a spatial frequency domain oxygenation imaging system. *J Biomed Opt*. 2011; 16(8):086015. [PubMed: 21895327]
21. Ponticorvo A, Taydas E, Mazhar A, Scholz T, Kim HS, Rimler J, Evans GR, Cuccia DJ, Durkin AJ. Quantitative assessment of partial vascular occlusions in a swine pedicle flap model using spatial frequency domain imaging. *Biomed Opt Express*. 2013; 4(2):298–306. [PubMed: 23412357]
22. Cheon YW, Lee WJ, Rah DK. Objective and quantitative evaluation of scar color using the L*a*b* color coordinates. *J Craniofac Surg*. 2010; 21(3):679–684. [PubMed: 20485028]
23. Li-Tsang CW, Lau JC, Liu SK. Validation of an objective scar pigmentation measurement by using a spectrophotometer. *Burns*. 2003; 29(8):779–784. [PubMed: 14636751]
24. Rah DK, Kim SC, Lee KH, Park BY, Kim DW. Objective evaluation of treatment effects on port-wine stains using L*a*b* color coordinates. *Plast Reconstr Surg*. 2001; 108(4):842–847. [PubMed: 11547137]
25. Kim SC, Kim DW, Hong JP, Rah DK. A quantitative evaluation of pigmented skin lesions using the L*a*b* color coordinates. *Yonsei Med J*. 2000; 41(3):333–339. [PubMed: 10957887]
26. Miyamoto K, Takiwaki H, Hillebrand GG, Arase S. Development of a digital imaging system for objective measurement of hyperpigmented spots on the face. *Skin Res Technol*. 2002; 8(4):227–235. [PubMed: 12423541]
27. Leon K, Mery D, Pedreschi F, Leon J. Color measurement in L*a*b* units from RGB digital images. *Food Research International*. 2006; 39(10):1084–1091.
28. Hill B, Roger T, Vorhagen FW. Comparative analysis of the quantization of color spaces on the basis of the CIELAB color-difference formula. *Acm Transactions on Graphics*. 1997; 16(2):109–154.
29. Stokes M, Fairchild MD, Berns RS. Precision Requirements for Digital Color Reproduction. *Acm Transactions on Graphics*. 1992; 11(4):406–422.
30. Ikeda H, Dai W, Higaki Y. A Study on Colorimetric Errors Caused by Quantizing Color Information. *IEEE T Instrum Meas*. 1992; 41(6):845–849.
31. Mahy M, Vanmellaert B, Vaneycken L, Oosterlinck A. The Influence of Uniform Color Spaces on Color Image-Processing - a Comparative-Study of Cielab, Cieluv, and Atd. *Journal of Imaging Technology*. 1991; 17(5):232–243.
32. Vander Haeghen Y, Naeyaert JM, Lemahieu I, Philips W. An imaging system with calibrated color image acquisition for use in dermatology. *IEEE Trans Med Imaging*. 2000; 19(7):722–730. [PubMed: 11055787]

33. Forbes PD. Radiation effects in swine. I. Vascular supply of the skin and hair. USNRDL-TR-67-141. Res Dev Tech Rep. 1967:1–18. [PubMed: 5300775]
34. Cuccia DJ, Bevilacqua F, Durkin AJ, Ayers FR, Tromberg BJ. Quantitation and mapping of tissue optical properties using modulated imaging. *J Biomed Opt.* 2009; 14(2):024012. [PubMed: 19405742]
35. Mazhar A, Dell S, Cuccia DJ, Gioux S, Durkin AJ, Frangioni JV, Tromberg BJ. Wavelength optimization for rapid chromophore mapping using spatial frequency domain imaging. *J Biomed Opt.* 2010; 15(6):061716. [PubMed: 21198164]
36. Erickson TA, Mazhar A, Cuccia D, Durkin AJ, Tunnell JW. Lookup-table method for imaging optical properties with structured illumination beyond the diffusion theory regime. *J Biomed Opt.* 2010; 15(3):036013. [PubMed: 20615015]
37. Gioux S, Mazhar A, Cuccia DJ, Durkin AJ, Tromberg BJ, Frangioni JV. Three-dimensional surface profile intensity correction for spatially modulated imaging. *J Biomed Opt.* 2009; 14(3):034045. [PubMed: 19566337]
38. Nguyen JQ, Saager RB, Cuccia DJ, Kelly KM, Jakowatz J, Hsiang D, Durkin AJ. Effects of motion on optical properties in the spatial frequency domain. *J Biomed Opt.* 2011; 16(12):126009. [PubMed: 22191926]
39. Engel H, Huang JJ, Tsao CK, Lin CY, Chou PY, Brey EM, Henry SL, Cheng MH. Remote real-time monitoring of free flaps via smartphone photography and 3G wireless internet: A prospective study evidencing diagnostic accuracy. *Microsurgery.* 2011; 31(8):589–595. [PubMed: 22072583]
40. Varkey P, Tan NC, Giroto R, Tang WR, Liu YT, Chen HC. A picture speaks a thousand words: the use of digital photography and the Internet as a cost-effective tool in monitoring free flaps. *Ann Plast Surg.* 2008; 60(1):45–48. [PubMed: 18281795]




Fig 1. Photograph of a) top view and b) underside of typical swine pedicle flap preparation with instrumentation labeled. Time course of blood flow changes during a series of typical c) arterial and d) venous partial occlusions.




Fig 2.
Diagram of imaging system and feedback occlusion system.




Fig. 3. Color images, overlaid ΔE images, and overlaid ΔHbT images at different time points corresponding to different occlusion levels from a venous occlusion experiment that showed the most significant color changes.




Fig. 4. Color images, overlaid ΔE images, and overlaid ΔHbT images at different time points corresponding to different occlusion levels from a venous occlusion experiment that showed the least significant color changes.




Fig. 5.
The time courses of a) ΔE and b) HbT changes across all venous occlusion experiments.
Statistically significant changes at a given occlusion level are marked with an asterisk.




Fig. 6. Color images, overlaid ΔE images, and overlaid ΔstO_2 images at different time points corresponding to different occlusion levels from a typical arterial occlusion experiment.




Fig. 7.
 The time courses of a) ΔE and b) stO_2 changes across all arterial occlusion experiments. Statistically significant changes at a given occlusion level are marked with an asterisk.