

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Draft Genome Sequence for *Desulfovibrio africanus* Strain PCS

### Permalink

<https://escholarship.org/uc/item/18w8p8b6>

### Journal

Microbiology Resource Announcements, 1(2)

### ISSN

2576-098X

### Authors

Brown, Steven D

Utturkar, Sagar M

Arkin, Adam P

et al.

### Publication Date

2013-05-02

### DOI

10.1128/genomea.00144-13

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# Draft Genome Sequence for *Desulfovibrio africanus* Strain PCS

Steven D. Brown,<sup>a,b</sup> Sagar M. Utturkar,<sup>b</sup> Adam P. Arkin,<sup>c</sup> Adam M. Deutschbauer,<sup>c</sup> Dwayne A. Elias,<sup>a</sup> Terry C. Hazen,<sup>a,d</sup> Romy Chakraborty<sup>e</sup>

Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA<sup>a</sup>; Graduate School of Genome Science and Technology, University of Tennessee, Knoxville, Tennessee, USA<sup>b</sup>; Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA<sup>c</sup>; College of Engineering, University of Tennessee, Knoxville, Tennessee, USA<sup>d</sup>; Earth Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA<sup>e</sup>

***Desulfovibrio africanus* strain PCS is an anaerobic sulfate-reducing bacterium (SRB) isolated from sediment from Paleta Creek, San Diego, CA. Strain PCS is capable of reducing metals such as Fe(III) and Cr(VI), has a cell cycle, and is predicted to produce methylmercury. We present the *D. africanus* PCS genome sequence.**

Received 27 February 2013 Accepted 13 March 2013 Published 11 April 2013

**Citation** Brown SD, Utturkar SM, Arkin AP, Deutschbauer AM, Elias DA, Hazen TC, Chakraborty R. 2013. Draft genome sequence for *Desulfovibrio africanus* strain PCS. *Genome Announc.* 1(2):e00144-13. doi:10.1128/genomeA.00144-13.

**Copyright** © 2013 Brown et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Romy Chakraborty, [rchakraborty@lbl.gov](mailto:rchakraborty@lbl.gov).

Sulfate-reducing bacteria (SRB) are anaerobic microorganisms that play important roles in sulfur and carbon cycles in diverse environments (see reviews [1–3]). *Desulfovibrio africanus* strain PCS was isolated from a lactate/sulfate enrichment culture inoculated with sediment samples obtained from Paleta Creek, San Diego, CA. The isolated strain was Gram negative, motile, non-sporulating, and 99% similar by 16S rRNA gene sequencing to *Desulfovibrio africanus* subsp. *uniflagellum* (GenBank accession number EU659693) and *D. africanus* strain Walvis Bay (CP003221.1) (4), which is consistent with the species definition (5). *D. africanus* strains, including PCS, have different morphotypes associated with a cell cycle (6–10), and PCS incompletely oxidizes lactate, accumulating acetate as an end product.

*D. africanus* strains have been shown to methylate inorganic mercury [Hg(II)] to methylmercury (MeHg), a potent human neurotoxin (10–12). The capability to produce MeHg is found only in a subset of SRB and Fe(III)-reducing bacteria (IRB) (11–16). A 4.2-Mb complete genome sequence for *D. africanus* strain Walvis Bay (4, 17), which produces MeHg, has been reported (10, 12). Recently, genetic studies have shown that a two-gene cluster encoding a putative corrinoid-containing CO dehydrogenase/acetyl coenzyme A (acetyl-CoA) synthase, HgcA, and a 2[4Fe-4S] ferredoxin, HgcB, is required to produce MeHg in SRB and IRB (18). HgcA and HgcB are predicted to have roles as a methyl carrier and an electron donor, respectively. To date, strain PCS has not been tested for its ability to generate MeHg.

The genome sequence for strain PCS was generated using Illumina data, as described previously (19). Briefly, CLC Genomics Workbench (version 5.5) was used to trim 100-bp reads from a paired-end library for quality sequence data, and these were then assembled using Velvet (version 1.2.01) (20). The resulting assembly generated 45 DNA contigs for an estimated genome size of ~3.9 Mb. The maximum contig size was 609,036 bp, the average contig size was 87,322 bp, and the  $N_{50}$  was 140,584 bp. The average read depth was approximately 560× the estimated genome size. The draft genome sequence was annotated as previously described (21) and 3,561 candidate protein coding genes were predicted.

The PCS genome had a G+C content of 61.2%, which is similar to the 61.4% G+C content reported for strain Walvis Bay (4). Strain PCS shows 95% average nucleotide identity to strain Walvis Bay when the two genome sequences are compared using the JSpecies program (22). Strain PCS contains putative *hgcA* (PCS\_01240) and *hgcB* (PCS\_01242) genes that are ~97% and 98% identical, respectively, to their Walvis Bay counterparts at the nucleotide level. In both strains, a gene encoding a predicted radical S-adenosylmethionine (SAM) superfamily or Fe-S oxidoreductase protein is in a 3' position relative to *hgcA* and 5' relative to *hgcB*, a genetic organization that differs from those of other MeHg-producing bacteria like *D. desulfuricans* strain ND132 (16, 18). The *D. africanus* PCS genome sequence will facilitate further studies with this bacterium.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [AOSV00000000](https://www.ncbi.nlm.nih.gov/nuccore/AOSV00000000). The version described in this paper is the first version, [AOSV01000000](https://www.ncbi.nlm.nih.gov/nuccore/AOSV01000000).

## ACKNOWLEDGMENTS

We thank Loren Hauser and Doug Hyatt (ORNL) for examining potential *D. africanus hgcA* start sites.

The work conducted by ENIGMA-Ecosystems and Networks Integrated with Genes and Molecular Assemblies (<http://enigma.lbl.gov>), a Scientific Focus Area Program at Lawrence Berkeley National Laboratory, was supported by the Office of Science, Office of Biological and Environmental Research (BER), of the U.S. Department of Energy under contract number DE-AC02-05CH11231. This work was also supported through the BER Mercury Scientific Focus Area led by ORNL. Oak Ridge National Laboratory is managed by UT-Battelle, LLC, for the U.S. Department of Energy under contract DE-AC05-00OR22725.

## REFERENCES

1. Keller KL, Wall JD. 2011. Genetics and molecular biology of the electron flow for sulfate respiration in *Desulfovibrio*. *Front. Microbiol.* 2:135.
2. Muyzer G, Stams AJM. 2008. The ecology and biotechnology of sulphate-reducing bacteria. *Nat. Rev. Microbiol.* 6:441–454.
3. Odum JM, Peck HDJ. 1984. Hydrogenase, electron-transfer proteins, and

- energy coupling in the sulfate-reducing bacteria *Desulfovibrio*. *Annu. Rev. Microbiol.* 38:551–592.
4. Brown SD, Wall JD, Kucken AM, Gilmour CC, Podar M, Brandt CC, Teshima H, Detter JC, Han CS, Land ML, Lucas S, Han J, Pennacchio L, Nolan M, Pitluck S, Woyke T, Goodwin L, Palumbo AV, Elias DA. 2011. Genome sequence of the mercury-methylating and pleomorphic *Desulfovibrio africanus* strain Walvis Bay. *J. Bacteriol.* 193:4037–4038.
  5. Campbell LL, Kasprzycki MA, Postgate JR. 1966. *Desulfovibrio africanus* sp. n., a new dissimilatory sulfate-reducing bacterium. *J. Bacteriol.* 92:1122–1127.
  6. Chakraborty R, Joyner D, Wozel E, Holman H-YN, Lam S, Hazen TC. 2006. *Desulfovibrio* strain PCS, a novel metal reducing pleomorphic sulfate reducing bacterium, poster Q-166. Abstr. 106th Gen. Meet. Am. Soc. Microbiol. American Society for Microbiology, Washington, DC.
  7. Chakraborty R, Joyner DC, Wozel E, Holman HY, Hazen TC. 2006. *Desulfovibrio* strain PCS, a metal reducing pleomorphic sulfate reducing bacterium, abstr 1936. 11th Int. Symp. Microb. Ecol., Vienna, Austria, 20 to 25 August 2006.
  8. Chakraborty R, Tang YJ, Pingitore F, Keasling JD, Hazen T. 2008. Metabolic pathways in the pleomorphic metal-reducing organism *Desulfovibrio africanus* strain PCS, poster K-054. Abstr. 108th Gen. Meet. Am. Soc. Microbiol. American Society for Microbiology, Washington, DC.
  9. Jones HE. 1971. A re-examination of *Desulfovibrio africanus*. *Arch. Mikrobiol.* 80:78–86.
  10. Moberly JG, Miller CL, Brown SD, Biswas A, Brandt CC, Palumbo AV, Elias DA. 2012. Role of morphological growth state and gene expression in *Desulfovibrio africanus* strain Walvis Bay mercury methylation. *Environ. Sci. Technol.* 46:4926–4932.
  11. Ekstrom EB, Morel M, FM, Benoit JM. 2003. Mercury methylation independent of the acetyl-coenzyme a pathway in sulfate-reducing bacteria. *Appl. Environ. Microbiol.* 69:5414–5422.
  12. Gilmour CC, Elias DA, Kucken AM, Brown SD, Palumbo AV, Schadt CW, Wall JD. 2011. The sulfate-reducing bacterium *Desulfovibrio desulfuricans* ND132 as a model for understanding bacterial mercury methylation. *Appl. Environ. Microbiol.* 77:3938–3951.
  13. Fleming EJ, Mack EE, Green PG, Nelson DC. 2006. Mercury methylation from unexpected sources: molybdate-inhibited freshwater sediments and an iron-reducing bacterium. *Appl. Environ. Microbiol.* 72:457–464.
  14. Compeau GC, Bartha R. 1985. Sulfate-reducing bacteria: principal methylators of mercury in anoxic estuarine sediment. *Appl. Environ. Microbiol.* 50:498–502.
  15. King JK, Kostka JE, Frischer ME, Saunders FM. 2000. Sulfate-reducing bacteria methylate mercury at variable rates in pure culture and in marine sediments. *Appl. Environ. Microbiol.* 66:2430–2437.
  16. Brown SD, Gilmour CC, Kucken AM, Wall JD, Elias DA, Brandt CC, Podar M, Chertkov O, Held B, Bruce DC, Detter JC, Tapia R, Han CS, Goodwin LA, Cheng J-F, Pitluck S, Woyke T, Mikhailova N, Ivanova NN, Han J, Lucas S, Lapidus AL, Land ML, Hauser LJ, Palumbo AV. 2011. Genome sequence of the mercury-methylating strain *Desulfovibrio desulfuricans* ND132. *J. Bacteriol.* 193:2078–2079.
  17. Hurt RA, Brown SD, Podar M, Palumbo AV, Elias DA. 2012. Sequencing intractable DNA to close microbial genomes. *PLoS One* 7:e41295. doi:10.1371/journal.pone.0041295.
  18. Parks JM, Johs A, Podar M, Bridou R, Hurt RA, Smith JD, Tomanicek SJ, Qian Y, Brown SD, Brandt CC, Palumbo AV, Smith JC, Wall JD, Elias DA, Liang L. 2013. The genetic basis for bacterial mercury methylation. *Science* 339:1332–1335.
  19. Brown SD, Utturkar SM, Klingeman DM, Johnson CM, Martin SL, Land ML, Lu TYS, Tse-Yuan S, Schadt CW, Doktycz MJ, Pelletier DA. 2012. Twenty-one genome sequences from *Pseudomonas* species and 19 genome sequences from diverse bacteria isolated from the rhizosphere and endosphere of *Populus deltoides*. *J. Bacteriol.* 194:5991–5993.
  20. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
  21. Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Grechkin Y, Ratner A, Jacob B, Huang J, Williams P, Huntemann M, Anderson I, Mavromatis K, Ivanova NN, Kyrpides NC. 2012. IMG: the integrated microbial genomes database and comparative analysis system. *Nucleic Acids Res.* 40:D115–D122.
  22. Richter M, Rosselló-Móra R. 2009. Shifting the genomic gold standard for the prokaryotic species definition. *Proc. Natl. Acad. Sci. U. S. A.* 106:19126–19131.