

UC Irvine

UC Irvine Previously Published Works

Title

Correction to: Physical exercise augmented cognitive behaviour therapy for older adults with generalised anxiety disorder (PEXACOG): study protocol for a randomized controlled trial.

Permalink

<https://escholarship.org/uc/item/19w3f97g>

Journal

Trials, 21(1)

Authors

Stavestrands, Silje

Sirevåg, Kristine

Nordhus, Inger

et al.

Publication Date

2020-04-06

DOI

10.1186/s13063-020-04239-9

Peer reviewed

CORRECTION

Open Access



Correction to: Physical exercise augmented cognitive behaviour therapy for older adults with generalised anxiety disorder (PEXACOG): study protocol for a randomized controlled trial

Silje Haukenes Stavestrands^{1,2*}, Kristine Sirevåg^{1,2}, Inger Hilde Nordhus^{1,3}, Trond Sjøbø², Trygve Bruun Endal², Hans M. Nordahl^{4,9}, Karsten Specht¹, Åsa Hammar¹, Anne Halmøy^{5,11}, Egil W. Martinsen^{3,10}, Eva Andersson⁶, Helene Hjelmervik¹, Jan Mohlman⁷, Julian F. Thayer⁸ and Anders Hovland^{1,2}

Correction to: *Trials* (2019) 20:174
<https://doi.org/10.1186/s13063-019-3268-9>

Following the publication of our article [1], we have become aware of one error in the exclusion criteria, inconsistencies in Figs. 1 and 2, and a typo in the reference list.

Exclusion criterion # 10 is incorrect.

Correction:

(10) participating regularly in 60 min of moderate intensity physical exercise per week, divided across 2 or more bouts of physical exercise.

The physical exercise exclusion criteria are omitted from Fig. 1 in the original publication. The correct version of Fig. 1 is included here.

In Fig. 2, the interim testing contains three tests which are not conducted at this point of measure. These are The Five-Minute Pyramid test, The Biering-Sørensen Test, and the Dumbbell Arm Press Test. The correct version of Fig. 2 is included here.

We have discovered a typo in the reference for the Expectancy / Credibility Scale. The correct reference is:

Borkovec, T. D. and S. D. Nau (1972). “Credibility of analogue therapy rationales.” *Journal of Behavior Therapy and Experimental Psychiatry* 3(4): 257–260.

The original article can be found online at <https://doi.org/10.1186/s13063-019-3268-9>.

Author details

¹Faculty of Psychology, University of Bergen, Box 7800, NO-5020 Bergen, Norway. ²Sollvi DPS, Osvegen 15, NO-5228 Nesttun, Norway. ³Faculty of Medicine, University of Oslo, Box 1078, Blindern, NO-0316 Oslo, Norway. ⁴Department of Mental Health, Norwegian University of Science and Technology, Box 8905, NO-7491 Trondheim, Norway. ⁵Faculty of Medicine, K.G. Jebsen Centre for Neuropsychiatric Disorders, University of Bergen, Box 7800, NO-5020 Bergen, Norway. ⁶The Swedish School of Sport and Health Sciences, GIH, Box 5626, SE-114 86 Stockholm, Sweden. ⁷Department of Psychology, William Paterson University, 300 Pompton Road, Wayne, NJ 07470, USA. ⁸Department of Psychology, Ohio State University, 1835 Neil Avenue, Columbus, OH 43210, USA. ⁹St.Olavs Hospital HF, Nidaros DPS, Box 3250, Sluppen, NO-7006 Trondheim, Norway. ¹⁰Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway. ¹¹Kronstad DPS/Division of Psychiatry, Haukeland University Hospital, Box 1400, NO-5021 Bergen, Norway.

Published online: 06 April 2020

Reference

1. Stavestrands SH, et al. Physical exercise augmented cognitive behaviour therapy for older adults with generalised anxiety disorder (PEXACOG): study protocol for a randomized controlled trial. *Trials*. 2019;20:174. <https://doi.org/10.1186/s13063-019-3268-9>.

The original article can be found online at <https://doi.org/10.1186/s13063-019-3268-9>

* Correspondence: silje.h.stavestrands@uib.no

¹Faculty of Psychology, University of Bergen, Box 7800, NO-5020 Bergen, Norway

²Sollvi DPS, Osvegen 15, NO-5228 Nesttun, Norway

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 1 Study design. Visual presentation of the study design, including sample, assessments, and interventions. GAD, generalised anxiety disorder; CBT, cognitive behaviour therapy

TIMEPOINT	STUDY PERIOD									
	Pre-treatment			Treatment				Post-treatment		
	W -x	W -1	0	W 1-5	W 6	W 7-16	W 17	M 10	M 16	
	Enrolment	Baseline	Allocation	Initial phase: PE or AP	Interim testing	CBT + PE or AP	Post testing	6-m F-U	12-m F-U	
ENROLMENT:										
Eligibility screen	X									
Informed consent	X									
Allocation			X							
INTERVENTIONS:										
CBT + Physical Exercise (PE)										
CBT + Attention Placebo (AP)										
ASSESSMENTS:										
Interviews										
ADIS-IV (primary outcome measure)	X						X			
M.I.N.I. Interview	X									
Mini Mental State Examination	X									
Demography	X									
Questionnaires										
GAD-7	X	X		X	X	X	X	X	X	X
Beck Anxiety Inventory		X			X		X	X	X	X
PSWQ (primary outcome measure)		X			X		X	X	X	X
Geriatric Anxiety Inventory		X			X		X	X	X	X
Beck Depression Inventory - II		X			X		X	X	X	X
Quality of life inventory		X			X		X	X	X	X
Bergen Insomnia Scale		X			X		X	X	X	X
Expectancy / Credibility scale		X					X		X	X
IPAQ		X			X		X		X	X
Objective measures										
Actigraph - 7 days		X					X		X	X
Cortisol - 3 saliva samplers		X			X		X	X	X	X
DNA - saliva sample		X								
HRV		X			X		X	X	X	X
BDNF - blood sample		XX			XX		XX	XX	XX	XX
Eklblom-Bak Cycle test		X			X		X	X	X	X
Five-Minute Pyramid Test		X					X	X	X	X
Biering-Sørensen test		X					X	X	X	X
Timed sit-to-stand test		X			X		X	X	X	X
45° sit-up test		X			X		X	X	X	X
Dumbbell arm press test		X					X	X	X	X
Neuropsychological testing										
Vocabulary (WASI)		X					X		X	X
Matrix Reasoning (WASI)		X					X		X	X
Grooved Pegboard Test		X					X		X	X
Coding (WAIS-IV)		X					X		X	X
Numbers (WAIS-IV)		X					X		X	X
Wisconsin Card Sorting Test		X					X		X	X
D-KEFS (TMT, VF, CWIT)		X					X		X	X
California Verbal Learning Test		X					X		X	X
Subsample of 40 participants										
Magnetic resonance imaging		X					X		X	X

Fig. 2: SPIRIT Figure. Schedule of enrolment, interventions and assessments.
 Abbreviations: W Week; M Month; F-U Follow-up
 Note. XX indicates blood sampling at two timepoints, before and after Eklblom-Bak Cycle test and 20 mins of moderate intensity physical exercise.

Fig. 2 (See legend on next page.)

(See figure on previous page.)

Fig. 2 Standard protocol items: recommendation for interventional trials (SPIRIT) figure. Schedule of enrolment, interventions and assessments. Overview of the measures applied in the study. CBT, cognitive behaviour therapy; ADIS-IV, Anxiety Disorders Interview Schedule for DSM-IV; M.I.N.I., Mini International Neuropsychiatric Interview; PSWQ, Penn State Worry Questionnaire; GAD-7, Generalized Anxiety Disorder 7-item scale; IPAQ, International Physical Activity Questionnaire; HRV, heart rate variability; BDNF, brain-derived neurotropic factor; WASI, Wechsler Abbreviated Scale of Intelligence; WAIS-IV, Wechsler Adult Intelligence Scale – Fourth Edition; D-KEFS, Delis–Kaplan Executive Function System; TMT, Trail-Making Test; VF, Verbal Fluency; CWIT, Color Word Interference Test