

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

THE STUDY OF INVARIANT SURFACES AND THEIR BREAK-UP BY THE HAMILTON-JACOBI METHOD

### Permalink

<https://escholarship.org/uc/item/1bc4j3p9>

### Authors

Warnock, R.L.  
Ruth, R.D.

### Publication Date

1986-08-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

Physics Division

OCT 13 1986

LIBRARY AND  
DOCUMENTS SECTION

Presented at the 13th International Conference on  
High Energy Accelerators, Novosibirsk, USSR,  
August 7-11, 1986

THE STUDY OF INVARIANT SURFACES AND THEIR BREAK-UP  
BY THE HAMILTON-JACOBI METHOD

R.L. Warnock and R.D. Ruth

August 1986

**TWO-WEEK LOAN COPY**  
*This is a Library-Circulating Copy  
which may be borrowed for two weeks.*



LBL-21979  
c.2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE STUDY OF INVARIANT SURFACES AND THEIR BREAK-UP BY THE HAMILTON-JACOBI METHOD\*

R. L. WARNOCK

Lawrence Berkeley Laboratory, University of California, Berkeley, CA 94720

R. D. RUTH

Stanford Linear Accelerator Center, Stanford University, Stanford, CA 94305

Summary

We describe a method to compute invariant tori in phase space for classical non-integrable Hamiltonian systems. Our procedure is to solve the Hamilton-Jacobi equation stated as a system of equations for Fourier coefficients of the generating function. The system is truncated to a finite number of Fourier modes and solved numerically by Newton's method. The resulting canonical transformation serves to reduce greatly the non-integrable part of the Hamiltonian. In examples studied to date the convergence properties of the method are excellent, even near chaotic regions and on the separatrices of isolated broad resonances. We propose a criterion for breakup of invariant surfaces, namely the vanishing of the Jacobian of the canonical transformation to new angle variables. By comparison with results from tracking, we find in an example with two nearly overlapping resonances that this criterion can be implemented with sufficient accuracy to determine critical parameters for the breakup ('transition to chaos') to an accuracy of 5-10%.

The Hamilton-Jacobi Equation

We present results for a system with one degree of freedom having a periodic time-dependent Hamiltonian. The generalization to higher dimensions will be obvious. In angle-action variables the Hamiltonian is

$$H(\phi, J, \theta) = H_0(J) + V(\phi, J, \theta), \quad (1)$$

where  $\theta$  is the machine azimuth or 'time', and the perturbation  $V$  is periodic in  $\theta$  with period  $2\pi$ . We seek a canonical transformation  $(\phi, J) \mapsto (\psi, K)$  in the form<sup>1</sup>

$$J = K + G_\phi(\phi, K, \theta), \quad (2)$$

$$\psi = \phi + G_K(\phi, K, \theta), \quad (3)$$

such that the new Hamiltonian becomes a function of  $K$  alone. Subscripts denote partial derivatives. The Hamilton-Jacobi equation to determine the generator  $G$  is the requirement that the new Hamiltonian  $H$  indeed depend only on  $K$ ; namely

$$H_0(K + G_\phi) + V(\phi, K + G_\phi, \theta) + G_\theta = H_1(K). \quad (4)$$

We are interested in periodic solutions of (4) with the Fourier development

$$G(\phi, K, \theta) = \sum_{m,n} g_{mn}(K) e^{i(m\phi - n\theta)}. \quad (5)$$

We rearrange (4) by adding and subtracting terms so as to isolate terms linear in  $G_\phi$  and  $G_\theta$ . We then take the Fourier

transform for  $m \neq 0$  to cast Eq. (4) in the form

$$g = A(g), \quad (6)$$

where  $g = [g_{mn}]$  is a vector of Fourier coefficients and

$$A_{mn}(g) = \frac{i}{(\nu(K)m - n)} \frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^{2\pi} d\phi d\theta e^{-i(m\phi - n\theta)} \times \\ [H(\phi, K + G_\phi, \theta) - H_0(K) - \nu(K)G_\phi], \quad m \neq 0, \quad (7)$$

where  $\nu(K) = \partial H_0 / \partial K$ . To truncate the system (6) for numerical solution we restrict  $(m,n)$  to some bounded set  $B$  of integers, with  $m \neq 0$ , and put

$$G_\phi = \sum_{(m,n) \in B} i m g_{mn}(K) e^{i(m\phi - n\theta)}. \quad (8)$$

In an iterative solution of (6) the set  $B$  is selected so that at iterate  $(p+1)$  all  $A_{mn}(g^{(p)})$  with  $(m,n) \in B$  are greater than some preassigned small number; here  $g^{(p)}$  is the  $p^{\text{th}}$  iterate. Only the amplitudes  $g_{mn}$  for  $m \neq 0$  are required to calculate  $G_\phi$ ; the  $m = 0$  amplitude and also the function  $H_1(K)$  are determined from (4) *a posteriori*. Once  $G_\phi$  is known,  $J(\phi, \theta)$  may be plotted from Eq. (2). The action  $K$  is an input parameter.

The equation in the form (6) is suitable for the examples treated below, but not for typical accelerator problems involving short nonlinear lattice elements. For the latter, the Fourier analysis in  $\theta$  has slow convergence and should be avoided. Retaining the Fourier analysis in  $\phi$ , and using the periodic Green function for the operator  $i m \nu + \partial / \partial \theta$ , we can find an integral equation for the amplitudes  $g_m(K; \theta)$ . The system can be discretized to provide an equation for the variables  $g_m(K; \theta_i)$ ,  $m \geq 1$ , where the  $\theta_i$  are mesh points located only in the nonlinear elements of the lattice. The solution is automatically periodic in  $\theta$ . An alternative procedure is to treat the equation as a system of differential equations in  $\theta$ . The equation must be integrated only once around the accelerator with periodicity achieved by iteration, in analogy to nonlinear closed orbit calculations.

An Integrable Example

We show results from solving (6) - (8) by Newton's method, starting from  $g = 0$ . Our first example is a locally integrable case in which some of the invariant surfaces may be expressed analytically, namely the 4<sup>th</sup> order isolated resonance model with

$$H(\phi, J, \theta) = \nu_0 J + \alpha J^2 / 2 + \epsilon J^2 \cos(4\phi - \theta), \quad (9)$$

where  $\nu_0$ ,  $\alpha$ , and  $\epsilon$  are constants. We have successfully calculated a variety of qualitatively different curves (surfaces of section at fixed  $\theta$ ) for various choices of the parameters. For details about accuracy, rate of convergence, etc., see Ref. 2. The most difficult curves to compute are the separatrices around wide islands. The method works surprisingly well even for

\* Work supported by the Department of Energy, contracts DE-AC03-76SF00515 and DE-AC03-76SF00098.

such curves, as is seen in Figures 1 and 2. Fig. 1 shows separatrices computed in 9 iterations with 31 modes in the set  $B$ . The points in Figs. 1 and 2 are plotted in normalized phase space ( $\sqrt{J} \cos \phi$  vs.  $\sqrt{J} \sin \phi$ ) at  $\theta = 0$ . The inner separatrix (almost a square) and the outer separatrix (four lobes intersecting at right angles) are from two different calculations for two different values of  $K$ . Fig. 2 is a plot of curves from the exact analytic formulas for comparison. Similar results are obtained for nonintegrable Hamiltonians, in a region sufficiently close to a single resonance.

## The Two Resonance Model

Our second example is nonintegrable, and contains all the generic phenomena of nonlinear mechanics in  $1\frac{1}{2}$  or 2 dimensions. In a restricted region of phase space it should describe the essential features of one dimensional betatron motion in the presence of nonlinearities. The example is the two-resonance model with the Hamiltonian

$$H = \nu_0 J + \frac{1}{2} \alpha J^2 + \epsilon_1 J^{5/2} \cos(5\phi - 3\theta) + \epsilon_2 J^2 \cos(8\phi - 3\theta). \quad (10)$$

For small perturbation strengths  $\epsilon_1, \epsilon_2$  we compute an invariant curve (section of invariant surface at  $\theta = 0$ ) for a tune equal to the golden mean  $\nu_* = (\sqrt{5} - 1)/2$ , which is between the two resonances. Here we refer to the exact perturbed tune  $\nu_* = dH_1/dK$ , not the unperturbed tune  $\nu = \nu_0 + \alpha K$ . To maintain the perturbed tune at a preassigned value, we include the equation  $\nu_* = dH_1/dK$  as a constraint in the iteration (see Ref. 2). Having found an initial curve we then increase  $\epsilon_1, \epsilon_2$  (arbitrarily taking  $\epsilon_1 = 2\epsilon_2$ ) and look for the transition to chaotic behavior.

We choose  $\nu_0 = 0.5$ ,  $\alpha = 0.1$ , and consider a sequence of three cases with strengths and resonance widths  $\Delta J_1, \Delta J_2$  as follows:

- (i)  $\epsilon_1 = 2\epsilon_2 = 6 \times 10^{-5}$ ,  $\Delta J_1 = 0.049$ ,  $\Delta J_2 = 0.054$ ;
- (ii)  $\epsilon_1 = 2\epsilon_2 = 10^{-4}$ ,  $\Delta J_1 = 0.063$ ,  $\Delta J_2 = 0.070$ ;
- (iii)  $\epsilon_1 = 2\epsilon_2 = 1.25 \times 10^{-4}$ ,  $\Delta J_1 = 0.070$ ,  $\Delta J_2 = 0.078$ ;

By the Chirikov resonance overlap criterion,<sup>3</sup> the corresponding invariant curves should be close to breakup, since the resonance separation is  $J_{r_1} - J_{r_2} = 0.25$ .

To identify the transition to chaos as the  $\epsilon$ 's are increased, we propose the criterion that the Jacobian of Eq. (3) vanish at some  $(\phi, \theta)$ :

$$\partial\psi/\partial\phi = 1 + G_{\phi K} = \partial J/\partial K = 0. \quad (11)$$

At such a point it is in general impossible to solve uniquely for  $\phi$  as a function of  $\psi$ . Since  $\partial\psi/\partial\phi = \partial J/\partial K$  the heuristic picture is that two curves, differing infinitesimally in their  $K$  values, make contact one with the other.

In Figures 7, 9 and 11 we show the invariant curves in Cartesian plots of  $J(\phi, \theta = 0)$  for cases (i), (ii) and (iii) respectively, while in Figures 8, 10, and 12 we give the corresponding plots of  $\partial J/\partial K(\phi, \theta = 0)$ . The latter quantity allows us to test condition (11), since the minimum values of  $\partial J/\partial K$  are quite insensitive to  $\theta$ . The anticipated zeros of  $\partial J/\partial K$  are on the verge of appearance in Fig. 12.

In Figures 3, 4, and 6 we show enlargements of small portions of the invariant curves for cases (i), (ii) and (iii), together with points obtained by tracking from initial conditions on the appropriate curve. An orbit from a single initial condition was followed through  $N$  turns in  $\theta$ , with  $N = 4000, 4000$ , and 1500 for cases (i), (ii) and (iii) respectively. The good agreement

between tracking and computed curves indicated in Figures 3 and 4 is maintained over the full range of  $\phi$ . Chaotic behavior is evident in case (iii), but completely absent in case (ii). In Fig. 5 we show an intermediate case,  $\epsilon_1 = 2\epsilon_2 = 1.2 \times 10^{-4}$  tracked for 3000 turns, which is ambiguous. It might represent chaos or merely a high-order island chain not yet filled in. We believe that the scatter of points in Figures 5 and 6 is genuine, since we have checked accuracy in integration of Hamilton's equations by backtracking.

Comparing Figures 12, 5 and 6, one sees that condition (11) is first met at roughly that perturbation strength at which chaotic motion appears in tracking. Actually, the Hamilton-Jacobi results for  $\partial J/\partial K$  (but not those for  $J$ ) are slightly ambiguous for  $\epsilon_1 = 2\epsilon_2 > 10^{-4}$ , since at such large perturbations we encounter a limitation on the number of modes that can be accommodated while retaining convergence of Newton's method. Thus we cannot say precisely where (11) is first satisfied. A more precise determination of the transition should be possible by using a second canonical transformation or a modification of Newton's method. Assessing present results from tracking and  $\partial J/\partial K$  together, we estimate that the curve for the golden mean tune breaks up at  $\epsilon_1 = 2\epsilon_2 = (1.2 \pm 0.05) \times 10^{-4}$ .

Aside from comparisons to tracking, a stringent test of the method is to do a second canonical transformation, call it  $G^{(1)}$ , and see how large  $G_\psi^{(1)}$  is in comparison to  $G_\phi$ . This gives a measure of the residual distortion of the invariant surface, a correction to the main distortion of harmonic oscillator behavior obtained at the first step. We have computed  $G^{(1)}$  to lowest order, which is adequate for a good estimate. Taking absolute values averaged over angles, we find that  $(|G_\psi^{(1)}|)/(|G_\phi|)$  varies from  $2.8 \times 10^{-6}$  (at  $\epsilon_1 = 2\epsilon_2 = 6 \times 10^{-5}$ ) to  $4.1 \times 10^{-3}$  at  $\epsilon_1 = 2\epsilon_2 = 1.2 \times 10^{-4}$ . Since this ratio is so small compared to 1, it seems that our surfaces are good approximations to actual KAM surfaces, even near the transition to chaos.






A related approach under study is to construct symplectic maps for tracking by solving the Hamilton-Jacobi equation for a nonperiodic  $G$  under the initial condition  $G(\phi, K, 0) = 0$ . Then the new variables  $(\psi, K)$  are interpreted as initial conditions  $(\phi_0, J_0)$ , and Equations (2) and (3) provide a map  $(\phi_0, J_0) \mapsto (\phi, J)$  expressing time evolution of the system. It should be possible to construct a map for a full turn in  $\theta$ .

## Conclusions

We conclude that the Hamilton-Jacobi method provides a promising alternative to canonical perturbation theory and its modern variants. Unlike perturbation theory its algebraic complexity does not increase as more accuracy is demanded, and the required computer programs are quite simple. The generalization of (11) to higher dimensions, namely the condition  $\det(1 + G_{\phi K}) = 0$ , may provide a useful criterion for the transition to chaos in the full 5-dimensional phase space of betatron motion. In principle, successive canonical transformations computed by our method on progressively larger mode sets should provide a KAM algorithm<sup>1</sup> with enhanced convergence, leading to exact KAM tori. We give an extended account of this work in Ref. 2.

## REFERENCES



1. V. I. Arnold, "Mathematical Methods of Classical Mechanics", Springer, Berlin, 1978.
2. R.L. Warnock and R.D. Ruth, "Invariant Tori Through Direct Solution of the Hamilton-Jacobi Equation", SLAC-PUB-3865, LBL-21709, to be published.
3. B. V. Chirikov, Phys. Reports 52, No. 5 (1979) 264.



7-86

5507B1

Figures 1-2: Invariant curves for the 4<sup>th</sup> order single resonance model, by Newton solution of the Hamilton-Jacobi equation and analytic formula, respectively.  
 Figures 3-6: Small segments of invariant curves compared to points from tracking, for increasing resonance widths.



7-86

$\phi/2\pi$

$\phi/2\pi$

5507B2

Figures 7,9,11: Invariant KAM curves  $J(\phi, \theta = 0)$ .  
 Figures 8,10,12: The Jacobians  $\partial\psi/\partial\phi = \partial J/\partial\kappa$  at  $\theta = 0$ .  
 Cases (i) - (iii) are for increasing resonance widths.

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.



*LAWRENCE BERKELEY LABORATORY  
TECHNICAL INFORMATION DEPARTMENT  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720*