

UC Berkeley

UC Berkeley Previously Published Works

Title

A Structural Perspective on the Regulation of the Epidermal Growth Factor Receptor

Permalink

<https://escholarship.org/uc/item/1ch6v7wn>

Journal

Annual Review of Biochemistry, 84(1)

ISSN

0066-4154

Authors

Kovacs, Erika
Zorn, Julie Anne
Huang, Yongjian
et al.

Publication Date

2015-06-02

DOI

10.1146/annurev-biochem-060614-034402

Peer reviewed

A Structural Perspective on the Regulation of the Epidermal Growth Factor Receptor

Erika Kovacs,^{1,2,3} Julie Anne Zorn,^{1,2,3}
Yongjian Huang,^{1,2,3} Tiago Barros,^{1,2,3}
and John Kuriyan^{1,2,3,4,5}

Departments of ¹Molecular and Cell Biology and ⁴Chemistry, ²California Institute for Quantitative Biosciences, ³Howard Hughes Medical Institute, University of California, Berkeley, California 94720; email: kovacs@berkeley.edu, jazorn@berkeley.edu, yjhuang@berkeley.edu, tiago@berkeley.edu, kuriyan@berkeley.edu

⁵Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720

Annu. Rev. Biochem. 2015. 84:739–64

First published online as a Review in Advance on January 26, 2015

The *Annual Review of Biochemistry* is online at biochem.annualreviews.org

This article's doi:
10.1146/annurev-biochem-060614-034402

Copyright © 2015 by Annual Reviews.
All rights reserved

Keywords

receptor tyrosine kinase, ligand-induced dimerization, asymmetric dimer, oncogenic mutations, transmembrane coupling

Abstract

The epidermal growth factor receptor (EGFR) is a receptor tyrosine kinase that plays a critical role in the pathogenesis of many cancers. The structure of intact forms of this receptor has yet to be determined, but intense investigations of fragments of the receptor have provided a detailed view of its activation mechanism, which we review here. Ligand binding converts the receptor to a dimeric form, in which contacts are restricted to the receptor itself, allowing heterodimerization of the four EGFR family members without direct ligand involvement. Activation of the receptor depends on the formation of an asymmetric dimer of kinase domains, in which one kinase domain allosterically activates the other. Coupling between the extracellular and intracellular domains may involve a switch between alternative crossings of the transmembrane helices, which form dimeric structures. We also discuss how receptor regulation is compromised by oncogenic mutations and the structural basis for negative cooperativity in ligand binding.

Contents

INTRODUCTION	740
STRUCTURES OF THE EXTRACELLULAR MODULE: THE TETHERED MONOMER AND THE BACK-TO-BACK DIMER	743
STRUCTURES OF THE EXTRACELLULAR MODULE IN COMPLEX WITH THERAPEUTIC ANTIBODIES	745
KINASE ACTIVATION THROUGH THE FORMATION OF AN ASYMMETRIC DIMER OF KINASE DOMAINS	746
ONCOGENIC MUTATIONS IN THE KINASE DOMAIN	750
THE JUXTAMEMBRANE REGION AS AN ACTIVATING SEGMENT	751
TRANSMEMBRANE COUPLING	752
NEGATIVE COOPERATIVITY IN LIGAND BINDING TO THE EPIDERMAL GROWTH FACTOR RECEPTOR	754
STUDIES OF THE FULL-LENGTH RECEPTOR	755
THE OLIGOMERIC STATE OF THE RECEPTOR: LIGAND- INDEPENDENT DIMERS AND HIGHER-ORDER OLIGOMERS	757
CONCLUDING REMARKS	757

INTRODUCTION

The transition from unicellular life forms to multicellular ones occurred independently several times in evolution (1). An early and defining step in the evolutionary branch that led to animal life was the emergence of signaling systems that are built on top of core signaling components that arose earlier in life. These newer signaling systems include pathways that control intercellular signaling, cell–cell adhesion, and the development of the organism. Among the most important components of these systems are the tyrosine kinases, which signal through the generation of phosphotyrosine residues, molecular beacons that recruit proteins containing Src homology 2 (SH2) domains and phosphotyrosine-binding (PTB) domains (2).

Receptor tyrosine kinases are the outermost sentinels of the SH2-based signaling pathways. The extracellular modules of these receptors receive signals in the form of peptide hormones or cell-surface proteins, and they respond by activating intracellular tyrosine kinase domains (**Figure 1a**). Two other broad classes of receptors also trigger tyrosine kinase activity but do not contain kinase domains themselves. These are the cytokine receptors and various receptors involved in immune responses, such as the T cell, B cell, and Fc receptors (3).

The receptor tyrosine kinases, the cytokine receptors, and the immunological receptors have one architectural feature in common, which is that the protein subunits that make up the receptor complexes have only one transmembrane helix. This fact has caused the intact forms of these receptors to evade high-resolution structural analysis, perhaps because the plasma membrane plays a much more critical role in maintaining their structure than is the case for G protein-coupled receptors (GPCRs) or ion channels, which are now understood at an impressive level of molecular detail. Despite this limitation, structural, biochemical, and biophysical probing of the mechanisms of receptor tyrosine kinases has yielded an increasingly deep understanding of how these proteins work. In this review, we discuss what is known about the structure and function of one family of receptor tyrosine kinases, of which the epidermal growth factor receptor (EGFR) is the prototypical member.




Figure 1

Model for activation, domain architecture and evolutionary lineage of epidermal growth factor receptor (EGFR) family members. (a) Model for activation of EGFR family members. (b) Domain boundaries in EGFR family members. (c) Dendrogram of the evolution of EGFR family members. Her2 does not have a known ligand, and the kinase domain of Her3 is catalytically impaired. Panel c modified from Reference 140.

Human EGFR (also known as Her1/ErbB1, after the viral erythroblastoma gene) and its three close relatives, human epidermal growth factor receptors 2, 3, and 4 (Her2/ErbB2, also known as the neu oncogene; Her3/ErbB3; and Her4/ErbB3), control cell growth and differentiation (**Figure 1b**) (4–6). These receptors elicit potent mitogenic responses, and genetic abnormalities in these receptors represent one of the most prevalent defects in cancer cells (7). In fact, EGFR was the first cell-surface receptor to be recognized as an oncogene (8, 9). More than 30% of breast cancers, 60% of non-small-cell lung cancers, and 40% of glioblastomas either overexpress or contain activating mutations in EGFR family members (10–12). The link between this family of receptors and cancer progression led to the first antibody-based therapy for cancer, trastuzumab (Herceptin[®]), which targets the extracellular module of Her2 (13).

The four EGFR family members are among the ~60 receptor tyrosine kinases in the human genome (14). All receptor tyrosine kinases have an N-terminal extracellular ligand-binding module and a cytoplasmic tyrosine kinase catalytic domain linked by the single transmembrane helix. Despite this general similarity in their architecture, the extracellular modules of the principal subfamilies of these receptors are quite different (14). The transmembrane helices are also very divergent in sequence. Although the kinase domains of the receptor tyrosine kinases are grouped together in one major branch of the kinome (15), they differ in their mechanisms of activation.

The analysis of the genomes of organisms that diverged early along the evolutionary branch leading to the metazoan lineage shows that the choanoflagellates, unicellular organisms that are not metazoans, contain receptor tyrosine kinases but do not contain EGFR (16, 17). EGFR appears in organisms that are on the metazoan lineage, such as the sponges (16, 17), and its appearance may be the result of a fusion of an extracellular module with a cytoplasmic tyrosine kinase that occurred independently of the fusion events that generated other receptor tyrosine kinases. That EGFR may have arisen independently of other receptor tyrosine kinases is also suggested by the fact that the sequence of the EGFR kinase domain is closer to that of Ack1, a nonreceptor tyrosine kinase (18), than to that of other receptor tyrosine kinases. The creation of receptors by independent fusion events has also led to the family of receptors for transforming growth factor β (TGF- β) and bone morphogenetic protein (BMP) (19, 20). Like receptor tyrosine kinases, these receptors have extracellular ligand-binding domains that are connected to a cytoplasmic kinase domain by a single transmembrane helix, but their kinase domains have specificity for serine and threonine rather than for tyrosine (21).

Three gene-duplication events led to the four EGFR family members that are present in vertebrates (**Figure 1c**) (22). Gene duplication often causes one of the two resulting proteins to become degenerate, leading to functional differentiation among members of the same family (23); this phenomenon is observed in the EGFR family. Her2 has lost the capacity to bind ligands, and functions primarily by forming heterodimers with other family members (24). Her3 has lost robust kinase activity (25), so it also signals through heterodimerization (26).

The extracellular module that is characteristic of the EGFR family is a tandem duplication of two kinds of domains. The first and third domains (domains I and III) are compact and have a β -helical fold. The second and fourth domains (domains II and IV) are elongated and contain several cysteine-rich elements. The extracellular module is followed by a transmembrane helix and an intracellular module with a juxtamembrane segment, a kinase domain, and a C-terminal tail (**Figure 1a**) (4–6). Ligand binding to the extracellular module promotes dimerization, resulting in autophosphorylation of tyrosine residues in the intracellular C-terminal tail. This process leads to the recruitment of effector proteins containing SH2 and/or PTB domains to the phosphorylated tyrosine residues and the triggering of downstream signaling cascades. In addition to being a recruitment site for these effectors, the C-terminal tail has been implicated in kinase regulation (27).

The formation of dimers or higher-order oligomers is an essential step for the activation of EGFR family members (28–30). Although ligand binding promotes dimerization, enhanced expression of the receptors can also drive dimerization through mass action, which is important in certain cancers. Heterodimerization between EGFR family members is also an important aspect of their function and, as mentioned above, is especially important for Her3 activation because the kinase domain of that receptor is impaired (25). Although Her2 has an active kinase domain, it does not readily form homodimers under normal conditions, and it is activated by heterodimerization with ligand-bound partners, particularly Her3. Her4 forms homodimers and can signal from the plasma membrane, but it is also cleaved by membrane-associated proteases upon activation, which causes translocation of the kinase domain from the plasma membrane to intracellular compartments (31, 32).

Here, we review the current state of knowledge of the structure of EGFR family members, as well as the insights that structural information provides into the molecular details of the regulation and activation of these receptors. We cover a rather narrow subset of the extensive literature on EGFR, focusing only on topics that have a direct bearing on the three-dimensional structure of the receptor.

STRUCTURES OF THE EXTRACELLULAR MODULE: THE TETHERED MONOMER AND THE BACK-TO-BACK DIMER

The extracellular modules of all four EGFR family members have been crystallized and their structures determined either with ligand bound (for EGFR and Her4) or without ligand (for all four receptors), as well as in complex with antibodies or antibody mimics. Leahy (33) has written a particularly insightful review on the EGFR family, which can hardly be bettered in terms of the clarity with which it describes the structural principles underlying the construction and function of the extracellular modules.

Three key points emerge from these structural studies of the extracellular modules (33). First, dimerization of EGFR has an unexpected feature, which is that the dimer interface is formed entirely by the receptor itself, with the ligand bound on the outside. Second, there are only two principal classes of conformations in all the crystal structures. One corresponds to an “extended” form that, whether in a monomeric or a dimeric state, resembles the conformation of one protomer in the active dimer. The other conformation is a folded-over or “tethered” form, in which the dimerization element is buried within a monomer. Third, the restriction to two principal conformations of the extracellular modules arises because domains I and III form a relatively rigid unit, as do domains II and IV. As a consequence, the extracellular module appears to “click” into either the compact, tethered form or the extended form that is primed for dimerization (**Figure 2**).

Knowledge of the ligand-bound structure is based on the crystal structure of a construct of the extracellular module of human EGFR lacking almost the entire domain IV, in complex with TGF- α (34); the structure of the entire extracellular module of human EGFR, in complex with EGF (35, 36); and the structure of the human Her4 extracellular region, in complex with its ligand Neuregulin-1 β (37). Dimerization of the extracellular module is mediated principally by domain II. A rigid loop is inserted into the cysteine-rich repeats of this domain. This “dimerization arm” interacts with the corresponding element in the dimer partner. The domain II dimerization arm is completely occluded by intramolecular interactions with domain IV in the monomeric tethered conformation. The tethered conformation has been observed in the crystal structure of the isolated extracellular module of human EGFR, which includes an EGF molecule bound with very low affinity as well (38); in a crystal structure containing the entire extracellular region of human Her3 (39); and in a crystal structure containing the entire extracellular region of human Her4 (40). Ligand binding results in a huge conformational change, an $\sim 130^\circ$ rotation of domains I and II with respect to domains III and IV, which converts the extracellular module from a folded-over conformation to an extended one and generates a heart-shaped “back-to-back” dimer configuration in which the ligand is nestled between domains I and III of each subunit of the dimer (**Figure 2a**).

The structure of the extracellular module of Her2 has also been determined, and it is unique compared with those of other family members because it adopts the extended conformation without ligand binding. This extended conformation is evident in the crystal structure of the entire extracellular region of rat Her2 (13) and a truncated construct of human Her2 (**Figure 2b**) (41). Two key residues in the autoinhibitory domain IV contact region are not conserved (Gly563 and His565 of EGFR are replaced with proline and phenylalanine, respectively, in Her2), which could explain the absence of the autoinhibitory domain II–IV contact in Her2. The extracellular module of Her2 does not homodimerize in solution, perhaps because of subtle conformational differences between the extended extracellular module of Her2 and the extended conformation observed in dimeric EGFR.

The structure of the extracellular module of Her2 resembles that of *Drosophila melanogaster* EGFR (dEGFR). dEGFR is regulated by growth factor ligands, but a crystal structure shows that




Figure 2

Structures of the extracellular modules of the epidermal growth factor receptor (EGFR) family members. (a) The conformational change induced by ligand binding. (Left) The tethered conformation of EGFR [Protein Data Bank (PDB) identifier 1NQL; EGF bound at low pH was removed for clarity] rearranges to (right) the extended conformation of EGFR (PDB 3NJP) upon ligand binding. (b) Unliganded Her3 (PDB 1M6B) and Her4 (PDB 2AHX) can adopt a tethered conformation similar to that of EGFR, whereas Her2 (PDB 1N8Y) is in an extended conformation, even in the absence of ligand.

it, too, lacks the intramolecular tether present in human EGFR (42). Instead, a distinct set of autoinhibitory interactions between domains I and III holds unliganded dEGFR in an extended but inactive state, which could provide an alternative means of autoinhibition in the human Her2 extracellular module as well. A structure of a heterodimer including Her2 is not available at present.

The extracellular modules of EGFR family members are heavily glycosylated, containing nearly 40 kDa of sugar moieties (9). Twelve *N*-glycosylation sites have been identified in EGFR

(43–45), and early studies using inhibitors of *N*-glycosylation have shown that the glycosylation of EGFR is important for its translocation to the cell surface and maturation (46). Several recent studies have investigated specific glycosylation sites in EGFR (47–49). For example, the N420D mutant of EGFR is constitutively phosphorylated, and this ligand-independent activation is due to spontaneous oligomer formation (47). (We use a numbering system that does not count the 24-residue-long signal peptide. An alternative numbering scheme has residue numbers increased by 24.) Removal of *N*-glycosylation at Asn579 weakens the tethering interaction between domains II and IV, leading to a more relaxed extracellular module conformation and increased ligand-binding affinity (48).

STRUCTURES OF THE EXTRACELLULAR MODULE IN COMPLEX WITH THERAPEUTIC ANTIBODIES

EGFR and Her2 were among the very first receptors to be identified and associated with human tumors. Ever since then, these receptors have been targets for therapeutic intervention (7). Inhibitors of the EGFR family fall into two major classes: (*a*) monoclonal antibodies that target the extracellular module of the receptor and (*b*) small molecules that target the intracellular tyrosine kinase domain. Among the antibodies that have been approved by the US Food and Drug Administration (FDA), cetuximab and panitumumab target EGFR, whereas pertuzumab and trastuzumab target Her2.

Several crystal structures of therapeutic antibodies binding to EGFR and Her2 are now available (**Figure 3**) (13, 50, 51). Cetuximab binds to and blocks the ligand-binding site on EGFR domain III and sterically prevents EGFR from extending into its active conformation (51). Pertuzumab binds directly to the Her2 dimerization arm and thereby blocks receptor dimerization and activation (50). Trastuzumab binds to domain IV of Her2, proximal to the membrane, but this binding does not block either dimerization or activation of Her2 (13). Instead, trastuzumab inhibits the proteolytic cleavage of the Her2 extracellular module (52). In the plasma membrane, metalloproteinases are known to cleave the extracellular domains of many proteins, including EGFR family members (53, 54). This cleavage leaves behind the transmembrane helix and the tyrosine kinase domain, and such a construct is constitutively active.




Figure 3

Therapeutic antibodies target epidermal growth factor receptor (EGFR) and Her2 in versatile ways. Structures of (*a*) the EGFR–cetuximab [Protein Data Bank (PDB) identifier 1YY9], (*b*) Her2–pertuzumab (PDB 1S78), and (*c*) Her2–trastuzumab (PDB 1N8Z) complexes.

An important consideration in antibody therapeutics is whether an immune response will be elicited through recognition of the Fc region of the immunoglobulin G (IgG) molecule by the Fc receptor of effector cells to induce cytotoxicity, termed antibody-dependent cell-mediated cytotoxicity (55). Variation in the Fc region of antibodies influences the extent of cytotoxic effector responses and is an important component of trastuzumab action (55).

Nanobodies are small antigen-binding elements from the variable regions of camelid antibodies, which contain only a heavy chain. Nanobodies have been successfully developed to target a smaller, concave EGFR epitope, which is inaccessible to the larger, flatter monoclonal antibodies (56). In addition to extensive antibody-engineering research on EGFR, alternative methods to block the extracellular modules have been explored. These include the development of synthetic binding proteins (57) and RNA aptamers (58) that target the extracellular modules. These are valuable tools for investigations into the activation mechanism of these receptors (59).

KINASE ACTIVATION THROUGH THE FORMATION OF AN ASYMMETRIC DIMER OF KINASE DOMAINS

The first view of the structure of the EGFR kinase domain came from crystallization of the unphosphorylated form with the cancer drug erlotinib by scientists at Genentech (60). This structure revealed the canonical kinase fold in an active conformation (**Figure 4a**). Protein kinases contain a conserved Asp–Phe–Gly (DFG) motif at the base of the so-called activation loop or segment, a key regulatory element. The aspartate side chain is flipped out of the catalytic center in the inactive conformations of many protein kinases (the DFG-out conformation), preventing it from coordinating Mg-ATP, as it must for catalytic activity. In the erlotinib–EGFR complex, the DFG motif is in the active conformation, and the activation loop is open and properly configured to bind peptide substrate. Another important structural element of the kinase active site is helix α C, in the N lobe of the kinase, which packs closely against the rest of the kinase in the active conformation. In the erlotinib complex, helix α C is in the canonical active conformation, rotated inward toward the ATP-binding site. This conformation of the helix presents a conserved glutamate side chain (Glu738) for an ion-pairing interaction with a conserved lysine side chain (Lys721).

The active conformation of the unphosphorylated EGFR kinase domain was unexpected (60). Studies of the insulin receptor kinase domain had demonstrated a crucial connection between activation-loop phosphorylation and the adoption of the active conformation in that kinase

Figure 4

Kinase domain structures of the epidermal growth factor receptor (EGFR) family members. (a) The active [Protein Data Bank (PDB) identifier 2GS6] and the inactive (PDB 2GS7) conformation of the EGFR kinase domain. Helix α C is colored dark blue, the Asp–Phe–Gly (DFG) motif green, and the activation loop red. The active structure has an ATP analog–peptide conjugate bound, and the inactive structure has AMP–PNP bound (*yellow*). D, E, and F stand for the amino acid residues aspartate, glutamate, and phenylalanine, respectively. (b) The asymmetric dimer of the EGFR kinase domain (PDB 2GS6). The activator kinase is colored yellow, and the receiver (enzymatically active) kinase is colored blue. Residue contacts that are important on the activator and the receiver are highlighted. (c) A sequence alignment of the EGFR family members from human and mouse. Two regions containing the residues involved in the N- and C-lobe faces of the dimer interface are shown in the upper and lower panels, respectively. Identical residues are colored in red. Residues in the N- and C-lobe faces of the dimer interface are denoted by ovals and triangles, respectively. Blue and magenta represent residues in the dimer interface that are conserved among EGFR, Her2, and Her4 but not in Her3. (d) Structure of the Her3 kinase domain (PDB 3KEX) with a close-up of the active site. Residues resulting in a catalytically impaired kinase are labeled. The catalytic Asp813 in EGFR is replaced by Asn815 in Her3, the critical Glu738 in helix α C of EGFR is replaced with His740 in Her3, and Val737 and Thr738 in Her3 stabilize the inactive conformation of helix α C. (e) Structure of the L834R/T766M double-mutant EGFR kinase domain (PDB 4LL0) with a close-up of the active site. The bound inhibitor, PD168393, is colored yellow, and the mutations are labeled.

(61, 62). In addition, a large body of research on other kinases, particularly serine/threonine kinases, had led to the assumption that activation-loop phosphorylation is required for adoption of the active conformation.

The puzzle regarding the conformation of the EGFR kinase domain deepened when scientists at GlaxoSmithKline determined the structure of the unphosphorylated EGFR kinase domain bound to the cancer drug lapatinib, which revealed a very similar conformation to that of inactive Src family kinases and cyclin-dependent kinases (CDKs) (63). In this conformation, helix α C is swung outward from the N lobe and the activation loop forms a short helix (**Figure 4a**). Although it is possible that the particular conformation of the kinase domain observed in crystal structures is induced by the drugs, this seems unlikely because of the high affinity of binding. Indeed, erlotinib also binds to the inactive conformation of EGFR (64, 65). Mutational studies demonstrated that activation-loop phosphorylation is not an absolute requirement for EGFR activation (66). What, then, triggers the switch from the inactive to the active conformation of the EGFR kinase domain?

A breakthrough in our understanding of how EGFR activates came with the discovery that two EGFR kinase domains can interact in an asymmetric fashion so that one, termed the activator, switches on the other, termed the receiver, in an allosteric mechanism that does not rely on transphosphorylation (**Figure 4b**) (67). The clue to the activation mechanism was hidden in the original structure of EGFR bound to erlotinib (60), which showed an extensive interface between the N lobe of one kinase domain and the C lobe of the other in the crystal lattice. The mechanism by which the kinase domain of EGFR is activated resembles the activation of CDKs by cyclin. The C lobe of the activator kinase domain plays the role of the cyclin, even though it is not related structurally to the cyclins (68).

A key experiment showed that although the EGFR kinase domain has low activity in solution, its specific activity increases substantially when concentrated on lipid vesicles (67). This activation does not require the tyrosine in the activation loop. Instead, it requires the intact asymmetric dimerization interface, as revealed by mutagenesis of both the full-length EGFR in cells and the kinase domain in vitro. EGFR variants containing single mutations at the asymmetric dimer interface—such as the activator-impaired V924R mutant and the receiver-impaired I682Q mutant—do not respond to EGF binding. However, cotransfection of the activator-impaired and receiver-impaired mutants allows the formation of an intact activating interface between kinase domains, and results in comparable activity to the wild-type receptor.

The discovery of the asymmetric dimer provided a conceptual framework for understanding how the degenerate kinase domain of Her3 activates other members of the EGFR family, particularly Her2. Sequence alignments show that residues in Her3 that correspond to the C-lobe (activator) interface are very similar to the residues found at the corresponding location in other members in the family, whereas residues that correspond to the N-lobe (receiver) interface are divergent in Her3 (**Figure 4c**). These and other features of Her3 suggest that its kinase domain functions as an activator for other EGFR family members. Experimental evidence for an allosteric activation mechanism that is driven by protein–protein interaction rather than by phosphorylation came from a study on Her2–Her3 heterodimers that showed, using mass spectrometry, that conformational changes correlated with activation do not require tyrosine phosphorylation in the activation loop (69).

The structure of the Her4 kinase domain in an active conformation confirmed the importance of the asymmetric dimer (70). The structures of the kinase domains in the activator and receiver positions in the Her4 asymmetric dimer are nearly superimposable on the structure of the asymmetric dimer formed by the kinase domains of EGFR. The structure of the Her2 kinase domain

in a homodimeric configuration also revealed the formation of a very similar asymmetric dimer (71).

The potential significance of the asymmetric dimer was anticipated by Groenen et al. (72), who modeled the structure of the EGFR kinase domain based on that of the insulin receptor, and noticed that there are exposed hydrophobic patches in the distal surfaces of both the N and C lobes of the modeled EGFR kinase domain. These authors speculated that the two lobes might interact to stabilize the active conformation, which is a remarkable insight considering that the structure of the EGFR kinase domain was only determined 5 years later.

The hydrophobic surface on the N lobe of the EGFR kinase domain is specific to the active conformation, as the residues that form this surface are sequestered by conformational change in the inactive Src-CDK conformation. The hydrophobic surface in the C lobe of the activator, in contrast, is not in a region where conformational changes are observed in kinase domains, and it is likely to remain available as an interaction surface even in the inactive state. This surface is exploited by the feedback inhibitor Mig6. The crystal structure of a complex between the EGFR kinase domain and a fragment of Mig6 shows that Mig6 binds to the distal surface of the C lobe of the kinase domain, and inhibition is achieved by blocking the activating dimer interface (73).

The formation of asymmetric dimers by the kinase domains is important for substrate presentation in fibroblast growth factor receptors (74), and heterodimerization of kinase domains is a critical step in the activation of RAF kinases (75). These dimers involve completely different interfaces than that observed in EGFR, and sequence comparisons suggest that the activation mechanism described here for the EGFR family is likely to be unique for this family.

Long-timescale molecular dynamics simulations have shown in atomic detail how the kinase domain of EGFR transitions between the active and inactive states (76). The simulations suggest that local unfolding, or “cracking,” at the hinge region between the N and C lobes of the kinase is a necessary step in the transition, leading to a set of conformations in which there is room for a rearrangement of the activation loop. The intermediate conformations revealed by the simulations differ significantly from both the inactive and active structures determined crystallographically.

Among the EGFR family members, the kinase domain of Her3 is unique in that it lacks several conserved residues that are critical for kinase activity. The structure of the Her3 kinase domain, reported by Jura et al. (25) and by Shi et al. (77), shows how sequence changes have disrupted the canonical active conformation of kinases (**Figure 4d**). Key catalytic residues are missing, helix α C is shortened and distorted, and the activation loop is unable to take on the canonical active conformation. Nevertheless, the activator surface is intact, and Jura et al. (25) showed that Her3 can serve as an activator for EGFR. These authors also showed that the Her3 kinase domain has essentially no detectable kinase activity toward peptide substrates.

The ability of the Her3 kinase domain to activate that of Her2 allosterically has been demonstrated by use of carboxyl group footprinting mass spectrometry to analyze Her2 and Her3 kinase domain heterodimers (69). These experiments have shown that Her2 and Her3 kinase domains preferentially form asymmetric heterodimers, with Her3 and Her2 monomers occupying the activator and receiver kinase positions, respectively.

Shi et al. (77) reported the remarkable finding that Her3 has a very low, but detectable autophosphorylation rate when concentrated on vesicles. These authors used quantum mechanical calculations to delineate a reaction pathway for Her3-catalyzed phosphoryl transfer that does not require the conserved catalytic base and that can be catalyzed by the “inactive-like” conformation observed crystallographically. A modified model for Her3 signaling was proposed on the basis of this low level of Her3 autophosphorylation activity. In this model, Her2 “preactivates” Her3 through transphosphorylation, and because phosphorylated Her3 demonstrates increased kinase

activity, the authors argued that activated Her3 monomers can then form functional homodimers (78). Nevertheless, the specific activity of Her3 autophosphorylation is extremely low, and has been estimated to be ~1,000-fold lower than that of EGFR (25). A functional role for Her3 autophosphorylation remains to be established.

ONCOGENIC MUTATIONS IN THE KINASE DOMAIN

The clinical importance of EGFR and Her2 in cancer development has resulted in substantial interest in oncogenic mutations in the kinase domains (7). One of the most common of these mutations, the L834R substitution, is located in the activation loop of the kinase domain. The crystal structure of the L834R mutant (79) shows that it is in the active conformation, and its oncogenic property has been attributed to its ability to lock the enzyme in the active conformation. This finding explains the increased sensitivity of this mutant to tyrosine kinase inhibitors (80) that may prefer the active conformation (60, 79), such as gefitinib and erlotinib.

A recent study suggests that the oncogenicity of L834R is the result of enhancing dimerization, not necessarily just stabilizing an active conformation (81). Using long-timescale molecular dynamics simulations, Shan et al. (81) showed that the N-lobe dimerization interface of the kinase domain is intrinsically disordered and becomes ordered only upon dimerization. They demonstrated that the L834R mutation facilitates EGFR dimerization by suppressing this local disorder. Static light scattering, native gel analyses, and enzyme assays indicated that this mutation causes abnormally high activity by promoting EGFR dimerization, rather than by simply allowing activation without dimerization. Another molecular dynamics study suggests a reconciliation of both views, and proposes that both the change in the relative stability of the active versus the inactive state and the suppression of disorder seem to play a role in the activation of EGFR variants with the L834R mutation (82).

Unfortunately, resistance to EGFR inhibitors often arises due to a second mutation, T766M. Thr766 is the “gatekeeper” residue in the kinase domain, so called because the nature of the residue at this position is a key determinant of inhibitor specificity. Structural analyses of the T766M variant of EGFR suggest that it can accommodate inhibitors that target the active state (83). This hypothesis agrees with the results of molecular dynamics simulations, which suggest that the methionine interacts with hydrophobic residues surrounding the active site to stabilize a compact, active conformation (82). An unexpected finding is that the T766M mutation in EGFR increases the affinity of the oncogenic L834R variant for ATP by more than an order of magnitude (83). The increased affinity for ATP relative to EGFR inhibitors is likely the primary mechanism by which the T766M mutant confers drug resistance.

Red-Brewer et al. (84) demonstrated a so-called superreceptor activity of these mutant kinases, especially the L834R/T766M double mutant. EGFR kinase domains are thought to assume the activator or receiver role in a random manner, but the EGFR mutants found in lung cancer preferentially assume the receiver position in the presence of wild-type EGFR or Her2, leading to hyperphosphorylation of the wild-type activator. The crystal structure of the T766M/L834R double mutant shows that the kinase domain adopts an active conformation, with an intact asymmetric dimer interface (**Figure 4e**) (84). Red-Brewer et al. argued that the double mutant adopts the receiver position because the energetic cost of inducing the active conformation in the L834R/T766M mutant is lower relative to wild-type receptors. Consequently, in a mixed population of wild-type and oncogenic EGFR, the mutants preferentially assume the active, receiver position. Thus, if wild-type receptors are expressed together with mutant EGFR, the net activity of the mutant will increase, suggesting that wild-type EGFRs have a critical role in tumorigenesis alongside their mutated counterparts.

THE JUXTAMEMBRANE REGION AS AN ACTIVATING SEGMENT

The intracellular juxtamembrane segment plays crucial regulatory roles in several receptor kinases. Structures of the type I TGF- β receptor kinase domain, a serine/threonine kinase, showed that the unphosphorylated juxtamembrane segment interacts with helix α C, holding it in an inactive conformation that is further buttressed by the FK506-binding protein FKBP12 (85). This autoinhibition is released by phosphorylation of serine residues within the juxtamembrane segment. This mechanism bears superficial resemblance to that observed in Ephrin receptors, in which the juxtamembrane segment also holds the kinase in an inactive conformation by interacting with helix α C (86). This theme is repeated in kinases such as FLT3 (87) and c-Kit (88). Given these findings, it came as a surprise that deletion of the juxtamembrane region results in loss of phosphorylation in the EGFR tail (89), a finding that suggests a crucial role for this segment in activation of the kinase.

The EGFR juxtamembrane segment consists of two major portions: JM-A (residues 645–663) and JM-B (residues 664–682). The role of the JM-B segment was determined independently and simultaneously by two groups. Red-Brewer et al. (90) determined the structure of a construct of EGFR that included the kinase domain and the juxtamembrane segment. Jura et al. (91) reinterpreted crystal lattice contacts in a structure of the Her4 kinase domain that included the JM-B segment (92). In both structures, the JM-B segment forms a clamp or a latch that reaches across from the N lobe of the receiver kinase domain in an asymmetric dimer to engage the C lobe of the activator kinase domain (**Figure 5a**).

Red-Brewer et al. (90) performed alanine-scanning mutagenesis and showed that mutations in the C-terminal 19 residues of the juxtamembrane segment abolish EGFR activation. Phosphorylation of T654 and T669 destabilizes, while the V665M lung cancer mutation stabilizes, the active dimer conformation by reducing and increasing the strength of the juxtamembrane latch




Figure 5

Structures of the juxtamembrane latch and the transmembrane helices of the epidermal growth factor receptor (EGFR). (a) The asymmetric dimer of EGFR, with the juxtamembrane (JM) latch colored green [Protein Data Bank (PDB) identifier 3G0P]. (b) An NMR structure of the transmembrane and JM-A helices of EGFR in lipid bicelles (PDB 2M20). Interactions between the N-terminal GxxxG-like motifs in the transmembrane segments and the LRLL motifs in the juxtamembrane segments are colored yellow.

interaction, respectively. By using *in vitro* kinase assays, Jura et al. (91) demonstrated that the juxtamembrane region dimerizes and activates the EGFR kinase, and that it is required on both the activator and the receiver. By designing several mutations in the juxtamembrane segment, based on the Her4 structure, these authors provided evidence that the JM-B segment latches the activated kinase domain to the activator. They also presented the structure of an alternative, symmetric, and inactive EGFR kinase dimer in which the formation of the activating juxtamembrane latch is prevented by the C-terminal tail, suggesting a means to prevent ligand-independent activation.

In the structure of the EGFR kinase domain determined by Red-Brewer et al. (90), the JM-A segment packs against another kinase domain, but this crystal contact is unlikely to be maintained in the intact receptor at the plasma membrane. On the basis of results obtained by NMR, Jura et al. (91) speculated that the JM-A segments of the two kinases instead form an antiparallel helical dimer in the active configuration of the receptor.

Replacement of the juxtamembrane segment with an unstructured linker in the context of the full-length receptor abolishes phosphorylation of EGFR, without measurable effects on receptor dimerization or ligand binding, which supports the critical role of the juxtamembrane region in the formation of the asymmetric dimer (93). Scheck et al. (94) demonstrated the formation of an antiparallel coiled coil within JM-A by fluorescence spectroscopy. They proposed that this conformational transition is functionally coupled to receptor activation by EGF, whereas TGF- α binding is communicated to the intracellular domains through formation of an alternative helical interface. This idea suggests that the juxtamembrane segment may differentially relay the signal initiated by binding of different ligands.

Apart from serving as a latch between asymmetric kinase dimers, the juxtamembrane segment has been proposed to interact with negatively charged lipids in the membrane and also with calmodulin (95), mediate feedback signals through threonine phosphorylation (96), and play a role in sorting and recycling (97). The juxtamembrane segment has also been implicated in negative cooperativity in ligand binding by EGFR, which we discuss below.

TRANSMEMBRANE COUPLING

The transmembrane segment of EGFR is a 24-residue-long single α -helix. The structure of the transmembrane segment of the Her2 homodimer (98), and later that of the EGFR–Her2 heterodimer (99), has been studied by NMR in lipid bicelles. In both cases, the transmembrane segments associate through N-terminal GxxxG-like motifs. These motifs have been identified as general dimerization motifs for transmembrane helices on the basis of experiments on the glycoporphin A transmembrane segment (100, 101), and they have been implicated in the self-association of transmembrane helices in EGFR family members (102). The observed dimeric structures of the transmembrane helices in lipid bicelles explain the effect of some oncogenic mutations, such as the I655V and the V659E substitutions in the transmembrane helix of Her2. Both residues participate in stabilizing dimerization through the N-terminal GxxxG-like motifs (98, 99). The I655V mutation may stabilize homodimerization through the replacement of a large side chain by a smaller one, and the V659E substitution may do so because of intermolecular hydrogen bonding of the glutamate side chain.

The structure of the transmembrane and the cytoplasmic juxtamembrane segment of EGFR in lipid bicelles has been determined by NMR, aided by molecular dynamics simulations (**Figure 5b**) (103, 104). This structure explained how the configuration of the transmembrane helices in the EGFR dimer can couple to the conformation of the juxtamembrane segments, and how these conformations are compatible with the asymmetric kinase dimer. The structure of the

transmembrane segment reveals a helical dimer that is consistent with the Her2 structures determined by Bocharov et al. (98) and Mineev et al. (99). The C-terminal ends of the transmembrane helices are separated by ~ 20 Å, which provides the appropriate spacing for an antiparallel interaction between the JM-A helices. This interface is formed by LRLL motifs in the juxtamembrane segment. Mutating all four of the small residues in the N-terminal dimerization interface in the transmembrane helix to isoleucine (T624I/G625I/G628I/A629I) results in significant inhibition of EGFR in cellular assays, providing experimental evidence for the importance of the N-terminal association between transmembrane helices in receptor activation (103).

The transmembrane helices of EGFR and Her2 contain a second dimerization motif toward their C-terminal ends. Fleishman et al. (105) have speculated that the presence of both N-terminal and C-terminal dimerization motifs generates a transmembrane switch, in which the dimer of transmembrane helices toggles between two configurations, with the N-terminal (extracellular) ends close together in one and the C-terminal (intracellular) ends close together in the other. These authors proposed that these configurations correspond to the active and inactive states of the receptor, respectively. Using NMR and molecular dynamics simulations, Endres et al. (103) and Arkhipov et al. (104) showed that stabilizing interactions through the C-terminal dimerization motif, by an I640E mutation, leads to the disruption of the N-terminal interface between the transmembrane helices and to the disruption of the antiparallel JM-A interaction. Using cell-based assays, these authors demonstrated that the I640E mutation in the intact receptor impairs EGF-dependent activation at low surface densities. Other researchers have suggested that the two conserved GxxxG-like motifs also play a role in mediating and/or stabilizing homo- and heterodimers (106, 107).

According to the model proposed by Endres et al. (103) and Arkhipov et al. (104), the extracellular modules in ligand-bound dimers assume a configuration favoring dimerization of the transmembrane helices near their N-terminal ends, dimerization of the juxtamembrane segments, and formation of asymmetric kinase dimers. In ligand-free dimers, by holding apart the N-terminal ends of the transmembrane helices, the extracellular modules instead favor C-terminal dimerization of the transmembrane helices; juxtamembrane segment dissociation and membrane burial; and formation of symmetric, inactive kinase dimers. **Figure 7a** (below) shows our current understanding of what the active state of the full-length receptor might look like.

Analyses of the monomer and dimer forms of EGFR by chemical and disulfide cross-linking suggested that EGFR has a preformed dimeric structure without bound ligand, and that ligand binding induces rotation of the extracellular juxtamembrane region—hence the transmembrane segment, which reorients the cytoplasmic module and results in activation of the kinase domains (108). Using an elegant protein engineering approach, Bell et al. (109) demonstrated similar rotational coupling in Her2. They designed a series of transmembrane helix mutants that sequentially move two glutamate residues, within a simplified transmembrane segment, across the entire transmembrane region. The movement of this dimerization motif is expected to rotate the kinase domains $\sim 103^\circ$ per residue. Rotation of this interface does not affect dimerization but leads to a periodic oscillation in kinase activation.

Lu et al. (36) have argued that the transmembrane helix plays a more passive role than is indicated by the preceding discussion. Analyses of disulfide cross-linking experiments indicate that EGF-induced dimerization of the transmembrane helices involves a less extensive interface than that found in glycophorin A and integrin, two receptors that dimerize in the absence of activation. Systematic mutagenesis of residues in the transmembrane helix of EGFR to leucine and phenylalanine shows that no single mutation disrupts transmembrane signaling. However, as shown by Das et al. (103), multiple mutations are required to disrupt this interface, supporting the model of active transmembrane coupling.

NEGATIVE COOPERATIVITY IN LIGAND BINDING TO THE EPIDERMAL GROWTH FACTOR RECEPTOR

A puzzling aspect of the binding of EGF to EGFR in cellular assays is that the Scatchard plot is not linear, as would be expected for independent binding sites on the receptor with no cooperativity. Instead, the Scatchard plot is curved in a concave-up manner. This feature of the binding isotherm was interpreted in terms of two populations of EGFR in cells (110). This interpretation suggests that a small population of the receptors (~10%) have very high affinity for the ligand ($K_d \sim 50$ pM), whereas the majority of receptors have much lower affinity ($K_d \sim 3$ nM). It was proposed that EGF binds with high affinity to the dimeric form of the receptor, but with low affinity to the monomeric form (28). Another interpretation is that spatial segregation of populations of the receptor underlies the heterogeneity of binding (111, 112).

Investigators in the field now consider the concave-up Scatchard plot as being due instead to negative cooperativity in dimeric receptors. Macdonald & Pike (113) have made extensive measurements of EGF binding to EGFR-bearing cells. They generated stable cell lines expressing EGFR under an inducible promoter, which allowed control of the level of EGFR expression. Using this system, they measured the binding of radiolabeled EGF to these cells as a function of EGFR expression level. For the wild-type receptor, the binding isotherms shift from left to right with increasing EGFR density; that is, the apparent binding becomes weaker at higher EGFR densities on the cell surface. Modeling of the data suggests that the affinity of EGF for the second site on the EGFR dimer (2.9 nM) is substantially less than the affinity for the first site (190 pM); that is, the dimer exhibits negative cooperativity.

Pike and colleagues (114) also obtained information about the structural requirements of negative cooperativity by performing binding analyses on cells expressing increasing levels of various mutant forms of EGFR. These findings link the extracellular juxtamembrane region, particularly the region responsible for the tethering interaction in domain IV (residues 561–585), to negative cooperativity. The intracellular juxtamembrane portion of the receptor is also involved in the generation of negative cooperativity, given that deletion of the kinase domain as well as the C-terminal tail yielded a receptor with negative cooperativity, whereas deletion of the entire intracellular module led to a receptor that showed no cooperativity (115).

By performing a more detailed analysis of various intracellular juxtamembrane mutants, Pike and colleagues (116) linked the stepwise binding of two ligands to kinase activation. In this model, the binding of EGF to the first site on the dimer induces the formation of one asymmetric kinase dimer; then the binding of EGF to the second site is required to disrupt the initial asymmetric dimer and allow the formation of the reciprocal asymmetric dimer. Thus, some of the energy of binding to the second site is used to reorient the first asymmetric dimer, leading to a lower binding affinity and the observed negative cooperativity. The implications of such a model have been studied extensively (117–120).

One of the difficulties of explaining negative cooperativity in human EGFR is that the isolated extracellular module does not show negative cooperativity *in vitro*. In contrast, the isolated extracellular module of the *Drosophila* receptor, dEGFR, exhibits negative cooperativity in binding to its ligand, Spitz, when the binding is analyzed *in vitro* using purified proteins. A breakthrough in understanding occurred with the structural and biophysical analysis of dEGFR by Lemmon and colleagues (121). The crystal structure of a singly liganded asymmetric extracellular dimer of dEGFR reveals the structural basis for negative cooperativity in that receptor (**Figure 6**).

Before interacting with Spitz, the ligand-binding sites are identical in preformed dimers of dEGFR. The first (highest-affinity) binding event yields the singly liganded, asymmetric dimer because Spitz “wedges” itself between domains I and III in one subunit of the dimer and pushes




Figure 6

Structural basis for negative cooperativity in ligand binding to *Drosophila* epidermal growth factor receptor (dEGFR). The unliganded [Protein Data Bank (PDB) identifier 3I2T], singly liganded (PDB 3LTG), and doubly liganded (PDB 3LTF) dEGFR dimer are depicted. The structure of the doubly liganded human EGFR dimer (PDB 3NJP) is shown on the right. Modified from Reference 121.

them apart. This process distorts domain II and forces a substantial reorientation of the dimerization arm, allowing formation of a more extensive, asymmetric dimer interface. In this asymmetric dEGFR dimer, domain II in the unoccupied receptor is structurally restrained and can no longer bend to allow Spitz to wedge itself fully into the unoccupied ligand-binding site without disrupting the extensive asymmetric interface. Thus, binding of Spitz to the second site is weaker, leading to negative cooperativity.

Although structures of human EGFR with only one ligand bound have not been determined, Tynan et al. (122) suggest that in humans, such asymmetry is brought forth by interactions with the plasma membrane. Fluorescence resonance energy transfer imaging reveals that a high-affinity ligand-binding human EGFR conformation is consistent with the extracellular region aligned flat on the plasma membrane, in such a way that the second binding site is occluded. Molecular dynamics simulations also suggest that asymmetrical interactions with the membrane may be responsible for this negative cooperativity (123). Such simulations further indicate that heterodimers formed by extracellular modules of Her2 and other EGFR family members assume an asymmetric conformation similar to that of *Drosophila* EGFR dimers (124).

STUDIES OF THE FULL-LENGTH RECEPTOR

Two groups have succeeded in purifying nearly full-length recombinant EGFR (lacking most of the C-terminal tail) in detergent micelles (125, 126). This approach allows the biochemical characterization of ligand-induced activation of the intact receptor, previously feasible only by indirect methods in cells. Wang et al. (127) studied the two most common non-small-cell lung cancer mutations in EGFR: L834R and a deletion mutant, $\Delta 722-726$. They found both to be at least as active as the EGF-bound receptor, and that activity was still strongly coupled to asymmetric kinase dimer formation. Previous studies of the isolated kinase domains had indicated that the L834R mutation was sufficient to promote full constitutive activity, and the authors pointed out




Figure 7

Full-length epidermal growth factor receptor (EGFR) and its oligomerization states. (a) A proposed composite model of full-length EGFR based on the structures of individual modules [Protein Data Bank (PDB) identifier 3NJP for the extracellular module, PDB 2M20 for the transmembrane–juxtamembrane A (JM-A) helices, and PDB ID 2GS6 for the kinase domains]. (b) Schematics for possible oligomerization states of EGFR in cells.

that such discrepancies underscore the limitations of attempting to understand the regulation of this complex receptor by studying isolated fragments.

Negative-stain electron microscopy has been used to visualize the coupling between the extracellular and intracellular modules of EGFR at low resolution (128, 129). The unliganded receptor adopts a monomeric and tethered extracellular module conformation. Intriguingly, the EGF-bound receptor adopts two distinct dimer conformations: a rodlike kinase domain structure consistent with the asymmetric dimer and a more globular intracellular region consistent with a more symmetric association. These two types of dimers coexist with a monomeric form. Inhibitors that stabilize the active or inactive conformation of the kinase active site, as well as various mutations, shift the equilibrium among the three observed forms. The coupling of one conformation of an activated receptor extracellular module to multiple kinase-domain

arrangements suggests that the linkage between the extracellular and intracellular regions of EGFR is unexpectedly flexible. It is important to note, however, that these experiments have been carried out on detergent-solubilized EGFR, which could substantially alter the behavior of the receptor compared with that in lipid bilayers.

THE OLIGOMERIC STATE OF THE RECEPTOR: LIGAND-INDEPENDENT DIMERS AND HIGHER-ORDER OLIGOMERS

Numerous studies suggest that the classical model of unliganded monomer to EGF-bound dimer transition might be too simple (**Figure 7b**). Several groups have been addressing the question of what the resting state of the receptor is—whether it is predominantly a monomer or, rather, an unliganded dimer (108, 130–136). Very briefly, the main conclusion arising from these reports is that in the basal state EGFR exists predominantly as a monomer, while it is in equilibrium with ligand-independent dimers. Clustering before ligand binding is rare for EGFR, but it might be a characteristic of Her2. The fraction of EGFR that dimerizes before ligand binding depends primarily on receptor surface densities and the cell types used.

In a related question regarding the oligomeric state of EGFR, Burgess and colleagues (137–139) argue that a dimer–tetramer transition is required for receptor activation. Their data are consistent with a significant fraction of liganded EGFR tetramers. They propose that such tetramers comprise two dimers juxtaposed in a side-by-side (or slightly staggered) arrangement (137) and that tetrameric EGFR is the main signaling unit (138, 139).

CONCLUDING REMARKS

Structural analyses of the extracellular and intracellular modules of EGFR have provided invaluable insight into how this receptor, central in cancer development, functions. Crystallography combined with biochemical, biophysical, and cell biology experiments have revealed the detailed mechanisms of how ligand binding leads to conformational change and how the extracellular module dimerizes through an interface completely mediated by the receptor itself. These approaches demonstrated how the EGFR kinase domains activate through an allosteric mechanism involving asymmetric kinase dimers. NMR provided insight into transmembrane–juxtamembrane coupling, and we now understand the structural basis for negative cooperativity in ligand binding in *Drosophila* EGFR.

Despite all these advances, we are still lacking a complete understanding of how the full-length receptor functions. This is mainly due to the technical difficulty of crystallizing single-pass transmembrane proteins, of which few, if any, structures are currently available. New avenues in membrane protein crystallography and electron microscopy, as well as long-timescale molecular dynamics simulations, are promising directions that might provide a breakthrough in the field. There are also several mechanistic questions that cannot be tackled from a purely structural perspective, and for which the integration of sophisticated cell biology approaches will be necessary. These remaining issues include understanding the oligomerization state of the receptor, both in the resting state and as the active signaling unit; understanding the precise mechanism of tail phosphorylation; and determining whether the asymmetry present at the kinase level is translated into differential phosphorylation of the activator and the receiver tail. We believe that these efforts will ultimately enable us to fully understand the complex regulation of this receptor family, knowledge that will be essential for improving cancer therapeutics directed against this class of receptors.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We thank Mark Lemmon, Linda Pike, Dan Leahy, and all members of our group for helpful discussions. This review was partially supported by the National Cancer Institute (grant number 2-R01-CA096504-06). J.A.Z. is supported by a postdoctoral fellowship from the National Institutes of Health, National Cancer Institute (F32 CA177087-02). Y.H. is a Howard Hughes Medical Institute International Student Research Fellow.

LITERATURE CITED

1. Ruiz-Trillo I, Burger G, Holland PWH, King N, Lang BF, et al. 2007. The origins of multicellularity: a multi-taxon genome initiative. *Trends Genet.* 23:113–18
2. Lim WA, Pawson T. 2010. Phosphotyrosine signaling: evolving a new cellular communication system. *Cell* 142:661–67
3. Hubbard SR, Till JH. 2000. Protein tyrosine kinase structure and function. *Annu. Rev. Biochem.* 69:373–98
4. Avraham R, Yarden Y. 2011. Feedback regulation of EGFR signalling: decision making by early and delayed loops. *Nat. Rev. Mol. Cell Biol.* 12:104–17
5. Endres NF, Engel K, Das R, Kovacs E, Kuriyan J. 2011. Regulation of the catalytic activity of the EGF receptor. *Curr. Opin. Struct. Biol.* 21:777–84
6. Lemmon MA, Schlessinger J, Ferguson KM. 2014. The EGFR family: not so prototypical receptor tyrosine kinases. *Cold Spring Harb. Perspect. Biol.* 6:a020768
7. Arteaga CL, Engelman JA. 2014. ERBB receptors: from oncogene discovery to basic science to mechanism-based cancer therapeutics. *Cancer Cell* 25:282–303
8. Downward J, Yarden Y, Mayes E, Scrase G, Totty N, et al. 1984. Close similarity of epidermal growth factor receptor and *v-erb-B* oncogene protein sequences. *Nature* 307:521–27
9. Ullrich A, Coussens L, Hayflick JS, Dull TJ, Gray A, et al. 1984. Human epidermal growth factor receptor cDNA sequence and aberrant expression of the amplified gene in A431 epidermoid carcinoma cells. *Nature* 309:418–25
10. Sharma SV, Bell DW, Settleman J, Haber DA. 2007. Epidermal growth factor receptor mutations in lung cancer. *Nat. Rev. Cancer* 7:169–81
11. Westphal M, Meima L, Szonyi E, Lofgren J, Meissner H, et al. 1997. Heregulins and the ErbB-2/3/4 receptors in gliomas. *J. Neuro-Oncol.* 35:335–46
12. Harari D, Yarden Y. 2000. Molecular mechanisms underlying ErbB2/HER2 action in breast cancer. *Oncogene* 19:6102–14
13. Cho HS, Mason K, Ramyar KX, Stanley AM, Gabelli SB, et al. 2003. Structure of the extracellular region of HER2 alone and in complex with the herceptin Fab. *Nature* 421:756–60
14. Lemmon MA, Schlessinger J. 2010. Cell signaling by receptor tyrosine kinases. *Cell* 141:1117–34
15. Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S. 2002. The protein kinase complement of the human genome. *Science* 298:1912–34
16. Richter DJ, King N. 2013. The genomic and cellular foundations of animal origins. *Annu. Rev. Genet.* 47:509–37
17. Nichols SA, Dirks W, Pearse JS, King N. 2006. Early evolution of animal cell signaling and adhesion genes. *PNAS* 103:12451–56
18. Yokoyama N, Miller WT. 2003. Biochemical properties of the Cdc42-associated tyrosine kinase ACK1. Substrate specificity, autophosphorylation, and interaction with Hck. *J. Biol. Chem.* 278:47713–23

19. Kang JS, Liu C, Derynck R. 2009. New regulatory mechanisms of TGF- β receptor function. *Trends Cell Biol.* 19:385–94
20. Massague J. 2012. TGF β signalling in context. *Nat. Rev. Mol. Cell Biol.* 13:616–30
21. Hanks SK, Hunter T. 1995. Protein kinases 6. The eukaryotic protein kinase superfamily: kinase (catalytic) domain structure and classification. *FASEB J.* 9:576–96
22. Stein RA, Staros JV. 2006. Insights into the evolution of the ErbB receptor family and their ligands from sequence analysis. *BMC Evol. Biol.* 6:79
23. Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate genes. *Science* 290:1151–55
24. Yarden Y, Sliwkowski MX. 2001. Untangling the ErbB signalling network. *Nat. Rev. Mol. Cell Biol.* 2:127–37
25. Jura N, Shan Y, Cao X, Shaw DE, Kuriyan J. 2009. Structural analysis of the catalytically inactive kinase domain of the human EGF receptor 3. *PNAS* 106:21608–13
26. Berger MB, Mendrola JM, Lemmon MA. 2004. ErbB3/HER3 does not homodimerize upon neuregulin binding at the cell surface. *FEBS Lett.* 569:332–36
27. Pines G, Huang PH, Zwang Y, White FM, Yarden Y. 2010. EGFRvIV: a previously uncharacterized oncogenic mutant reveals a kinase autoinhibitory mechanism. *Oncogene* 29:5850–60
28. Yarden Y, Schlessinger J. 1987. Self-phosphorylation of epidermal growth factor receptor: evidence for a model of intermolecular allosteric activation. *Biochemistry* 26:1434–42
29. Yarden Y, Schlessinger J. 1987. Epidermal growth factor induces rapid, reversible aggregation of the purified epidermal growth factor receptor. *Biochemistry* 26:1443–51
30. Honegger AM, Kris RM, Ullrich A, Schlessinger J. 1989. Evidence that autophosphorylation of solubilized receptors for epidermal growth factor is mediated by intermolecular cross-phosphorylation. *PNAS* 86:925–29
31. Vecchi M, Baulida J, Carpenter G. 1996. Selective cleavage of the heregulin receptor ErbB-4 by protein kinase C activation. *J. Biol. Chem.* 271:18989–95
32. Cheng QC, Tikhomirov O, Zhou W, Carpenter G. 2003. Ectodomain cleavage of ErbB-4: characterization of the cleavage site and m80 fragment. *J. Biol. Chem.* 278:38421–27
33. Leahy DJ. 2004. Structure and function of the epidermal growth factor (EGF/ErbB) family of receptors. *Adv. Protein Chem.* 68:1–27
34. Garrett TP, McKern NM, Lou M, Elleman TC, Adams TE, et al. 2002. Crystal structure of a truncated epidermal growth factor receptor extracellular domain bound to transforming growth factor α . *Cell* 110:763–73
35. Ogiso H, Ishitani R, Nureki O, Fukai S, Yamanaka M, et al. 2002. Crystal structure of the complex of human epidermal growth factor and receptor extracellular domains. *Cell* 110:775–87
36. Lu C, Mi LZ, Grey MJ, Zhu J, Graef E, et al. 2010. Structural evidence for loose linkage between ligand binding and kinase activation in the epidermal growth factor receptor. *Mol. Cell Biol.* 30:5432–43
37. Liu P, Cleveland TE 4th, Bouyain S, Byrne PO, Longo PA, Leahy DJ. 2012. A single ligand is sufficient to activate EGFR dimers. *PNAS* 109:10861–66
38. Ferguson KM, Berger MB, Mendrola JM, Cho HS, Leahy DJ, Lemmon MA. 2003. EGF activates its receptor by removing interactions that autoinhibit ectodomain dimerization. *Mol. Cell* 11:507–17
39. Cho HS, Leahy DJ. 2002. Structure of the extracellular region of HER3 reveals an interdomain tether. *Science* 297:1330–33
40. Bouyain S, Longo PA, Li S, Ferguson KM, Leahy DJ. 2005. The extracellular region of ErbB4 adopts a tethered conformation in the absence of ligand. *PNAS* 102:15024–29
41. Garrett TP, McKern NM, Lou M, Elleman TC, Adams TE, et al. 2003. The crystal structure of a truncated ErbB2 ectodomain reveals an active conformation, poised to interact with other ErbB receptors. *Mol. Cell* 11:495–505
42. Alvarado D, Klein DE, Lemmon MA. 2009. ErbB2 resembles an autoinhibited invertebrate epidermal growth factor receptor. *Nature* 461:287–91
43. Smith KD, Davies MJ, Bailey D, Renouf DV, Hounsell EF. 1996. Analysis of the glycosylation patterns of the extracellular domain of the epidermal growth factor receptor expressed in Chinese hamster ovary fibroblasts. *Growth Factors* 13:121–32

44. Sato C, Kim JH, Abe Y, Saito K, Yokoyama S, Kohda D. 2000. Characterization of the N-oligosaccharides attached to the atypical Asn-X-Cys sequence of recombinant human epidermal growth factor receptor. *J. Biochem.* 127:65-72
45. Zhen Y, Caprioli RM, Staros JV. 2003. Characterization of glycosylation sites of the epidermal growth factor receptor. *Biochemistry* 42:5478-92
46. Gamou S, Shimizu N. 1988. Glycosylation of the epidermal growth factor receptor and its relationship to membrane transport and ligand binding. *J. Biochem.* 104:388-96
47. Tsuda T, Ikeda Y, Taniguchi N. 2000. The Asn-420-linked sugar chain in human epidermal growth factor receptor suppresses ligand-independent spontaneous oligomerization. Possible role of a specific sugar chain in controllable receptor activation. *J. Biol. Chem.* 275:21988-94
48. Whitson KB, Whitson SR, Red-Brewer ML, McCoy AJ, Vitali AA, et al. 2005. Functional effects of glycosylation at Asn-579 of the epidermal growth factor receptor. *Biochemistry* 44:14920-31
49. Fernandes H, Cohen S, Bishayee S. 2001. Glycosylation-induced conformational modification positively regulates receptor-receptor association: a study with an aberrant epidermal growth factor receptor (EGFRvIII/ΔEGFR) expressed in cancer cells. *J. Biol. Chem.* 276:5375-83
50. Franklin MC, Carey KD, Vajdos FF, Leahy DJ, de Vos AM, Sliwkowski MX. 2004. Insights into ErbB signaling from the structure of the ErbB2-pertuzumab complex. *Cancer Cell* 5:317-28
51. Li S, Schmitz KR, Jeffrey PD, Wiltzius JJ, Kussie P, Ferguson KM. 2005. Structural basis for inhibition of the epidermal growth factor receptor by cetuximab. *Cancer Cell* 7:301-11
52. Baselga J, Albanell J, Molina MA, Arribas J. 2001. Mechanism of action of trastuzumab and scientific update. *Semin. Oncol.* 28:4-11
53. Vecchi M, Carpenter G. 1997. Constitutive proteolysis of the ErbB-4 receptor tyrosine kinase by a unique, sequential mechanism. *J. Cell Biol.* 139:995-1003
54. Codony-Servat J, Albanell J, Lopez-Talavera JC, Arribas J, Baselga J. 1999. Cleavage of the HER2 ectodomain is a pervanadate-activable process that is inhibited by the tissue inhibitor of metalloproteases-1 in breast cancer cells. *Cancer Res.* 59:1196-201
55. Clynes RA, Towers TL, Presta LG, Ravetch JV. 2000. Inhibitory Fc receptors modulate in vivo cytotoxicity against tumor targets. *Nat. Med.* 6:443-46
56. Schmitz KR, Bagchi A, Roovers RC, van Bergen en Henegouwen PMP, Ferguson KM. 2013. Structural evaluation of EGFR inhibition mechanisms for nanobodies/VHH domains. *Structure* 21:1214-24
57. Hackel BJ, Neil JR, White FM, Witttrup KD. 2012. Epidermal growth factor receptor downregulation by small heterodimeric binding proteins. *Protein Eng. Des. Sel.* 25:47-57
58. Chen CH, Chernis GA, Hoang VQ, Landgraf R. 2003. Inhibition of heregulin signaling by an aptamer that preferentially binds to the oligomeric form of human epidermal growth factor receptor 3. *PNAS* 100:9226-31
59. Zhang Q, Park E, Kani K, Landgraf R. 2012. Functional isolation of activated and unilaterally phosphorylated heterodimers of ERBB2 and ERBB3 as scaffolds in ligand-dependent signaling. *PNAS* 109:13237-42
60. Stamos J, Sliwkowski MX, Eigenbrot C. 2002. Structure of the epidermal growth factor receptor kinase domain alone and in complex with a 4-anilinoquinazoline inhibitor. *J. Biol. Chem.* 277:46265-72
61. Hubbard SR. 1997. Crystal structure of the activated insulin receptor tyrosine kinase in complex with peptide substrate and ATP analog. *EMBO J.* 16:5572-81
62. Hubbard SR, Wei L, Ellis L, Hendrickson WA. 1994. Crystal structure of the tyrosine kinase domain of the human insulin receptor. *Nature* 372:746-54
63. Wood ER, Truesdale AT, McDonald OB, Yuan D, Hassell A, et al. 2004. A unique structure for epidermal growth factor receptor bound to GW572016 (lapatinib): relationships among protein conformation, inhibitor off-rate, and receptor activity in tumor cells. *Cancer Res.* 64:6652-59
64. Gajiwala KS, Feng J, Ferre R, Ryan K, Brodsky O, et al. 2013. Insights into the aberrant activity of mutant EGFR kinase domain and drug recognition. *Structure* 21:209-19
65. Park JH, Liu Y, Lemmon MA, Radhakrishnan R. 2012. Erlotinib binds both inactive and active conformations of the EGFR tyrosine kinase domain. *Biochem. J.* 448:417-23

66. Gotoh N, Tojo A, Hino M, Yazaki Y, Shibuya M. 1992. A highly conserved tyrosine residue at codon 845 within the kinase domain is not required for the transforming activity of human epidermal growth factor receptor. *Biochem. Biophys. Res. Commun.* 186:768–74
67. Zhang X, Gureasko J, Shen K, Cole PA, Kuriyan J. 2006. An allosteric mechanism for activation of the kinase domain of epidermal growth factor receptor. *Cell* 125:1137–49
68. Jura N, Zhang X, Endres NF, Seeliger MA, Schindler T, Kuriyan J. 2011. Catalytic control in the EGF receptor and its connection to general kinase regulatory mechanisms. *Mol. Cell* 42:9–22
69. Collier TS, Diraviyam K, Monsey J, Shen W, Sept D, Bose R. 2013. Carboxyl group footprinting mass spectrometry and molecular dynamics identify key interactions in the HER2–HER3 receptor tyrosine kinase interface. *J. Biol. Chem.* 288:25254–64
70. Qiu C, Tarrant MK, Choi SH, Sathyamurthy A, Bose R, et al. 2008. Mechanism of activation and inhibition of the HER4/ErbB4 kinase. *Structure* 16:460–67
71. Aertgeerts K, Skene R, Yano J, Sang BC, Zou H, et al. 2011. Structural analysis of the mechanism of inhibition and allosteric activation of the kinase domain of HER2 protein. *J. Biol. Chem.* 286:18756–65
72. Groenen LC, Walker F, Burgess AW, Treutlein HR. 1997. A model for the activation of the epidermal growth factor receptor kinase involvement of an asymmetric dimer? *Biochemistry* 36:3826–36
73. Zhang X, Pickin KA, Bose R, Jura N, Cole PA, Kuriyan J. 2007. Inhibition of the EGF receptor by binding of MIG6 to an activating kinase domain interface. *Nature* 450:741–44
74. Bae JH, Boggon TJ, Tome F, Mandiyan V, Lax I, Schlessinger J. 2010. Asymmetric receptor contact is required for tyrosine autophosphorylation of fibroblast growth factor receptor in living cells. *PNAS* 107:2866–71
75. Hu J, Stites EC, Yu H, Germino EA, Meharena HS, et al. 2013. Allosteric activation of functionally asymmetric RAF kinase dimers. *Cell* 154:1036–46
76. Shan Y, Arkhipov A, Kim ET, Pan AC, Shaw DE. 2013. Transitions to catalytically inactive conformations in EGFR kinase. *PNAS* 110:7270–75
77. Shi F, Telesco SE, Liu Y, Radhakrishnan R, Lemmon MA. 2010. ErbB3/HER3 intracellular domain is competent to bind ATP and catalyze autophosphorylation. *PNAS* 107:7692–97
78. Steinkamp MP, Low-Nam ST, Yang S, Lidke KA, Lidke DS, Wilson BS. 2014. ErbB3 is an active tyrosine kinase capable of homo- and heterointeractions. *Mol. Cell. Biol.* 34:965–77
79. Yun CH, Boggon TJ, Li Y, Woo MS, Greulich H, et al. 2007. Structures of lung cancer–derived EGFR mutants and inhibitor complexes: mechanism of activation and insights into differential inhibitor sensitivity. *Cancer Cell* 11:217–27
80. Carey KD, Garton AJ, Romero MS, Kahler J, Thomson S, et al. 2006. Kinetic analysis of epidermal growth factor receptor somatic mutant proteins shows increased sensitivity to the epidermal growth factor receptor tyrosine kinase inhibitor, erlotinib. *Cancer Res.* 66:8163–71
81. Shan Y, Eastwood MP, Zhang X, Kim ET, Arkhipov A, et al. 2012. Oncogenic mutations counteract intrinsic disorder in the EGFR kinase and promote receptor dimerization. *Cell* 149:860–70
82. Sutto L, Gervasio FL. 2013. Effects of oncogenic mutations on the conformational free-energy landscape of EGFR kinase. *PNAS* 110:10616–21
83. Yun CH, Mengwasser KE, Toms AV, Woo MS, Greulich H, et al. 2008. The T790M mutation in EGFR kinase causes drug resistance by increasing the affinity for ATP. *PNAS* 105:2070–75
84. Red-Brewer M, Yun CH, Lai D, Lemmon MA, Eck MJ, Pao W. 2013. Mechanism for activation of mutated epidermal growth factor receptors in lung cancer. *PNAS* 110:e3595–604
85. Huse M, Chen YG, Massague J, Kuriyan J. 1999. Crystal structure of the cytoplasmic domain of the type I TGF β receptor in complex with FKBP12. *Cell* 96:425–36
86. Wybenga-Groot LE, Baskin B, Ong SH, Tong J, Pawson T, Sicheri F. 2001. Structural basis for autoinhibition of the Ephb2 receptor tyrosine kinase by the unphosphorylated juxtamembrane region. *Cell* 106:745–57
87. Griffith J, Black J, Faerman C, Swenson L, Wynn M, et al. 2004. The structural basis for autoinhibition of FLT3 by the juxtamembrane domain. *Mol. Cell* 13:169–78
88. Mol CD, Dougan DR, Schneider TR, Skene RJ, Kraus ML, et al. 2004. Structural basis for the autoinhibition and STI-571 inhibition of c-Kit tyrosine kinase. *J. Biol. Chem.* 279:31655–63

89. Thiel KW, Carpenter G. 2007. Epidermal growth factor receptor juxtamembrane region regulates allosteric tyrosine kinase activation. *PNAS* 104:19238–43
90. Red-Brewer M, Choi SH, Alvarado D, Moravcevic K, Pozzi A, et al. 2009. The juxtamembrane region of the EGF receptor functions as an activation domain. *Mol. Cell* 34:641–51
91. Jura N, Endres NF, Engel K, Deindl S, Das R, et al. 2009. Mechanism for activation of the EGF receptor catalytic domain by the juxtamembrane segment. *Cell* 137:1293–307
92. Wood ER, Shewchuk LM, Ellis B, Brignola P, Brashear RL, et al. 2008. 6-Ethynylthieno[3,2-d]- and 6-ethynylthieno[2,3-d]pyrimidin-4-anilines as tunable covalent modifiers of ErbB kinases. *PNAS* 105:2773–78
93. He L, Hristova K. 2012. Consequences of replacing EGFR juxtamembrane domain with an unstructured sequence. *Sci. Rep.* 2:854
94. Scheck RA, Lowder MA, Appelbaum JS, Schepartz A. 2012. Bipartite tetracysteine display reveals allosteric control of ligand-specific EGFR activation. *ACS Chem. Biol.* 7:1367–76
95. McLaughlin S, Smith SO, Hayman MJ, Murray D. 2005. An electrostatic engine model for autoinhibition and activation of the epidermal growth factor receptor (EGFR/ErbB) family. *J. Gen. Physiol.* 126:41–53
96. Hunter T, Ling N, Cooper JA. 1984. Protein kinase C phosphorylation of the EGF receptor at a threonine residue close to the cytoplasmic face of the plasma membrane. *Nature* 311:480–83
97. Hobert ME, Kil SJ, Medof ME, Carlin CR. 1997. The cytoplasmic juxtamembrane domain of the epidermal growth factor receptor contains a novel autonomous basolateral sorting determinant. *J. Biol. Chem.* 272:32901–9
98. Bocharov EV, Mineev KS, Volynsky PE, Ermolyuk YS, Tkach EN, et al. 2008. Spatial structure of the dimeric transmembrane domain of the growth factor receptor ErbB2 presumably corresponding to the receptor active state. *J. Biol. Chem.* 283:6950–56
99. Mineev KS, Bocharov EV, Pustovalova YE, Bocharova OV, Chupin VV, Arseniev AS. 2010. Spatial structure of the transmembrane domain heterodimer of ErbB1 and ErbB2 receptor tyrosine kinases. *J. Mol. Biol.* 400:231–43
100. Lemmon MA, Treutlein HR, Adams PD, Brunger AT, Engelman DM. 1994. A dimerization motif for transmembrane α -helices. *Nat. Struct. Biol.* 1:157–63
101. Russ WP, Engelman DM. 2000. The GxxxG motif: a framework for transmembrane helix–helix association. *J. Mol. Biol.* 296:911–19
102. Mendrola JM, Berger MB, King MC, Lemmon MA. 2002. The single transmembrane domains of ErbB receptors self-associate in cell membranes. *J. Biol. Chem.* 277:4704–12
103. Endres NF, Das R, Smith AW, Arkhipov A, Kovacs E, et al. 2013. Conformational coupling across the plasma membrane in activation of the EGF receptor. *Cell* 152:543–56
104. Arkhipov A, Shan Y, Das R, Endres NF, Eastwood MP, et al. 2013. Architecture and membrane interactions of the EGF receptor. *Cell* 152:557–69
105. Fleishman SJ, Schlessinger J, Ben-Tal N. 2002. A putative molecular-activation switch in the transmembrane domain of ErbB2. *PNAS* 99:15937–40
106. Gerber D, Sal-Man N, Shai Y. 2004. Two motifs within a transmembrane domain, one for homodimerization and the other for heterodimerization. *J. Biol. Chem.* 279:21177–82
107. Escher C, Cymer F, Schneider D. 2009. Two GxxxG-like motifs facilitate promiscuous interactions of the human ErbB transmembrane domains. *J. Mol. Biol.* 389:10–16
108. Moriki T, Maruyama H, Maruyama IN. 2001. Activation of preformed EGF receptor dimers by ligand-induced rotation of the transmembrane domain. *J. Mol. Biol.* 311:1011–26
109. Bell CA, Tynan JA, Hart KC, Meyer AN, Robertson SC, Donoghue DJ. 2000. Rotational coupling of the transmembrane and kinase domains of the Neu receptor tyrosine kinase. *Mol. Biol. Cell* 11:3589–99
110. King AC, Cuatrecasas P. 1982. Resolution of high and low affinity epidermal growth factor receptors. Inhibition of high affinity component by low temperature, cycloheximide, and phorbol esters. *J. Biol. Chem.* 257:3053–60
111. Mayawala K, Vlachos DG, Edwards JS. 2005. Heterogeneities in EGF receptor density at the cell surface can lead to concave up scatchard plot of EGF binding. *FEBS Lett.* 579:3043–47
112. den Hartigh JC, van Bergen en Henegouwen PMP, Verkleij AJ, Boonstra J. 1992. The EGF receptor is an actin-binding protein. *J. Cell Biol.* 119:349–55

113. Macdonald JL, Pike LJ. 2008. Heterogeneity in EGF-binding affinities arises from negative cooperativity in an aggregating system. *PNAS* 105:112–17
114. Adak S, DeAndrade D, Pike LJ. 2011. The tethering arm of the EGF receptor is required for negative cooperativity and signal transduction. *J. Biol. Chem.* 286:1545–55
115. Macdonald-Obermann JL, Pike LJ. 2009. The intracellular juxtamembrane domain of the epidermal growth factor (EGF) receptor is responsible for the allosteric regulation of EGF binding. *J. Biol. Chem.* 284:13570–76
116. Adak S, Yang KS, Macdonald-Obermann J, Pike LJ. 2011. The membrane-proximal intracellular domain of the epidermal growth factor receptor underlies negative cooperativity in ligand binding. *J. Biol. Chem.* 286:45146–55
117. Yang KS, Ilagan MX, Piwnica-Worms D, Pike LJ. 2009. Luciferase fragment complementation imaging of conformational changes in the epidermal growth factor receptor. *J. Biol. Chem.* 284:7474–82
118. Macdonald-Obermann JL, Piwnica-Worms D, Pike LJ. 2012. Mechanics of EGF receptor/ErbB2 kinase activation revealed by luciferase fragment complementation imaging. *PNAS* 109:137–42
119. Li Y, Macdonald-Obermann J, Westfall C, Piwnica-Worms D, Pike LJ. 2012. Quantitation of the effect of ErbB2 on epidermal growth factor receptor binding and dimerization. *J. Biol. Chem.* 287:31116–25
120. Macdonald-Obermann JL, Pike LJ. 2014. Different EGF receptor ligands show distinct kinetics and biased or partial agonism for homodimer and heterodimer formation. *J. Biol. Chem.* 289:26178–88
121. Alvarado D, Klein DE, Lemmon MA. 2010. Structural basis for negative cooperativity in growth factor binding to an EGF receptor. *Cell* 142:568–79
122. Tynan CJ, Roberts SK, Rolfe DJ, Clarke DT, Loeffler HH, et al. 2011. Human epidermal growth factor receptor (EGFR) aligned on the plasma membrane adopts key features of *Drosophila* EGFR asymmetry. *Mol. Cell. Biol.* 31:2241–52
123. Arkhipov A, Shan Y, Kim ET, Shaw DE. 2014. Membrane interaction of bound ligands contributes to the negative binding cooperativity of the EGF receptor. *PLOS Comput. Biol.* 10:e1003742
124. Arkhipov A, Shan Y, Kim ET, Dror RO, Shaw DE. 2013. Her2 activation mechanism reflects evolutionary preservation of asymmetric ectodomain dimers in the human EGFR family. *eLife* 2:e00708
125. Qiu C, Tarrant MK, Boronina T, Longo PA, Kavran JM, et al. 2009. In vitro enzymatic characterization of near full length EGFR in activated and inhibited states. *Biochemistry* 48:6624–32
126. Mi LZ, Grey MJ, Nishida N, Walz T, Lu C, Springer TA. 2008. Functional and structural stability of the epidermal growth factor receptor in detergent micelles and phospholipid nanodiscs. *Biochemistry* 47:10314–23
127. Wang Z, Longo PA, Tarrant MK, Kim K, Head S, et al. 2011. Mechanistic insights into the activation of oncogenic forms of EGF receptor. *Nat. Struct. Mol. Biol.* 18:1388–93
128. Mi LZ, Lu C, Li Z, Nishida N, Walz T, Springer TA. 2011. Simultaneous visualization of the extracellular and cytoplasmic domains of the epidermal growth factor receptor. *Nat. Struct. Mol. Biol.* 18:984–89
129. Lu C, Mi LZ, Schürpf T, Walz T, Springer TA. 2012. Mechanisms for kinase-mediated dimerization of the epidermal growth factor receptor. *J. Biol. Chem.* 287:38244–53
130. Gadella TW Jr, Jovin TM. 1995. Oligomerization of epidermal growth factor receptors on A431 cells studied by time-resolved fluorescence imaging microscopy. A stereochemical model for tyrosine kinase receptor activation. *J. Cell Biol.* 129:1543–58
131. Martín-Fernández M, Clarke DT, Tobin MJ, Jones SV, Jones GR. 2002. Preformed oligomeric epidermal growth factor receptors undergo an ectodomain structure change during signaling. *Biophys. J.* 82:2415–27
132. Sako Y, Minoghchi S, Yanagida T. 2000. Single-molecule imaging of EGFR signalling on the surface of living cells. *Nat. Cell Biol.* 2:168–72
133. Low-Nam ST, Lidke KA, Cutler PJ, Roovers RC, van Bergen en Henegouwen PMP, et al. 2011. ErbB1 dimerization is promoted by domain co-confinement and stabilized by ligand binding. *Nat. Struct. Mol. Biol.* 18:1244–49
134. Nagy P, Claus J, Jovin TM, Arndt-Jovin DJ. 2010. Distribution of resting and ligand-bound ErbB1 and ErbB2 receptor tyrosine kinases in living cells using number and brightness analysis. *PNAS* 107:16524–29
135. Saffarian S, Li Y, Elson EL, Pike LJ. 2007. Oligomerization of the EGF receptor investigated by live cell fluorescence intensity distribution analysis. *Biophys. J.* 93:1021–31

136. Chung I, Akita R, Vandlen R, Toomre D, Schlessinger J, Mellman I. 2010. Spatial control of EGF receptor activation by reversible dimerization on living cells. *Nature* 464:783–87
137. Clayton AH, Walker F, Orchard SG, Henderson C, Fuchs D, et al. 2005. Ligand-induced dimer–tetramer transition during the activation of the cell surface epidermal growth factor receptor A multidimensional microscopy analysis. *J. Biol. Chem.* 280:30392–99
138. Clayton AH, Orchard SG, Nice EC, Posner RG, Burgess AW. 2008. Predominance of activated EGFR higher-order oligomers on the cell surface. *Growth Factors* 26:316–24
139. Kozler N, Barua D, Henderson C, Nice EC, Burgess AW, et al. 2014. Recruitment of the adaptor protein Grb2 to EGFR tetramers. *Biochemistry* 53:2594–604
140. Gómez A, Volf JN, Hornung U, Schartl M, Wellbrock C. 2004. Identification of a second *egfr* gene in *Xiphophorus* uncovers an expansion of the epidermal growth factor receptor family in fish. *Mol. Biol. Evol.* 21:266–75



Contents

It Seems Like Only Yesterday <i>Charles C. Richardson</i>	1
Veritas per structuram <i>Stephen C. Harrison</i>	37
Nuclear Organization <i>Yosef Gruenbaum</i>	61
The Balbiani Ring Story: Synthesis, Assembly, Processing, and Transport of Specific Messenger RNA–Protein Complexes <i>Petra Björk and Lars Wieslander</i>	65
Functions of Ribosomal Proteins in Assembly of Eukaryotic Ribosomes In Vivo <i>Jesús de la Cruz, Katrin Karbstein, and John L. Woolford Jr.</i>	93
Lamins: Nuclear Intermediate Filament Proteins with Fundamental Functions in Nuclear Mechanics and Genome Regulation <i>Yosef Gruenbaum and Roland Foissner</i>	131
Regulation of Alternative Splicing Through Coupling with Transcription and Chromatin Structure <i>Shiran Naftelberg, Ignacio E. Schor, Gil Ast, and Alberto R. Kornblihtt</i>	165
DNA Triplet Repeat Expansion and Mismatch Repair <i>Ravi R. Iyer, Anna Pluciennik, Marek Napierala, and Robert D. Wells</i>	199
Nuclear ADP-Ribosylation and Its Role in Chromatin Plasticity, Cell Differentiation, and Epigenetics <i>Michael O. Hottiger</i>	227
Application of the Protein Semisynthesis Strategy to the Generation of Modified Chromatin <i>Matthew Holt and Tom Muir</i>	265
Mechanisms and Regulation of Alternative Pre-mRNA Splicing <i>Yeon Lee and Donald C. Rio</i>	291
The Clothes Make the mRNA: Past and Present Trends in mRNP Fashion <i>Guramrit Singh, Gabriel Pratt, Gene W. Yeo, and Melissa J. Moore</i>	325

Biochemical Properties and Biological Functions of FET Proteins <i>Jacob C. Schwartz, Thomas R. Cech, and Roy R. Parker</i>	355
Termination of Transcription of Short Noncoding RNAs by RNA Polymerase II <i>Karen M. Arndt and Daniel Reines</i>	381
PIWI-Interacting RNA: Its Biogenesis and Functions <i>Yuka W. Iwasaki, Mikiko C. Siomi, and Harubiko Siomi</i>	405
The Biology of Proteostasis in Aging and Disease <i>Johnathan Labbadia and Richard I. Morimoto</i>	435
Magic Angle Spinning NMR of Proteins: High-Frequency Dynamic Nuclear Polarization and ^1H Detection <i>Yongchao Su, Loren Andreas, and Robert G. Griffin</i>	465
Cryogenic Electron Microscopy and Single-Particle Analysis <i>Dominika Elmlund and Hans Elmlund</i>	499
Natural Photoreceptors as a Source of Fluorescent Proteins, Biosensors, and Optogenetic Tools <i>Daria M. Shcherbakova, Anton A. Shemetov, Andrii A. Kaberniuk, and Vladislav V. Verkhusha</i>	519
Structure, Dynamics, Assembly, and Evolution of Protein Complexes <i>Joseph A. Marsh and Sarah A. Teichmann</i>	551
Mechanisms of Methicillin Resistance in <i>Staphylococcus aureus</i> <i>Sharon J. Peacock and Gavin K. Paterson</i>	577
Structural Biology of Bacterial Type IV Secretion Systems <i>Vidya Chandran Darbari and Gabriel Waksman</i>	603
ATP Synthase <i>Wolfgang Junge and Nathan Nelson</i>	631
Structure and Energy Transfer in Photosystems of Oxygenic Photosynthesis <i>Nathan Nelson and Wolfgang Junge</i>	659
Gating Mechanisms of Voltage-Gated Proton Channels <i>Yasushi Okamura, Yuichiro Fujiwara, and Souhei Sakata</i>	685
Mechanisms of ATM Activation <i>Tanya T. Paull</i>	711
A Structural Perspective on the Regulation of the Epidermal Growth Factor Receptor <i>Erika Kovacs, Julie Anne Zorn, Yongjian Huang, Tiago Barros, and John Kuriyan</i> ...	739

Chemical Approaches to Discovery and Study of Sources and Targets of Hydrogen Peroxide Redox Signaling Through NADPH Oxidase Proteins <i>Thomas F. Brewer, Francisco J. Garcia, Carl S. Onak, Kate S. Carroll, and Christopher J. Chang</i>	765
Form Follows Function: The Importance of Endoplasmic Reticulum Shape <i>L.M. Westrate, J.E. Lee, W.A. Prinz, and G.K. Voeltz</i>	791
Protein Export into Malaria Parasite-Infected Erythrocytes: Mechanisms and Functional Consequences <i>Natalie J. Spillman, Josh R. Beck, and Daniel E. Goldberg</i>	813
The Twin-Arginine Protein Translocation Pathway <i>Ben C. Berks</i>	843
Transport of Sugars <i>Li-Qing Chen, Lily S. Cheung, Liang Feng, Widmar Tanner, and Wolf B. Frommer</i>	865
A Molecular Description of Cellulose Biosynthesis <i>Josua T. McNamara, Jacob L.W. Morgan, and Jochen Zimmer</i>	895
Cellulose Degradation by Polysaccharide Monooxygenases <i>William T. Beeson, Van V. Vu, Elise A. Span, Christopher M. Phillips, and Michael A. Marletta</i>	923
Physiology, Biomechanics, and Biomimetics of Hagfish Slime <i>Douglas S. Fudge, Sarah Schorno, and Shannon Ferraro</i>	947
Indexes	
Cumulative Index of Contributing Authors, Volumes 80–84	969
Cumulative Index of Article Titles, Volumes 80–84	973

Errata

An online log of corrections to *Annual Review of Biochemistry* articles may be found at
<http://www.annualreviews.org/errata/biochem>